

**AIMS**African Institute for
Mathematical Sciences
RWANDA

Lecture 2: Probability II

Introduction to Probability and Statistics

evans.gouno@univ-ubs.fr

- 1 σ -field
- 2 Conditional Probability
- 3 Independence
- 4 Random variables
- 5 Distribution of r.v.'s

σ -field

Probability will be defined on a collection of events that should be a σ -field.

Definition

A collection $\mathcal{F}$ of subsets of Ω is called a σ -**field** if it satisfies the following conditions:

- $\emptyset \in \mathcal{F}$ or $\Omega \in \mathcal{F}$,
- if $A \in \mathcal{F}$ then $\bar{A} \in \mathcal{F}$.
- if $A_1, A_2, \dots \in \mathcal{F}$ then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$,

Probability measure

Definition

A probability measure P on $(\Omega, \mathcal{F})$ is a function $P : \mathcal{F} \rightarrow [0, 1]$ satisfying

- 1 $P(A) \geq 0$, for all $A \in \mathcal{F}$.
- 2 $P(\Omega) = 1$.
- 3 If $A_1, A_2, \dots$ is a collection of disjoint members of $\mathcal{F}$ ($A_i \cap A_j = \emptyset$ for all pairs satisfying $i \neq j$), then

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i).$$

The triple $(\Omega, \mathcal{F}, P)$, is called a **probability space**.

The axiom 3 is called the **countable additivity**.

Examples

Example 1:

A coin is tossed once.

We take $\Omega = \{H, T\}$ and $\mathcal{F} = \{\emptyset, \{H\}, \{T\}, \Omega\}$.

A possible probability measure P is given by:

$$P(\emptyset) = 0, \quad P(H) = p, \quad P(T) = 1 - p, \quad P(\Omega) = 1,$$

where $p \in [0, 1]$.

Examples

Example 2:

Let consider $\Omega = [0, 1]$

$\mathcal{F} = \mathcal{B}([0, 1])$, the Borel σ -algebra on $[0, 1]$.

➡ this is the smallest σ -algebra containing all open intervals $(a, b) \subseteq [0, 1]$.

Let A be an interval from $\mathcal{B}([0, 1])$

$P(A) = \text{Upper bound of } A - \text{Lower bound of } A = \text{length of } A$.

➡ Lebesgue measure.

e.g. $P((a, b]) = b - a, \forall x \in [0, 1], P(\{x\}) = 0$.

Properties

- $P(\emptyset) = 0$.

Properties

- $P(\emptyset) = 0$.
- For every A , $P(\overline{A}) = 1 - P(A)$.

Properties

- $P(\emptyset) = 0$.
- For every A , $P(\overline{A}) = 1 - P(A)$.
- Let A and B be two events such that $A \subseteq B$.
Then $P(A) \leq P(B)$.

Properties

- $P(\emptyset) = 0$.
- For every A , $P(\overline{A}) = 1 - P(A)$.
- Let A and B be two events such that $A \subseteq B$.
Then $P(A) \leq P(B)$.
- For every A , $P(A) \leq 1$.

Properties

- $P(\emptyset) = 0$.
- For every A , $P(\overline{A}) = 1 - P(A)$.
- Let A and B be two events such that $A \subseteq B$.
Then $P(A) \leq P(B)$.
- For every A , $P(A) \leq 1$.
- **Finite Additivity.**
Let $A_1, \dots, A_n$ be disjoint events.
Then $P(A_1 \cup \dots \cup A_n) = P(A_1) + \dots + P(A_n)$.

Properties

- **Boole's Inequality.** For any sequence $A_1, A_2, A_3 \dots$

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty} P(A_i).$$

Proof:

Properties

- **Boole's Inequality.** For any sequence $A_1, A_2, A_3 \dots$

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty} P(A_i).$$

Proof: Let $B_1 = A_1$ and $B_n = \bar{A}_1 \cap \dots \cap \bar{A}_{n-1} \cap A_n$.

We have $\bigcup_{n=1}^{\infty} B_n = \bigcup_{n=1}^{\infty} A_n$ and $\bigcap_{n=1}^{\infty} B_n = \emptyset$.

Then, from the definition,

$$P\left(\bigcup_{n=1}^{\infty} A_n\right) = P\left(\bigcup_{n=1}^{\infty} B_n\right) = \sum_{n=1}^{\infty} P(B_n).$$

But $B_n \subseteq A_n$ therefore $P(B_n) \leq P(A_n)$ and the result follows.

Properties

Lemma

Let A and B be two events,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

Properties

Lemma

Let A and B be two events,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$

Proof

$A \cup B = (A \cap \bar{B}) \cup (A \cap B) \cup (\bar{A} \cap B)$. (disjoint events)

$P(A \cup B) = P(A \cap \bar{B}) + P(A \cap B) + P(\bar{A} \cap B)$. (additivity)

But $A = (A \cap \bar{B}) \cup (A \cap B)$ and $B = (B \cap \bar{A}) \cup (B \cap A)$.

Therefore $P(A \cap \bar{B}) = P(A) - P(A \cap B)$

and $P(B \cap \bar{A}) = P(B) - P(A \cap B)$.

Then

$$P(A \cup B) = P(A) - \cancel{P(A \cap B)} + \cancel{P(A \cap B)} + P(B) - P(A \cap B).$$

Properties

Exercise

Let A, B, C be 3 events.

Consider $E_1 = A \cap \overline{B} \cap \overline{C}$ and $E_2 = A \cap (B \cup C)$.

We have seen that E_1 and E_2 are disjoint and that $E_1 \cup E_2 = A$.

Suppose that $P(A) = 0.6$, $P(B) = 0.4$, $P(C) = 0.3$, $P(A \cap B) = 0.2$, $P(B \cap C) = 0.1$, $P(A \cap C) = 0.1$ and $P(A \cap B \cap C) = 0.05$.

Calculate $P(E_1)$ and $P(E_2)$

Properties

Solution

- $P(E_2) = P(A \cap (B \cup C)) = P((A \cap B) \cup (A \cap C))$.

From the previous lemma, we have:

$$\begin{aligned} P((A \cap B) \cup (A \cap C)) &= P(A \cap B) + P(A \cap C) - P(A \cap B \cap C) \\ &= 0.2 + 0.1 - 0.05 = \boxed{0.25} \end{aligned}$$

- We know that $E_1 \cup E_2 = A$ and $E_1 \cap E_2 = \emptyset$ then

$$P(E_1 \cup E_2) = P(E_1) + P(E_2) \iff P(A) = P(E_1) + P(E_2)$$

$$\text{And, } P(E_1) = P(A) - P(E_2) = 0.6 - 0.25 = \boxed{0.35}$$

Properties

Poincaré's theorem:

Let $A_1, \dots, A_n$ be events. Then

$$\begin{aligned} P(A_1 \cup A_2 \cup \dots \cup A_n) &= \sum_{i=1}^n P(A_i) - \sum_{i < j} P(A_i \cap A_j) \\ &+ \sum_{i < j < k} P(A_i \cup A_j \cap A_k) - \dots \\ &+ (-1)^{n+1} P(A_1 \cap \dots \cap A_n). \end{aligned}$$

Conditional Probability

Consider two events A and B . Suppose that the event B occurs.

What can we say about the chance of occurrence for event A ?

What is the probability of A given B ?

Conditional Probability

Consider two events A and B . Suppose that the event B occurs.

What can we say about the chance of occurrence for event A ?

What is the probability of A given B ?

Definition

If $P(B) > 0$ the conditional probability of A given B is

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)}.$$

• **Remark:** $P(A \cap B) = P(A \mid B)P(B)$.

Conditional Probability

Lemma

For any events A and B such that $0 < P(B) < 1$,

$$P(A) = P(A \mid B)P(B) + P(A \mid \bar{B})P(\bar{B}).$$

Proof

Conditional Probability

Lemma

For any events A and B such that $0 < P(B) < 1$,

$$P(A) = P(A \mid B)P(B) + P(A \mid \bar{B})P(\bar{B}).$$

Proof

$A = (A \cap B) \cup (A \cap \bar{B})$. $A \cap B$ and $A \cap \bar{B}$ are disjoint.

Then $P(A) = P(A \cap B) + P(A \cap \bar{B})$.

But $P(A \cap B) = P(A \mid B)P(B)$ and $P(A \cap \bar{B}) = P(A \mid \bar{B})P(\bar{B})$.

Therefore

$$P(A) = P(A \mid B)P(B) + P(A \mid \bar{B})P(\bar{B}).$$

Conditional Probability

Generalization:

The Law of Total Probability

Let $B_1, \dots, B_n$ be a partition of Ω . Then, for any event A ,

$$P(A) = \sum_{i=1}^n P(A \mid B_i)P(B_i).$$

Bayes' Theorem

Let $B_1, \dots, B_n$ be a partition of Ω such that $P(B_i) > 0$ for each i .
If $P(A) > 0$ then, for each $i = 1, \dots, n$,

$$P(B_i \mid A) = \frac{P(A \mid B_i)P(B_i)}{\sum_{j=1}^n P(A \mid B_j)P(B_j)}.$$

Conditional Probability

Exercise

Larry divides his email into three categories $A_1 = \text{"spam"}$, $A_2 = \text{"low priority"}$ and $A_3 = \text{"high priority"}$.

From previous experience, he finds that $P(A_1) = 0.7$, $P(A_2) = 0.2$ and $P(A_3) = 0.1$.

Let B be the event that the email contains the word "free".

From previous experience, Larry knows that $P(B | A_1) = 0.9$, $P(B | A_2) = 0.01$ and $P(B | A_3) = 0.01$. Larry receives an email with the word "free".

What is the probability that it is "spam" ?

Conditional Probability

Solution

By definition: $P(A_1 | B) = \frac{P(A_1 \cap B)}{P(B)}.$

$$P(A_1 \cap B) = P(B | A_1)P(A_1) = 0.9 \times 0.7 = 0.63.$$

By the Law of Total Probability:

$$\begin{aligned} P(B) &= P(B | A_1)P(A_1) + P(B | A_2)P(A_2) + P(B | A_3)P(A_3) \\ &= 0.9 \times 0.7 + 0.01 \times 0.2 + 0.01 \times 0.1 = 0.633 \end{aligned}$$

● Conclusion:

$$P(A_1 | B) = \frac{0.63}{0.633} = 0.9952$$

Independence

The conditional probability characterizes the fact the occurrence of an event B modifies the chance of occurrence of A . When it is not the case that is to say the probability remains unchanged $P(A | B) = P(A)$, we said that A and B are independent.

Definition

Two events are said to be independent if

$$P(A \cap B) = P(A)P(B).$$

More generally, a family $\{A_i : i \in I\}$ is called independent if

$$P\left(\bigcap_{i \in J} A_i\right) = \prod_{i \in J} P(A_i)$$

for all finite subsets J of I .

Independence

Exercise

Toss a fair coin n times. Let A be the event “*At least one head occurs*”.
Let T_j be the event that tails occurs on the j^{th} toss.
Use T_j , $j = 1, \dots, n$ and their properties to compute the probability of A ?

Independence

Exercise

Toss a fair coin n times. Let A be the event “At least one head occurs”.

Let T_j be the event that tails occurs on the j^{th} toss.

Use $T_j, j = 1, \dots, n$ and their properties to compute the probability of A ?

Solution:

$P(A) = 1 - P(\bar{A}) = 1 - P(\text{no heads at all}) = 1 - P(\text{all tails})$.

The event $\{\text{all tails}\}$ is $T_1 \cap T_2 \cap \dots \cap T_n$ that is tail occurs on the first toss and on the second toss and so on until the n^{th} toss.

The tosses are independent thus: $P(T_1 \cap T_2 \cap \dots \cap T_n) = \prod_{i=1}^n P(T_i)$.

$T_j = \{H \text{ or } T, (j-1) \text{ times}, T, H \text{ or } T, (n-j) \text{ times}\}$.

$\#T_j = 2^{j-1} \times 2^{n-j} = 2^{n-1}$ and $\#\Omega = 2^n$, then $P(T_j) = 1/2$ and:

$$P(A) = 1 - \frac{1}{2^n}.$$

Independence

Exercise

Two people are taken turns trying to sink a basketball into a net. Person 1 succeeds with probability $1/3$ while person 2 succeeds with probability $1/4$.

What is the probability that person 1 succeeds before person 2?

Hint: Consider the possible sequences of successive games with F for the event “the player fails” and S for “the player succeeds”.

Independence

Solution

Suppose that Person 1 starts the game before Person 2.
Denote E the event of interest.

Person	1	2	1	2	1	2	...	1	2	1	...	Probability
First game	S											1/3
Second game	F	F	S									$2/3 \times 3/4 \times 1/3$
Third game	F	F	F	F	S							$2/3 \times 3/4 \times 2/3 \times 3/4 \times 1/3$
							$\vdots$					$\vdots$
k^{th} game	F	F	F	F	F		...	F	F	S		$(2/3 \times 3/4)^{k-1} \times 1/3$
$\vdots$												$\vdots$

$$P(E) = \sum_{k=1}^{+\infty} (2/3 \times 3/4)^{k-1} \times 1/3 = \sum_{k=1}^{+\infty} (1/2)^{k-1} \times 1/3 = \frac{1}{3} \left(\frac{1}{1 - 1/2} \right) = 2/3.$$

Random variables

Definition

A **random variable** (r.v.) X is a mapping that assigns a real number to each outcome:

$$\begin{aligned} X : \Omega &\longrightarrow V \\ \omega &\longmapsto x = X(\omega) \end{aligned}$$

A random variable can be

- **discrete**, that is $V \subseteq \mathbb{N}$. (ex : $\{0, 1\}$, $\{1, 2, 3, 4\}$, $\mathbb{N}$, etc.)
- **continuous**, that is $V \subseteq \mathbb{R}$ (ex : $[0, 1]$ ou $\mathbb{R}$, etc.).

Examples of random variable

- **discrete:**

- number of failures of a system component in a given time period,
- number of $\{Head\}$ obtained after flipping a coin n times,
- number of dots shown after rolling two six-sided dice.

- **continue:**

- times to failure of a device,
- necessary time to perform a given task,
- weights, heights of individuals in a population.
- measurement errors.

Examples of random variable

- Random experiment: Tossing a coin.

- Possible outcomes: *Tail* or *Head*.

- Universe:

$$\Omega = \{Tail, Head\}$$

- Associated random variable: $X(Tail) = 0$ et $X(Head) = 1$.

- $V = \{0, 1\}$.

- $Tail = X^{-1}(0)$ and $Head = X^{-1}(1)$.

Examples of random variables

- Random experiment: Rolling two dice simultaneously.

- Universe:

$$\Omega = \left\{ \left(\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} \right), \left(\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} \right), \dots, \left(\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} \right), \dots, \left(\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} \right) \right\}$$

- Random variable: X , sum of the number of dots on the face of the two dice.

- For Example:

$$X \left(\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array} \right) = 4.$$

- $V = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$.

Measuring events

- r.v. versus event

$$\begin{aligned} X^{-1}(4) &= \{\omega \in \Omega \mid X(\omega) = 4\} \\ &= \left\{ \left(\begin{array}{|c|} \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \\ \hline \end{array} \right), \left(\begin{array}{|c|} \hline \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \right), \left(\begin{array}{|c|} \hline \bullet \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \\ \hline \end{array} \right) \right\}. \end{aligned}$$

- In practise, we denote:

$$P\{X^{-1}(4)\} = P\{X(\omega) = 4\} = P(X = 4),$$

- $P(X = x), x \in V$ is called **the distribution of X** or **the law of X** .

Distribution

Exercise

Toss an unfair coin (that is $P(H) \neq P(T)$).

What is the distribution of X the r.v. associated to the outcomes?

Distribution

Exercise

Toss an unfair coin (that is $P(H) \neq P(T)$).

What is the distribution of X the r.v. associated to the outcomes?

- $P(H) = P(X = 1) = p$.
- $P(T) = P(X = 0) = 1 - p$.

$$x \in \{0, 1\}, P(X = x) = (1 - p)^{1-x} p^x.$$

Distribution of a random variable

Exercise

What is the distribution of the r.v. X , sum of the dots obtained when rolling two dice simultaneously ?

Distribution of a random variable

Description of Ω

Die 1

						
Die 2	 2	3	4	5	6	7
	3	4	5	6	7	8
	4	5	6	7	8	9
	5	6	7	8	9	10
	6	7	8	9	10	11
	7	8	9	10	11	12

Distribution of a random variable

The size of Ω is 36.

To compute $P(X = x)$, we count the number of couples such that the summation of the two dice dots is equal to x and we divide by $|\Omega| = 36$.

For example, $P(X = 5) = \frac{4}{36} = \frac{1}{9}$.

Distribution of a random variable

The size of Ω is 36.

To compute $P(X = x)$, we count the number of couples such that the summation of the two dice dots is equal to x and we divide by $|\Omega| = 36$.

For example, $P(X = 5) = \frac{4}{36} = \frac{1}{9}$.

Thus, the distribution of X is:

x	2	3	4	5	6
$P(X = x)$	1/36	1/18	1/12	1/9	5/36

x	7	8	9	10	11	12
$P(X = x)$	1/6	5/36	1/9	1/12	1/18	1/36



Distribution of a discrete r.v.

When a random variable is discrete, its law or distribution is described by the quantities $P(X = x)$, $x \in V$.

We have:

$$P(X = x) \in [0, 1], \forall x \in V \text{ et } \sum_{x \in V} P(X = x) = 1.$$

Distribution of a discrete r.v.

Exercise

Let us consider a population with a large number of individuals that may possess the character A .

Let p be the proportion of individuals that possess A .

We draw a sample of size n from the whole population.

Consider the random variable X , number of individuals in the sample possessing A .

What is the distribution of X ?

Hint: Consider the r.v. $Y_i = 1$ if the individual i has the character A , 0 otherwise and find the relationship between X and Y_i , $i = 1, \dots, n$.

Distribution of a discrete r.v.

Solution

Consider a r.v. Y_i such that $Y_i = 1$ if the individual i from the sample possesses the character A and $Y_i = 0$ otherwise.

From the text, $P(Y_i = 1) = p$ and $P(Y_i = 0) = 1 - p$.

$$\begin{aligned} \{X=x\} &= \{ \text{Amongst the } n \text{ individuals exactly } x \text{ possess the character } A \} \\ &= \{x \text{ r.v. } Y_i \text{ are equal to } 1 \text{ and } (n-x) \text{ r.v. are equal to } 0\}. \end{aligned}$$

There are several possible sequences of 0,1 containing exactly x times 1 and $(n-x)$ times 0.

Counting these sequences, we obtain: $C_n^x = \frac{n!}{x!(n-x)!}$

(Write, count and compare the sequences with $n = 5$ and $x = 3$).