Topic 6: Differentiability and Marginal Reasoning

(Why economists care about slopes, not just optima)

---

1. Why differentiability enters now

Up to now, we have focused on two fundamental questions:

1. **Does an optimal choice exist?**
2. **What does the set of optimal choices look like?**

We now turn to a different question:

> **How do optimal choices change when the environment changes?**

This is the core question of **comparative statics**.

To answer it, economists often rely on *local approximations*:

- small changes in prices,
- small changes in income,
- small changes in parameters.

Differentiability is the mathematical structure that makes such reasoning precise.

---

2. From global behavior to local behavior

Recall the generic optimization problem

$$\max_{x \in X} f(x).$$

So far, our analysis has been **global**:

- existence relies on compactness,
- uniqueness relies on strict concavity,
- stability relies on hemicontinuity.

Differentiability allows us to zoom in and ask:

> What happens to $f(x)$ when we move x *a little bit*?

This shift — from global structure to local behavior — is what makes marginal analysis possible.

---

3. Differentiability: the basic idea

Let $f : \mathbb{R}^n \to \mathbb{R}$.

We say that f is **differentiable at** x if there exists a linear map[1]

$$\mathrm{D}f(x) \colon \mathbb{R}^n \to \mathbb{R}$$

such that

$$f(x+h) \approx f(x) + Df(x)[h]$$

for small vectors h.

What does it mean:

> $Df(x)[h]$ is the predicted change in f if I take a small step h away from x.

This definition, however, uses a vector h which is important. So it says, basically: If you give me a direction in which to walk, and (what we will later call) the *gradient* (the linear extrapolation of the change in the function value in our desired direction) at your starting point, then I can tell you where you end up in ''height'' **relative to your starting point** even without knowing the precise function or the starting point.

Important: Differentiability does *not* mean that the function is linear. It means that **locally**, it can be well approximated by something linear.

Sidenote: The relation $\approx$ is one that is used in various ways in mathematics always trying to mean something like the left hand side is close to the right hand side. You may correctly ask, ''what is the metric in which we should measure that closeness?'' And in general I have no answer. In this particular case, the notion of closeness we have in mind behind this is that as we let $h \to 0$ the error we are making vanishes somewhat continuously. The more classical (and also more precise) definition of differentiability would be that $f : U \to \mathbb{R}$ is differentiable at $x \in U$ if

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h}$$

[1] A linear map is a function that has the following property: $L(ax + by) = aL(x) + bL(y)$ for any vectors x and y and scalars a and b

exists. However, the one we have works for multidimensions[2] and carries more economic meaning, and once we talk about it, the notion of $\approx$ we mean here, will be clear.

---

4. How good is "approximately"? Little-o and big-O

Let us now be a bit more precises in *which* sense $f(x+h) \approx f(x) + Df(x)[h]$, that is, in *which* sense is the left-hand side close to the right-hand side? The answer is a piece of notation you will meet constantly—here, in econometrics, in macro, etc.

The idea is simple. We are never really interested in the size of an approximation error on its own. We are interested in how small the error is **relative** to the stuff we are interested in, in particular if we make stepsizes small.

Write the error as $r(h)$, so that

$$f(x+h) = f(x) + Df(x)[h] + r(h).$$

We say that

$$r(h) = o(\|h\|) \qquad \text{if} \qquad \lim_{h \to 0} \frac{r(h)}{\|h\|} = 0,$$

and that

$$r(h) = O(\|h\|) \qquad \text{if} \qquad \frac{|r(h)|}{\|h\|} \text{ stays bounded as } h \to 0.$$

In words, this is a description: $> r(h) = o(\|h\|)$: the error is *negligible next to the step*. Shrink the step and the error shrinks strictly faster. So if h is small enough our approximation is as precise as we want it to be. $> > r(h) = O(\|h\|)$: the error is *at worst proportional to the step*. It does not blow up, but it need not become negligible either.

So little-o is a statement about a race, big-O a statement about a ceiling. Little-o is the stronger claim: o implies O, never the other way round.

A word on the equals sign, because it is a genuine abuse of notation that everyone commits and nobody apologises for. Writing $r(h) = o(\|h\|)$ does not say that r equals some particular object called $o(\|h\|)$. It says r *belongs to the class* of functions that die faster than $\|h\|$. Read it as "is of smaller order than", not as

[2] Look up Frechet derivatives for the more precise definition of the unidimensional one I have given in this sidenote

"equals". The one move you may not make is to cancel such a statement across an equation. Thus, doing algebra with $o(\cdot)$ and $O(\cdot)$ is fishy. People still do it, but one shouldn't take that literally.

Differentiability, restated

With this vocabulary the definition of Section 3 becomes precise and the $\approx$ disappears:

> f is **differentiable at** x if there is a linear map $Df(x)$ with
>
> $$f(x+h) = f(x) + Df(x)[h] + o(\|h\|).$$

This is what the sidenote promised. And the approximation is not merely "good enough for practical purposes"; it is good in an exact sense, because $Df(x)$ is the *only* linear map whose error dies faster than the step itself. That uniqueness is what entitles us to say *the* derivative rather than *a* derivative.

Why insist on all this? Because "small" is not a property a number has on its own. It is a property a number has *relative to something else*. A one-cent error is nothing next to a price change of a euro and enormous next to a price change of a hundredth of a cent. Little-o makes the comparison explicit, and thereby turns marginal reasoning from a hope into a theorem.[3]

A quick economic reading. Let $\pi(k)$ be a firm's profit at input level k, and raise the input by a small amount h:

$$\pi(k+h) = \pi(k) + \pi'(k)\,h + o(h).$$

The term $\pi'(k)h$ is the marginal calculation an economist actually performs. The $o(h)$ is the *licence* to perform it: whatever we threw away is not just small, it is small compared to the very change we are studying, so once h is small enough it cannot overturn the sign of the marginal effect. Without that licence, "marginal profit is positive, so raise the input" would be a guess rather than an argument.

---

5. Partial derivatives and gradients

When f is differentiable, the linear map $Df(x)$ can be represented by a vector of **partial derivatives**

[3] You will meet stochastic cousins of these symbols in econometrics, written o_p and O_p, where "vanishes" means convergence in probability rather than an ordinary limit. The bookkeeping idea is identical; only the notion of vanishing changes. Nothing here is stochastic — o and O are statements about functions, not about random variables.

$$\nabla f(x) = \left(\frac{\partial f}{\partial x_1}(x), \ldots, \frac{\partial f}{\partial x_n}(x) \right).$$

The gradient vector encodes:

- the direction of steepest ascent,
- the marginal effect of each component.

What is a partial derivative?

> The partial derivative with respect to x_i measures how the objective changes when we increase x_i slightly, holding all other components fixed.
>
> The gradient collects all partial derivatives, that is the changes in each direction.

From the gradient we know how we would walk on the landscape given by the function in each direction by making *linear combinations* of the partial derivatives.

The ''holding everything else fixed'' language the gradient encapsulates is fundamental in economic reasoning (we often call it ceteris paribus or c.p. reasoning).

---

6. Marginal interpretation in economics

Marginal reasoning appears everywhere in microeconomics:

- **Marginal utility**: how utility changes with one more unit of a good,
- **Marginal cost**: how cost changes with one more unit of output,
- **Marginal willingness to pay**: how much an agent is willing to give up for a small increase.

All of these are formalized using partial derivatives. In fact, often that is a shortcut we take because differentiability implies continuity which means that 'small changes' are possible in the first place. As economists we have decided to often assume continuity, because it makes the statements clearer and the error in translating back to economics is not too big. However, one has to, of course, always be vigilant whether such an assumption that only serves the economist, is costly in terms of the interpretation.

Reading a derivative backwards

Topic 1 said when a mapping can be undone. We can also undo derivatives.

Let f be differentiable and strictly monotone, so that f^{-1} exists, and write

$y = f(x)$. Differentiating the identity $f\big(f^{-1}(y)\big) = y$ on both sides gives

$$f'\big(f^{-1}(y)\big) \cdot \big(f^{-1}\big)'(y) = 1, \qquad \text{so} \qquad \big(f^{-1}\big)'(y) = \frac{1}{f'\big(f^{-1}(y)\big)}.$$

The formula needs $f' \neq 0$ at the point in question, for the reason written into it: otherwise we are dividing by zero. Think of a flat graph. The inverse would imply an infinite slope, because it becomes a vertical line.

Note, however, that generally, only the slope is lost, not the inverse itself. Take $f(x) = x^3$: it is strictly increasing, so $f^{-1}(y) = y^{1/3}$ exists everywhere, but $f'(0) = 0$ and $\big(f^{-1}\big)'(y) = \frac{1}{3}y^{-2/3} \to \infty$ as $y \to 0$. R

In economics this is intuitive. Inverse demand $p(\cdot)$ and demand $x(\cdot)$ are the same relation read in two directions, so their slopes are reciprocals:

$$p'(x) = \frac{1}{x'(p)}.$$

A demand that reacts sharply to price (large $|x'|$) is an inverse demand that reacts mildly to quantity (small $|p'|$), and the other way round. Neither statement contains more information than the other; which one is convenient depends only on whether the model treats price or quantity as the thing being chosen.

Why is it great to have differentiability? It gives us a disciplined way to talk about ''small changes'' without ambiguity.

---

7. First-order conditions (interior solutions)

Suppose:

- $X \subseteq \mathbb{R}^n$ is open,
- f is differentiable,
- x^* is an interior maximizer.

Then a necessary condition for optimality is

$$\nabla f(x^*) = 0.$$

What this means:

> If the function has our differentiability properties, then at an optimum, we must be in a (locally) flat area.

The reason is (hopefully) clear: If we are at the hill top, anywhere we go, we must go down. However, any path that *crosses* the hilltop from somewhere lower to somewhere lower must first go up and then down again. Differentiability gives us that everywhere we can 'think linearly' if our next step is small enough.

But then, it must be that when switching from going uphill to downhill, we eventually pass a part where the linear approximation is flat (even if only for very short). And this passing must happen exactly at the hill top.

The condition above is called a **first-order condition**.

It does *not* guarantee optimality by itself.
It identifies **candidates** for optimality (so called critical points).

So if we are looking for an optimum in a smooth problem like the one described above, it is *first order* for us to find points that satisfy that condition. Once we have those, we can worry about which of those to eliminate.

---

8. Why first-order conditions are not enough

A point where $\nabla f(x) = 0$ could be:

- a maximum,
- a minimum,
- a saddle point.

A Maximum is what we want, a hill top. A minimum is the lowest point in the valley and the opposite of what we want. A saddle point is something tricky. It is the lowest point of a valley in some direction, but the hill top in another. So standing on the saddle point, means it goes down in some direction, and up in others (like at that one point at the back of a saddle), in some it may be even flat. It is, however, clearly not a hill top.

Moreover, we need to distinguish between local and global optimality. If we are on a hill top in a mountain range, that means we are at a maximum (it goes down in all directions, locally), but nothing says that this is the highest hill top.

The first-order condition tells us nothing about these things.

This is why first-order conditions must be combined with **global structure**:

- concavity of the objective ensures that any critical point is a global maximum,
- differentiability ensures that no jumps or kinks occur that bring about points that can trump the critical points without being critical points themselves
- convexity of the choice set ensures that no local traps exist where we are not allowed to go to.

This connects directly back to Topics 3 and 4.

---

9. Second-order intuition (informal)

Second-order information captures **curvature**.

If f is twice differentiable, curvature is summarized by the **Hessian matrix**

$$D^2 f(x).$$

It is a square table with n rows and n columns that collects all second-order partial derivatives. The entry in row i and column j is

$$\frac{\partial^2 f}{\partial x_i \, \partial x_j}(x),$$

the change in the marginal effect of x_j when x_i increases a little. If these second derivatives are continuous, the order of differentiation does not matter, so the entry in row i, column j equals the entry in row j, column i. A matrix with this property is called **symmetric**.[4] For a production function, symmetry says something that is not obvious at all: one more unit of input j raises the marginal product of input i by exactly as much as one more unit of input i raises the marginal product of input j.

We will not work with Hessians much as objects in their own right, but with their help one can verify optimality of a critical point (yet often this is not very elegant). The next section shows exactly how.

The key intuition is:

> Concavity means the function bends downward everywhere.

Under global concavity of the objective:

- first-order conditions become sufficient,
- local optima are global.

---

10. Taylor approximation: buying a better error term

Above we showed that curvature is summarized by the Hessian, and that with its help one can verify optimality of a critical point. But we had not been precise. To do that, we use Taylor's theorem (often called Taylor approximation) is how. It is also the tool that makes precise a trade we have been making implicitly all along: one more derivative assumed, one more term in the expansion, one order less error.

[4] The result is known as Young's theorem (sometimes also named after Schwarz or Clairaut).

The statement

Let f be twice continuously differentiable near x. Then

$$f(x+h) = f(x) + \nabla f(x) \cdot h + \tfrac{1}{2}\, h^\top D^2 f(x)\, h + o(\|h\|^2).$$

In English:

> Step a little way h from x. The function value changes by a linear term, plus a quadratic term built from the Hessian, plus an error that is negligible even next to $\|h\|^2$.

Set this beside the first-order statement from Section 4:

$$f(x+h) = f(x) + \nabla f(x) \cdot h + o(\|h\|).$$

The pattern is the whole point of the theorem, so let us reiterate:

> Each further derivative you are willing to assume buys you one further term in the expansion, and pushes the error down by one order in $\|h\|$.


It is a real trade, and it is not free. The price is smoothness of f. If f is merely differentiable you get the linear term and $o(\|h\|)$, and that is all you get.

It is worth seeing this claim rather than only reading it. Take a utility function $u(x) = \sqrt{x}$ and expand it at $x_0 = 4$. Since $u(4) = 2$, $u'(4) = \frac{1}{4}$ and $u''(4) = -\frac{1}{32}$, the two approximations are

$$L(x) = 2 + \tfrac{1}{4}(x-4) \qquad \text{and} \qquad Q(x) = 2 + \tfrac{1}{4}(x-4) - \tfrac{1}{64}(x-4)^2.$$

Panel (a) is the qualitative picture: at x_0 all three curves agree, and as we walk away the straight line is the first to part company while the parabola keeps hugging the curve. Panel (b) says the same thing in a form you can check. Fix a tolerance $\varepsilon = 0.1$ and ask how far from x_0 each approximation stays inside it. The line manages $x \in [1.87,\ 6.93]$, the parabola $[1.06,\ 8.39]$ — a range about 1.45 times as wide.

Look also at the shape of the two error curves near x_0, because that is the theorem itself drawn: the linear error leaves zero like $\|h\|^2$, the quadratic error like $\|h\|^3$. That is what "one order less error" buys. It is also why the advantage grows as you demand more: tighten the tolerance to $\varepsilon = 0.02$ and the ratio of the two ranges widens to about 1.9.



(a) Both approximations touch $\sqrt{x}$ at x_0. The line leaves first.

(b) The error, measured. The band where Q stays within ε is 1.45 times wider.

Figure 1: Linear and quadratic approximation of the square root at $x_0 = 4$

Why it is true

The one-dimensional case carries the entire idea. Fix x and h and define

$$g(t) := f(x + th), \qquad t \in [0, 1],$$

so we walk along the segment from x to $x + h$ and record the height as we go. Then g is a function of one variable with $g(0) = f(x)$ and $g(1) = f(x + h)$, and the chain rule gives $g'(0) = \nabla f(x) \cdot h$ and $g''(0) = h^\top D^2 f(x)\, h$ — the two terms of the expansion, already assembled.

So the multidimensional statement *is* the one-dimensional statement about g, and the one-dimensional statement is the mean value theorem applied one order further than usual: control g'' on the segment and keep honest track of what is left over.

Let's go over this. The existence of the second derivative *at* x is what lets us write the quadratic term at all, and—this is worth knowing—it is already enough on its own for the leftover to be $o(\|h\|^2)$. Assuming a little more, namely that $D^2 f$ is *continuous* near x, is the hypothesis we stated above. It costs nothing in any application we will meet, and it buys a cleaner proof and the exact remainder form in the footnote below. What we may *not* drop is the second derivative itself: with only one derivative there is no quadratic term to write, and the best we can say remains the $o(\|h\|)$ statement of Section 4.[5]

[5]Taylor's theorem is used freely in MWG but not stated there as a numbered result. A precise statement and proof is in any calculus text. If you prefer the error written as an exact quantity rather than an order statement, the standard alternative is the Lagrange remainder $\frac{1}{2} h^\top D^2 f(\xi)\, h$ for some unknown point ξ on the segment between x and $x + h$. Same theorem, different bookkeeping: an unknown point instead of an unknown small quantity.

Why that helps with second-order conditions

Now let x^* be a critical point, so $\nabla f(x^*) = 0$. The linear term vanishes and Taylor leaves

$$f(x^* + h) - f(x^*) = \tfrac{1}{2}\, h^\top D^2 f(x^*)\, h + o(\|h\|^2).$$

Everything about *local* optimality is now visible in one line: the change in height is, up to a negligible error, a quadratic form in the direction we walk.

- Negative for every $h \neq 0$ (the Hessian is negative definite): every direction goes down, so x^* is a **local maximum**.
- Positive for every $h \neq 0$: every direction goes up, a **local minimum**.
- Negative in some directions and positive in others: the saddle of Section 8 — and now we can say precisely which directions are the "down" ones and which the "up" ones, because we read them off the quadratic form.

This is the exact version of the hill-top story told in Section 7.

Notice where the $o(\|h\|^2)$ does its work. It is what allows the quadratic term to *decide* the comparison: because the error dies faster than $\|h\|^2$, a strictly signed quadratic form cannot be overturned by it once h is small enough. And if the quadratic form is only *semi*definite — zero in some direction — then along that direction the error is no longer negligible next to it, and second-order information genuinely does not settle the question. That is not a failure of the method. It is the method telling us the truth, namely that we have to look further.

Naming the cases: definite and semidefinite matrices

The sign patterns we just went through have names, and the names are used for any square matrix, not only for Hessians.

Let A be a matrix with n rows and n columns, and write a_{ij} for the entry in row i and column j. For a vector $h \in \mathbb{R}^n$, the number

$$h^\top A\, h = \sum_{i=1}^{n} \sum_{j=1}^{n} a_{ij}\, h_i h_j$$

is called the **quadratic form** of A evaluated at h. With $A = D^2 f(x^*)$ it is, up to the factor $\frac{1}{2}$, the second-order term above. The matrix A is

- **negative semidefinite** if $h^\top A\, h \leq 0$ for every h,
- **negative definite** if $h^\top A\, h < 0$ for every $h \neq 0$,
- **positive semidefinite**, respectively **positive definite**, if the same holds with $\geq$, respectively $>$.

These definitions are statements about the second-order term only. A function whose Hessian at a critical point is negative semidefinite but not negative definite can still rise in a direction where the form is zero: $f(x) = x^4$ at $x = 0$ does.

For $n = 1$ the matrix is a single number a and the quadratic form is ah^2. Negative definite then just means $a < 0$, and negative semidefinite means $a \leq 0$. The definitions extend "the sign of a number" to tables of numbers. A matrix whose quadratic form is positive in some directions and negative in others is none of the four; that was the saddle.[6]

Two things can be read off directly, and both get used.

The diagonal. Let h be the vector with a 1 in position i and zeros elsewhere. Then $h^\top A\, h = a_{ii}$. So every diagonal entry of a negative semidefinite matrix satisfies $a_{ii} \leq 0$. The converse fails: nonpositive diagonal entries do not make a matrix negative semidefinite. Take $f(k_1, k_2) = k_1 k_2$. Its Hessian has zeros on the diagonal and ones off the diagonal, so the quadratic form is $2h_1 h_2$, which is positive for $h = (1, 1)$ and negative for $h = (1, -1)$. Each input on its own has a constant marginal product, and still the Hessian is not negative semidefinite, because the two inputs reinforce each other. By the result stated next, $k_1 k_2$ is therefore not concave.

Concavity. Section 9 said that concavity means the function bends downward everywhere. Here is the precise version. Let f be twice continuously differentiable on an open convex set X. Then

> f is concave if and only if $D^2 f(x)$ is negative semidefinite at every $x \in X$.

Why? Walk along a line, as in the Taylor argument. For fixed x and h, the function $g(t) = f(x + th)$ of the single variable t has second derivative $g''(t) = h^\top D^2 f(x + th)\, h$. A function on a convex set is concave exactly when it is concave along every segment in the set; this is where the convexity of X enters, and openness lets us differentiate in every direction h. And a twice differentiable function of one variable is concave exactly when its second derivative is nowhere positive.

If $D^2 f(x)$ is negative definite at every x, then f is *strictly* concave, meaning that Jensen's inequality (Topic 3, Section 8) holds strictly for any two distinct points and any $\lambda \in (0, 1)$. The converse of this last statement fails: $f(x) = -x^4$ is strictly concave, yet $f''(0) = 0$. For convex functions, replace "negative" by "positive" throughout.[7]

[6] The definitions do not ask A to be symmetric, and in demand theory you will meet a matrix that is shown to be negative semidefinite before anyone knows whether it is symmetric. For the quadratic form only the symmetric part of A matters: swapping the roles of i and j in the double sum shows that A and the matrix with entries $\frac{1}{2}(a_{ij} + a_{ji})$ have the same quadratic form.

[7] MWG defines negative (semi)definiteness in Mathematical Appendix M.D (Definition M.D.1) and proves the characterization of concave functions as Theorem M.C.2.

This also completes the classification of critical points. At an interior local maximum the Hessian *must* be negative semidefinite: if the quadratic form were positive in some direction h, the expansion above would send us uphill for small steps in that direction. So negative semidefinite is necessary, negative definite at a critical point is sufficient, and the gap between the two is exactly the semidefinite case described above.

An example. Take the Cobb–Douglas function $f(k_1, k_2) = k_1^{1/2} k_2^{1/2}$ for $k_1, k_2 > 0$, read as a production function or as a utility function. Differentiating twice gives

$$D^2 f(k) = \frac{f(k)}{4} \begin{pmatrix} -\dfrac{1}{k_1^2} & \dfrac{1}{k_1 k_2} \\ \dfrac{1}{k_1 k_2} & -\dfrac{1}{k_2^2} \end{pmatrix},$$

and therefore

$$h^\top D^2 f(k)\, h = -\frac{f(k)}{4} \left(\frac{h_1}{k_1} - \frac{h_2}{k_2} \right)^2 \leq 0.$$

The Hessian is symmetric, as it must be, and negative semidefinite, so f is concave. It is not negative definite: the quadratic form is zero whenever $h_1/k_1 = h_2/k_2$, that is, when we move in the direction of k itself. That is no accident. Scaling both inputs by the same factor scales output by that factor (Topic 1, Section 10 called this homogeneity of degree one), so along a ray through the origin f is a straight line and has no curvature to show.

An economic example: the cost of a small mistake

Take a firm choosing an input k at unit cost with output price p, so that profit is $\pi(k) = pf(k) - k$, and let k^* be an interior optimum, meaning $pf'(k^*) = 1$. What does it cost the firm to miss the optimum slightly and choose $k^* + h$? Taylor answers at once:

$$\pi(k^* + h) - \pi(k^*) = \underbrace{0}_{\text{first order}} + \tfrac{1}{2}\, pf''(k^*)\, h^2 \;+\; o(h^2).$$

The first-order term is zero *because* k^* was chosen optimally. So the cost of a small mistake is second order: halve the mistake and you quarter the loss.

This is one of the most heavily used facts in applied microeconomics. It is why small optimization errors are usually treated as harmless, and it is the same vanishing first-order term that will make the envelope theorem work in Topic 7. It is also a statement we could not even have phrased two sections ago: "the loss is of order h^2" *is* the content, and o is what lets us say it.

11. Differentiability and constraints (preview)

Most economic problems involve constraints:

$$\max_{x \in X} f(x).$$

When constraints bind, optimality conditions involve:

- trade-offs,
- shadow prices,
- Lagrange multipliers.

We will introduce this machinery later.

For now, the key message is:

> Differentiability allows us to express trade-offs precisely.

12. An economic example: marginal utility

Consider a utility function $u(x_1, x_2)$.

The **partial derivative**

$$\frac{\partial u}{\partial x_1}(x)$$

measures the marginal utility of good 1 at bundle x.[8]

Comparing marginal utilities across goods explains:

- substitution,
- demand responses,
- interior versus corner solutions.

Without differentiability, these ideas lose precision.

There is another form of differentiation, which is the **total derivative**. The way to think about this is to think about the partial derivative as exactly what it is, the change in one direction of the choice vector. So in the above example, $\frac{\partial u}{\partial x_1}(x)$ means we keep x_2 unchanged and only change x_1 (marginally). But of course, there are many ways to change things "slightly." We could walk in direction 1, in direction 2 or any weighted combination of it. The *total derivative* tells us what

[8] These are the standard derivatives we know from undergraduates and school.

that means for our function. And it is (because of the linear map we defined in the beginning) a remarkably simple object. So here it is[9]

$$\mathrm{d}u(x) = \frac{\partial u}{\partial x_1}(x)\mathrm{d}x_1 + \frac{\partial u}{\partial x_2}(x)\mathrm{d}x_2$$

The d here is a way to think about small changes which (however) have a relative size. So if we go in the x_1 direction only, then $\mathrm{d}x_2 = 0$ is 0 and $\mathrm{d}x_1 = 1$.[10]

Differentiating along a ray: Euler's formula

Topic 1, Section 10 introduced homogeneity: f is homogeneous of degree r if $f(\lambda x) = \lambda^r f(x)$ for every $\lambda > 0$. That is a statement about scaling. With derivatives it becomes a statement about marginal effects.

Let f be differentiable and homogeneous of degree r. Fix a point x and treat both sides of the identity as functions of the single variable λ. They are the same function of λ, so they have the same derivative. On the left, the chain rule gives $\nabla f(\lambda x) \cdot x$: we walk along the ray through x, the same device as $g(t) = f(x + th)$ in Section 10. On the right we get $r\lambda^{r-1} f(x)$. Now set $\lambda = 1$:

$$\nabla f(x) \cdot x = \sum_{i=1}^{n} \frac{\partial f}{\partial x_i}(x)\, x_i = r\, f(x).$$

This is **Euler's formula** which says

> "For a differentiable function that is homogeneous of degree r, the marginal effects, each weighted by the level of its own variable, add up to r times the value of the function."

Two examples show why economists use it all the time.

Degree one: paying inputs their marginal product. Let $f(k_1, \ldots, k_n)$ be a differentiable production function with constant returns to scale, so $r = 1$. Euler's formula reads

$$\sum_{i=1}^{n} \frac{\partial f}{\partial k_i}(k)\, k_i = f(k).$$

If every unit of every input is paid that input's marginal product, the payments add up to exactly the output. Nothing is left over, and nothing is missing. For the Cobb–Douglas function of Section 10, $\frac{\partial f}{\partial k_i}(k) = \frac{f(k)}{2k_i}$, so each of the two inputs receives half of the output.

[9] This is just spelling out the definition of derivative!!

[10] Important: While $\mathrm{d}x_1$ is an object we can move around (multiply by $\mathrm{d}x_1$) for example, ∂x_1 is **not,** instead, here the object is the entire $\frac{\partial u}{\partial x_1}(x)$.

Degree zero: demand. Topic 1, Section 10 showed that demand does not react when all prices and wealth are scaled together. Suppose the demand for good ℓ is a differentiable function $x_\ell(p, w)$ of the prices $p_1, \ldots, p_L$ and wealth w. With $r = 0$ the right-hand side of Euler's formula is zero:

$$\sum_{j=1}^{L} \frac{\partial x_\ell}{\partial p_j}(p, w)\, p_j + \frac{\partial x_\ell}{\partial w}(p, w)\, w = 0.$$

The price effects and the wealth effect on any one good are tied together: whoever knows all but one of them knows the last. You will meet this identity again as one of the restrictions that consumer theory places on demand.[11]

---

13. Why this matters for comparative statics

Differentiability underlies a bunch of concepts we will see in Microeconomics:

- Slutsky decompositions,
- envelope theorems (Topic 7),
- local comparative statics.

It allows us to replace

> ''What happens when something changes?''

with

> ''What is the derivative with respect to that change?''

Topic 7 applies this logic to optimized values. It also introduces the KKT conditions needed when an optimum lies on a constraint, where the unconstrained condition $\nabla f = 0$ is no longer appropriate.

This is a powerful simplification — but one that rests on strong assumptions.

---

14. Where this appears in Mas-Colell, Whinston, and Green (MWG)

The main references are:

- **MWG, Chapter 3** (consumer theory)
- **MWG, Mathematical Appendix M.A** (matrix notation for derivatives), **M.B** (homogeneous functions and Euler's formula), **M.C** (concave and quasiconcave functions), and **M.D** (definite and semidefinite matrices)

[11] MWG states and proves Euler's formula as Theorem M.B.2 in Mathematical Appendix M.B, and applies it to demand in Section 2.E.

- **MWG, Chapter 5** (duality and marginal conditions)

Differentiability is often used implicitly; these notes make the logic explicit.

---

15. What you should take away

After this topic, you should be comfortable with:

- interpreting derivatives as local approximations,
- reading o and O as statements about how fast an error vanishes relative to the step,
- reading gradients as vectors of marginal effects,
- understanding when first-order conditions apply,
- using a second-order expansion to classify a critical point,
- telling a negative definite matrix from a negative semidefinite one, and reading a negative semidefinite Hessian as concavity,
- turning homogeneity into Euler's formula,
- differentiating an inverse, and reading demand and inverse demand as reciprocal slopes,
- seeing how differentiability complements convexity.

Differentiability does not replace global reasoning.
It refines it.