

REVIEW · IMMUNOLOGY

How do mRNA vaccines work?

REFERENCES	TOKENS	VERIFICATION	COMPILED
45 verified	≈6.1k	Crossref	26 Aug 2026

SUMMARY

An mRNA vaccine gives cells a short-lived recipe for one pathogen protein, not the pathogen itself. A lipid shell gets the recipe into cells and also helps alert the immune system; antigen production then recruits antibodies, memory B cells and T cells. The platform has delivered strong protection from severe respiratory disease, but protection from infection wanes, variant matching is imperfect, and myocarditis is a rare risk concentrated in young males after dose 2.

01 The platform is a transient, programmable protein factory

- The payload is messenger RNA (mRNA) inside a lipid nanoparticle (LNP): the particle protects RNA, promotes cell entry and cytoplasmic release, and ribosomes translate the sequence without the RNA entering the nucleus or becoming DNA. Chaudhary et al. 2021, Hou et al. 2021, Pardi & Krammer 2024
- Ionizable lipids bind RNA during particle formation but are closer to neutral in the body, improving delivery over permanently charged carriers. Cullis & Felgner 2024, Hou et al. 2021
- The encoded antigen is designed: COVID-19 vaccines used prefusion-stabilized spike, building on earlier coronavirus structural work, so new targets can reuse the manufacturing platform. Corbett et al. 2020, Pardi & Krammer 2024

02 Modified RNA and purification made useful expression possible

- Unmodified RNA activates innate sensors: substituting modified nucleosides suppressed signalling through Toll-like receptors 3, 7 and 8 and reduced cytokine release from human dendritic cells. Karikó et al. 2005
- Removing double-stranded RNA contaminants by high-performance liquid chromatography eliminated detectable interferon and inflammatory-cytokine induction in the tested cells and raised translation **10- to 1,000-fold**. Karikó et al. 2011
- Translation fidelity is a real design constraint: N1-methylpseudouridine caused +1 ribosomal frameshifting and immune recognition of off-target products in mice and people, although no adverse outcome was reported and synonymous recoding reduced the signal. Mulrone et al. 2024

03 The carrier is both delivery vehicle and immune alarm

- In mice, the tested LNP formulation acted as an adjuvant even without antigen-encoding RNA, driving T-follicular-helper and germinal-centre responses through its ionizable lipid and interleukin-6; the exact pathway and magnitude cannot be assumed for every LNP or for humans. Alameh et al. 2021, Verbeke et al. 2022
- Mouse tissue profiling found vaccine mRNA chiefly in draining-node monocytes, macrophages and migratory dendritic cells at day 1; MDA5-type I interferon signalling supported antigen-specific CD8 T-cell expansion, while several other proposed pathways were dispensable in that model. Li et al. 2022

04 Lymph nodes convert a brief signal into durable immune memory

- Early human trials found strong neutralizing antibodies and TH1-skewed CD4 and CD8 T-cell responses, while CD8 cells expanded after the first dose before neutralizing antibodies had fully matured. Sahin et al. 2020, Oberhardt et al. 2021
- Serial lymph-node sampling found spike-specific germinal-centre B cells for at least **12 weeks** and T-follicular-helper cells for about **6 months**, although these intensive studies involved small, relatively young cohorts. Turner et al. 2021, Mudd et al. 2022
- Across 1,540 tracked B-cell clones, somatic mutation rose **3.5-fold over six months**; affinity-matured memory B cells and bone-marrow plasma cells persisted even as circulating antibody declined. Kim et al. 2022
- The delivery-to-memory pathway and its evidence boundaries are summarized in Figure 1.




Figure 1. A temporary RNA recipe builds immune memory.

- **Shows:** A lipid nanoparticle delivers modified mRNA, ribosomes make antigen in the cytoplasm, immune cells coordinate in a draining lymph node, and antibodies plus memory B and T cells remain.
- **Evidence boundary:** Delivery, temporary expression and persistent human lymph-node responses are observed; the detailed sensing pathway and lipid-nanoparticle adjuvant effect rest mainly on mouse experiments.
- **Sources:** [Pardi & Krammer 2024](#), [Alameh et al. 2021](#), [Li et al. 2022](#), [Turner et al. 2021](#), [Kim et al. 2022](#)

05 Antibodies predict protection, but platform performance depends on pathogen and outcome

- The pivotal placebo-controlled COVID-19 trials reported **95.0%** efficacy (95% credible interval 90.3-97.6; n=43,448) for BNT162b2 and **94.1%** (95% CI 89.3-96.8; n=30,420) for mRNA-1273 against symptomatic disease before major immune-escape variants; six-month BNT162b2 efficacy was 91.3% (89.0-93.2) and 96.7% (80.3-99.9) against severe disease. [Polack et al. 2020](#), [Baden et al. 2021](#), [Thomas et al. 2021](#)
- In mRNA-1273 recipients, estimated vaccine efficacy rose from 78% (95% CI 54-89) at a neutralizing titre of 10 to 96% (94-98) at 1,000; a 24-study meta-analysis also linked neutralization loss to reduced protection across variants. Correlation does not mean antibodies mediate every component of protection. [Gilbert et al. 2022](#), [Cromer et al. 2022](#)
- The platform generalizes but does not guarantee success: one-dose mRNA-1345 prevented RSV lower-respiratory disease by 83.7% (95.88% CI 66.0-92.2; n=35,541), whereas an influenza formulation produced stronger A but weaker B responses than licensed vaccine. [Wilson et al. 2023](#), [Ma et al. 2025](#), [Kandinov et al. 2025](#), [Whitaker et al. 2023](#)

VACCINE	EVIDENCE	MAIN RESULT	IMPORTANT BOUNDARY
BNT162b2	Placebo RCT, n=43,448	95.0% efficacy against symptomatic COVID-19	Ancestral-virus era Polack et al. 2020
mRNA-1273	Placebo RCT, n=30,420	94.1% efficacy; 30 severe cases, all placebo	Ancestral-virus era Baden et al. 2021
mRNA-1345 RSV	Placebo RCT, n=35,541, age at least 60	83.7% against lower-respiratory disease with at least 2 signs	Median follow-up 112 days Wilson et al. 2023
ARCT-154 self-amplifying RNA	Placebo RCT, 5 micrograms	56.6% against any and 95.3% against severe COVID-19	Mostly Delta; short follow-up Hô et al. 2024

06 Protection wanes fastest where the airway and variants matter most

- Across 78 estimates, effectiveness fell **21.0 percentage points** (95% CI 13.9-29.8) against infection from month 1 to 6, versus **10.0 points** (6.1-15.4) against severe disease; only 3 of 18 included studies had low overall risk of bias. [Feikin et al. 2022](#)
- The anatomical pattern fits: vaccinated people had strong blood responses but lower airway neutralization and no detectable spike-specific B or T cells in bronchoalveolar lavage, unlike convalescents. This small study supports, but does not by itself prove, a mechanism for breakthrough infection. [Tang et al. 2022](#)

- Variant matching helps incompletely. A BA.1-bivalent booster raised BA.1 titres over the ancestral booster but did not test effectiveness; later BQ.1.1 and XBB.1 escaped much BA.5-bivalent-dose neutralization. [Chalkias et al. 2022](#), [Kurahade et al. 2023](#)
- A 28-study synthesis found bivalent boosters added about **28-31% relative effectiveness** against infection and **58-62%** against severe outcomes versus earlier-dose comparators, with substantial heterogeneity. [Song et al. 2024](#)
- XBB.1.5-vaccine effectiveness in 53.4 million adults fell by month 5 to 26.7% against infection, 52.3% against hospitalization and 69.4% against death; JN.1 replacement reduced it further. Antibody responses remained biased toward epitopes shared with the ancestral strain, with limited truly XBB-specific recruitment in some people. [Ma et al. 2026](#), [Johnston et al. 2024](#)
- The two outcome-specific declines and their supported explanations are separated in [Figure 2](#).




Figure 2. Protection wanes faster against infection than severe disease.

- **Shows:** Average decline from month 1 to month 6 was 21.0 percentage points against infection and 10.0 points against severe disease in a pre-Omicron synthesis.
- **Evidence boundary:** Equal row sizes avoid a false magnitude encoding; airway immunity and variant escape are supported explanations, but the studies do not quantify how much each mechanism contributes.
- **Sources:** [Feikin et al. 2022](#), [Tang et al. 2022](#), [Kurahade et al. 2023](#), [Ma et al. 2026](#)

07 Myocarditis is rare, real, and sharply concentrated by age, sex, dose and product

- Active surveillance of 11.8 million mRNA doses found no prespecified serious-outcome signal and anaphylaxis at about **5 per million doses**; a separate matched national study found most examined adverse events were not elevated but confirmed a myocarditis excess. [Klein et al. 2021](#), [Barda et al. 2021](#)
- A 99-million-person multinational study reproduced the myocarditis/pericarditis signal after mRNA vaccination; its observed-versus-expected design remains sensitive to background-rate assumptions. [Faksova et al. 2024](#)
- In 39.6 million adolescent doses, pooled myopericarditis was **43.5 per million** (95% CI 30.8-61.6): **66.0** in males versus **10.1** in females, and **60.4** after dose 2 versus **16.6** after dose 1. [Guo et al. 2023](#)

- Nordic registers showed that males aged 16-24 had **55.5 excess cases per million** after BNT162b2 dose 2 and **183.9 per million** after mRNA-1273 dose 2; English data likewise found the highest relative risk after mRNA-1273 dose 2 and a smaller booster signal. [Karlstad et al. 2022](#), [Stowe et al. 2023](#)
- Infection is not a uniform comparator: across all ages, infection-associated myocarditis risk was higher, but among men younger than 40 the estimated excess after mRNA-1273 dose 2 was 97 per million versus 16 per million after a positive test. [Patone et al. 2022](#)
- The sex, dose and product concentration—and the incompatible denominators—are shown in [Figure 3](#).




Figure 3. Myocarditis risk is concentrated rather than uniform.

- **Shows:** Panel a separates pooled adolescent rates by sex and dose; panel b separates Nordic excess events in males aged 16–24 by vaccine product.
- **Evidence boundary:** The panels use different populations, denominators and axes, so comparisons are valid within a panel but marker positions cannot be compared across panels.
- **Sources:** [Guo et al. 2023](#), [Karlstad et al. 2022](#), [Patone et al. 2022](#), [Faksova et al. 2024](#)

08 Speed and programmability remain constrained by stability, delivery and evidence gaps

- Purification, particle mixing, scale-up and cold-chain stability remain bottlenecks. [Rosa et al. 2021](#), [Hou et al. 2021](#)
- Freeze-drying preserved one mRNA-LNP formulation for 12 weeks at room temperature and 24 weeks at 4 degrees C in preclinical tests, but another study found post-drying RNA retention and translation depended strongly on the ionizable lipid; thermostability is therefore formulation-specific. [Muramatsu et al. 2022](#), [Lamoot et al. 2023](#)

- [Self-amplifying RNA](#) includes replication machinery and can reduce dose: 5-microgram ARCT-154 protected against severe Delta-era COVID-19, and as a fourth-dose booster it produced non-inferior ancestral and higher BA.4/5 neutralization than BNT162b2 at day 28; neither result establishes long-term superiority. [Blakney et al. 2021](#), [Hô et al. 2024](#), [Oda et al. 2024](#)

References

45 · VERIFIED VIA CROSSREF

- Alameh et al. 2021** Lipid nanoparticles enhance the efficacy of mRNA and protein subunit vaccines by inducing robust T follicular helper cell and humoral responses. *Immunity*. <https://doi.org/10.1016/j.immuni.2021.11.001>
- Baden et al. 2021** Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa2035389>
- Barda et al. 2021** Safety of the BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Setting. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa2110475>
- Blakney et al. 2021** An Update on Self-Amplifying mRNA Vaccine Development. *Vaccines*. <https://doi.org/10.3390/vaccines9020097>
- Chalkias et al. 2022** A Bivalent Omicron-Containing Booster Vaccine against Covid-19. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa2208343>
- Chaudhary et al. 2021** mRNA vaccines for infectious diseases: principles, delivery and clinical translation. *Nature Reviews Drug Discovery*. <https://doi.org/10.1038/s41573-021-00283-5>
- Corbett et al. 2020** SARS-CoV-2 mRNA vaccine design enabled by prototype pathogen preparedness. *Nature*. <https://doi.org/10.1038/s41586-020-2622-0>
- Cromer et al. 2022** Neutralising antibody titres as predictors of protection against SARS-CoV-2 variants and the impact of boosting: a meta-analysis. *The Lancet Microbe*. [https://doi.org/10.1016/s2666-5247\(21\)00267-6](https://doi.org/10.1016/s2666-5247(21)00267-6)
- Cullis & Felgner 2024** The 60-year evolution of lipid nanoparticles for nucleic acid delivery. *Nature Reviews Drug Discovery*. <https://doi.org/10.1038/s41573-024-00977-6>
- Faksova et al. 2024** COVID-19 vaccines and adverse events of special interest: A multinational Global Vaccine Data Network (GVDN) cohort study of 99 million vaccinated individuals. *Vaccine*. <https://doi.org/10.1016/j.vaccine.2024.01.100>
- Feikin et al. 2022** Duration of effectiveness of vaccines against SARS-CoV-2 infection and COVID-19 disease: results of a systematic review and meta-regression. *The Lancet*. [https://doi.org/10.1016/s0140-6736\(22\)00152-0](https://doi.org/10.1016/s0140-6736(22)00152-0)
- Gilbert et al. 2022** Immune correlates analysis of the mRNA-1273 COVID-19 vaccine efficacy clinical trial. *Science*. <https://doi.org/10.1126/science.abm3425>
- Guo et al. 2023** Incidence of myopericarditis after mRNA COVID-19 vaccination: A meta-analysis with focus on adolescents aged 12–17 years. *Vaccine*. <https://doi.org/10.1016/j.vaccine.2023.05.049>
- Hou et al. 2021** Lipid nanoparticles for mRNA delivery. *Nature Reviews Materials*. <https://doi.org/10.1038/s41578-021-00358-0>
- Hô et al. 2024** Safety, immunogenicity and efficacy of the self-amplifying mRNA ARCT-154 COVID-19 vaccine: pooled phase 1, 2, 3a and 3b randomized, controlled trials. *Nature Communications*. <https://doi.org/10.1038/s41467-024-47905-1>
- Johnston et al. 2024** Immunological imprinting shapes the specificity of human antibody responses against SARS-CoV-2 variants. *Immunity*. <https://doi.org/10.1016/j.immuni.2024.02.017>
- Kandinov et al. 2025** An mRNA-based seasonal influenza vaccine in adults: Results of two phase 3 randomized clinical trials and correlate of protection analysis of hemagglutination inhibition titers. *Human Vaccines & Immunotherapeutics*. <https://doi.org/10.1080/21645515.2025.2484088>
- Karikó et al. 2005** Suppression of RNA Recognition by Toll-like Receptors: The Impact of Nucleoside Modification and the Evolutionary Origin of RNA. *Immunity*. <https://doi.org/10.1016/j.immuni.2005.06.008>
- Karikó et al. 2011** Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA. *Nucleic Acids Research*. <https://doi.org/10.1093/nar/gkr695>
- Karlstad et al. 2022** SARS-CoV-2 Vaccination and Myocarditis in a Nordic Cohort Study of 23 Million Residents. *JAMA Cardiology*. <https://doi.org/10.1001/jamacardio.2022.0583>
- Kim et al. 2022** Germinal centre-driven maturation of B cell response to mRNA vaccination. *Nature*. <https://doi.org/10.1038/s41586-022-04527-1>
- Klein et al. 2021** Surveillance for Adverse Events After COVID-19 mRNA Vaccination. *JAMA*. <https://doi.org/10.1001/jama.2021.15072>
- Kurhade et al. 2023** Low neutralization of SARS-CoV-2 Omicron BA.2.75.2, BQ.1.1 and XBB.1 by parental mRNA vaccine or a BA.5 bivalent booster. *Nature Medicine*. <https://doi.org/10.1038/s41591-022-02162-x>
- Lamoot et al. 2023** Successful batch and continuous lyophilization of mRNA LNP formulations depend on cryoprotectants and ionizable lipids. *Biomaterials Science*. <https://doi.org/10.1039/d2bm02031a>
- Li et al. 2022** Mechanisms of innate and adaptive immunity to the Pfizer-BioNTech BNT162b2 vaccine. *Nature Immunology*. <https://doi.org/10.1038/s41590-022-01163-9>
- Ma et al. 2025** Immune correlates analysis of mRNA-1345 RSV vaccine efficacy clinical trial. *Nature Communications*. <https://doi.org/10.1038/s41467-025-61153-x>
- Ma et al. 2026** Effectiveness of the monovalent XBB.1.5 COVID-19 vaccines: A systematic review and meta-analysis. *Journal of Microbiology, Immunology and Infection*. <https://doi.org/10.1016/j.jmii.2025.07.002>
- Mudd et al. 2022** SARS-CoV-2 mRNA vaccination elicits a robust and persistent T follicular helper cell response in humans. *Cell*. <https://doi.org/10.1016/j.cell.2021.12.026>

Mulrone et al. 2024 N1-methylpseudouridylation of mRNA causes +1 ribosomal frameshifting. *Nature*. <https://doi.org/10.1038/s41586-023-06800-3>

Muramatsu et al. 2022 Lyophilization provides long-term stability for a lipid nanoparticle-formulated, nucleoside-modified mRNA vaccine. *Molecular Therapy*. <https://doi.org/10.1016/j.ymthe.2022.02.001>

Oberhardt et al. 2021 Rapid and stable mobilization of CD8+ T cells by SARS-CoV-2 mRNA vaccine. *Nature*. <https://doi.org/10.1038/s41586-021-03841-4>

Oda et al. 2024 Immunogenicity and safety of a booster dose of a self-amplifying RNA COVID-19 vaccine (ARCT-154) versus BNT162b2 mRNA COVID-19 vaccine: a double-blind, multicentre, randomised, controlled, phase 3, non-inferiority trial. *The Lancet Infectious Diseases*. [https://doi.org/10.1016/s1473-3099\(23\)00650-3](https://doi.org/10.1016/s1473-3099(23)00650-3)

Pardi & Krammer 2024 mRNA vaccines for infectious diseases — advances, challenges and opportunities. *Nature Reviews Drug Discovery*. <https://doi.org/10.1038/s41573-024-01042-y>

Patone et al. 2022 Risk of Myocarditis After Sequential Doses of COVID-19 Vaccine and SARS-CoV-2 Infection by Age and Sex. *Circulation*. <https://doi.org/10.1161/circulationaha.122.059970>

Polack et al. 2020 Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa2034577>

Rosa et al. 2021 mRNA vaccines manufacturing: Challenges and bottlenecks. *Vaccine*. <https://doi.org/10.1016/j.vaccine.2021.03.038>

Sahin et al. 2020 COVID-19 vaccine BNT162b1 elicits human antibody and TH1 T cell responses. *Nature*. <https://doi.org/10.1038/s41586-020-2814-7>

Song et al. 2024 A systematic review and meta-analysis on the effectiveness of bivalent mRNA booster vaccines against Omicron variants. *Vaccine*. <https://doi.org/10.1016/j.vaccine.2024.04.049>

Stowe et al. 2023 Risk of myocarditis and pericarditis after a COVID-19 mRNA vaccine booster and after COVID-19 in those with and without prior SARS-CoV-2 infection: A self-controlled case series analysis in England. *PLOS Medicine*. <https://doi.org/10.1371/journal.pmed.1004245>

Tang et al. 2022 Respiratory mucosal immunity against SARS-CoV-2 after mRNA vaccination. *Science Immunology*. <https://doi.org/10.1126/sciimmunol.add4853>

Thomas et al. 2021 Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine through 6 Months. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa2110345>

Turner et al. 2021 SARS-CoV-2 mRNA vaccines induce persistent human germinal centre responses. *Nature*. <https://doi.org/10.1038/s41586-021-03738-2>

Verbeke et al. 2022 Innate immune mechanisms of mRNA vaccines. *Immunity*. <https://doi.org/10.1016/j.immuni.2022.10.014>

Whitaker et al. 2023 mRNA vaccines against respiratory viruses. *Current Opinion in Infectious Diseases*. <https://doi.org/10.1097/qco.0000000000000948>

Wilson et al. 2023 Efficacy and Safety of an mRNA-Based RSV PreF Vaccine in Older Adults. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa2307079>