



BULLETS · NUTRITION

Which health benefits of the Mediterranean diet are actually supported?

REFERENCES	TOKENS	VERIFICATION	COMPILED
97 verified	≈17.5k	Crossref	27 Aug 2026

SUMMARY

The clearest benefit is cardiovascular: supported Mediterranean-diet programmes reduce major events in people at elevated risk, usually by a modest absolute amount. Diabetes prevention is probable, and higher adherence is consistently associated with longer survival. Weight, liver fat, and metabolic markers improve modestly; cancer, dementia, and depression findings are promising but rely much more on observational studies, small trials, or indirect outcomes.

01 The intervention is a family of plant-forward patterns, not a fixed menu

- The recurring core is vegetables, fruit, legumes, nuts, minimally refined cereals, olive oil, and fish, with less red or processed meat and sweets. Dairy, alcohol, quantities, and scoring vary; adherence scores can reward different foods or population medians.^{1,2,3,4}
- The 14-item Mediterranean Diet Adherence Screener (MEDAS) rapidly measures reported behaviour, not biological dose.⁵

- The Seven Countries Study supplied the observational hypothesis that later trials tested; it was not causal proof.⁶
- Review quality varies: among 24 cardiovascular syntheses, complete compliance with adapted methodological criteria ranged from 8% to 75%.⁷
- The recurring core, variable definitions, and unsupported shortcuts are separated in Figure 1.




Figure 1. A Mediterranean diet is a pattern, not one fixed menu.

- **Shows:** The recurring food-group pattern, the elements that vary across studies, and four interpretations that the intervention evidence does not support.
- **Evidence boundary:** The replacement arrow is pattern-level synthesis, not a serving prescription or proof that one component causes the outcomes.
- **Sources:** ^{1,2,4,8}



02 Corrected PREDIMED and two secondary-prevention trials support fewer cardiovascular events

- The original 2013 PREDIMED report is retracted and is not evidence here. After a statistical audit exposed unusual baseline distributions, investigators documented household enrolment and site-level allocation departures and issued a corrected 2018 republication.^{9,10}
- In the corrected analysis (7,447 high-risk adults; median 4.8 y), the primary composite occurred less often with extra-virgin olive oil (hazard ratio (HR) 0.69, 95% confidence interval (CI) 0.53–0.91) and nuts (0.72, 0.54–0.95) than with low-fat advice. Excluding 1,588 participants with known or suspected allocation departures produced essentially the same estimates.¹⁰

- In the Lyon Diet Heart Study after myocardial infarction (n=605), cardiac death or nonfatal infarction occurred in 14 Mediterranean-pattern participants and 44 controls over 46 months; the magnitude is unusually large and has not been reproduced.¹¹
- CORDIOPREV randomized 1,002 patients with coronary disease for 7 y: major events occurred in 87 Mediterranean-diet and 111 low-fat participants, with adjusted hazard ratios around 0.72–0.75.¹²

TRIAL	POPULATION	COMPARATOR	MAIN RESULT	EVIDENCE KEY
PREDIMED republica- tion	High cardiovascular risk; n=7,447	Low-fat advice	Major events: HR 0.69 with olive oil; 0.72 with nuts	10
Lyon Diet Heart	Recent myocardial infarction; n=605	Prudent Western diet	Cardiac death/nonfatal infarction: 14 vs 44	11
CORDIOPREV	Established coronary disease; n=1,002	Low-fat diet	87 vs 111 major events; HR about 0.72–0.75	12

- The exact trial results—and the reason their magnitudes should not be pooled visually—are shown in Figure 2.




Figure 2. Three endpoint trials support cardiovascular benefit.

- **Shows:** Corrected PREDIMED hazard ratios and confidence intervals, plus the reported Lyon Diet Heart and CORDIOPREV event counts.
- **Evidence boundary:** Populations, comparators, endpoints, and follow-up differ; the 2013 PREDIMED report is retracted, and only the corrected 2018 republication is plotted.
- **Sources:** ^{10,11,12,13}



03 Evidence syntheses confirm benefit over minimal support, not clear superiority over every healthy diet

- A network meta-analysis of 40 trials (35,548 participants at increased risk) found Mediterranean programmes superior to minimal intervention for all-cause mortality (odds ratio 0.72), cardiovascular mortality (0.55), stroke (0.65), and nonfatal infarction (0.48); at intermediate baseline risk this meant about 17 fewer deaths and 17 fewer infarctions per 1,000 over 5 y.¹³
- The same analysis found no convincing mortality or infarction advantage over low-fat programmes, which also outperformed minimal advice.¹³
- Earlier randomized-trial pooling found fewer major vascular events (risk ratio 0.63, 95% CI 0.53–0.75) but no all-cause mortality reduction (1.00, 0.86–1.15); excluding one unreliable trial weakened several secondary results.¹⁴
- Cochrane included 30 trials and 12,461 participants but judged most comparisons low or moderate certainty because interventions, comparators, reporting, and event counts differed.¹⁵
- Independent reviews reproduce lower cardiovascular event rates but stress that primary-prevention evidence leans heavily on PREDIMED and programmes often include counselling, free foods, or exercise.^{16,17}
- The 2026 Italian-guideline syntheses again found lower primary and recurrent cardiovascular risk, but combined randomized and cohort evidence and reported certainty ranging from low to high by outcome.^{18,19}
- Compared with low-fat diets, Mediterranean patterns yield small favourable changes in weight, blood pressure, glucose, and lipids; between-diet differences are much smaller than comparisons with usual diet.²⁰

04 Longer survival is a consistent association, but not a randomized mortality result

- In the foundational Greek cohort (22,043 adults), each 2-point adherence increase was associated with 25% lower total mortality; in 74,607 older Europeans the corresponding association was 8%, illustrating how estimates shrink across settings.^{21,22}
- Early pooling of 12 cohorts (1.57 million participants) estimated all-cause mortality risk ratio 0.91 per 2 adherence points; later meta-analyses reached similar 9%–10% reductions per 2 points.^{23,24,25}
- The newest broad synthesis included 54 cohorts, 1.83 million participants, and 346,034 deaths: risk ratio 0.96 (95% CI 0.95–0.97) per 1-point increase, with moderate certainty.²⁶
- Two large US health-professional cohorts linked the top adherence category to total mortality HR 0.82 and cardiovascular disease HR 0.83; these are carefully adjusted but remain observational.^{27,28}

- Among 25,315 US women followed for up to 25 y, the highest adherence group had mortality HR 0.77; metabolic, inflammatory, lipoprotein, insulin-resistance, and body-mass pathways explained only part of the association.²⁹
- Associations also appear in people with diabetes (mortality HR 0.63 per 2 points), yet healthier adherers differ in activity, smoking, education, and care access.³⁰
- Biomarker-based adherence predicted mortality better than repeated food questionnaires in an older Italian cohort, but imperfect specificity could not eliminate confounding.³¹

05 Weight and metabolic risk factors improve, mostly by small amounts

- A 50-study synthesis associated the pattern with lower metabolic-syndrome risk and pooled changes of roughly –2.4 mm Hg systolic pressure, –6 mg/dL triglycerides, –4 mg/dL glucose, and less than 1 cm waist circumference.³²
- Across 57 controlled trials (36,983 participants), direction favoured the Mediterranean diet for 18 of 28 metabolic measures, but comparator and intervention heterogeneity was often high.³³
- In DIRECT (n=322), 2-y weight loss was 4.4 kg with Mediterranean, 2.9 kg with low-fat, and 4.7 kg with low-carbohydrate diets; thus the pattern beat low fat but not low carbohydrate.³⁴
- Across 121 trials, most named diets produced modest 6-mo losses that faded by 12 mo; Mediterranean programmes retained some cardiovascular risk-factor improvement but no unique weight-loss advantage.³⁵
- PREDIMED-Plus combined a 30% energy reduction, activity, and intensive behavioural support. In a 1,521-person imaging substudy, it reduced total and visceral fat and slowed relative lean-mass loss over 3 y, but diet alone cannot receive the credit.³⁶
- A polyphenol-enriched green variant reduced visceral fat by 14.1% versus 6.0% with standard Mediterranean and 4.2% with healthy guidance; 88% of participants were men in a monitored workplace.³⁷
- A 2024 review of olive-oil-enriched Mediterranean trials found no consistent cardiometabolic benefit beyond a very small triglyceride reduction, countering claims that extra oil alone reproduces the whole pattern.⁸
- In pre-existing metabolic disease, 69-study pooling found modest improvements in body mass index, waist, low-density lipoprotein (LDL) cholesterol, triglycerides, glucose, insulin resistance, and C-reactive protein (CRP), while glycated haemoglobin (HbA1c) and lean-mass findings remained inconsistent.³⁸

**06 Type 2 diabetes has unusually coherent trial, cohort, and biomarker evidence**

- In the diabetes-free PREDIMED subgroup, Mediterranean diets reduced incident diabetes relative to low-fat advice; annual adherence analysis gave HR 0.80 (95% CI 0.70–0.92) per 2 MEDAS points and 0.46 (0.25–0.83) for highest versus lowest adherence.^{39,40}
- PREDIMED-Plus later found that adding energy restriction and activity reduced 6-y diabetes incidence by 31% relative to an unrestricted Mediterranean diet (9.5% vs 12.0%), again testing a multicomponent programme rather than diet alone.⁴¹
- A Mediterranean biomarker score trained in a feeding trial predicted diabetes HR 0.71 (0.65–0.77) per standard deviation in EPIC-InterAct (22,202 participants), compared with 0.90 for self-report; specificity and residual confounding remain possible.⁴²

- In 112,493 UK Biobank participants, the top Mediterranean-lifestyle quartile had diabetes HR 0.70, but the exposure deliberately included physical activity, rest, and social habits.⁴³
- Systematic reviews of prospective studies estimate 19%–23% lower incidence with higher adherence. In established diabetes, randomized evidence shows modest improvements rather than remission by diet alone.^{44,45}
- A 2025 synthesis of 11 randomized controlled trials found HbA1c –0.31 percentage points, fasting glucose –0.85 mmol/L, and body mass index –0.83 kg/m² versus control diets.⁴⁶

CARDIOMETABOLIC CLAIM	BEST ESTIMATE	INTERPRETATION	EVIDENCE KEY
Incident diabetes, PREDIMED-Plus	9.5% vs 12.0% over 6 y	Additional benefit from energy restriction + activity	41
HbA1c in established diabetes	–0.31 percentage points	Small but clinically useful adjunct	46
Weight in DIRECT	–4.4 kg at 2 y	Better than low fat; similar to low carbohydrate	34
Olive-oil-enriched trials	Mostly null	Oil supplementation is not the whole pattern	8

07 Cancer associations are broad, but causal evidence is thin

- An umbrella review rated support strongest for overall cancer incidence and mortality, while most site-specific findings were only suggestive or weak.⁴⁷
- The largest updated synthesis (117 studies; 3.20 million participants) associated high adherence with cancer mortality risk ratio 0.87, colorectal cancer 0.83, and breast cancer 0.94; most site-specific certainty was low or very low.⁴⁸
- Cancer meta-analyses converge on inverse observational associations, but varying scores, retrospective data, and lifestyle confounding limit causal interpretation.⁴⁹
- Dose-response pooling supports lower colorectal risk, while a separate postmenopausal breast-cancer cohort/meta-analysis found protection concentrated in estrogen-receptor-negative disease; neither is randomized.^{50,51}
- PREDIMED reported fewer invasive breast cancers in the olive-oil arm (hazard ratio 0.32, 95% CI 0.13–0.79), but this secondary analysis contained only 35 cases.⁵²
- Olive-oil intake itself was associated with cardiovascular, diabetes, and mortality benefits but not cancer (risk ratio 0.94, 0.86–1.03), weakening single-ingredient explanations.⁵³
- A breast-cancer umbrella review likewise found consistent direction but predominantly observational evidence, leaving prevention trials as the missing test.⁵⁴

08 Dementia cohorts are positive; the largest dedicated cognitive trial was null

- Meta-analyses of observational studies associate higher adherence with cognitive impairment HR 0.82, dementia 0.89, and Alzheimer disease 0.70, but report heterogeneity and possible publication bias.^{55,56}
- UK Biobank followed 60,298 adults for 9.1 y: high adherence predicted dementia HR 0.77 and absolute risk 1.18% versus 1.73%, independent of measured genetic risk.⁵⁷
- Earlier synthesis found mild cognitive impairment risk ratio 0.75 and Alzheimer disease 0.71 in cohorts; two small trials yielded episodic-memory standardized mean difference 0.20.^{58,59}
- A PREDIMED substudy reported better global cognition after 6.5 y, but it was small and inherited the parent trial's allocation concerns.⁶⁰
- In contrast, the 3-y MIND trial (604 older adults; calorie-restricted intervention and control) found a global-cognition difference of 0.035 standard deviations (95% CI –0.022 to 0.092) and no meaningful magnetic resonance imaging (MRI) difference.⁶¹
- A MIND-diet meta-analysis remains favourable because cohort weights dominate, while a 2024 PREVENT analysis found no association with cognition or neuroimaging.^{62,63}

**09 Depression trials point in the same direction, but effect size is unstable**

- Prospective dietary-pattern studies associate high adherence with roughly one-third lower incident depression, but reverse causation and social-pattern confounding remain plausible.^{64,65}
- SMILES randomized 67 adults for 12 wk: the diet arm improved Montgomery-Åsberg Depression Rating Scale scores 7.1 points more than social support; remission was 32% versus 8%.⁶⁶
- HELFIMED (n=152) combined Mediterranean advice with fish-oil supplementation and group delivery, producing greater symptom improvement than social groups; the design cannot isolate the diet.⁶⁷
- AMMEND randomized 72 young men and found larger symptom reductions over 12 wk, extending the signal to another small population-specific trial.⁶⁸
- Five-trial pooling (1,507 participants) estimated symptom standardized mean difference -0.53 , but heterogeneity was 87%, the prediction interval crossed zero, and certainty was low.⁶⁹
- A 104-study mental-health review found broadly favourable associations, but only 16 studies were intervention-al.⁷⁰

10 Liver fat falls with calorie-restricted variants, without clear superiority to low-fat diets

- An ad libitum Mediterranean diet and a low-fat diet both reduced liver fat in a 12-wk randomized trial, with no decisive between-diet advantage.⁷¹
- In DIRECT, hepatic-fat reduction statistically mediated part of the Mediterranean diet's cardiometabolic advantage over low fat, but mediation does not identify which foods caused it.⁷²
- DIRECT-PLUS (n=294) found relative liver-fat reductions of 38.9% with green Mediterranean, 19.6% with standard Mediterranean, and 12.2% with healthy guidance at 18 mo; all groups received activity support and the enriched arm included green tea, walnuts, and Mankai.⁷³
- Earlier randomized-trial pooling suggested improvements in steatosis and insulin resistance, whereas enzyme effects were small and inconsistent.^{74,75}
- A 2024 head-to-head meta-analysis concluded Mediterranean and low-fat diets were similarly effective for short-term liver fat and enzymes, a direct contradiction to claims of unique superiority.⁷⁶
- An Asian-adapted trial supports transferability for fatty-liver management, but changes the tested pattern and leaves long-term outcomes unknown.⁷⁷

11 Measured biology is plausible and responsive, but does not substitute for clinical outcomes

- Trial meta-analysis finds small improvements in endothelial function and inflammatory markers, though effects differ by assay, population, and comparator.^{78,79}
- A 2026 synthesis of 33 trials (3,476 participants) found reductions in high-sensitivity CRP, interleukin-6, and interleukin-17, but not conventional CRP, interleukin-10, tumour-necrosis factor, or total antioxidant capacity.⁸⁰
- Mechanistic reviews therefore support several converging pathways—fat quality, polyphenols, fibre fermentation, inflammation, endothelial function, insulin sensitivity—without identifying one necessary ingredient.⁸¹
- In an 8-wk isocaloric trial (n=82), the diet lowered total cholesterol and altered bile acids, the microbiome, and metabolome without changing body weight; glucose, insulin, and CRP were null.⁸²
- A 1-y older-adult intervention (microbiome subset n=612) shifted microbial taxa and functions that covaried with lower frailty and inflammation, while a separate obesity trial showed that two healthy diets altered gut communities and insulin sensitivity.^{83,84}
- Plasma metabolomic profiles partly tracked adherence and cardiovascular risk, providing objective exposure measures rather than proof that the metabolites mediate events.⁸⁵
- Baseline bile acids and microbes may modify lipid and adiposity responses; PREDIMED-Plus also links diet-induced microbiome and metabolome shifts with risk factors. These are secondary analyses, not independent endpoint trials.^{86,87}

12 Translation depends on culture, cost, support, and what replaces meat or alcohol

- Outside the Mediterranean, food availability, cooking, resources, and adherence scores change; cultural adaptation is part of the intervention.^{88,89}
- Score agreement is limited even within one study population, so an effect per point is not automatically portable between instruments or countries.⁹⁰
- MedEx-UK demonstrated that a theory-informed programme could improve adherence and remain acceptable in older adults at dementia risk, but the intervention supplied behavioural maintenance and activity support.⁹¹
- Price, access, cooking skills, family norms, and knowledge impede adherence; free foods, counselling, and frequent contact make trials more supportive than routine life.⁹²
- Economic modelling in secondary prevention suggests Mediterranean diet plus activity can be cost-effective, but estimates inherit assumptions about adherence and event reduction.⁹³



- A sustainability review (35 studies, mostly models) classified the pattern as environmentally favourable in 91% of comparisons, with results driven mainly by lower ruminant-meat intake; water footprints and local production can reverse particular comparisons.^{94,95}
- Traditional scores reward moderate alcohol, but modern interpretation should not encourage non-drinkers to start: cancer and addiction risk make wine an optional historical component, not a therapeutic ingredient.⁹⁶
- Dietary-pattern mortality reviews still identify a much larger observational than randomized evidence base, so residual healthy-user bias remains a central limit even after extensive adjustment.⁹⁷
- The intervention package, adherence conditions, and causal stopping point are traced in Figure 3.




Figure 3. Benefits belong to the supported programme as delivered.

- **Shows:** Trials evaluate a replacement pattern with varying support; real-world conditions influence adherence before risk factors, imaging outcomes, or clinical events are measured.
- **Evidence boundary:** The pathway is programme logic, not a quantified mediation model; it does not isolate one ingredient, prove unsupported diet alone, or guarantee identical effects after adaptation.
- **Sources:** 10,91,92,94,41

13 The evidence supports a hierarchy of claims, not a universal health halo

OUTCOME	BEST-SUPPORTED CONCLUSION	MAIN UNCERTAINTY
Major cardiovascular events	Probably reduced in higher-risk adults, especially versus minimal advice	Absolute benefit and superiority to other high-quality diets ¹³
Type 2 diabetes	Probably prevented; glycaemia improves modestly in established disease	Added effects of calorie restriction, activity, and intensive support ⁴¹
All-cause mortality	Consistently associated with 4% lower risk per adherence point	No mortality trial capable of confirming the cohort estimate ²⁶
Weight and metabolic markers	Small, clinically useful changes	Not uniquely better than low-carbohydrate or low-fat diets ³⁵
Cancer	Lower incidence/mortality associations for several sites	Mostly observational; one small secondary randomized signal ⁴⁸
Cognition	Cohorts favourable; dedicated 3-y trial null	Duration, population risk, active comparator, residual confounding ⁶¹
Depression	Small trials favourable	Low certainty, high heterogeneity, weak blinding ⁶⁹



OUTCOME	BEST-SUPPORTED CONCLUSION	MAIN UNCERTAINTY
Fatty liver	Liver fat improves with energy-restricted variants	Similar short-term efficacy to low-fat diets; few hard outcomes ⁷⁶

- The most defensible interpretation is selective: use the pattern as a supported cardiovascular and diabetes-prevention programme, not as proof that olive oil, wine, or any single food prevents every chronic disease.
- For outcomes beyond cardiometabolic disease, larger preregistered trials with active comparators, objective adherence measures, diverse populations, and clinical rather than surrogate endpoints would change confidence most.
- The uneven directness and certainty across outcomes are mapped in Figure 4.




Figure 4. The evidence is strongest for cardiometabolic outcomes.

- **Shows:** Outcomes are arranged by human evidence directness and certainty, from clinical endpoint trials through consistent but less-direct evidence to limited or surrogate outcomes.
- **Evidence boundary:** This is a narrative-review synthesis, not a formal GRADE score; position, marker area, and colour do not encode effect size or individual benefit.
- **Sources:** ^{13,41,26,48,61}

References

97 · VERIFIED VIA CROSSREF

1. Davis C, Bryan J, Hodgson J, Murphy K (2015) Definition of the Mediterranean Diet; A Literature Review. *Nutrients*. <https://doi.org/10.3390/nu7115459>
2. Bach-Faig A, Berry EM, Lairon D, Reguant J, Trichopoulos A, Dernini S, Medina FX, Battino M, Belahsen R, Miranda G, Serra-Majem L (2011) Mediterranean diet pyramid today. Science and cultural updates. *Public Health Nutrition*. <https://doi.org/10.1017/s1368980011002515>
3. Bach A, Serra-Majem L, Carrasco JL, Roman B, Ngo J, Bertomeu I, Obrador B (2006) The use of indexes evaluating the adherence to the Mediterranean diet in epidemiological studies: a review. *Public Health Nutrition*. <https://doi.org/10.1079/phn2005936>
4. Hutchins-Wiese HL, Bales CW, Porter Starr KN (2022) Mediterranean diet scoring systems: understanding the evolution and applications for Mediterranean and non-Mediterranean countries. *British Journal of Nutrition*. <https://doi.org/10.1017/s0007114521002476>
5. Schröder H, Fitó M, Estruch R, Martínez-González MA, Corella D, Salas-Salvadó J, Lamuela-Raventós R, Ros E, Salaverria I, Fiol M, Lapetra J, Vinyoles E, Gómez-Gracia E, Lahoz C, Serra-Majem L, Pintó X, Ruiz-Gutiérrez V, Covas MI (2011) A Short Screener Is Valid for Assessing Mediterranean Diet Adherence among Older Spanish Men and Women. *The Journal of Nutrition*. <https://doi.org/10.3945/jn.110.135566>
6. Menotti A, Puddu PE (2015) How the Seven Countries Study contributed to the definition and development of the Mediterranean diet concept: A 50-year journey. *Nutrition, Metabolism and Cardiovascular Diseases*. <https://doi.org/10.1016/j.numecd.2014.12.001>
7. Huedo-Medina TB, Garcia M, Bihuniak JD, Kenny A, Kerstetter J (2016) Methodologic quality of meta-analyses and systematic reviews on the Mediterranean diet and cardiovascular disease outcomes: a review. *The American Journal of Clinical Nutrition*. <https://doi.org/10.3945/ajcn.115.112771>
8. Keshani M, Sadeghi N, Tehrani SD, Ahmadi AR, Sharma M (2024) Mediterranean diet enriched with olive oil shows no consistent benefits on cardiometabolic and anthropometric parameters: a systematic review with meta-analysis of randomized controlled trials. *European Journal of Nutrition*. <https://doi.org/10.1007/s00394-024-03478-9>
9. Carlisle JB (2017) Data fabrication and other reasons for non-random sampling in 5087 randomised, controlled trials in anaesthetic and general medical journals. *Anaesthesia*. <https://doi.org/10.1111/anae.13938>
10. Estruch R, Ros E, Salas-Salvadó J, Covas MI, Corella D, Arós F, Gómez-Gracia E, Ruiz-Gutiérrez V, Fiol M, Lapetra J, Lamuela-Raventós RM, Serra-Majem L, Pintó X, Basora J, Muñoz MA, Sorlí JV, Martínez JA, Fitó M, Gea A, Hernán MA, Martínez-González MA (2018) Primary Prevention of Cardiovascular Disease with a Mediterranean Diet Supplemented with Extra-Virgin Olive Oil or Nuts. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa1800389>
11. de Lorgeril M, Salen P, Martin JL, Monjaud I, Delaye J, Mamelle N (1999) Mediterranean Diet, Traditional Risk Factors, and the Rate of Cardiovascular Complications After Myocardial Infarction. *Circulation*. <https://doi.org/10.1161/01.cir.99.6.779>



12. Delgado-Lista J, Alcalá-Díaz JF, Torres-Peña JD, Quintana-Navarro GM, Fuentes F, García-Ríos A, Ortiz-Morales AM, González-Requero AI, Pérez-Caballero AI, Yubero-Serrano EM, Rangel-Zuñiga OA, Camargo A, Rodríguez-Cantalejo F, López-Segura F, Badimon L, Ordovas JM, Pérez-Jiménez F, Pérez-Martínez P, López-Miranda J, Alcalá-Díaz JF, Almada Peña Y, Aranda E, Arenas de Larriba AP, Badimon L, Badimon JJ, Blanco-Molina A, Blanco-Rojó R, Bolívar-Muñoz J, Caballero-Villarraso J, Camargo A, Chica J, Corina A, Criado-García J, Cruz-Teno C, Daponte-Codina A, de Teresa Galvan E, Delgado-Casado N, Delgado-Lista J, Estruch R, Fernandez JM, Fernandez-Gandara C, Fuentes-Jimenez F, Garcia-Carpintero Fernandez-Pacheco S, Garcia-Rios A, Gomez-Delgado F, Gomez-Garduño A, Gomez-Luna P, Gomez-Luna MJ, Gonzalez-Guardia L, Gonzalez-Requero AI, Gutierrez-Mariscal FM, Haro-Mariscal CM, Jimenez-Lucena R, Jimenez-Morales AI, Leon-Acuña A, Lopez-Miranda J, Lopez-Segura F, Marin-Hinojosa C, Meneses Alvarez ME, Mesa-Luna D, Moya-Garrido MN, Muñoz-Carvajal I, Navarro-Martos V, Ochoa JJ, Ordovas JM, Ortiz-Minuesa JA, Ortiz-Morales AM, Pan M, Peña-Orihuela P, Pérez-Caballero AI, Pérez-Corral I, Pérez-Jiménez F, Pérez-Martínez P, Pi-Sunyer FX, Quintana-Navarro GM, Ramírez-Lara I, Rangel-Zuñiga OA, Rodríguez-Artalejo F, Rodríguez-Cantalejo J, Romero MA, Roncero-Ramos I, Ruano-Ruiz JA, Ruiz de Castroviejo J, Sanchez-Villegas P, Suarez de Lezo J, Suarez de Lezo J, Torres-Peña JD, Vals-Delgado C, Valverde R, Visioli F, Yubero-Serrano EM (2022) Long-term secondary prevention of cardiovascular disease with a Mediterranean diet and a low-fat diet (CORDIOPREV): a randomised controlled trial. *The Lancet*. [https://doi.org/10.1016/s0140-6736\(22\)290012-2](https://doi.org/10.1016/s0140-6736(22)290012-2)
13. Karam G, Agarwal A, Sadeghirad B, Jalink M, Hitchcock CL, Ge L, Kiflen R, Ahmed W, Zea AM, Milenkovic J, Chedrawe MA, Rabassa M, El Dib R, Goldenberg JZ, Guyatt GH, Boyce E, Johnston BC (2023) Comparison of seven popular structured dietary programmes and risk of mortality and major cardiovascular events in patients at increased cardiovascular risk: systematic review and network meta-analysis. *BMJ*. <https://doi.org/10.1136/bmj-2022-072003>
14. Liyanage T, Ninomiya T, Wang A, Neal B, Jun M, Wong MG, Jardine M, Hillis GS, Perkovic V (2016) Effects of the Mediterranean Diet on Cardiovascular Outcomes—A Systematic Review and Meta-Analysis. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0159252>
15. Rees K, Takeda A, Martin N, Ellis L, Wijesekara D, Vepa A, Das A, Hartley L, Stranges S (2019) Mediterranean-style diet for the primary and secondary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.cd009825.pub3>
16. Bloomfield HE, Koeller E, Greer N, MacDonald R, Kane R, Wilt TJ (2016) Effects on Health Outcomes of a Mediterranean Diet With No Restriction on Fat Intake. *Annals of Internal Medicine*. <https://doi.org/10.7326/m16-0361>
17. Sebastian SA, Padda I, Johal G (2024) Long-term impact of mediterranean diet on cardiovascular disease prevention: A systematic review and meta-analysis of randomized controlled trials. *Current Problems in Cardiology*. <https://doi.org/10.1016/j.cpcardiol.2024.102509>
18. Volpe R, Ciccone MM, Pala B, Barbarano F, Camastra S, Caprio M, Casirati A, Ferrera A, Galletti F, Greatti A, Mollica MP, Paolillo S, Parretti D, Nucci D, Veronese N, Fontana L, Maggi S, Onder G, Silano M, Zanetti M, Gianfredi V, Ricci M, Volpe M, Maggi S, Volpe M, Zanetti M, Onder G, Veronese N, Gianfredi V, Nucci D, Silano M, Rossi L, Antonella B, Paolo C, Maria CA, Matteo CM, Giovambattista D, Ferruccio G, Emilia G, Antonella L, Anna POS, Francesco L, Francesco M, Ida MM, Gerardo M, Barbara P, Domenico R, Elena S, Massimiliano S, Immacolata SM, Ersilia T, Andrea U, Roberto V, Giovanni Z, Elena A, Fiorella A, Letizia B, Federico B, Alberto B, Ilari BM, Carola B, Stefania C, Ilaria C, Massimiliano C, Irene C, Amanda C, Elisa C, Anastasia C, Martina C, Martina C, Giuseppe CAF, Laura C, Marco C, Stefano C, Lucilla C, Chiara D, Andrea D, Francesco DL, Lorenzo E, Maurizio F, Enrica F, Martina F, Armando F, Lucia G, Cristina G, Alessandro G, Paola G, Federica L, Fabrizia L, Elisa M, Alessandro M, Pina MM, Alessia M, Alisha M, Giovanna M, Barbara P, Stefania P, Damiano P, Cristina PM, Costanza P, Saverio RF, Cristiana R, Francesco RG, Alessia S, Piero S, Sara S, Dario TG, Ilaria T, Filippo V, Fernanda V, Marilena V, Salvatore Z, Morsella A, Limongi F, Ragusa FS, Laviano A, Sieber C (2026a) Effectiveness of Mediterranean diet for the primary prevention of cardiovascular diseases: A systematic review and meta-analysis featured in the Italian National Guidelines "La Dieta Mediterranea". *Nutrition*. <https://doi.org/10.1016/j.nut.2025.113038>
19. Volpe R, Ciccone MM, Pala B, Barbarano F, Camastra S, Caprio M, Casirati A, Ferrera A, Galletti F, Greatti A, Mollica MP, Paolillo S, Parretti D, Nucci D, Veronese N, Fontana L, Maggi S, Onder G, Silano M, Zanetti M, Gianfredi V, Ricci M, Volpe M (2026b) Efficacy of Mediterranean diet for the prevention of cardiovascular disease in patients: A systematic review and meta-analysis featured in the Italian National Guidelines "La Dieta Mediterranea". *Nutrition*. <https://doi.org/10.1016/j.nut.2025.113053>
20. Nordmann AJ, Suter-Zimmermann K, Bucher HC, Shai I, Tuttle KR, Estruch R, Briel M (2011) Meta-Analysis Comparing Mediterranean to Low-Fat Diets for Modification of Cardiovascular Risk Factors. *The American Journal of Medicine*. <https://doi.org/10.1016/j.amjmed.2011.04.024>
21. Trichopoulos A, Costacou T, Bamia C, Trichopoulos D (2003) Adherence to a Mediterranean Diet and Survival in a Greek Population. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa025039>
22. Trichopoulos A, Orfanos P, Norat T, Bueno-de-Mesquita B, Ocké MC, Peeters PH, van der Schouw YT, Boeing H, Hoffmann K, Boffetta P, Nagel G, Masala G, Krogh V, Panico S, Tumino R, Vineis P, Bamia C, Naska A, Benetou V, Ferrari P, Slimani N, Pera G, Martínez-García C, Navarro C, Rodríguez-Barranco M, Dorronsoro M, Spencer EA, Key TJ, Bingham S, Khaw KT, Kesse E, Clavel-Chapelon F, Boutron-Ruault MC, Berglund G, Wirfalt E, Hallmans G, Johansson I, Tjønneland A, Olsen A, Overvad K, Hundborg HH, Riboli E, Trichopoulos D (2005) Modified Mediterranean diet and survival: EPIC-elderly prospective cohort study. *BMJ*. <https://doi.org/10.1136/bmj.38415.644155.8f>
23. Sofi F, Cesari F, Abbate R, Gensini GF, Casini A (2008) Adherence to Mediterranean diet and health status: meta-analysis. *BMJ*. <https://doi.org/10.1136/bmj.a1344>
24. Eleftheriou D, Benetou V, Trichopoulos A, La Vecchia C, Bamia C (2018) Mediterranean diet and its components in relation to all-cause mortality: meta-analysis. *British Journal of Nutrition*. <https://doi.org/10.1017/s0007114518002593>
25. Soltani S, Jayedi A, Shab-Bidar S, Becerra-Tomás N, Salas-Salvado J (2019) Adherence to the Mediterranean Diet in Relation to All-Cause Mortality: A Systematic Review and Dose-Response Meta-Analysis of Prospective Cohort Studies. *Advances in Nutrition*. <https://doi.org/10.1093/advances/nmz041>
26. Nucci D, Ragusa FS, Mazza E, Veronese N, Gianfredi V, Volpe M, Maggi S, Onder G, Sieber C, Zanetti M, Silano M, Chernykh A, Lorenzo FD, Romiti GF, Schirò P, Ungar A, Maggi S, Volpe M, Zanetti M, Onder G, Veronese N, Gianfredi V, Nucci D, Silano M, Rossi L, Antonella B, Paolo C, Maria CA, Matteo CM, Giovambattista D, Ferruccio G, Emilia G, Antonella L, Francesco L, Massimiliano S, Immacolata SM, Ersilia T, Andrea U, Roberto V, Giovanni Z, Elena A, Fiorella A, Letizia B, Federico B, Alberto B, Ilari BM, Carola B, Stefania C, Ilaria C, Massimiliano C, Irene C, Amanda C, Elisa C, Anastasia C, Martina C, Martina C, Giuseppe CAF, Laura C, Marco C, Stefano C, Lucilla C, Chiara D, Andrea D, Francesco DL, Lorenzo E, Maurizio F, Enrica F, Martina F, Armando F, Lucia G, Cristina G, Alessandro G, Paola G, Federica L, Fabrizia L, Elisa M, Alessandro M, Pina MM, Alessia M, Alisha M, Giovanna M, Barbara P, Stefania P, Damiano P, Cristina PM, Costanza P, Saverio RF, Cristiana R, Francesco RG, Alessia S, Piero S, Sara S, Dario TG, Ilaria T, Filippo V, Fernanda V, Marilena V, Salvatore Z, Morsella A, Limongi F, Ragusa FS, Fontana L, Laviano A, Sieber C (2026) Adherence to the Mediterranean diet and all-cause mortality: A systematic review and meta-analysis within the Italian National Guidelines "La Dieta Mediterranea". *Nutrition*. <https://doi.org/10.1016/j.nut.2026.113189>
27. Shan Z, Wang F, Li Y, Baden MY, Bhupathiraju SN, Wang DD, Sun Q, Rexrode KM, Rimm EB, Qi L, Tabung FK, Giovannucci EL, Willett WC, Manson JE, Qi Q, Hu FB (2023) Healthy Eating Patterns and Risk of Total and Cause-Specific Mortality. *JAMA Internal Medicine*. <https://doi.org/10.1001/jamainternmed.2022.6117>
28. Shan Z, Li Y, Baden MY, Bhupathiraju SN, Wang DD, Sun Q, Rexrode KM, Rimm EB, Qi L, Willett WC, Manson JE, Qi Q, Hu FB (2020) Association Between Healthy Eating Patterns and Risk of Cardiovascular Disease. *JAMA Internal Medicine*. <https://doi.org/10.1001/jamainternmed.2020.2176>
29. Ahmad S, Moorthy MV, Lee IM, Ridker PM, Manson JE, Buring JE, Demler OV, Mora S (2024) Mediterranean Diet Adherence and Risk of All-Cause Mortality in Women. *JAMA Network Open*. <https://doi.org/10.1001/jamanetworkopen.2024.14322>
30. Bonaccio M, Di Castelnuovo A, Costanzo S, Persichillo M, De Curtis A, Donati MB, de Gaetano G, Iacoviello L, on behalf of the MOLI-SANI study Investigators (2016) Adherence to the traditional Mediterranean diet and mortality in subjects with diabetes. Prospective results from the MOLI-SANI study. *European Journal of Preventive Cardiology*. <https://doi.org/10.1177/2047487315569409>
31. Hidalgo-Liberona N, Meroño T, Zamora-Ros R, Rabassa M, Semba R, Tanaka T, Bandinelli S, Ferrucci L, Andres-Lacueva C, Cherubini A (2021) Adherence to the Mediterranean diet assessed by a novel dietary biomarker score and mortality in older adults: the InCHIANTI cohort study. *BMC Medicine*. <https://doi.org/10.1186/s12916-021-02154-7>
32. Kistorini CM, Milionis HJ, Esposito K, Giugliano D, Goudevenos JA, Panagiotakos DB (2011) The Effect of Mediterranean Diet on Metabolic Syndrome and its Components. *Journal of the American College of Cardiology*. <https://doi.org/10.1016/j.jacc.2010.09.073>
33. Papadaki A, Nolen-Doerr E, Mantzoros CS (2020) The Effect of the Mediterranean Diet on Metabolic Health: A Systematic Review and Meta-Analysis of Controlled Trials in Adults. *Nutrients*. <https://doi.org/10.3390/nu12113342>
34. Shai I, Schwarzfuchs D, Henkin Y, Shahar DR, Witkow S, Greenberg I, Golan R, Fraser D, Bolotin A, Vardi H, Tangi-Rozental O, Zuk-Ramot R, Sarusi B, Brickner D, Schwartz Z, Sheiner E, Marko R, Katorza E, Thiery J, Fiedler GM, Blüher M, Stumvoll M, Stampfer MJ (2008) Weight Loss with a Low-Carbohydrate, Mediterranean, or Low-Fat Diet. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa0708681>
35. Ge L, Sadeghirad B, Ball GDC, da Costa BR, Hitchcock CL, Svendrovski A, Kiflen R, Quadri K, Kwon HY, Karamouzian M, Adams-Webber T, Ahmed W, Damannhoury S, Zeraatkar D, Nikolakopoulou A, Tsuyuki RT, Tian J, Yang K, Guyatt GH, Johnston BC (2020) Comparison of dietary macronutrient patterns of 14 popular named dietary programmes for weight and cardiovascular risk factor reduction in adults: systematic review and network meta-analysis of randomised trials. *BMJ*. <https://doi.org/10.1136/bmj.m696>
36. Konieczna J, Ruiz-Canela M, Galmes-Panades AM, Abete I, Babio N, Fiol M, Martín-Sánchez V, Estruch R, Vidal J, Buil-Cosiales P, García-Gavilán JF, Moñino M, Marcos-Delgado A, Casas R, Olbryta R, Fitó M, Hu FB, Martínez-González MA, Martínez JA, Romaguera D, Salas-Salvado J (2023) An Energy-Reduced Mediterranean Diet, Physical Activity, and Body Composition. *JAMA Network Open*. <https://doi.org/10.1001/jamanetworkopen.2023.37994>
37. Zelicha H, Kloting N, Kaplan A, Yaskolka Meir A, Rinott E, Tsaban G, Chassidim Y, Blüher M, Ceglarek U, Isermann B, Stumvoll M, Quayson RN, von Bergen M, Engelmann B, Rolle-Kampczyk UE, Haange SB, Tuohy KM, Diotallevi C, Shelef I, Hu FB, Stampfer MJ, Shai I (2022) The effect of high-polyphenol Mediterranean diet on visceral adiposity: the DIRECT PLUS randomized controlled trial. *BMC Medicine*. <https://doi.org/10.1186/s12916-022-02525-8>
38. Muscogiuri G, Maiorino MI, Paolini B, Aversano F, Buscemi C, Cappiello I, Caruso I, Ceriani E, Chimienti M, Cicero FAG, Cintoni M, Desideri G, D'Eusebio C, Medea G, Patti MC, Randazzo C, Troiano E, Vitale M, Zimmiti S, Veronese N, Gianfredi V, Volpe M, Ricci M, Maggi S, Onder G, Silano M, Nucci D, Zanetti M, Fontana L, Casirati A, Leonardi F (2026) Mediterranean diet for the management of pre-existing metabolic diseases: Evidence from a systematic review and meta-analysis featured in the Italian national guidelines "La Dieta Mediterranea". *Nutrition*. <https://doi.org/10.1016/j.nut.2025.112975>
39. Salas-Salvado J, Bulló M, Estruch R, Ros E, Covas MI, Ibarrola-Jurado N, Corella D, Arós F, Gómez-Gracia E, Ruiz-Gutiérrez V, Romaguera D, Lapetra J, Lamuela-Raventós RM, Serra-Majem L, Pintó X, Basora J, Muñoz MA, Sorlí JV, Martínez-González MA (2014) Prevention of Diabetes With Mediterranean Diets. *Annals of Internal Medicine*. <https://doi.org/10.7326/m13-1725>
40. Martínez-González MA, Montero P, Ruiz-Canela M, Toledo E, Estruch R, Gómez-Gracia E, Li J, Ros E, Arós F, Hernández A, Corella D, Fiol M, Lapetra J, Serra-Majem L, Pintó X, Cofán M, Sorlí JV, Babio N, Márquez-Sandoval YF, Castañer O, Salas-Salvado J (2023) Yearly attained adherence to Mediterranean diet and incidence of diabetes in a large randomized trial. *Cardiovascular Diabetology*. <https://doi.org/10.1186/s12933-023-01994-2>

41. Ruiz-Canela M, Corella D, Martínez-González MA, Babio N, Martínez JA, Forga L, Alonso-Gómez AM, Wärnberg J, Vioque J, Romaguera D, López-Miranda J, Estruch R, Santos-Lozano JM, Serra-Majem L, Bueno-Cavanillas A, Tur JA, Martín-Sánchez V, Riera-Mestre A, Delgado-Rodríguez M, Matía-Martin P, Vidal J, Vázquez C, Damiel L, Buil-Cosiales P, Shyam S, Sorli JV, Castañer O, García-Rios A, Torres-Collado L, Gómez-Gracia E, Zulet MA, Konieczna J, Casas R, Cano-Ibáñez N, Tojal-Sierra L, Bernal-López RM, Toledo E, García-Gavilán J, Fernández-Carrión R, Goday A, Arenas-Larriva AP, González-Palacios S, Schröder H, Ros E, Fitó M, Hu FB, Tinahones FJ, Salas-Salvado J (2025) Comparison of an Energy-Reduced Mediterranean Diet and Physical Activity Versus an Ad Libitum Mediterranean Diet in the Prevention of Type 2 Diabetes. *Annals of Internal Medicine*. <https://doi.org/10.7326/annals-25-00388>

42. Sobiecki JG, Imamura F, Davis CR, Sharp SJ, Koulman A, Hodgson JM, Guevara M, Schulze MB, Zheng JS, Agnoli C, Bonet C, Colorado-Yohar SM, Fagherazzi G, Franks PW, Gundersen TE, Jannasch F, Kaaks R, Katzke V, Molina-Montes E, Nilsson PM, Palli D, Panico S, Papier K, Rolandsson O, Sacerdote C, Tjønneland A, Tong TYN, van der Schouw YT, Danesh J, Buterworth AS, Riboli E, Murphy KJ, Wareham NJ, Forouhi NG (2023) A nutritional biomarker score of the Mediterranean diet and incident type 2 diabetes: Integrated analysis of data from the MedLey randomised controlled trial and the EPIC-InterAct case-cohort study. *PLOS Medicine*. <https://doi.org/10.1371/journal.pmed.1004221>

43. Maroto-Rodríguez J, Ortolá R, Carballo-Casla A, Iriarte-Campo V, Salinero-Frío MA, Rodríguez-Artalejo F, Sotos-Prieto M (2023) Association between a mediterranean lifestyle and Type 2 diabetes incidence: a prospective UK biobank study. *Cardiovascular Diabetology*. <https://doi.org/10.1186/s12933-023-01999-x>

44. Schwingshackl L, Missbach B, König J, Hoffmann G (2015) Adherence to a Mediterranean diet and risk of diabetes: a systematic review and meta-analysis. *Public Health Nutrition*. <https://doi.org/10.1017/s1368980014001542>

45. Esposito K, Maiorino MI, Bellastella G, Chiodini P, Panagiotakos D, Giugliano D (2015) A journey into a Mediterranean diet and type 2 diabetes: a systematic review with meta-analyses. *BMJ Open*. <https://doi.org/10.1136/bmjopen-2015-008222>

46. Wu MJ, Hung CH, Yong SB, Ching GS, Hsu HJ (2025) Impact of the Mediterranean Diet on Glycemic Control, Body Mass Index, Lipid Profile, and Blood Pressure in Type 2 Diabetes: A Meta-Analysis of Randomized Controlled Trials. *Nutrients*. <https://doi.org/10.3390/nu17243908>

47. Dinu M, Pagliai G, Casini A, Sofi F (2018) Mediterranean diet and multiple health outcomes: an umbrella review of meta-analyses of observational studies and randomised trials. *European Journal of Clinical Nutrition*. <https://doi.org/10.1038/ejcn.2017.58>

48. Morze J, Danielewicz A, Przybyłowicz K, Zeng H, Hoffmann G, Schwingshackl L (2021) An updated systematic review and meta-analysis on adherence to mediterranean diet and risk of cancer. *European Journal of Nutrition*. <https://doi.org/10.1007/s00394-020-02346-6>

49. Schwingshackl L, Schwedhelm C, Galbete C, Hoffmann G (2017) Adherence to Mediterranean Diet and Risk of Cancer: An Updated Systematic Review and Meta-Analysis. *Nutrients*. <https://doi.org/10.3390/nu9101063>

50. Zhong Y, Zhu Y, Li Q, Wang F, Ge X, Zhou G, Miao L (2020) Association between Mediterranean diet adherence and colorectal cancer: a dose-response meta-analysis. *The American Journal of Clinical Nutrition*. <https://doi.org/10.1093/ajcn/nqaa083>

51. van den Brandt PA, Schulpen M (2017) Mediterranean diet adherence and risk of postmenopausal breast cancer: results of a cohort study and meta-analysis. *International Journal of Cancer*. <https://doi.org/10.1002/ijc.30654>

52. Toledo E, Salas-Salvado J, Donat-Vargas C, Buil-Cosiales P, Estruch R, Ros E, Corella D, Fitó M, Hu FB, Arós F, Gómez-Gracia E, Romaguera D, Ortega-Calvo M, Serra-Majem L, Pintó X, Schröder H, Basora J, Sorlí JV, Bulló M, Serra-Mir M, Martínez-González MA (2015) Mediterranean Diet and Invasive Breast Cancer Risk Among Women at High Cardiovascular Risk in the PREDIMED Trial. *JAMA Internal Medicine*. <https://doi.org/10.1001/jamainternmed.2015.4838>

53. Martínez-González MA, Sayón-Orea C, Bullón-Vela V, Bes-Rastrollo M, Rodríguez-Artalejo F, Yusta-Boyo MJ, García-Solano M (2022) Effect of olive oil consumption on cardiovascular disease, cancer, type 2 diabetes, and all-cause mortality: A systematic review and meta-analysis. *Clinical Nutrition*. <https://doi.org/10.1016/j.clnu.2022.10.001>

54. González-Palacios Torres C, Barrios-Rodríguez R, Muñoz-Bravo C, Toledo E, Dierssen T, Jiménez-Moleón JJ (2023) Mediterranean diet and risk of breast cancer: An umbrella review. *Clinical Nutrition*. <https://doi.org/10.1016/j.clnu.2023.02.012>

55. Fekete M, Varga P, Ungvari Z, Fekete JT, Buda A, Szappanos Á, Lehoczki A, Mózes N, Grosso G, Godos J, Menyhart O, Munkácsy G, Tarantini S, Yabluchanskiy A, Ungvari A, Györfy B (2025) The role of the Mediterranean diet in reducing the risk of cognitive impairment, dementia, and Alzheimer's disease: a meta-analysis. *GeroScience*. <https://doi.org/10.1007/s11357-024-01488-3>

56. Nucci D, Sommariva A, Degoni LM, Gallo G, Mancarella M, Ntarelli F, Savoia A, Catalini A, Ferranti R, Pregliasco FE, Castaldi S, Gianfridi V (2024) Association between Mediterranean diet and dementia and Alzheimer disease: a systematic review with meta-analysis. *Aging Clinical and Experimental Research*. <https://doi.org/10.1007/s40520-024-02718-6>

57. Shannon OM, Ranson JM, Gregory S, Macpherson H, Milte C, Lentjes M, Mulligan A, McEvoy C, Griffiths A, Matu J, Hill TR, Adamson A, Siervo M, Minihane AM, Muniz-Tererra G, Ritchie C, Mathers JC, Llewellyn DJ, Stevenson E (2023) Mediterranean diet adherence is associated with lower dementia risk, independent of genetic predisposition: findings from the UK Biobank prospective cohort study. *BMC Medicine*. <https://doi.org/10.1186/s12916-023-02772-3>

58. Fu J, Tan LJ, Lee JE, Shin S (2022) Association between the mediterranean diet and cognitive health among healthy adults: A systematic review and meta-analysis. *Frontiers in Nutrition*. <https://doi.org/10.3389/fnut.2022.946361>

59. Singh B, Parsaik AK, Mielke MM, Erwin PJ, Knopman DS, Petersen RC, Roberts RO (2014) Association of Mediterranean Diet with Mild Cognitive Impairment and Alzheimer's Disease: A Systematic Review and Meta-Analysis. *Journal of Alzheimer's Disease*. <https://doi.org/10.3233/jad-130830>

60. Martínez-Lapiscina EH, Clavero P, Toledo E, Estruch R, Salas-Salvado J, San Julián B, Sanchez-Tainta A, Ros E, Valls-Pedret C, Martínez-González MA (2013) Mediterranean diet improves cognition: the PREDIMED-NAVARRA randomised trial. *Journal of Neurology, Neurosurgery & Psychiatry*. <https://doi.org/10.1136/nnnp-2012-304792>

61. Barnes LL, Dhana K, Liu X, Carey VJ, Ventrelle J, Johnson K, Hollings CS, Bishop L, Laranjo N, Stubbs BJ, Reilly X, Agarwal P, Zhang S, Grodstein F, Tangney CC, Holland TM, Aggarwal NT, Arfanakis K, Morris MC, Sacks FM (2023) Trial of the MIND Diet for Prevention of Cognitive Decline in Older Persons. *New England Journal of Medicine*. <https://doi.org/10.1056/nejmoa2302368>

62. Chen H, Dhana K, Huang Y, Huang L, Tao Y, Liu X, Melo van Lent D, Zheng Y, Ascherio A, Willett W, Yuan C (2023) Association of the Mediterranean Dietary Approaches to Stop Hypertension Intervention for Neurodegenerative Delay (MIND) Diet With the Risk of Dementia. *JAMA Psychiatry*. <https://doi.org/10.1001/jamapsychiatry.2023.0800>

63. Gregory S, Buller-Peralta I, Bridgeman K, Góngora VDL, Dounavi M, Low A, Ntalianis G, O'Brien J, Parra MA, Ritchie CW, Ritchie K, Shannon OM, Stevenson EJ, Muniz-Terrera G (2024) The Mediterranean diet is not associated with neuroimaging or cognition in middle-aged adults: a cross-sectional analysis of the PREVENT dementia programme. *European Journal of Neurology*. <https://doi.org/10.1111/ene.16345>

64. Lassale C, Batty GD, Baghadi A, Jacka F, Sánchez-Villegas A, Kivimäki M, Akbaraly T (2019) Healthy dietary indices and risk of depressive outcomes: a systematic review and meta-analysis of observational studies. *Molecular Psychiatry*. <https://doi.org/10.1038/s41380-018-0237-8>

65. Shafiei F, Salari-Moghaddam A, Larjani B, Esmailzadeh A (2019) Adherence to the Mediterranean diet and risk of depression: a systematic review and updated meta-analysis of observational studies. *Nutrition Reviews*. <https://doi.org/10.1093/nutrit/nuy070>

66. Jacka FN, O'Neill A, Opie R, Itsiopoulos C, Cotton S, Mohebbi M, Castle D, Dash S, Mihalopoulos C, Chatterton ML, Brazionis L, Dean OM, Hodge AM, Berk M (2017) A randomised controlled trial of dietary improvement for adults with major depression (the 'SMILES' trial). *BMC Medicine*. <https://doi.org/10.1186/s12916-017-0791-y></



82. Meslier V, Laiola M, Roager HM, De Filippis F, Roume H, Quinquis B, Giacco R, Mennella I, Ferracane R, Pons N, Pasolli E, Rivellese A, Dragsted LO, Vitaglione P, Ehrlich SD, Ercolini D (2020) Mediterranean diet intervention in overweight and obese subjects lowers plasma cholesterol and causes changes in the gut microbiome and metabolome independently of energy intake. *Gut*. <https://doi.org/10.1136/gutjnl-2019-320438>
83. Ghosh TS, Rampelli S, Jeffery IB, Santoro A, Neto M, Capri M, Giampieri E, Jennings A, Candela M, Turroni S, Zoetendal EG, Hermes GDA, Elodie C, Meunier N, Brugere CM, Pujos-Guillot E, Berendsen AM, De Groot LCPGM, Feskens EJM, Kaluza J, Pietruszka B, Bielak MJ, Comte B, Maijo-Ferre M, Nicoletti C, De Vos WM, Fairweather-Tait S, Cassidy A, Brigidi P, Franceschi C, O'Toole PW (2020) Mediterranean diet intervention alters the gut microbiome in older people reducing frailty and improving health status: the NU-AGE 1-year dietary intervention across five European countries. *Gut*. <https://doi.org/10.1136/gutjnl-2019-319654>
84. Haro C, Montes-Borrego M, Rangel-Zúñiga OA, Alcalá-Díaz JF, Gómez-Delgado F, Pérez-Martínez P, Delgado-Lista J, Quintana-Navarro GM, Tinahones FJ, Landa BB, López-Miranda J, Camargo A, Pérez-Jiménez F (2016) Two Healthy Diets Modulate Gut Microbial Community Improving Insulin Sensitivity in a Human Obese Population. *The Journal of Clinical Endocrinology & Metabolism*. <https://doi.org/10.1210/jc.2015-3351>
85. Li J, Guasch-Ferré M, Chung W, Ruiz-Canela M, Toledo E, Corella D, Bhupathiraju SN, Tobias DK, Tabung FK, Hu J, Zhao T, Turman C, Feng YCA, Clish CB, Mucci L, Eliassen AH, Costenbader KH, Karlson EW, Wolpin BM, Ascherio A, Rimm EB, Manson JE, Qi L, Martínez-González MÁ, Salas-Salvadó J, Hu FB, Liang L (2020) The Mediterranean diet, plasma metabolome, and cardiovascular disease risk. *European Heart Journal*. <https://doi.org/10.1093/eurheartj/ehaa209>
86. Gao P, Rinott E, Dong D, Mei Z, Wang F, Liu Y, Kamer O, Yaskolka Meir A, Tuohy KM, Blüher M, Stumvoll M, Stampfer MJ, Shai I, Wang DD (2024) Gut microbial metabolism of bile acids modifies the effect of Mediterranean diet interventions on cardiometabolic risk in a randomized controlled trial. *Gut Microbes*. <https://doi.org/10.1080/19490976.2024.2426610>
87. García-Gavilán JF, Atzeni A, Babio N, Liang L, Belzer C, Vioque J, Corella D, Fitó M, Vidal J, Moreno-Indias I, Torres-Collado L, Coltell O, Toledo E, Clish C, Hernando J, Yun H, Hernández-Cacho A, Jeanfavre S, Dennis C, Gómez-Pérez AM, Martínez MA, Ruiz-Canela M, Tinahones FJ, Hu FB, Salas-Salvadó J (2024) Effect of 1-year lifestyle intervention with energy-reduced Mediterranean diet and physical activity promotion on the gut metabolome and microbiota: a randomized clinical trial. *The American Journal of Clinical Nutrition*. <https://doi.org/10.1016/j.ajcnut.2024.02.021>
88. Hoffman R, Gerber M (2013) Evaluating and adapting the Mediterranean diet for non-Mediterranean populations: A critical appraisal. *Nutrition Reviews*. <https://doi.org/10.1111/nure.12040>
89. Woodside J, Young IS, McKinley MC (2022) Culturally adapting the Mediterranean Diet pattern – a way of promoting more 'sustainable' dietary change? *British Journal of Nutrition*. <https://doi.org/10.1017/s0007114522001945>
90. Olmedo-Requena R, González-Donquiles C, Dávila-Batista V, Romaguera D, Castelló A, Molina de la Torre AJ, Amiano P, Dierssen-Sotos T, Guevara M, Fernández-Tardón G, Lozano-Lorca M, Alguacil J, Peiró R, Huerta JM, Gracia-Lavedan E, Aragonés N, Fernández-Villa T, Solans M, Gómez-Acebo I, Castaño-Vinyals G, Kogevinas M, Pollán M, Martín V (2019) Agreement among Mediterranean Diet Pattern Adherence Indexes: MCC-Spain Study. *Nutrients*. <https://doi.org/10.3390/nu11030488>
91. Jennings A, Shannon OM, Gillings R, Lee V, Elsworth R, Bundy R, Rao G, Hanson S, Hardeman W, Paddick SM, Siervo M, Aldred S, Mathers JC, Hornberger M, Miniham AM (2024) Effectiveness and feasibility of a theory-informed intervention to improve Mediterranean diet adherence, physical activity and cognition in older adults at risk of dementia: the MedEx-UK randomised controlled trial. *BMC Medicine*. <https://doi.org/10.1186/s12916-024-03815-z>
92. Biggi C, Biasini B, Ogrinc N, Strojnik L, Endrizzi I, Menghi L, Khémiri I, Mankai A, Slama FB, Jamoussi H, Riviou K, Elfazazi K, Rehman N, Scazzina F, Menozzi D (2024) Drivers and Barriers Influencing Adherence to the Mediterranean Diet: A Comparative Study across Five Countries. *Nutrients*. <https://doi.org/10.3390/nu16152405>
93. Bonekamp NE, Visseren FLJ, van der Schouw YT, van der Meer MG, Teraa M, Ruigrok YM, Geleijnse JM, Koopal C, UCC-SMART study group, Cramer MJ, Nathoe HM, van der Harst P, van de Meer MG, de Borst GJ, Teraa M, Bots ML, van Smeden M, Emmelot-Vonk MH, de Jong PA, Lely AT, van der Kaaij NP, Kappelle LJ, Ruigrok YM, Verhaar MC, Dorresteyn JAN, Visseren FLJ (2024) Cost-effectiveness of Mediterranean diet and physical activity in secondary cardiovascular disease prevention: results from the UCC-SMART cohort study. *European Journal of Preventive Cardiology*. <https://doi.org/10.1093/eurjpc/zwae123>
94. Lorca-Camara V, Bosque-Prous M, Bes-Rastrollo M, O'Callaghan-Gordo C, Bach-Faig A (2024) Environmental and Health Sustainability of the Mediterranean Diet: A Systematic Review. *Advances in Nutrition*. <https://doi.org/10.1016/j.advnut.2024.100322>
95. Dernini S, Berry E, Serra-Majem L, La Vecchia C, Capone R, Medina F, Aranceta-Bartrina J, Belahsen R, Burlingame B, Calabrese G, Corella D, Donini L, Lairon D, Meybeck A, Pekcan A, Piscopo S, Yngve A, Trichopoulos A (2017) Med Diet 4.0: the Mediterranean diet with four sustainable benefits. *Public Health Nutrition*. <https://doi.org/10.1017/s13688980016003177>
96. Martínez-González MA (2024) Should we remove wine from the Mediterranean diet?: a narrative review. *The American Journal of Clinical Nutrition*. <https://doi.org/10.1016/j.ajcnut.2023.12.020>
97. English LK, Ard JD, Bailey RL, Bates M, Bazzano LA, Boushey CJ, Brown C, Butera G, Callahan EH, de Jesus J, Mattes RD, Mayer-Davis EJ, Novotny R, Obbagy JE, Rahavi EB, Sabate J, Snetselaar LG, Stoody EE, Van Horn LV, Venkatraman S, Heymsfield SB (2021) Evaluation of Dietary Patterns and All-Cause Mortality. *JAMA Network Open*. <https://doi.org/10.1001/jamanetworkopen.2021.22277> ■