



POPSCI · CHEMICAL ECOLOGY

Inside the scent trail of a “mosquito magnet”

REFERENCES

13 verified

TOKENS

≈2.5k

VERIFICATION

Crossref

COMPILED

27 Aug 2026

Why do mosquitoes seem to feast on one person and ignore the next? The difference is real, but the best clues point not to sweet blood or a single magic molecule, but to the shifting chemical signature above our skin.

For more than three years, researchers asked *Aedes aegypti* mosquitoes to choose between nylon sleeves worn on people’s forearms. One person’s scent beat every rival’s almost every time; their attraction score was more than 100 times those of the two least attractive people. The pecking order held across months, mosquito strains and repeated samples.¹

That experiment turns a familiar complaint into a measurable trait: some people really are persistent mosquito magnets. Yet a bite begins long before the proboscis meets blood. A mosquito must first locate a living animal, decide that it is the right species, then pick among nearby individuals. The emerging answer is less like a blood-type label and more like a scent trail assembled from breath, warmth, skin chemistry and microbes—with each mosquito species weighting the clues differently.^{2,3}

01 First, find something alive

Carbon dioxide from breath can wake up a mosquito’s search, while body odor and heat help guide the approach. But these channels overlap. *Aedes aegypti* engineered without a working carbon-dioxide receptor still showed only a slight reduction in attraction to live humans in a large semi-field test, apparently because odor and heat could compensate.⁴

In a large outdoor flight cage in Zambia, *Anopheles gambiae* barely visited human-temperature targets that lacked carbon dioxide or body odor. Add breath and scent, however, and the mosquitoes chose not only a human target but one sleeping person over another.⁵ The mosquito is not following a single beacon. It is cross-checking a bundle of signals.

02 The acids in your invisible halo

The strongest trail currently leads to skin odor. In the sleeve study, highly attractive people carried larger amounts of several carboxylic acids—a broad family of compounds present in human skin emissions—and the association appeared again in a validation group. Diluting the most attractive person’s sleeve erased the mosquitoes’ preference, although the study could not show that any one acid was sufficient on its own.¹ Follow-up receptor experiments confirmed that acids linked to human attraction activate acid-sensitive mosquito receptors.⁶

The Zambia study landed nearby from a much more naturalistic direction: people preferred by *Anopheles gambiae* emitted relatively more butyric, isobutyric and isovaleric acids, plus acetoin, a compound that skin microbes can produce.⁵ Still, chemistry is contextual. Four geographic populations of *Aedes aegypti* differed more than 300-fold in the dose of lactic acid needed to attract them, and lactic acid alone attracted fewer than 40% of mosquitoes versus more than 87% for whole human odor.⁷ A scent blend is a chord, not a note.

03 The microbes behind the perfume

Human skin bacteria transform sweat, oils and shed material into airborne molecules, making the skin microbiome a plausible upstream source of individual scent. Among 48 people, those highly attractive to *Anopheles gambiae* had more culturable skin bacteria but lower bacterial diversity than poorly attractive people.⁸ A later study of *Anopheles coluzzii* also linked particular bacterial groups and odor compounds with high or low attractiveness.⁹

These are associations, not proof that rearranging a person’s microbes would rearrange the biting queue. Reviews of the field identify the microbiome as promising but still entangled with host chemistry, environment and mosquito behavior.¹⁰ The perfume may be microbial in part; its recipe is not yet settled.

04 The plot twist: there is no universal mosquito magnet

A 2026 experiment offered odors from 119 people to three mosquito species. Each species produced a different human ranking and tracked different chemical features; even sex differences appeared only for *Aedes aegypti*.¹¹ What attracts a yellow-fever mosquito may not impress an Asian tiger mosquito or a southern house mosquito.

Human genetics may influence the raw material. In a small study of 18 identical and 19 non-identical female twin pairs, attraction to hand odor was more alike within identical pairs, yielding a heritability estimate of 0.62 (standard error 0.124). But the narrow, older cohort and wide uncertainty leave the responsible genes unknown.¹²

Blood type makes a cleaner story—and a weaker one. Laboratory *Aedes albopictus* preferred type-O blood when different blood groups were offered through artificial membrane feeders. But that design deliberately removed human odor and tested a meal after the host-finding problem had already been solved; it cannot show that mosquitoes are more likely to fly toward a person with type O.¹³



05 What the mosquito is really choosing

Return to those scented sleeves. Their owner's place in the ranking persisted because the sleeve carried a remarkably durable chemical calling card, not because a mosquito could inspect the blood beneath it. The best evidence says that acid-rich skin-odor blends help explain who gets noticed, while breath and heat guide the search and microbes

and genes may shape the blend. But “mosquito magnet” is a relationship, not a permanent human category: change the mosquito species, setting or odor mixture, and the winner can change too. The decisive next experiment is one that manipulates a specific part of human scent safely, then shows across several mosquito species and real-world settings that the biting order changes with it.

References

13 · VERIFIED VIA CROSSREF

1. De Obaldia ME, Morita T, Dedmon LC, Boehmler DJ, Jiang CS, Zeledon EV, Cross JR, Vossall LB (2022) Differential mosquito attraction to humans is associated with skin-derived carboxylic acid levels. *Cell*. <https://doi.org/10.1016/j.cell.2022.09.034>
2. Coutinho-Abreu IV, Riffell JA, Akbari OS (2022) Human attractive cues and mosquito host-seeking behavior. *Trends in Parasitology*. <https://doi.org/10.1016/j.pt.2021.09.012>
3. Blanken SL, Prudhomme O'Meara W, Hol FJH, Bousema T, Markwalter CF (2024) À la carte: how mosquitoes choose their blood meals. *Trends in Parasitology*. <https://doi.org/10.1016/j.pt.2024.05.007>
4. McMeniman CJ, Corfas RA, Matthews BJ, Ritchie SA, Vossall LB (2014) Multimodal Integration of Carbon Dioxide and Other Sensory Cues Drives Mosquito Attraction to Humans. *Cell*. <https://doi.org/10.1016/j.cell.2013.12.044>
5. Giraldo D, Rankin-Turner S, Corver A, Tauxe GM, Gao AL, Jackson DM, Simubali L, Book C, Stevenson JC, Thuma PE, McCoy RC, Gordus A, Mburu MM, Simulundu E, McMeniman CJ (2023) Human scent guides mosquito thermotaxis and host selection under naturalistic conditions. *Current Biology*. <https://doi.org/10.1016/j.cub.2023.04.050>
6. Ray G, Huff RM, Castillo JS, Bellantuono AJ, DeGennaro M, Pitts RJ (2023) Carboxylic acids that drive mosquito attraction to humans activate ionotropic receptors. *PLOS Neglected Tropical Diseases*. <https://doi.org/10.1371/journal.pntd.0011402>
7. Williams CR, Ritchie SA, Russell RC, Eiras AE, Kline DL, Geier M (2006) Geographic Variation in Attraction to Human Odor Compounds by *Aedes aegypti* Mosquitoes (Diptera: Culicidae): A Laboratory Study. *Journal of Chemical Ecology*. <https://doi.org/10.1007/s10886-006-9097-9>
- Verhulst NO, Qiu YT, Beijleveld H, Maliepaard C, Knights D, Schulz S, Berg-Lyons D, Lauber CL, Verduijn W, Haasnoot GW, Mumm R, Bouwmeester HJ, Claas FHJ, Dicke M, van Loon JJA, Takken W, Knight R, Smallegange RC (2011) Composition of Human Skin Microbiota Affects Attractiveness to Malaria Mosquitoes. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0028991>
- Showering A, Martinez J, Benavente ED, Gezan SA, Jones RT, Oke C, Tytheridge S, Pretorius E, Scott D, Allen RL, D'Alessandro U, Lindsay SW, Armour JAL, Pickett J, Logan JG (2022) Skin microbiome alters attractiveness to *Anopheles* mosquitoes. *BMC Microbiology*. <https://doi.org/10.1186/s12866-022-02502-4>
- Martinez J, Showering A, Oke C, Jones RT, Logan JG (2021) Differential attraction in mosquito-human interactions and implications for disease control. *Philosophical Transactions of the Royal Society B: Biological Sciences*. <https://doi.org/10.1098/rstb.2019.0811>
- Marrero KM, Castillo JS, Lucas-Barbosa D, Bellantuono AJ, Marrero MA, Cid D, Costa-da-Silva AL, Verhulst NO, DeGennaro M (2026) Individual humans are more attractive to certain mosquito species. *iScience*. <https://doi.org/10.1016/j.isci.2026.117006>
- Fernández-Grandon GM, Gezan SA, Armour JAL, Pickett JA, Logan JG (2015) Heritability of Attractiveness to Mosquitoes. *PLOS ONE*. <https://doi.org/10.1371/journal.pone.0122716>
- Bursali F, Şimşek FM (2023) Blood Feeding Preferences of Laboratory-Reared *Aedes albopictus* for Human Blood Groups and Its Effect on Their Fertility. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi*. <https://doi.org/10.9775/kvfd.2023.29881> ■