

Buoyancy fluxes of sea-ice leads in the Arctic

Josué Martínez-Moreno¹, Camille Lique¹, Claude Talandier¹, someone else?

¹Laboratoire d’Océanographie Physique et Spatiale (LOPS), University of Brest/IFREMER/IRD/CNRS,
Brest, France

Key Points:

- Buoyancy fluxes in Linear Kinematic Features vary seasonally
- Buoyancy fluxes in Linear Kinematic Features can be 20-30% larger than in the ice-pack.
- The presence of Linear Kinematic Features has the potential transform water masses in the ocean surface.

Corresponding author: Josué Martínez-Moreno, josue.martinez.moreno@ifremer.fr

Abstract

The Arctic sea-ice, in particular the ice-pack, acts as an insulator between the atmosphere and the ocean below. Linear kinematic features (LKF), or leads, common in Arctic sea ice serves as openings that allow exchanges between ocean and atmosphere, for example: heat, salt, moisture, and gases. Therefore, leads can influence heat exchanges, ice melt or growth, ocean dynamics, and the atmospheric boundary layer. The scale of leads varies from a few meters to kilometers in width, and from few kilometers up to hundreds of kilometers in length. Their formation is driven by the winds and ocean currents. Existing literature has suggested that leads play a significant role in the heat budget of the surface of the ocean and likely impact water mass transformation within the Arctic. However, due to the lack of observations and high resolution models resolving LKFs, an estimate of the magnitude of the impact in the ocean has been elusive. Here, we explore the magnitude and impact in the ocean surface caused by the LKFs using SEDNA, a high resolution pan-Arctic hindcast ($1/60^\circ$; NEMO + SI3 configuration). We show that despite the LKF only account for a small area of the sea-ice cover (5%), they contribute up to 20-30% more buoyancy than the ice-pack at the surface. Furthermore, of the total surface water mass transformation in the Arctic ice cover, LKFs explain approximately 10% of it. LKFs make a significant contribution despite their small area coverage; however, the presented estimate shows a large local impact but to a lesser degree in the Arctic Ocean than previously hypothesized.

1 Introduction

Arctic sea-ice plays a crucial role in regulating Earth's climate by acting as a natural insulator between the atmosphere and the ocean below (Wettlaufer et al., 1997; Untersteiner, 1961). This insulating property acts as a thermal barrier, limiting heat loss from the ocean to the atmosphere and minimizing evaporation, thereby maintaining the stability of the polar environment (Ruffieux et al., 1995). A ubiquitous feature of Arctic sea ice is the formation and persistence of linear kinematic features (LKF), commonly known as leads. These features are elongated openings in the sea ice cover that expose the ocean surface to the atmosphere (Lüpkes et al., 2008).

Leads vary in scale from a few meters to kilometers in width and span distances of a few kilometers up to hundreds of kilometers (Tschudi et al., 1998). Their formation mechanisms are diverse, ranging from mechanical to thermodynamic processes (Ram-

pal et al., 2016; Hutchings et al., 2005). Mechanically, sea ice deformation and fracturing are caused by winds and ocean currents (Richter-Menge et al., 2002). Winds are thought to be the main forcing of the sea ice deformation, where persistent winds stress result in sea ice fracturing and eventually leads (Linow & Dierking, 2017; Hutter et al., 2018). Although ocean currents have a smaller contribution to the sea ice deformation compared to wind forcing, ocean drag at the bottom of the ice acts to balance the wind-induced sea ice velocities (Weiss & Marsan, 2004). Additionally, observations show that ocean currents may play a bigger role in dictating the fracturing patterns of sea ice in regions with strong currents and bathymetric features (Willmes et al., 2023). Alongside these mechanical forces, thermodynamic processes such as rapid variation of the air or ocean temperature can result in the fracturing of sea ice (Lewis, 1998). Once a lead is formed, warm air temperatures can instigate localized melting, maintaining the lead open. Conversely, during colder periods, rapid freezing facilitate the closure of a lead. The intricate interplay between these thermodynamic processes and dynamic forces dictates the evolution and persistence of leads.

Leads are crucial since they act as gateways between the atmosphere and the ocean, exchanging heat, moisture, and gases (Michaelis & Lüpkes, 2022; Andreas et al., 2002). Depending on the season and the atmospheric forcing, this direct contact between the ocean water and atmosphere can result in locally intensified heat fluxes at the ocean-atmosphere interface (Marcq & Weiss, 2012; Maykut, 1986), localized ice melt or growth, and modify the ocean dynamics (Bourgault et al., 2020). For example, in winter, heat exchange from the ocean to the cold atmosphere, in addition to ice freezing within the leads, results in local convective processes in both the ocean and atmosphere (Lüpkes et al., 2008; McPhee et al., 2005). A change in the surface temperature and salinity due to the heat and salt flux at the ocean-ice -atmosphere interfaces transforms locally the surface water masses (Walsh, 1982). Leads have a undeniable local impact on the ocean and it has been hypothesized that leads may play a significant role in the Arctic by modulating the stratification, altering ocean circulation, and transformation of water masses (Reiser et al., 2020; Smith et al., 1990). However, the magnitude and influence of the LKFs compared to the surrounding ice pack in the Arctic basin is yet to be quantified.

To investigate and quantify the impact of the LKFs in the Arctic Ocean, we use a really high-resolution pan-Arctic hindcast, SEDNA (Talandier & Lique, 2023). Then, we compute a mask of the linear kinematic features during the year 2014, as well as the




Figure 1. Linear kinematic features from SEDNA. a) Linear kinematic mean frequency during winter between 2011 and 2015. b) Snapshot of the total deformation for the XX of XX. c) True color image from Sentinel 3 on the 12th of April 2022 for the magenta area in panel a. d) Mean ice concentration from SEDNA (800m resolution) on the 21st of April 2014. e) Total deformation used to identifies the linear kinematic features. f) Identified linear kinematic features on the 21st of April 2014.

buoyancy fluxes and water mass transformation for the identified LKFs, and compare them to the ice pack (See Methods for details of the calculations and numerical model).

Observation vs Models

Climate models and high resolution sea ice models are unable to reproduce LKFs due to the lack of resolution to capture the small footprint of LKFs in the Arctic (Wang et al., 2016). SEDNA is capable of overcoming this limitation due to its spatial resolution of $\sim 800m$ on average in the Arctic. The climatology of the winter lead probability, i.e. the likelihood of a lead occurring for any given day during winter, is shown in

Figure 1a. There are several distinct spatial patterns that match those from observations (Willmes et al., 2023). For example, we observe increased lead frequency in the shelves, along bathymetric features, and in regions with large kinetic energy. Several hotspots can be identified along the sea shelves and in the Chukchi Sea, highlighting the model’s ability to reproduce LKF dynamics. **Should we something more robust to show that the model does a good job?**

SEDNA is capable of reproducing the overall pattern of the observed winter lead distribution in the Arctic shelves, the Greenland Sea, Barents Sea, Kara Sea and Laptev Sea. However, observations show a prominent signature in the Beaufort Sea that is not captured in our simulation using the same winter definition. Lagging the winter definition by a month, allows for a better representation of the LKFs in the Beaufort Sea. A snapshot of the total deformation is shown in Figure 1b, where the local maxima in the deformation correspond to the linear kinematic features identified by the algorithm described in the Methods section. A closer examination, focusing north of Greenland in panels c, d, e, and f of Figure 1, reveals similarities between the spatial pattern of typically observed leads on the 12th of April 2022 from a true color satellite image (Fig. 1c) and the ice concentration of SEDNA for the 21st of April 2014. Note that the sea ice concentration in the identified leads shown in Fig.1e-f remains higher than 80% concentration; this is a consequence of the visco-plastic nature of the sea ice model and the daily averaged output that smooths the presence of LKFs in the fields. Furthermore, in our simulation, but in fact, in all numerical models using a visco-plastic rheology, it has been observed that they do not reproduce the typical fracturing angles as in observations (). Despite this, the important similarities between the observations and model lead probability allow us for the first time to estimate quantitatively the impact LKF have in the ocean.

In situ observations have shown that LKFs in winter can have heat fluxes up to two orders of magnitude larger than those of the surrounding ice covered ocean (). Presumably impacting the buoyancy flux and the circulation of the Arctic Ocean. Figure 2a and e show snapshots of the buoyancy flux, revealing the typical patterns of the Arctic. In winter (Fig. 2a) we observe an overall negative buoyancy flux, a consequence of the freezing of sea ice and thus brine rejection. Meanwhile, in summer, there is an overall positive buoyancy flux due to the melting and freshening of the Arctic (Fig. 2b). Overall, in these two maps we observe that inside the ice covered region (magenta contour), the

fluxes are spatially homogeneous, except for the MIZ and the LKFs. Regarding the LKFs, we observe that these features have larger buoyancy fluxes than the surroundings ice covered ocean.

In figure 2, we plot two transects on the 1st of January 2014 and 30th of June 2014 showing the heat flux (panels b, f), freshwater heat-equivalent flux (panels c and g), and ice thickness and buoyancy flux (panels d and h). Examining the 1st of January 2014, we observe a small heat flux in the LKFs (Fig.2 b); blue shaded areas) that does not impact the heat flux at the ocean-atmosphere interface because the ocean surface is at freezing point. Consequently, the extra heat flux is used for sea ice growth. In contrast, the salt flux (Fig.2c) shows two peaks with twice the magnitude of the surrounding ice cover collocated with the identified LKFs. A positive freshwater flux corresponds to a salinification of the ocean surface due to sea ice growth and a negative buoyancy flux that increases the density of the water at the surface (Fig.2d). As expected, there is a decrease in ice thickness where the LKFs are identified. On the 25th of June (Fig.2f), we observe an increase in the heat fluxes within some of the identified LKFs to up to $\approx 25W/m^2$. Meanwhile, the freshwater flux shows a decrease within the LKFs, indicating a freshening of the surface due to sea ice melt. This change is captured by the positive buoyancy flux, decreasing the density at the surface (Fig.2g and h). Similarly to the winter snapshot, we note a decrease in ice thickness of more than $1m$ within the LKFs. It is important to note that some peaks in the buoyancy flux align with the edge of the detected leads, likely due to the enhanced refreezing or melting that occur at the edge of the LKFs. Furthermore, not all the identified LKFs exhibit significant ocean-atmosphere fluxes. For these features, we explored snapshots of the simulation a few days later and noted that the LKFs are starting to open, and thus the ocean and ice still behave as they do in the ice pack (**Check this idea?**).

As in observations, we detect heat fluxes larger in the LKFs than in the surrounding ice pack in both summer and winter, however, the freshwater heat-equivalent flux due to melting and freezing is significantly larger than the heat fluxes associated with the LKFs. By identifying the daily mask of the LKFs, we explore the evolution of the buoyancy fluxes over the full year of 2014 in the next section.




Figure 2. Maps of buoyancy flux and transects of downward heat flux, freshwater heat-equivalent flux, ice thickness, and mixed layer depth for the 1st of January and 25th of June 2014. a) Buoyancy flux in Winter the 1st of January 2014. Transect on the 1st of January 2014 of b) downward heat flux, c) freshwater heat-equivalent flux, d) ice thickness and mixed layer depth. e) Buoyancy flux in Summer the 30th of June 2014. Transect on the 25th of June 2014 of b) downward heat flux, c) freshwater heat-equivalent flux, d) ice thickness (magenta) and buoyancy flux (green). Note that the transect location and orientation changes between the winter and summer dates. Magenta contour in panels a and e correspond to the 0% concentration contour of sea ice. Vertical blue bars shows the mask of LKFs in the transect.

Impact of LKF in the ocean buoyancy

We now consider the spatially averaged impact of LKFs on ocean buoyancy. We start by looking at the percentage area covered by three distinct masks of the sea ice covered Arctic Ocean: i) the daily identified LKF, ii) the marginal ice zone (15-80% sea ice concentration), and iii) the ice pack (80-100% sea ice concentration). The areas covered by the different masks show a seasonal cycle; during summer the percentage of the pack ice cover area drops from approximately 90% to 50% and the marginal ice zone increases from 10% to approximately 50%. Meanwhile, the percentage of the area covered by LKFs never exceeds more than 5% of the sea ice covered Arctic and its seasonal cycle reaches a maximum in the summer season, since thinner ice can break more easily.

To demonstrate the role of leads in buoyancy flux, we multiply the buoyancy flux and area weighted them by the respective masks of the sea ice cover; the MIZ, the ice pack, and the LKF. The buoyancy flux shows an important seasonal cycle, where positive values correspond to melting season of sea ice, thus forming lighter waters in the ocean surface, and negative values correspond to the freezing season of sea ice, thus forming denser waters. The buoyancy flux in the MIZ forms generally lighter waters (Fig.3b), this suggests that in the MIZ there is a preferential melting of sea ice, where both the haline and thermal components of the buoyancy flux act to lighten the ocean surface (Fig.3c and d). Over the seasonal cycle, the pack ice changes sign from winter to summer (Fig.3b), this sign change corresponds to a shift from ice formation in winter (formation of denser waters) to ice melting in summer (formation of lighter waters), where the primary contribution to the buoyancy flux arises from the haline buoyancy component. Analogous to the ice pack, the buoyancy in the LKF has a seasonal cycle and it shows a similar temporal evolution as the ice pack. Although LKFs cover less than 5% of the total Arctic ice cover area, over certain periods of time the LKF buoyancy flux can be 10%-20% larger than that of the ice pack.

To further quantify the role of the ice pack and the LKF on the overturning circulation of the Arctic, we use the water mass transformation framework (WMT; Walin, 1982). This approach gives us information on the formation of lighter or denser water due to changes in the surface fluxes. In agreement with previous studies, the surface freshwater flux dominates Arctic water mass transformation. The yearly mean WMT transformation of the ice pack (Fig. 4a) is characterized by a broad region of strong positive




Figure 3. Time-series during 2014 of a) the area cover, b) buoyancy fluxes, c) haline buoyancy flux component, and d) thermal buoyancy flux component for linear kinematic features, transitional ice zone (ice concentration 0-80%), and pack ice (ice concentration >80%).

transformation (losing buoyancy) in the density range of $22.5 < \gamma_n < 27.5$, reaching a maximum of 1.5 Sv, and a negative transformation (gaining buoyancy) below $\gamma_n < 22.5$. The yearly mean WMT in LKF also has a positive peak within the $22.5 < \gamma_n < 27.5$ reaching up to 0.15 Sv, in other words 10% of the yearly water mass transformation in the ice covered Arctic.

The daily WMT of the ice pack and LKF are shown in Figure 4b and c respectively. Note that the magnitude of the WMT in the LKF is 10% of that in the ice pack. In both we observe a seasonal cycle. In the winter months, a positive transformation occurs meaning a loss of buoyancy, due to brine rejection due to sea ice growth. In the transition between summer and winter, we reach almost zero transformation due to the fact that both




Figure 4. Mean and daily Water mass transformation over 2014. a) Mean water mass transformation over 2014 for the ice pack and LKF. Hovmoller diagram of the water mass transformation (Sv) for the b) Pack Ice and c) linear kinematic features. Pane d) is the ratio of water mass transformation between the LKF and the Pack Ice. Note the magnitude change in the colorbar of panels a), b), and c).

denser and lighter waters are been formed as ice freezes and melts. In summer, a negative transformation or a gain of buoyancy occurs due to the melting of sea ice. More importantly, the WMT in the ice pack and the LKF show a similar seasonal behavior, where some small differences can be observed in the WMT ratio between the LKF and ice pack. The major differences in the density classes can be observed in $\gamma_n < 22.5$ and $\gamma_n > 27.5$, although this signal is likely to arise in LKF due to their capacity to form fresher water and denser water when the ocean is in direct contact with the atmosphere, the magnitude of the WMT in these density classes are negligible.

2 Conclusions

Linear kinematic features are ubiquitous in the sea ice covered oceans that allow direct exchanges between the ocean and the atmosphere, whereby fresher and denser water can be formed at the ocean surface depending on the atmospheric forcing. Generally, the importance of these fluxes at the ocean-atmosphere interface have been speculated to be significantly important for the Arctic dynamics (). The present study assesses the impact of LKFs in the ocean surface by diagnosing the buoyancy fluxes using high-resolution hindcast (SEDNA). SEDNA reproduces the LKF spatial distribution over most of the Arctic, with exception of the Beaufort Sea. This mismatch is caused by a one-month

delay in the simulated LKF. In other words, there is a temporal shift in the simulated LKF frequency and those of observed. This mismatch could have important implications in the Arctic circulation, thus further research is needed to better understand this temporal shift in the formation of leads.

Although LKFs have a significant local impact, they contribute only a small fraction to the overall buoyancy flux and water mass transformation of the Arctic. Despite their observable impact on the direct exchanges between the ocean and the atmosphere, LKFs make up only a fraction of the total sea ice cover in the Arctic. Similarly, in terms of water mass transformation, LKFs account for a limited portion of the overall transformation process. While their presence can lead to localized changes in the density of surface waters due to brine rejection and ice melting, their spatial extent and duration are not sufficient to significantly alter the ocean dynamics of the Arctic. Our findings show that LKF cover a small area of the ice-covered Arctic and they have a contribution of up to 10% of the buoyancy flux and water mass transformation.

LKFs generated through shear have been shown to have a significant impact on ocean surface dynamics. For example, McPhee et al. (2005) observed a change in the heat fluxes in the ocean subsurface up to two orders of magnitude larger compared to the surrounding areas. Although our definition to identify LKFs include this process, we do not observe such signatures in our simulation. Therefore, further research is required to better understand different LKF generation mechanisms and their potential impact in the Arctic Ocean.

Our results may also be sensitive to the specific sea ice and vertical mixing parameterizations used in our model, particularly those relating to ice/ocean fluxes, such as brine rejection, enhanced vertical diffusivity, and vertical exchanges. Finally, it remains to be shown whether LKFs can locally impact regions of the Arctic basin where the sea ice and stratification may allow for a more significant impact on the ocean below LKFs.

3 Open Research

4 Methods

Model description

Here we employ the NEMO numerical platform (release 4.0.5) to establish a $1/60^\circ$ pan-Arctic ocean-sea ice configuration named SEDNA. The ocean core (NEMO-OCE)

solves the primitive equations (PE) using finite differences on an Arakawa C-grid mesh, while the ocean is coupled to the SI3 sea-ice component (NEMO-SI3). The pan-Arctic domain covers the Bering Strait on the Pacific side and extends southward to 56°N in the Subpolar North Atlantic gyre and to the Baltic Sea entrance. The ocean model incorporates a linear free surface, utilizes a 3rd order flux form scheme for momentum advection, and employs a 4th order Flux Corrected Transport (FCT) for tracer advection. Additionally, the sea-ice representation includes a 5-category model with the Elasto-Visco-Plastic (EVP) rheology activated and a land-fast capability to better simulate sea-ice behavior in specific areas.

The NCAR bulk formula from Large & Yeager (2009) to calculate air-sea fluxes and uses hourly ERA5 atmospheric data for near-surface variables. The three lateral boundaries are constrained with daily GLORYS12V1 Re-analysis data, and monthly freshwater fluxes from river discharges are combined with Greenland land ice melt data to supply SEDNA. The simulation extends over seven years (2009 to 2015) and starts from rest, utilizing initial conditions based on the World Ocean Atlas 2009 temperature/salinity and mean January 2009 ice state from PIOMAS re-analysis. Preliminary assessment against available observations and re-analyses data shows that the simulation reasonably captures key ocean and sea-ice characteristics.

LKF identification

The LKF from SEDNA were identified by using the algorithm proposed by Hutter et al. (2019). This method uses the total deformation rate defined as:

$$T_d = \left[\underbrace{\left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right)^2}_{Divergence} + \underbrace{\left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right)^2 + \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} \right)^2}_{Shear} \right]^{1/2}, \quad (1)$$

where u and v are the sea ice velocities. The deformation rate varies spatially depending on the background deformation and the sea ice properties (), where the largest deformation rates occur along the boundaries of ice flows where the LKF are found. Therefore, to identify the LKFs it is required to identify the local maxima using edge detection (DoG filter) in the deformation rate field. After the edge detection, using binary map we obtain a mask with the original width of identified LKF. This mask is then used to compute the statistics presented in this study. After this step, Hutter et al. (2019) reduce the LKFs width to a single pixel to then track the LKF over time. Since our study

focus on the ocean-atmosphere interactions, we only use the identification algorithm to produce a mask of LKFs during 2014 for the Arctic, where we retain the original width computed from the deformation rate. This mask is then used to quantify and contrast the impact of LKFs compared to the ice-pack.

Buoyancy flux and water mass transformation

- Describe MLD definition
- Describe buoyancy flux and decomposition between thermal and haline components.

Acknowledgments

References

- Andreas, E. L., Guest, P. S., Persson, P. O. G., Fairall, C. W., Horst, T. W., Moritz, R. E., & Semmer, S. R. (2002). Near-surface water vapor over polar sea ice is always near ice saturation. *Journal of Geophysical Research: Oceans*, 107(C10). doi: 10.1029/2000jc000411
- Bourgault, P., Straub, D., Duquette, K., Nadeau, L.-P., & Tremblay, B. (2020). Vertical Heat Fluxes beneath Idealized Sea Ice Leads in Large-Eddy Simulations: Comparison with Observations from the SHEBA Experiment. *Journal of Physical Oceanography*, 50(8), 2189–2202. doi: 10.1175/jpo-d-19-0298.1
- Hutchings, J. K., Heil, P., & Hibler, W. D. (2005). Modeling Linear Kinematic Features in Sea Ice. *Monthly Weather Review*, 133(12), 3481–3497. doi: 10.1175/mwr3045.1
- Hutter, N., Losch, M., & Menemenlis, D. (2018). Scaling Properties of Arctic Sea Ice Deformation in a High-Resolution Viscous-Plastic Sea Ice Model and in Satellite Observations. *Journal of Geophysical Research: Oceans*, 123(1), 672–687. doi: 10.1002/2017jc013119
- Hutter, N., Zampieri, L., & Losch, M. (2019). Leads and ridges in Arctic sea ice from RGPS data and a new tracking algorithm. *The Cryosphere*, 13(2), 627–645. doi: 10.5194/tc-13-627-2019
- Lewis, J. K. (1998). Thermomechanics of pack ice. *Journal of Geophysical Research:*

- 293 *Oceans*, 103(C10), 21869–21882. doi: 10.1029/98jc01256
- 294 Linow, S., & Dierking, W. (2017). Object-Based Detection of Linear Kinematic Fea-
 295 tures in Sea Ice. *Remote Sensing*, 9(5), 493. doi: 10.3390/rs9050493
- 296 Lüpkes, C., Vihma, T., Birnbaum, G., & Wacker, U. (2008). Influence of leads in
 297 sea ice on the temperature of the atmospheric boundary layer during polar night.
 298 *Geophysical Research Letters*, 35(3). doi: 10.1029/2007gl032461
- 299 Marcq, S., & Weiss, J. (2012). Influence of sea ice lead-width distribution on turbu-
 300 lent heat transfer between the ocean and the atmosphere. *The Cryosphere*, 6(1),
 301 143–156. doi: 10.5194/tc-6-143-2012
- 302 Maykut, G. A. (1986). The surface heat and mass balance. In N. Untersteiner (Ed.),
 303 *The geophysics of sea ice* (pp. 395–463). Boston, MA: Springer US. doi: 10.1007/
 304 978-1-4899-5352-0_6
- 305 McPhee, M. G., Kwok, R., Robins, R., & Coon, M. (2005). Upwelling of arctic
 306 pycnocline associated with shear motion of sea ice. *Geophysical Research Letters*,
 307 32(10). doi: 10.1029/2004GL021819
- 308 Michaelis, J., & Lüpkes, C. (2022). The Impact of Lead Patterns on Mean Profiles
 309 of Wind, Temperature, and Turbulent Fluxes in the Atmospheric Boundary Layer
 310 over Sea Ice. *Atmosphere*, 13(1), 148. doi: 10.3390/atmos13010148
- 311 Rampal, P., Bouillon, S., Ólason, E., & Morlighem, M. (2016). neXtSIM: a new La-
 312 grangian sea ice model. *The Cryosphere*, 10(3), 1055–1073. doi: 10.5194/tc-10
 313 -1055-2016
- 314 Reiser, F., Willmes, S., & Heinemann, G. (2020). A New Algorithm for Daily Sea
 315 Ice Lead Identification in the Arctic and Antarctic Winter from Thermal-Infrared
 316 Satellite Imagery. *Remote Sensing*, 12(12), 1957. doi: 10.3390/rs12121957
- 317 Richter-Menge, J. A., McNutt, S. L., Overland, J. E., & Kwok, R. (2002). Re-
 318 lating arctic pack ice stress and deformation under winter conditions. *Jour-*
 319 *nal of Geophysical Research: Oceans*, 107(C10), SHE 15–1–SHE 15–13. doi:
 320 10.1029/2000jc000477
- 321 Ruffieux, D., Persson, P. O. G., Fairall, C. W., & Wolfe, D. E. (1995). Ice pack and
 322 lead surface energy budgets during leadex 1992. *Journal of Geophysical Research:*
 323 *Oceans*, 100(C3), 4593–4612. doi: 10.1029/94JC02485
- 324 Smith, S. D., Muench, R. D., & Pease, C. H. (1990). Polynyas and leads: An
 325 overview of physical processes and environment. *Journal of Geophysical Research:*

- 326 *Oceans*, 95(C6), 9461–9479. doi: 10.1029/jc095ic06p09461
- 327 Talandier, C., & Lique, C. (2023, February). *Sedna-delta*. Zenodo. Retrieved from
 328 <https://doi.org/10.5281/zenodo.7656924> doi: 10.5281/zenodo.7656924
- 329 Tschudi, M., Curry, J., & Maslanik, J. (1998). Airborne observations of leads in the
 330 Beaufort Sea. *IGARSS '98. Sensing and Managing the Environment. 1998 IEEE*
 331 *International Geoscience and Remote Sensing. Symposium Proceedings. (Cat.*
 332 *No.98CH36174)*, 2, 986–988 vol.2. doi: 10.1109/igarss.1998.699648
- 333 Untersteiner, N. (1961). On the mass and heat budget of arctic sea ice. *Archiv*
 334 *für Meteorologie, Geophysik und Bioklimatologie, Serie A*, 12(2), 151–182. doi: 10
 335 .1007/bf02247491
- 336 Walin, G. (1982). On the relation between sea-surface heat flow and thermal circu-
 337 lation in the ocean. *Tellus*, 34(2), 187–195. doi: 10.1111/j.2153-3490.1982.tb01806
 338 .X
- 339 Wang, Q., Danilov, S., Jung, T., Kaleschke, L., & Wernecke, A. (2016). Sea ice leads
 340 in the Arctic Ocean: Model assessment, interannual variability and trends. *Geo-*
 341 *physical Research Letters*, 43(13), 7019–7027. doi: 10.1002/2016gl068696
- 342 Weiss, J., & Marsan, D. (2004). Scale properties of sea ice deformation and fractur-
 343 ing. *Comptes Rendus Physique*, 5(7), 735–751. doi: 10.1016/j.crhy.2004.09.005
- 344 Wettlaufer, J. S., Worster, M. G., & Huppert, H. E. (1997). the phase evolution of
 345 Young Sea Ice. *Geophysical Research Letters*, 24(10), 1251–1254. doi: 10.1029/
 346 97gl00877
- 347 Willmes, S., Heinemann, G., & Schnaase, F. (2023). Patterns of wintertime Arctic
 348 sea ice leads and their relation to winds and ocean currents. *The Cryosphere Dis-*
 349 *cussions*, 2023, 1–23. doi: 10.5194/tc-2023-22