



## GT03

### Geometric Structures in Manifolds

#### Estructuras Geométricas en Variedades Diferenciables

##### Organizers

##### Organizadores

##### Antolatzaileak

**Guillermo Sánchez Arellano**

(Universidad Complutense de Madrid)

**Pablo Nicolás Martínez**

(Centre de Recerca Matemàtica)

**Benedetta Facciotti**

(Universitat Politècnica de Catalunya)

**Robert Cardona Aguilar**

(Universitat de Barcelona)

##### Description

##### Descripción


##### Deskribapena

*The main object of study of differential topology are differentiable manifolds. Their behavior is in many instances too flexible, so they are commonly endowed with a geometric structure which simplifies and enriches their study. Such structures can be of various types: Riemannian metrics, symplectic forms, contact distributions, complex structures, spin structures, foliations... All of them behave differently, but one can commonly observe relations between techniques from different areas.*

*Here we provide a link between geometers from different fields mainly focused on symplectic and contact geometry, as well as in their applications to different areas such as topology and dynamical systems.*

El comportamiento de las variedades diferenciables puede resultar demasiado flexible en algunas ocasiones, por lo que se les suele agregar algún tipo de estructura geométrica que enriquezca y simplifique su estudio. Algunos ejemplos son: métricas riemannianas, formas simplécticas, distribuciones de contacto, estructuras complejas, estructuras de spin o foliaciones. Aunque cada una presenta diferentes comportamientos, en ocasiones pueden encontrarse relaciones entre las técnicas empleadas en cada área.

Con esta sesión tratamos de poner en contacto a geómetras de diferentes áreas, enfocándonos especialmente en las geometrías simpléctica y de contacto y en sus aplicaciones a la topología y a los sistemas dinámicos

MSC Codes	Códigos MSC	MSC Kodeak
	57R17 (primary)  57R15; 37J39 (secondary)	
Slots	Bloques	Blokeak
	1.A (Aula 0.19S); 1.B (Aula 0.19S); 1.C (Aula 0.19S)	
QR Code	Código QR	QR Kodea
		
Session Schedule	Horario de la Sesión	Saioaren Ordutegia

L13   17:30-17:50   0.19S <i>Hodge structures on configuration spaces of orbits</i> <b>Alejandro Calleja</b> (UCM-ICMAT)		
L13   18:00-18:20   0.19S <i>Complex and symplectic geometry in dimension 8</i> <b>Luis Pizarro Golvano</b> (Universidad de Zaragoza)		
L13   18:30-18:50   0.19S <i>The cosymplectic Chern--Hamilton conjecture</i> <b>Søren István Adorján Dyhr</b> (Universitat Politècnica de Catalunya & Centre de Recerca Matemàtica)		
L13   19:00-19:20   0.19S <i>On integrable contact systems and bi-Hamiltonian structures</i> <b>Asier López-Gordón</b> (Institute of Mathematics of the Polish Academy of Sciences)		

M14 | 15:00-15:20 | 0.19S

*Prelegendrian submanifolds in elliptic distribution*

**Wei Zhou** (UCM-ICMAT)

M14 | 15:30-15:50 | 0.19S

*Geometric formalism of Poisson-Poincaré reduction*

**Miguel A. Berbel** (Universidad Pontificia Comillas)

M14 | 16:00-16:20 | 0.19S

*Symplectic invariant connections*

**José Luis Carmona Jiménez** (IMAR)

M14 | 16:30-16:50 | 0.19S

*A Gentle Introduction to Generalized Riemannian Geometry*

**Jaime Pedregal Pastor** (Utrecht University)

M14 | 17:30-17:50 | 0.19S

*Floer Theory for the 3-Body Problem*

**Jagna Wi niewska** (Universitat Politècnica de Catalunya)

M14 | 18:00-18:20 | 0.19S

*Morse Theory Applied to Topological Quantum Field Theories*

**Enrique Aycart Maldonado** (Complutense University of Madrid)

M14 | 18:30-18:50 | 0.19S

*A sketch of the Atiyah-Singer index theorem and its extensions*

**Josep Fontana McNally** (Universitat Politècnica de Catalunya)

**Monday 13****17:30-17:50****[Room 0.19S]****Lunes 13****17:30-17:50****[Aula 0.19S]****Astelehena 13****17:30-17:50****[Gela 0.19S]*****Hodge structures on configuration spaces of orbits*****Alejandro Calleja**

(UCM-ICMAT)

Given an algebraic variety  $X$  with an action of an algebraic group  $G$ , we define the  $n$ -th configuration space of orbits as the set of  $n$ -tuples of points of  $X$  such that the orbits through  $G$  of these points are pairwise disjoint. In this talk we introduce these spaces and their applications to Knot Theory. We will also show how we can study the Hodge structure of these spaces by relating it to the one of  $X$ .

[arXiv:2403.07765](https://arxiv.org/abs/2403.07765)**Monday 13****18:00-18:20****[Room 0.19S]****Lunes 13****18:00-18:20****[Aula 0.19S]****Astelehena 13****18:00-18:20****[Gela 0.19S]*****Complex and symplectic geometry in dimension 8*****Luis Pizarro Golvano**

(Universidad de Zaragoza)

The objective of this talk is to ask some questions about complex non-Kähler geometry and symplectic geometry by working with some examples of homogeneous spaces that are built from a nilpotent Lie group. This type of homogeneous spaces are known as nilmanifolds and they are the key in order to build examples and counter examples in problems of geometric structures in manifolds. In addition, nilmanifolds allow you to study geometry using linear algebra and Lie Groups theory.

**Monday 13**  
**18:30-18:50**  
**[Room 0.19S]**

**Lunes 13**  
**18:30-18:50**  
**[Aula 0.19S]**

**Astelehena 13**  
**18:30-18:50**  
**[Gela 0.19S]**

*The cosymplectic Chern--Hamilton conjecture*

**Søren István Adorján Dyhr**

(Universitat Politècnica de Catalunya & Centre de Recerca Matemàtica)

We study a functional on compatible metrics on compact, 3-dimensional, cosymplectic manifolds. It generalizes the functional studied by Chern and Hamilton in 1984 for contact manifolds.

We classify which manifolds admit critical metrics by computing local forms and making explicit constructions: they are either co-Kähler or mapping tori of the 2-torus by maps conjugate to hyperbolic toral automorphisms. In the first case, the Reeb vector field is Killing, in the second it is Anosov.

Joint work with Ángel González-Prieto, Eva Miranda, and Daniel Peralta-Salas.

**Monday 13**  
**19:00-19:20**  
**[Room 0.19S]**

**Lunes 13**  
**19:00-19:20**  
**[Aula 0.19S]**

**Astelehena 13**  
**19:00-19:20**  
**[Gela 0.19S]**

*On integrable contact systems and bi-Hamiltonian structures*

**Asier López-Gordón**

(Institute of Mathematics of the Polish Academy of Sciences)

We have recently proven a Liouville-Arnol'd theorem for contact Hamiltonian systems. In this setting, one has a  $(2n+1)$ -dimensional contact manifold with  $n+1$  functions in involution w.r.t. the Jacobi bracket, and the invariant submanifolds are  $n$ -codimensional and coisotropic. In a work in progress, we utilize Jacobi-Nijenhuis structures and extend the theory of bi-Hamiltonian systems to obtain those functions in involution in a contact manifold.

Joint work with Leonardo Colombo, Manuel de León, María Emma Eyrea Irazú, and Manuel Lainz.

[arXiv:2302.12061](https://arxiv.org/abs/2302.12061)

Tuesday 14

15:00-15:20

[Room 0.19S]

Martes 14

15:00-15:20

[Aula 0.19S]

Asteartea 14

15:00-15:20

[Gela 0.19S]

*Prelegendrian submanifolds in elliptic distribution***Wei Zhou**

(UCM-ICMAT)

The maximally non-involutive  $(4, 6)$ -distribution has two classes: elliptic and hyperbolic. The latter was shown to exhibit an h-principle by Javier Martínez and Álvaro del Pino. For the elliptic case, there is a natural association with a contact 7-manifold, an  $S^1$ -fibration. In this talk, we introduce the counterpart to Legendrian submanifolds in elliptic distributions and discuss the potential existence of non-isotopic PreLegendrian submanifolds in  $\mathbb{R}^6$ .

Joint work with Álvaro del Pino.

Tuesday 14

15:30-15:50

[Room 0.19S]

Martes 14

15:30-15:50

[Aula 0.19S]

Asteartea 14

15:30-15:50

[Gela 0.19S]

*Geometric formalism of Poisson-Poincaré reduction***Miguel A. Berbel**

(Universidad Pontificia Comillas)

This talk presents the covariant bracket formulation of first-order Hamiltonian field theories, introducing specific polysymplectic and multisymplectic structures on the configuration bundle. When the physical system has a symmetry, a  $G$ -principal connection can be used to reduce these geometric structures and simplify its description. We focus on the Poisson–Poincaré reduction and related canonical structures that may play a crucial role in this reduction process.

**Tuesday 14**  
**16:00-16:20**  
**[Room 0.19S]**

**Martes 14**  
**16:00-16:20**  
**[Aula 0.19S]**

**Asteartea 14**  
**16:00-16:20**  
**[Gela 0.19S]**

***Symplectic invariant connections***  
**José Luis Carmona Jiménez**  
(IMAR)

Homogeneous spaces are differentiable manifolds with a transitive action of a Lie group. The Ambrose-Singer Theorem characterizes Riemannian homogeneous spaces through an invariant tensor that satisfies covariant equations and is central to the Tricerri-Vanhecke program, which focuses on Riemannian homogeneous manifolds. However, it is limited to metric cases.

We present a generalization of the Ambrose-Singer Theorem for non-metric instances and apply this to symplectic homogeneous manifolds.

Joint work with Marco Castrillón López.

[arXiv:2001.06254](https://arxiv.org/abs/2001.06254)

**Tuesday 14**  
**16:30-16:50**  
**[Room 0.19S]**

**Martes 14**  
**16:30-16:50**  
**[Aula 0.19S]**

**Asteartea 14**  
**16:30-16:50**  
**[Gela 0.19S]**

***A Gentle Introduction to Generalized Riemannian Geometry***  
**Jaime Pedregal Pastor**  
(Utrecht University)

Generalized geometry has proven to be a powerful unifying framework for different geometries such as complex, symplectic, Poisson and the like. The theory of generalized metrics has led to further links to other geometries, e.g. bihermitian geometry, which are also of interest in supersymmetric sigma models in string theory. In this talk we will give a gentle introduction to generalized geometry, focusing in particular on the Riemannian aspect of the theory.

*Tuesday 14*  
*17:30-17:50*  
*[Room 0.19S]*

**Martes 14**  
**17:30-17:50**  
**[Aula 0.19S]**

**Asteartea 14**  
**17:30-17:50**  
**[Gela 0.19S]**

*Floer Theory for the 3-Body Problem*

**Jagna Wi niewska**

(Universitat Politècnica de Catalunya)

In my talk I will explain how to use tools from symplectic geometry and Hamiltonian dynamics to tackle the following problem: Can we send a rocket between any two points in the gravitational field of the Moon and the Earth, using the engines only at the beginning and at the end of the journey?

*Tuesday 14*  
*18:00-18:20*  
*[Room 0.19S]*

**Martes 14**  
**18:00-18:20**  
**[Aula 0.19S]**

**Asteartea 14**  
**18:00-18:20**  
**[Gela 0.19S]**

*Morse Theory Applied to Topological Quantum Field Theories*

**Enrique Aycart Maldonado**

(Complutense University of Madrid)

Topological Quantum Field Theories (TQFTs) are powerful algebro-geometric tools that provide us with a new way of computing topological and algebraic invariants of closed manifolds. In particular, cobordism categories play a fundamental role in the formulation of TQFTs, providing a geometric framework that connects category theory, topology and theoretical physics. We will focus on the application of Morse theory to the study of the two-dimensional case and the associated bicategory of bordisms.



*Tuesday 14*  
*18:30-18:50*  
*[Room 0.19S]*

**Martes 14**  
**18:30-18:50**  
**[Aula 0.19S]**

**Asteartea 14**  
**18:30-18:50**  
**[Gela 0.19S]**

*A sketch of the Atiyah-Singer index theorem and its extensions*

**Josep Fontana McNally**

(Universitat Politècnica de Catalunya)

The Atiyah-Singer index theorem brings together ideas from analysis, geometry, and topology, and generalizes many important results such as the Gauss-Bonnet and Riemann-Roch theorems. In this talk, we present the key objects in the statement, sketch one of the different proofs that have been given, and describe an extension to manifolds with boundary. We discuss how this last extension is the starting point of  $b$ -geometry, a framework to work with singular differential forms.