

Name- Ankit Meena

Roll.No- 2003121002

I am Ankit Meena, Roll.No. - 2003121002  
here by declare that during the course of  
this exam have not used any means of  
communication through phone, chat or any messaging  
app or social media app to discuss regarding  
this exam with any human or any bot.



08-03-2021



Problem: 1

(a)

we know  $I_d(s) = I_d(0) e^{-\tau_d}$  (A)

The relation in terms of flux can be written -

$S_d(s) = S_d(0) e^{-\tau_d(h)}$  (B)

where  $S_d(s)$  = flux at distance  $s$

$S_d(0)$  = flux without absorption

So apparent magnitude related to flux  $\rightarrow$

$$m = -2.5 \log S + \text{const.}$$

$$S \propto 10^{-0.4m}$$

Now  $\frac{S_d}{S_{d,0}} = 10^{-0.4(m-m_0)}$

$$\frac{S_d}{S_{d,0}} = e^{-\tau_d} = 10^{-\log \tau_d}$$

then

$$A_d = m - m_0 = -2.5 \log \left( \frac{S_d}{S_{d,0}} \right) = 2.5 \ln \tau_d$$

$A_d = 1.086 \tau_d$  A.P.



1(b) Use the Tully-Fisher relation to estimate the  $d_{pc} \rightarrow$   
we know the H-I line width  $w \sim 70 \text{ km/s}$

I can plot the absolute magnitude  $M$ . So

I can derive the distance to the galaxy using  
the standard relation -

$$M = m - 5 \log_{10} d_{pc} + 5 - A \quad (1)$$

~~In given galaxy for H-band~~

$$M_H = -9.50 + \log_{10} w - 2.50 = -21.67$$

where -  $d_{pc}$  = distance in parsec

$A$  = dust attenuation along the  
line of sight

To convert the measured H-I line width  $w$   
into an absolute magnitude  $M$  in the H-Band

$M_H$  (in given units)

$$M_H = -9.50 (\log_{10} w - 2.50) - 21.67 \quad (2)$$



Anshu

Ankit Meena

2003/21002

4

where  $w \approx 70 \text{ km/s}$

∴ we are dealing with flux measurement in the infrared, where the effects of dust attenuation are typically very small

then neglect the attenuation correction  $A$  in equation ① -

then 
$$M = m - 5 \log_{10} D_{pc} + 5$$

⇒ 
$$D_{pc} = 10^{\frac{m_H - M_H - 5}{-5}} \quad \text{--- ②}$$

from equation ②  $w \approx 70 \text{ km/s}$

$$m_H \approx -9.50 \cdot (\log_{10} 70 - 2.50) - 21.67 \approx -15.44$$

$$m_H \approx -15.44$$

from equation ③ put  $m_H \approx -15.44$ ,  $m_H = 15.1$

then 
$$D_{pc} = 10^{\frac{-15.44 + 15.1 - 5}{-5}} \approx 12.8 \text{ Mpc}$$

$$D_{pc} = 12.8 \text{ Mpc}$$



Ankit

Ankit Meena

2003121002

5

I(d)

Fundamental plane for elliptical galaxies:-

⇒ we get a relation between the surface brightness and effective radius

~~Equation (1)~~

$$R_e \propto \langle I_e \rangle^{-0.83} \quad \text{--- (A)}$$

where  $\langle I_e \rangle$  = average surface brightness

effective radius:-

$$L = 2\pi R_e^2 \langle I_e \rangle \quad \text{--- (B)}$$

from (A) & (B) -

$$L \propto R_e^2 \propto \langle I_e \rangle \langle I_e \rangle^{-0.66}$$

$$\text{so } \langle I_e \rangle \propto L^{-1.5} \quad \text{--- (C)}$$



Anurag

Ankit Meena

2003/21002

6

Electrical galaxies distribution in the 3-D  
parameter space ( $R_e$ ,  $\langle I_e \rangle$ ,  $\sigma_0$ ) is located  
close to a plane defined by -

$$R_e \approx \sigma_0^{1.4} \langle I_e \rangle^{0.85} \quad (1)$$

where  $\sigma_0$  = central velocity  
dispersion.

log form of  $\langle R_e \rangle$  -

$$\log R_e \approx 0.34 \langle M \rangle_c + 1.4 \log \sigma_0 + \text{const.} \quad (2)$$

$\sigma_0$

where  $\langle M \rangle_c$  is average surface  
brightness in

$R_e$

Equation (2) defines a plane in 3-D parameter space  
known as fundamental plane.



Amit

Ankit Meena

2003121002

(7)

Problem 2 →

(a)

given in questions -

⇒ Coma cluster has a radial velocity = 6750 km/s.

⇒  $R_e$  → effective radius = 50'

⇒ 3-dimensional velocity dispersion = 900 km/s

$R_{gravitational radius} = 2.5 R_{effective radius}$

we know mass estimate

$$M = \frac{R_{gr} \cdot \langle v^2 \rangle}{G}$$

$$\text{Viral mass} \Rightarrow M_{vir} = \frac{3 R_{vir} \sigma_v^2}{G}$$

where  $\sigma_v$  = velocity dispersion

$$M_{vir} = \frac{4\pi}{3} \cdot \Delta c \cdot \rho_{cr} \cdot R_{vir}^3$$

$$\Delta c \approx 200 \text{ \& } \rho_{cr} = 10^{-26} \text{ kg/m}^3$$



Ankit Meena

Ankit Meena

2003/21002

8

so equality the two selection ~~condition~~  $\rightarrow$

$$\frac{4\pi}{3} \times 200 \times 10^{-26} \times R_{vir}^3 = \frac{\sigma_{vir} \times (900 \times 10^3)^2}{57}$$

$$\sigma_{vir}^2 = \frac{(900 \times 10^3)^2 \times 3}{200 \times 4\pi \times 10^{-26} \times 57}$$

$$\sigma_{vir} = 1.45 \times 10^{27}$$

so using estimate mass (galaxy cluster).

$$M_N = \frac{R_v \sigma_v^2}{57}$$

$$M_N = \frac{(900 \times 10^3)^2 \times 1.45 \times 10^{27}}{6.67 \times 10^{11}}$$

$$M_{vir} = 1.76 \times 10^{27}$$



Ankit

Ankit Meena

2003121002

9

3(c)

given in question -

$$\text{displacement } z = 0.15 z_0$$

$$\Rightarrow z(t=0) = 0.15 z_0$$

$$\Rightarrow z(t) \text{ (now) } = z_0$$

$$\S \frac{m g(t=0)}{m g(t)} = \frac{50}{13}$$

using  $\rightarrow$

$$z(t) = z(t=0) + p \ln \left[ \frac{m g(t=0)}{m g(t)} \right]$$

put above value -

$$z_0 = 0.15 z_0 + p \ln \left[ \frac{50}{13} \right]$$

$$z_0 = 0.15 z_0 + p (1.345)$$

$$p = \frac{z_0 - 0.15 z_0}{1.345} = 0.637 z_0$$

=



Ankit Meena

Ankit Meena  
2003121002

(10)

4  
(d)

$\Rightarrow$  Synchrotron emission from a source ~~is~~ when

we have polarization in linear so -

$$\frac{I_{\perp}}{I_{\parallel}} = \frac{\int_0^{\infty} [F(x) + G(x)] dx}{\int_0^{\infty} [F(x) - G(x)] dx}$$

where   $I_{\perp}$  = perpendicular component of intensity

$I_{\parallel}$  = parallel component of the intensity,

the function obtained along  $I_{\parallel}(\omega)$  &  $I_{\perp}(\omega)$   
where  $G(x)$  is obtained by subtracting  $I_{\parallel}(\omega)$   
from  $I_{\perp}(\omega)$ . then -

$$\text{Also } \int_0^{\infty} x^{\mu} f(x) dx = \frac{2^{\mu+1}}{(\mu+2)} \Gamma\left(\frac{\mu}{2} + \frac{7}{3}\right) \pi\left(\frac{\mu}{2} + \frac{7}{3}\right)$$

where  $\Gamma$  = gamma function



Dr. Ankit Meena

Ankit Meena  
2003/2/002

(11)

$$\Rightarrow \int_0^{\infty} x^N G(x) dx = 2^N \Gamma\left(\frac{N}{2} + \frac{1}{3}\right) \Gamma\left(\frac{N}{2} + \frac{2}{3}\right)$$

then setting  $N=0$  we get-

$$\frac{I_{\perp}}{I_{\parallel}} = \frac{\Gamma(7/3) \Gamma(2/3) + \Gamma(4/3) \Gamma(2/3)}{\Gamma(7/3) \Gamma(2/3) - \Gamma(4/3) \Gamma(2/3)}$$

$$\frac{I_{\perp}}{I_{\parallel}} = \frac{\Gamma(2/3) \cdot \{\Gamma(7/3) + \Gamma(4/3)\}}{\Gamma(2/3) \{\Gamma(7/3) - \Gamma(4/3)\}}$$

$$\therefore \Gamma(7/3) = \frac{4}{3} \Gamma(4/3)$$

$$= \frac{\frac{4}{3} + 1}{\frac{4}{3} - 1} \frac{\Gamma(4/3)}{\Gamma(4/3)} = 7$$

$$\frac{I_{\perp}}{I_{\parallel}} = 7 \Rightarrow 7:1$$

Scattered polarization is defined -

$$\pi = \frac{I_{\perp}(\omega) - I_{\parallel}(\omega)}{I_{\perp}(\omega) + I_{\parallel}(\omega)} = \frac{G(x)}{F(x)}$$



Answer

Ankit neeraj  
2003/2002

12

Now if the electron have energy spectrum.

$$N(E) = K E^{-\phi} dE$$

So then

$$\pi = \frac{\int_0^{\infty} G(n) x^{(p-3)/2} dn}{\int_0^{\infty} f(n) x^{(p-3)/2} dn}$$

$$\pi = \frac{p+1}{4} \frac{\Gamma(p/4 + 7/12)}{\Gamma(p/4 + 19/12)} = \frac{p+1}{p+7/3}$$

$$\pi = \frac{p+1}{p+7/3}$$

~~So for a typical value of  $p=2$  then  $\pi = \frac{9}{13} \approx 70\%$   
So this is maximum value of linear polarization  
in all type of electron selection.~~

for  $p=2$ , then  $\pi = \frac{9}{13} \approx 70\%$



Anil K. Meena

Ankit Meena

2003/21002

13

Problem: 5g

5 (b) Superluminal motion has been found to occur in many of celestial bodies with peculiar events.

For some active galaxies, the distribution around the line  $B_{app} = \delta$

where  $B_{app}$  = apparent transverse velocity

$\delta$  = Doppler factor.

a weak correlation of  $B_{app}$  with  $\delta$  in


constant  $B_{app}$  is positively correlated with Lorentz factor  $\gamma$  or  $\gamma^2 (1+z)^{-1}$

⇒ In astronomy superluminal motion is the apparently faster than light motion seen in some radio galaxies.

$$L_{app} = \delta^4 L$$

$$\delta = \gamma / (1 - \beta \cos \theta)$$

$$B_{app} = \beta \sin \theta / (1 - \beta \cos \theta)$$





Anubhav

Anubhav  
2003121002

14

5(b)

Flat spectrum radio quasars and narrow-line Seyfert 1 (NLS1)  $\rightarrow$

$\Rightarrow$  AGNs can be divided according to the radio power as radio-loud and radio-quiet. Objects  $\sim$  Almost 10% of the AGNs are radio loud. ~~all~~ all others fall in the category of radio-quiet objects.

① Radio loud AGNs or radio galaxies have emission line spectra similar to the Seyferts - 1 but they are extremely bright on the radio wavelength source.

② AGNs with the broad permitted lines. Emission from hot, high-velocity gas that is near the black hole. as type-2 AGNs in the radio-quiet group. These include the Seyfert - 1 galaxies and high-luminosity radio-quiet quasars.



Ankit

Ankit meena

2003/2/002

15

5(d)

M87 - black hole image →

⇒ Supermassive black holes are bathed in glowing gas that is heated to billions of degrees. Because their extreme gravity traps light, so black holes cast a shadow on this bright emission. The shadow is surrounded by bright photon ring.

⇒ The photon ring is composed of series of increasingly sharp subrings. Each subring is produced by photons that traveled around the black hole ~~image~~  $\frac{n}{2}$  times. Before.

reaching the observer. So these subrings stack up to give the full image.

