

# THE SKEW-NORMAL APPROXIMATION OF THE BINOMIAL DISTRIBUTION

Joyce Tipping

Truman State University

Spring 2011

# INTRODUCTION

## DEFINITION (BINOMIAL)

Let  $X$  represent the number of successes in a sequence of  $n$  independent Bernoulli trials with probability of success  $p$  on each trial.

# INTRODUCTION

## DEFINITION (BINOMIAL)

Let  $X$  represent the number of successes in a sequence of  $n$  independent Bernoulli trials with probability of success  $p$  on each trial.

$X$  has discrete pdf

$$f_X(x) = \binom{n}{x} p^x q^{n-x}$$

# INTRODUCTION

## DEFINITION (BINOMIAL)

Let  $X$  represent the number of successes in a sequence of  $n$  independent Bernoulli trials with probability of success  $p$  on each trial.

$X$  has discrete pdf

$$f_X(x) = \binom{n}{x} p^x q^{n-x}$$

and cdf

$$F_X(x) = P(X \leq x) = \sum_{k=0}^x f_X(k).$$

# INTRODUCTION

The binomial cdf is easy to calculate for small  $n$  ...

# INTRODUCTION

The binomial cdf is easy to calculate for small  $n$  ...

But as  $n$  gets larger, it becomes increasingly difficult.

# INTRODUCTION

The binomial cdf is easy to calculate for small  $n$  ...

But as  $n$  gets larger, it becomes increasingly difficult.

For example ...

# INTRODUCTION

When  $n = 3$ ,

$$F(1) = \binom{3}{1} p^1 q^2 + \binom{3}{0} p^0 q^3$$



# INTRODUCTION

When  $n = 3$ ,

$$F(1) = \binom{3}{1} p^1 q^2 + \binom{3}{0} p^0 q^3$$

When  $n = 25$ ,

$$\begin{aligned} F(12) = & \binom{25}{12} p^{12} q^{13} + \binom{25}{11} p^{11} q^{14} + \binom{25}{10} p^{10} q^{15} + \binom{25}{9} p^9 q^{16} \\ & + \binom{25}{8} p^8 q^{17} + \binom{25}{7} p^7 q^{18} + \binom{25}{6} p^6 q^{19} + \binom{25}{5} p^5 q^{20} \\ & + \binom{25}{4} p^4 q^{21} + \binom{25}{3} p^3 q^{22} + \binom{25}{2} p^2 q^{23} + \binom{25}{1} p^1 q^{24} \\ & + \binom{25}{0} p^0 q^{25} \end{aligned}$$

# INTRODUCTION

A common technique is to use the normal distribution as an approximation:

$$F_X(x) \approx \Phi\left(\frac{x + 0.5 - \mu}{\sigma}\right),$$

where  $\mu = np$ ,  $\sigma = \sqrt{np(1-p)}$ , and  $\Phi$  is the standard normal cdf.

# INTRODUCTION

A common technique is to use the normal distribution as an approximation:

$$F_X(x) \approx \Phi\left(\frac{x + 0.5 - \mu}{\sigma}\right),$$

where  $\mu = np$ ,  $\sigma = \sqrt{np(1-p)}$ , and  $\Phi$  is the standard normal cdf.

When does this work well? ...

# INTRODUCTION

A common technique is to use the normal distribution as an approximation:


$$F_X(x) \approx \Phi\left(\frac{x + 0.5 - \mu}{\sigma}\right),$$

where  $\mu = np$ ,  $\sigma = \sqrt{np(1-p)}$ , and  $\Phi$  is the standard normal cdf.

When does this work well? ... In a nutshell, when the binomial is symmetric.


# INTRODUCTION

The binomial is symmetric when  $p = 0.5$




# INTRODUCTION

The binomial is symmetric when  $p = 0.5$  or  $n$  is very large.



# INTRODUCTION


However, when  $n$  is medium and  $p$  is extreme ...



the binomial is very skewed ...

# INTRODUCTION

However, when  $n$  is medium and  $p$  is extreme ...




and the normal approximation doesn't work very well.



# INTRODUCTION

However, when  $n$  is medium and  $p$  is extreme ...



Introducing ... the skew-normal distribution.

# OUTLINE

Today's itinerary:

1. Skew-Normal distribution – basic properties

# OUTLINE

Today's itinerary:

1. Skew-Normal distribution – basic properties
2. Method of Moments – derive an approximation

# OUTLINE

Today's itinerary:

1. Skew-Normal distribution – basic properties
2. Method of Moments – derive an approximation
3. Accuracy – examine the accuracy of our approximation

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

## DEFINITION (SKEW-NORMAL)

Let  $Y$  be a skew-normal distribution, with location parameter  $\mu \in \mathbb{R}$ , scale parameter  $\sigma > 0$ , and shape parameter  $\lambda \in \mathbb{R}$ . Then  $Y$  has pdf

$$f(x|\mu, \sigma, \lambda) = \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi\left(\frac{\lambda(x - \mu)}{\sigma}\right), \quad x \in \mathbb{R},$$

where  $\phi$  is the standard normal pdf and  $\Phi$  is the standard normal cdf.

We write  $Y \sim SN(\mu, \sigma, \lambda)$ .

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

So where did this funny-looking pdf come from? ...

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

So where did this funny-looking pdf come from? ...

## LEMMA

*If  $f_0$  is a one-dimensional probability density function symmetric about 0, and  $G$  is a one-dimensional distribution function such that  $G'$  exists and is a density symmetric about 0, then*

$$f(z) = 2 \cdot f_0(z) \cdot G\{w(z)\} \quad (-\infty < z < \infty)$$

*is a density function for any odd function  $w(\cdot)$ . (Lemma 1, ?)*

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

So where did this funny-looking pdf come from? ...

## LEMMA

*If  $f_0$  is a one-dimensional probability density function symmetric about 0, and  $G$  is a one-dimensional distribution function such that  $G'$  exists and is a density symmetric about 0, then*

$$f(z) = 2 \cdot f_0(z) \cdot G\{w(z)\} \quad (-\infty < z < \infty)$$

*is a density function for any odd function  $w(\cdot)$ . (Lemma 1, ?)*

► Kernel



# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

So where did this funny-looking pdf come from? ...

## LEMMA

*If  $f_0$  is a one-dimensional probability density function symmetric about 0, and  $G$  is a one-dimensional distribution function such that  $G'$  exists and is a density symmetric about 0, then*

$$f(z) = 2 \cdot f_0(z) \cdot G\{w(z)\} \quad (-\infty < z < \infty)$$

*is a density function for any odd function  $w(\cdot)$ . (Lemma 1, ?)*

- ▶ Kernel
- ▶ CDF

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

Basic properties:

$$E(Y) = \mu + b\delta\sigma$$

$$E(Y^2) = \mu^2 + 2b\delta\mu\sigma + \sigma^2$$

$$E(Y^3) = \mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3$$

$$\text{Var}(Y) = \sigma^2(1 - b^2\delta^2)$$

$$\text{where } b = \sqrt{\frac{2}{\pi}} \text{ and } \delta = \frac{\lambda}{\sqrt{1 + \lambda^2}}. \text{ (?)}$$

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when  $\lambda = 0$ ?

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when  $\lambda = 0$ ?

$$f(x|\mu, \sigma, \lambda = 0) = \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0)$$

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when  $\lambda = 0$ ?

$$\begin{aligned}f(x|\mu, \sigma, \lambda = 0) &= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0) \\&= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot 0.5\end{aligned}$$

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when  $\lambda = 0$ ?

$$\begin{aligned}f(x|\mu, \sigma, \lambda = 0) &= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0) \\&= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot 0.5 \\&= \frac{1}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right)\end{aligned}$$

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when  $\lambda = 0$ ?

$$\begin{aligned}f(x|\mu, \sigma, \lambda = 0) &= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0) \\&= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot 0.5 \\&= \frac{1}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \\&= \frac{1}{\sqrt{2\pi}\sigma} \cdot \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right),\end{aligned}$$

# THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when  $\lambda = 0$ ?

$$\begin{aligned}f(x|\mu, \sigma, \lambda = 0) &= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0) \\&= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot 0.5 \\&= \frac{1}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \\&= \frac{1}{\sqrt{2\pi}\sigma} \cdot \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right),\end{aligned}$$

which is the pdf of the normal distribution  $(\mu, \sigma)$ .



# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## DEFINITION (STANDARD SKEW-NORMAL)

The  $SN(0, 1, \lambda)$  distribution is called the standard skew-normal and has pdf

$$f_Z(x|\lambda) = 2 \cdot \phi(x) \cdot \Phi(\lambda x), \quad x \in \mathbb{R}.$$

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## DEFINITION (STANDARD SKEW-NORMAL)

The  $SN(0, 1, \lambda)$  distribution is called the standard skew-normal and has pdf

$$f_Z(x|\lambda) = 2 \cdot \phi(x) \cdot \Phi(\lambda x), \quad x \in \mathbb{R}.$$

Similar to the normal and standard normal,  $Z = \frac{Y - \mu}{\sigma}$  and  $Y = \sigma Z + \mu$ .

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (1)

*If  $Z \sim SN(0, 1, \lambda)$ , then  $(-Z) \sim SN(0, 1, -\lambda)$ .*

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (1)

*If  $Z \sim SN(0, 1, \lambda)$ , then  $(-Z) \sim SN(0, 1, -\lambda)$ .*

*Short proof:*

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (1)

*If  $Z \sim SN(0, 1, \lambda)$ , then  $(-Z) \sim SN(0, 1, -\lambda)$ .*

*Short proof:*

$$f_{(-Z)}(x) = f_Z(-x)$$

*Q.E.D.*

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (1)

*If  $Z \sim SN(0, 1, \lambda)$ , then  $(-Z) \sim SN(0, 1, -\lambda)$ .*

*Short proof:*

$$\begin{aligned}f_{(-Z)}(x) &= f_Z(-x) \\ &= 2 \cdot \phi(-x) \cdot \Phi(-\lambda x)\end{aligned}$$

*Q.E.D.*

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (1)

*If  $Z \sim SN(0, 1, \lambda)$ , then  $(-Z) \sim SN(0, 1, -\lambda)$ .*

*Short proof:*

$$\begin{aligned}f_{(-Z)}(x) &= f_Z(-x) \\&= 2 \cdot \phi(-x) \cdot \Phi(-\lambda x) \\&= 2 \cdot \phi(x) \cdot \Phi(-\lambda x),\end{aligned}$$

*Q.E.D.*

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (1)

If  $Z \sim SN(0, 1, \lambda)$ , then  $(-Z) \sim SN(0, 1, -\lambda)$ .

*Short proof:*

$$\begin{aligned}f_{(-Z)}(x) &= f_Z(-x) \\&= 2 \cdot \phi(-x) \cdot \Phi(-\lambda x) \\&= 2 \cdot \phi(x) \cdot \Phi(-\lambda x),\end{aligned}$$


which is the pdf of  $SN(0, 1, -\lambda)$ .

*Q.E.D.*



# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

*Property 1:*  $-SN(0, 1, \lambda) \sim SN(0, 1, -\lambda)$



# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (2)

*If  $Z \sim SN(0, 1, \lambda)$ , then  $Z^2 \sim \chi_1^2$  (chi-square with 1 degree of freedom).*

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (2)

*If  $Z \sim SN(0, 1, \lambda)$ , then  $Z^2 \sim \chi_1^2$  (chi-square with 1 degree of freedom).*

*Short proof:*

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (2)

*If  $Z \sim SN(0, 1, \lambda)$ , then  $Z^2 \sim \chi_1^2$  (chi-square with 1 degree of freedom).*

*Short proof:*

Lemma 1 comes with a handy result, (? , page 161):

If  $Y \sim f_0$  and  $Z \sim f$ , then  $|Y| \stackrel{d}{=} |Z|$ , where the notation  $\stackrel{d}{=}$  denotes equality in distribution.

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (2)

*If  $Z \sim SN(0, 1, \lambda)$ , then  $Z^2 \sim \chi_1^2$  (chi-square with 1 degree of freedom).*

*Short proof:*

Lemma 1 comes with a handy result, (? , page 161):

If  $Y \sim f_0$  and  $Z \sim f$ , then  $|Y| \stackrel{d}{=} |Z|$ , where the notation  $\stackrel{d}{=}$  denotes equality in distribution.

Let  $X \sim N(0, 1)$ . Since  $X^2 \sim \chi_1^2$  and  $|X| \stackrel{d}{=} |Z|$ , then  $Z^2 \sim \chi_1^2$ .  
*Q.E.D.*

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (3)

As  $\lambda \rightarrow \pm\infty$ ,  $SN(0, 1, \lambda)$  tends to the half normal distribution,  $\pm|N(0, 1)|$ .

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (3)


As  $\lambda \rightarrow \pm\infty$ ,  $SN(0, 1, \lambda)$  tends to the half normal distribution,  $\pm|N(0, 1)|$ .

Let  $X \sim |N(0, 1)|$ . Then

$$f_X(x) = \begin{cases} 0 & \text{when } -\infty < x \leq 0 \\ 2\phi & \text{when } 0 < x < \infty \end{cases}.$$

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL


*Property 3:*  $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$  as  $\lambda \rightarrow \infty$ :






# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

*Property 3:*  $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$  as  $\lambda \rightarrow \infty$ :




# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

*Property 3:*  $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$  as  $\lambda \rightarrow \infty$ :




# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

*Property 3:*  $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$  as  $\lambda \rightarrow \infty$ :




# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

*Property 3:*  $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$  as  $\lambda \rightarrow \infty$ :




# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

*Property 3:*  $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$  as  $\lambda \rightarrow -\infty$ :




# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

*Property 3:*  $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$  as  $\lambda \rightarrow -\infty$ :




# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

*Property 3:*  $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$  as  $\lambda \rightarrow -\infty$ :



# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL


*Property 3:*  $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$  as  $\lambda \rightarrow -\infty$ :





# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

*Property 3:*  $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$  as  $\lambda \rightarrow -\infty$ :



# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (4)

*The moment generating function of  $SN(0, 1, \lambda)$  is*

$$M(t|\lambda) = 2 \cdot \Phi(\delta t) \cdot e^{t^2/2},$$

*where  $\delta = \frac{\lambda}{\sqrt{1 + \lambda^2}}$  and  $t \in (-\infty, \infty)$ .*

# THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

## PROPERTY (4)

*The moment generating function of  $SN(0, 1, \lambda)$  is*

$$M(t|\lambda) = 2 \cdot \Phi(\delta t) \cdot e^{t^2/2},$$

*where  $\delta = \frac{\lambda}{\sqrt{1 + \lambda^2}}$  and  $t \in (-\infty, \infty)$ .*

According to Equation 5 in ?, the mgf of  $SN(\mu, \sigma, \lambda)$  is

$$M(t) = E\{e^{tY}\} = 2 \cdot \exp\left(\mu t + \frac{\sigma^2 t^2}{2}\right) \cdot \Phi(\delta \sigma t).$$

Our result follows.

# METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

# METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

# METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

1. Find the first three moments about the mean (central moments) of the binomial and the skew-normal.

# METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

1. Find the first three moments about the mean (central moments) of the binomial and the skew-normal.

What are central moments?:  $E(X)$ ,  $E([X - E(X)]^2)$ ,  $E([X - E(X)]^3)$ .

# METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

1. Find the first three moments about the mean (central moments) of the binomial and the skew-normal.

What are central moments?:  $E(X)$ ,  $E([X - E(X)]^2)$ ,  $E([X - E(X)]^3)$ .

2. Set them equal to each other.



# METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

1. Find the first three moments about the mean (central moments) of the binomial and the skew-normal.

What are central moments?:  $E(X)$ ,  $E([X - E(X)]^2)$ ,  $E([X - E(X)]^3)$ .

2. Set them equal to each other.
3. Take  $n$  and  $p$  to be constants; solve for  $\mu$ ,  $\sigma$ , and  $\lambda$ .

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Let's start with the binomial ...

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

The first two central moments are just the mean and variance:

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

The first two central moments are just the mean and variance:

$$E(B) = np$$

$$E([B - E(B)]^2) = \text{Var}(B) = np(1 - p)$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

The third one takes some elbow grease.

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

The third one takes some elbow grease.

First we'll need to find  $E(B^2)$  and  $E(B^3)$ .

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Recall that  $\text{Var}(B) = E(B^2) - [E(B)]^2$ .

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Recall that  $\text{Var}(B) = E(B^2) - [E(B)]^2$ . Thus

$$E(B^2) = \text{Var}(B) + [E(B)]^2$$



# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Recall that  $\text{Var}(B) = E(B^2) - [E(B)]^2$ . Thus

$$\begin{aligned} E(B^2) &= \text{Var}(B) + [E(B)]^2 \\ &= np(1-p) + n^2p^2 \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Recall that  $\text{Var}(B) = E(B^2) - [E(B)]^2$ . Thus

$$\begin{aligned} E(B^2) &= \text{Var}(B) + [E(B)]^2 \\ &= np(1-p) + n^2p^2 \\ &= np - np^2 + n^2p^2. \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

We will get  $E(B^3)$  via the third factorial moment,  $E[B(B-1)(B-2)]$ .

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$E[B(B-1)(B-2)]$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} &E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \frac{n!}{x! (n-x)!} p^x q^{n-x} \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \frac{n!}{x! (n-x)!} p^x q^{n-x} \\ &= \sum_{x=3}^n \frac{n!}{(x-3)! (n-x)!} p^x q^{n-x} \end{aligned}$$



# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \frac{n!}{x! (n-x)!} p^x q^{n-x} \\ &= \sum_{x=3}^n \frac{n!}{(x-3)! (n-x)!} p^x q^{n-x} \\ &= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)! (n-x)!} p^{x-3} q^{n-x} \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let  $y = x - 3$ ; then  $x = y + 3$ , and  $x = 3, x = n \Rightarrow y = 0, y = n - 3$ :

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let  $y = x - 3$ ; then  $x = y + 3$ , and  $x = 3, x = n \Rightarrow y = 0, y = n - 3$ :

$$= n(n-1)(n-2)p^3 \cdot \sum_{y=0}^{n-3} \frac{(n-3)!}{y!(n-(y+3))!} p^y q^{n-(y+3)}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let  $y = x - 3$ ; then  $x = y + 3$ , and  $x = 3, x = n \Rightarrow y = 0, y = n - 3$ :

$$= n(n-1)(n-2)p^3 \cdot \sum_{y=0}^{n-3} \frac{(n-3)!}{y!(n-(y+3))!} p^y q^{n-(y+3)}$$

$$= n(n-1)(n-2)p^3 \cdot \underbrace{\sum_{y=0}^{n-3} \frac{(n-3)!}{y!((n-3)-y)!} p^y q^{(n-3)-y}}_{\text{[pdf of } \textit{Bin}(n-3, p) \text{ summed over its domain]} = 1}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let  $y = x - 3$ ; then  $x = y + 3$ , and  $x = 3, x = n \Rightarrow y = 0, y = n - 3$ :

$$= n(n-1)(n-2)p^3 \cdot \sum_{y=0}^{n-3} \frac{(n-3)!}{y!(n-(y+3))!} p^y q^{n-(y+3)}$$

$$= n(n-1)(n-2)p^3 \cdot \underbrace{\sum_{y=0}^{n-3} \frac{(n-3)!}{y!((n-3)-y)!} p^y q^{(n-3)-y}}_{\text{[pdf of } \textit{Bin}(n-3, p) \text{ summed over its domain]} = 1}$$

$$= n(n-1)(n-2)p^3$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let  $y = x - 3$ ; then  $x = y + 3$ , and  $x = 3, x = n \Rightarrow y = 0, y = n - 3$ :

$$= n(n-1)(n-2)p^3 \cdot \sum_{y=0}^{n-3} \frac{(n-3)!}{y!(n-(y+3))!} p^y q^{n-(y+3)}$$

$$= n(n-1)(n-2)p^3 \cdot \underbrace{\sum_{y=0}^{n-3} \frac{(n-3)!}{y!((n-3)-y)!} p^y q^{(n-3)-y}}_{\text{[pdf of } \textit{Bin}(n-3, p) \text{ summed over its domain]} = 1}$$

$$= n(n-1)(n-2)p^3$$

$$= n^3 p^3 - 3n^2 p^3 + 2np^3$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get  $E(B^3)$ , we expand the left side of the previous equation:



# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get  $E(B^3)$ , we expand the left side of the previous equation:

$$E[B(B-1)(B-2)]$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get  $E(B^3)$ , we expand the left side of the previous equation:

$$\begin{aligned} E[B(B-1)(B-2)] \\ = E[B^3 - 3B^2 + 2B] \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get  $E(B^3)$ , we expand the left side of the previous equation:

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= E[B^3 - 3B^2 + 2B] \\ &= E(B^3) - 3E(B^2) + 2E(B) \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get  $E(B^3)$ , we expand the left side of the previous equation:

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= E[B^3 - 3B^2 + 2B] \\ &= E(B^3) - 3E(B^2) + 2E(B) \\ &= E(B^3) - 3(np - np^2 + n^2p^2) + 2np \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get  $E(B^3)$ , we expand the left side of the previous equation:

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= E[B^3 - 3B^2 + 2B] \\ &= E(B^3) - 3E(B^2) + 2E(B) \\ &= E(B^3) - 3(np - np^2 + n^2p^2) + 2np \\ &= E(B^3) - 3np + 3np^2 - 3n^2p^2 + 2np \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get  $E(B^3)$ , we expand the left side of the previous equation:

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= E[B^3 - 3B^2 + 2B] \\ &= E(B^3) - 3E(B^2) + 2E(B) \\ &= E(B^3) - 3(np - np^2 + n^2p^2) + 2np \\ &= E(B^3) - 3np + 3np^2 - 3n^2p^2 + 2np \\ &= E(B^3) + 3np^2 - 3n^2p^2 - np \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Left side:  $E(B^3) + 3np^2 - 3n^2p^2 - np$

Right side:  $n^3p^3 - 3n^2p^3 + 2np^3$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Left side:  $E(B^3) + 3np^2 - 3n^2p^2 - np$

Right side:  $n^3p^3 - 3n^2p^3 + 2np^3$

Set them equal and solve for  $E(B^3)$ :



# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Left side:  $E(B^3) + 3np^2 - 3n^2p^2 - np$

Right side:  $n^3p^3 - 3n^2p^3 + 2np^3$

Set them equal and solve for  $E(B^3)$ :

$$E(B^3) + 3np^2 - 3n^2p^2 - np = n^3p^3 - 3n^2p^3 + 2np^3$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Left side:  $E(B^3) + 3np^2 - 3n^2p^2 - np$

Right side:  $n^3p^3 - 3n^2p^3 + 2np^3$

Set them equal and solve for  $E(B^3)$ :

$$E(B^3) + 3np^2 - 3n^2p^2 - np = n^3p^3 - 3n^2p^3 + 2np^3$$

$$\Rightarrow E(B^3) = n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$E\left([B - E(B)]^3\right)$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E \left( [B - E(B)]^3 \right) \\ &= E \left( B^3 - 3B^2 E(B) + 3B[E(B)]^2 - [E(B)]^3 \right) \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E \left( [B - E(B)]^3 \right) \\ &= E \left( B^3 - 3B^2 E(B) + 3B[E(B)]^2 - [E(B)]^3 \right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \\ &= (n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np) \\ &\quad - 3(np - np^2 + n^2p^2)(np) + 2(np)^3 \end{aligned}$$



# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \\ &= (n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np) \\ &\quad - 3(np - np^2 + n^2p^2)(np) + 2(np)^3 \\ &= \cancel{n^3p^3} - \cancel{3n^2p^3} + 2np^3 - 3np^2 + \cancel{3n^2p^2} + np \\ &\quad - \cancel{3n^2p^2} + \cancel{3n^2p^3} - \cancel{3n^3p^3} + \cancel{2n^3p^3} \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \\ &= (n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np) \\ &\quad - 3(np - np^2 + n^2p^2)(np) + 2(np)^3 \\ &= \cancel{n^3p^3} - \cancel{3n^2p^3} + 2np^3 - 3np^2 + \cancel{3n^2p^2} + np \\ &\quad - \cancel{3n^2p^2} + \cancel{3n^2p^3} - \cancel{3n^3p^3} + \cancel{2n^3p^3} \\ &= 2np^3 - 3np^2 + np \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \\ &= (n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np) \\ &\quad - 3(np - np^2 + n^2p^2)(np) + 2(np)^3 \\ &= \cancel{n^3p^3} - \cancel{3n^2p^3} + 2np^3 - 3np^2 + \cancel{3n^2p^2} + np \\ &\quad - \cancel{3n^2p^2} + \cancel{3n^2p^3} - \cancel{3n^3p^3} + \cancel{2n^3p^3} \\ &= 2np^3 - 3np^2 + np \\ &= np(p-1)(2p-1) \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Let's restate our results:

$$E(B) = np,$$

$$E([B - E(B)]^2) = np(1 - p),$$

$$E([B - E(B)]^3) = np(p - 1)(2p - 1)$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Now lets move on to skew-normal ...

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the first and second central moments are the mean and variance.

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the first and second central moments are the mean and variance.

$$E(Y) = \mu + b\delta\sigma$$
$$\text{Var}(Y) = \sigma^2(1 - b^2\delta^2)$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:



# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$E([Y - E(Y)]^3)$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned} E([Y - E(Y)]^3) \\ = E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned} & E([Y - E(Y)]^3) \\ &= E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \\ &= (\mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3) \\ &\quad - 3(\mu^2 + 2b\delta\mu\sigma + \sigma^2)(\mu + b\delta\sigma) + 2(\mu + b\delta\sigma)^3 \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned} & E([Y - E(Y)]^3) \\ &= E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \\ &= (\mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3) \\ &\quad - 3(\mu^2 + 2b\delta\mu\sigma + \sigma^2)(\mu + b\delta\sigma) + 2(\mu + b\delta\sigma)^3 \\ &= \cancel{\mu^3} + \cancel{3b\delta\mu^2\sigma} + \cancel{3\mu\sigma^2} + \cancel{3b\delta\sigma^3} - b\delta^3\sigma^3 - \cancel{3\mu^3} - \cancel{3b\delta\mu^2\sigma} \\ &\quad - \cancel{6b\delta\mu^2\sigma} - \cancel{6b^2\delta^2\mu\sigma^2} - \cancel{3\mu\sigma^2} - \cancel{3b\delta\sigma^3} + 2\mu^3 + 6b\delta\mu^2\sigma \\ &\quad + \cancel{6b^2\delta^2\mu\sigma^2} + 2b^3\delta^3\sigma^3 \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned} & E([Y - E(Y)]^3) \\ &= E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \\ &= (\mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3) \\ &\quad - 3(\mu^2 + 2b\delta\mu\sigma + \sigma^2)(\mu + b\delta\sigma) + 2(\mu + b\delta\sigma)^3 \\ &= \cancel{\mu^3} + \cancel{3b\delta\mu^2\sigma} + \cancel{3\mu\sigma^2} + \cancel{3b\delta\sigma^3} - b\delta^3\sigma^3 - \cancel{3\mu^3} - \cancel{3b\delta\mu^2\sigma} \\ &\quad - \cancel{6b\delta\mu^2\sigma} - \cancel{6b^2\delta^2\mu\sigma^2} - \cancel{3\mu\sigma^2} - \cancel{3b\delta\sigma^3} + \cancel{2\mu^3} + \cancel{6b\delta\mu^2\sigma} \\ &\quad + \cancel{6b^2\delta^2\mu\sigma^2} + 2b^3\delta^3\sigma^3 \\ &= 2b^3\delta^3\sigma^3 - b\delta^3\sigma^3 \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned} & E([Y - E(Y)]^3) \\ &= E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \\ &= (\mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3) \\ &\quad - 3(\mu^2 + 2b\delta\mu\sigma + \sigma^2)(\mu + b\delta\sigma) + 2(\mu + b\delta\sigma)^3 \\ &= \cancel{\mu^3} + \cancel{3b\delta\mu^2\sigma} + \cancel{3\mu\sigma^2} + \cancel{3b\delta\sigma^3} - b\delta^3\sigma^3 - \cancel{3\mu^3} - \cancel{3b\delta\mu^2\sigma} \\ &\quad - \cancel{6b\delta\mu^2\sigma} - \cancel{6b^2\delta^2\mu\sigma^2} - \cancel{3\mu\sigma^2} - \cancel{3b\delta\sigma^3} + 2\mu^3 + 6b\delta\mu^2\sigma \\ &\quad + \cancel{6b^2\delta^2\mu\sigma^2} + 2b^3\delta^3\sigma^3 \\ &= 2b^3\delta^3\sigma^3 - b\delta^3\sigma^3 \\ &= b\delta^3\sigma^3(2b^2 - 1) \end{aligned}$$

# METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Our results, restated:

$$E(Y) = \mu + b\delta\sigma = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}}$$

$$E([Y - E(Y)]^2) = \sigma^2(1 - b^2\delta^2) = \sigma^2 \left( 1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2} \right)$$

$$E([Y - E(Y)]^3) = b\delta^3\sigma^3(2b^2 - 1) = \sigma^3 \sqrt{\frac{2}{\pi}} \left( \frac{\lambda}{\sqrt{1 + \lambda^2}} \right)^3 \left( \frac{4}{\pi} - 1 \right)$$

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We're finally ready to derive our approximation!



# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We start by setting the central moments of the binomial equal to the central moments of the skew-normal.

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We start by setting the central moments of the binomial equal to the central moments of the skew-normal.

$$np = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}} \quad (1a)$$

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We start by setting the central moments of the binomial equal to the central moments of the skew-normal.

$$np = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}} \quad (1a)$$

$$np(1 - p) = \sigma^2 \left( 1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2} \right) \quad (1b)$$

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We start by setting the central moments of the binomial equal to the central moments of the skew-normal.

$$np = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}} \quad (1a)$$

$$np(1 - p) = \sigma^2 \left( 1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2} \right) \quad (1b)$$

$$np(p - 1)(2p - 1) = \sigma^3 \sqrt{\frac{2}{\pi}} \left( \frac{\lambda}{\sqrt{1 + \lambda^2}} \right)^3 \left( \frac{4}{\pi} - 1 \right) \quad (1c)$$

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

To get  $\lambda$ , divide the cube of (1b) by the square of (1c):

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

To get  $\lambda$ , divide the cube of (1b) by the square of (1c):

$$\frac{\sigma^6 \left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}\right)^3}{\sigma^6 \cdot \frac{2}{\pi} \left(\frac{\lambda}{\sqrt{1+\lambda^2}}\right)^6 \left(\frac{4}{\pi} - 1\right)^2} = \frac{n^3 p^3 (1-p)^3}{n^2 p^2 (p-1)^2 (2p-1)^2}$$

(2)

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

To get  $\lambda$ , divide the cube of (1b) by the square of (1c):

$$\begin{aligned} \frac{\sigma^6 \left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}\right)^3}{\sigma^6 \cdot \frac{2}{\pi} \left(\frac{\lambda}{\sqrt{1+\lambda^2}}\right)^6 \left(\frac{4}{\pi} - 1\right)^2} &= \frac{n^3 p^3 (1-p)^3}{n^2 p^2 (p-1)^2 (2p-1)^2} \\ \Rightarrow \frac{\left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}\right)^3}{\frac{2}{\pi} \left(\frac{\lambda^2}{1+\lambda^2}\right)^3 \left(\frac{4}{\pi} - 1\right)^2} &= \frac{np(1-p)}{(1-2p)^2} \end{aligned} \quad (2)$$

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Equation (2) can be solved for  $\lambda^2$ .



# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Equation (2) can be solved for  $\lambda^2$ .

Then take  $\lambda$  to be

$$\lambda = \{\text{sign of } (1 - 2p)\} \sqrt{\lambda^2}.$$

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Equation (2) can be solved for  $\lambda^2$ .

Then take  $\lambda$  to be

$$\lambda = \{\text{sign of } (1 - 2p)\} \sqrt{\lambda^2}.$$

Why?

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION


Equation (2) can be solved for  $\lambda^2$ .

Then take  $\lambda$  to be

$$\lambda = \{\text{sign of } (1 - 2p)\} \sqrt{\lambda^2}.$$


Why? Recall Property 3 ...

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When  $p < 0.5$ :


# METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When  $p < 0.5$ :

- ▶ The binomial skews right (weight shifts left) and approaches  $+|N(0, 1)| \rightarrow \lambda$  is positive


# METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When  $p < 0.5$ :


- ▶ The binomial skews right (weight shifts left) and approaches  $+|N(0, 1)| \rightarrow \lambda$  is positive
- ▶  $(1 - 2p)$  is positive

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When  $p > 0.5$ :

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION




When  $p > 0.5$ :

- ▶ The binomial skews left (weight shifts right) and approaches  $-|N(0, 1)| \rightarrow \lambda$  is negative



# METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When  $p > 0.5$ :

- ▶ The binomial skews left (weight shifts right) and approaches  $-|N(0, 1)| \rightarrow \lambda$  is negative
- ▶  $(1 - 2p)$  is negative

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Once you have  $\lambda$ , solve for  $\sigma$  and then  $\mu$ .

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Once you have  $\lambda$ , solve for  $\sigma$  and then  $\mu$ .

$$np(1 - p) = \sigma^2 \left( 1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2} \right) \Rightarrow \sigma = \sqrt{\frac{np(1 - p)}{1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2}}}$$

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Once you have  $\lambda$ , solve for  $\sigma$  and then  $\mu$ .

$$np(1-p) = \sigma^2 \left( 1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2} \right) \Rightarrow \sigma = \sqrt{\frac{np(1-p)}{1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}}}$$

$$np = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1+\lambda^2}} \Rightarrow \mu = np - \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1+\lambda^2}}$$

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

When  $p = 0.5$ , the binomial is symmetric, so  $\lambda = 0$ .

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

When  $p = 0.5$ , the binomial is symmetric, so  $\lambda = 0$ .

This takes us back to the usual normal approximation:

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

When  $p = 0.5$ , the binomial is symmetric, so  $\lambda = 0$ .

This takes us back to the usual normal approximation:

$$\sigma = \sqrt{\frac{np(1-p)}{1 - \frac{2}{\pi} \cdot \frac{0^2}{1+0^2}}} = \sqrt{\frac{np(1-p)}{1}} = \sqrt{np(1-p)}$$

# METHOD OF MOMENTS: DERIVING AN APPROXIMATION

When  $p = 0.5$ , the binomial is symmetric, so  $\lambda = 0$ .

This takes us back to the usual normal approximation:

$$\sigma = \sqrt{\frac{np(1-p)}{1 - \frac{2}{\pi} \cdot \frac{0^2}{1+0^2}}} = \sqrt{\frac{np(1-p)}{1}} = \sqrt{np(1-p)}$$
$$\mu = np - \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{0}{\sqrt{1+0^2}} = np - 0 = np$$



# METHOD OF MOMENTS: RESTRICTIONS

Unfortunately, though better than the normal approximation, our skew-normal method isn't universal.

# METHOD OF MOMENTS: RESTRICTIONS

Unfortunately, though better than the normal approximation, our skew-normal method isn't universal.

To be able to solve for  $\lambda$ , we must put a few restrictions on  $n$  and  $p$ .

# METHOD OF MOMENTS: RESTRICTIONS

$$\text{Let } u = \frac{\lambda^2}{1 + \lambda^2} \text{ and } v = 1/u = \frac{1 + \lambda^2}{\lambda^2}.$$

# METHOD OF MOMENTS: RESTRICTIONS

$$\text{Let } u = \frac{\lambda^2}{1 + \lambda^2} \text{ and } v = 1/u = \frac{1 + \lambda^2}{\lambda^2}.$$

Then we can rewrite (2):

$$\frac{\left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2}\right)^3}{\frac{2}{\pi} \left(\frac{\lambda^2}{1 + \lambda^2}\right)^3 \left(\frac{4}{\pi} - 1\right)^2}$$

# METHOD OF MOMENTS: RESTRICTIONS

$$\text{Let } u = \frac{\lambda^2}{1 + \lambda^2} \text{ and } v = 1/u = \frac{1 + \lambda^2}{\lambda^2}.$$

Then we can rewrite (2):

$$\frac{\left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2}\right)^3}{\frac{2}{\pi} \left(\frac{\lambda^2}{1 + \lambda^2}\right)^3 \left(\frac{4}{\pi} - 1\right)^2}$$

$\vdots$

(*magic*)

$\vdots$

# METHOD OF MOMENTS: RESTRICTIONS

$$\text{Let } u = \frac{\lambda^2}{1 + \lambda^2} \text{ and } v = 1/u = \frac{1 + \lambda^2}{\lambda^2}.$$

Then we can rewrite (2):

$$\begin{aligned} & \frac{\left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2}\right)^3}{\frac{2}{\pi} \left(\frac{\lambda^2}{1 + \lambda^2}\right)^3 \left(\frac{4}{\pi} - 1\right)^2} \\ & \quad \vdots \\ & \quad (magic) \\ & \quad \vdots \\ & \left(v - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right) = g(v) \end{aligned}$$

# METHOD OF MOMENTS: RESTRICTIONS

$$g(v) \text{ is increasing in } v = \frac{1 + \lambda^2}{\lambda^2} \geq 1.$$

# METHOD OF MOMENTS: RESTRICTIONS

$g(v)$  is increasing in  $v = \frac{1 + \lambda^2}{\lambda^2} \geq 1$ . Therefore:

$$\min_v g(v)$$



# METHOD OF MOMENTS: RESTRICTIONS

$g(v)$  is increasing in  $v = \frac{1 + \lambda^2}{\lambda^2} \geq 1$ . Therefore:

$$\min_v g(v) = g(1)$$

# METHOD OF MOMENTS: RESTRICTIONS

$g(v)$  is increasing in  $v = \frac{1 + \lambda^2}{\lambda^2} \geq 1$ . Therefore:

$$\min_v g(v) = g(1) = \left(1 - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right)$$

# METHOD OF MOMENTS: RESTRICTIONS

$g(v)$  is increasing in  $v = \frac{1 + \lambda^2}{\lambda^2} \geq 1$ . Therefore:

$$\min_v g(v) = g(1) = \left(1 - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right) = 1.009524 \approx 1$$

# METHOD OF MOMENTS: RESTRICTIONS

To be able to solve (2) for  $\lambda$ , we must have

$$\{\text{right hand side of (2)}\} \geq \{\text{min of left hand side of (2)}\}$$

(3)

# METHOD OF MOMENTS: RESTRICTIONS

To be able to solve (2) for  $\lambda$ , we must have

$$\{\text{right hand side of (2)}\} \geq \{\text{min of left hand side of (2)}\}$$

$$\frac{np(1-p)}{(1-2p)^2} \geq 1$$

(3)

# METHOD OF MOMENTS: RESTRICTIONS

To be able to solve (2) for  $\lambda$ , we must have

$$\{\text{right hand side of (2)}\} \geq \{\text{min of left hand side of (2)}\}$$

$$\frac{np(1-p)}{(1-2p)^2} \geq 1$$

$$np(1-p) \geq \frac{1}{(1-2p)^2}. \quad (3)$$

# METHOD OF MOMENTS: RESTRICTIONS

To be able to solve (2) for  $\lambda$ , we must have

$$\{\text{right hand side of (2)}\} \geq \{\text{min of left hand side of (2)}\}$$

$$\frac{np(1-p)}{(1-2p)^2} \geq 1$$

$$np(1-p) \geq \frac{1}{(1-2p)^2}. \quad (3)$$

From (3), we can answer two questions:

# METHOD OF MOMENTS: RESTRICTIONS

One:

Given  $p$ , what is the least  $n$  necessary?



# METHOD OF MOMENTS: RESTRICTIONS


One:

Given  $p$ , what is the least  $n$  necessary?

$$n \geq \frac{(1 - 2p)^2}{p(1 - p)}$$

# METHOD OF MOMENTS: RESTRICTIONS

Least possible  $n$ , given a fixed  $p$ :



# METHOD OF MOMENTS: RESTRICTIONS

Two:

Given  $n$ , what is the range of possible  $p$ 's?

# METHOD OF MOMENTS: RESTRICTIONS


Two:

Given  $n$ , what is the range of possible  $p$ 's?

$$\frac{1}{2} - \frac{1}{2}\sqrt{\frac{n}{n+4}} \leq p \leq \frac{1}{2} + \frac{1}{2}\sqrt{\frac{n}{n+4}}$$

# METHOD OF MOMENTS: RESTRICTIONS

Range of possible  $p$ , given a fixed  $n$ :



# DEMONSTRATING IMPROVED ACCURACY

We have an approximation!!

But is it more accurate? ...

# DEMONSTRATING IMPROVED ACCURACY

We have an approximation!!


But is it more accurate? ... Answer: Yes!

# DEMONSTRATING IMPROVED ACCURACY: VISUAL


The easiest way of gauging accuracy is by visual inspection.




# DEMONSTRATING IMPROVED ACCURACY: VISUAL




# DEMONSTRATING IMPROVED ACCURACY: VISUAL




# DEMONSTRATING IMPROVED ACCURACY: VISUAL




# DEMONSTRATING IMPROVED ACCURACY: VISUAL




# DEMONSTRATING IMPROVED ACCURACY: VISUAL




# DEMONSTRATING IMPROVED ACCURACY: VISUAL



# DEMONSTRATING IMPROVED ACCURACY: VISUAL




# DEMONSTRATING IMPROVED ACCURACY: VISUAL





# DEMONSTRATING IMPROVED ACCURACY: VISUAL




# DEMONSTRATING IMPROVED ACCURACY: MABS

A more numerical way of gauging accuracy is the *MABS*, defined by ? as

$$\text{MABS}(n, p) = \max_{k \in \{0, 1, \dots, n\}} \left| F_{B(n, p)}(k) - F_{\text{appr}(n, p)}(k + 0.5) \right|.$$


# DEMONSTRATING IMPROVED ACCURACY: MABS

MABS as a function of  $p$ , with fixed  $n = 25$ :




# DEMONSTRATING IMPROVED ACCURACY: MABS

MABS as a function of  $p$ , with fixed  $n = 100$ :




# DEMONSTRATING IMPROVED ACCURACY: MABS

MABS as a function of  $n$ , with fixed  $p = 0.05$ :



# DEMONSTRATING IMPROVED ACCURACY: MABS

MABS as a function of  $n$ , with fixed  $p = 0.1$ :



# RESOURCES

Estimations of  $SN(\mu, \sigma, \lambda)$  for  $Bin(n, p)$

|     |      | $n$                  |                      |                      |                       |                        |
|-----|------|----------------------|----------------------|----------------------|-----------------------|------------------------|
|     |      | 25                   | 50                   | 100                  | 250                   | 500                    |
| $p$ | 0.05 | (-0.11, 1.74, 4.56)  | (0.79, 2.30, 2.54)   | (2.85, 3.06, 1.86)   | (9.58, 4.52, 1.38)    | (21.32, 6.11, 1.15)    |
|     | 0.10 | (0.89, 2.20, 2.31)   | (2.97, 2.94, 1.74)   | (7.44, 3.94, 1.40)   | (21.53, 5.88, 1.10)   | (45.62, 8.01, 0.94)    |
|     | 0.15 | (2.02, 2.49, 1.79)   | (5.32, 3.34, 1.43)   | (12.25, 4.51, 1.19)  | (33.77, 6.77, 0.96)   | (70.30, 9.27, 0.82)    |
|     | 0.20 | (3.23, 2.67, 1.50)   | (7.76, 3.61, 1.24)   | (17.18, 4.89, 1.04)  | (46.18, 7.39, 0.85)   | (95.18, 10.16, 0.74)   |
|     | 0.25 | (4.49, 2.79, 1.29)   | (10.28, 3.78, 1.09)  | (22.20, 5.15, 0.93)  | (58.71, 7.83, 0.76)   | (120.22, 10.80, 0.67)  |
|     | 0.30 | (5.80, 2.85, 1.12)   | (12.86, 3.88, 0.95)  | (27.31, 5.32, 0.82)  | (71.34, 8.12, 0.68)   | (145.39, 11.24, 0.60)  |
|     | 0.35 | (7.17, 2.86, 0.96)   | (15.50, 3.92, 0.83)  | (32.49, 5.39, 0.72)  | (84.09, 8.28, 0.60)   | (170.70, 11.50, 0.53)  |
|     | 0.40 | (8.59, 2.83, 0.80)   | (18.23, 3.89, 0.70)  | (37.76, 5.39, 0.61)  | (96.96, 8.32, 0.51)   | (196.18, 11.60, 0.45)  |
|     | 0.45 | (10.12, 2.73, 0.61)  | (21.08, 3.79, 0.53)  | (43.21, 5.29, 0.47)  | (110.07, 8.23, 0.40)  | (221.93, 11.54, 0.35)  |
|     | 0.50 | (12.50, 2.50, 0.00)  | (25.00, 3.54, 0.00)  | (50.00, 5.00, 0.00)  | (125.00, 7.91, 0.00)  | (250.00, 11.18, 0.00)  |
|     | 0.55 | (14.88, 2.73, -0.61) | (28.92, 3.79, -0.53) | (56.79, 5.29, -0.47) | (139.93, 8.23, -0.40) | (278.07, 11.54, -0.35) |
|     | 0.60 | (16.41, 2.83, -0.80) | (31.77, 3.89, -0.70) | (62.24, 5.39, -0.61) | (153.04, 8.32, -0.51) | (303.82, 11.60, -0.45) |
|     | 0.65 | (17.83, 2.86, -0.96) | (34.50, 3.92, -0.83) | (67.51, 5.39, -0.72) | (165.91, 8.28, -0.60) | (329.30, 11.50, -0.53) |
|     | 0.70 | (19.20, 2.85, -1.12) | (37.14, 3.88, -0.95) | (72.69, 5.32, -0.82) | (178.66, 8.12, -0.68) | (354.61, 11.24, -0.60) |
|     | 0.75 | (20.51, 2.79, -1.29) | (39.72, 3.78, -1.09) | (77.80, 5.15, -0.93) | (191.29, 7.83, -0.76) | (379.78, 10.80, -0.67) |
|     | 0.80 | (21.77, 2.67, -1.50) | (42.24, 3.61, -1.24) | (82.82, 4.89, -1.04) | (203.82, 7.39, -0.85) | (404.82, 10.16, -0.74) |
|     | 0.85 | (22.98, 2.49, -1.79) | (44.68, 3.34, -1.43) | (87.75, 4.51, -1.19) | (216.23, 6.77, -0.96) | (429.70, 9.27, -0.82)  |
|     | 0.90 | (24.11, 2.20, -2.31) | (47.03, 2.94, -1.74) | (92.56, 3.94, -1.40) | (228.47, 5.88, -1.10) | (454.38, 8.01, -0.94)  |
|     | 0.95 | (25.11, 1.74, -4.56) | (49.21, 2.30, -2.54) | (97.15, 3.06, -1.86) | (240.42, 4.52, -1.38) | (478.68, 6.11, -1.15)  |

# RESOURCES

All values in my project were computed using a Python library, which is freely available online:

<http://github.com/joycetipping/skew-normal-capstone/>



# BIBLIOGRAPHY I