

CALCULATING A SKEW-NORMAL APPROXIMATION

Although easier with a computer program, calculating estimates for μ , σ , and λ by hand is perfectly possible.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (??):

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (??):

$$\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi} \right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2} \right) = \frac{np(1 - p)}{(1 - 2p)^2} \quad (1)$$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (??):

$$\underbrace{\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right)}_{f(\lambda)} = \frac{np(1 - p)}{(1 - 2p)^2} \quad (1)$$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (??):

$$\underbrace{\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right)}_{f(\lambda)} = \underbrace{\frac{np(1 - p)}{(1 - 2p)^2}}_{k_{n,p}} \quad (1)$$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (??):

$$\underbrace{\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right)}_{f(\lambda)} = \underbrace{\frac{np(1 - p)}{(1 - 2p)^2}}_{k_{n,p}} \quad (1)$$

The closed form solution to (1) is pretty hideous, so we'll take a numerical approach.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (??):

$$\underbrace{\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right)}_{f(\lambda)} = \underbrace{\frac{np(1 - p)}{(1 - 2p)^2}}_{k_{n,p}} \quad (1)$$

The closed form solution to (1) is pretty hideous, so we'll take a numerical approach.

Our goal is to find λ such that $f(\lambda)$ is within a certain margin of error (e) of $k_{n,p}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Recall that the sign of λ is determined independently of the value.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Recall that the sign of λ is determined independently of the value.

It is possible to show, by taking its derivative, that f is monotonically decreasing for positive λ .

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Recall that the sign of λ is determined independently of the value.

It is possible to show, by taking its derivative, that f is monotonically decreasing for positive λ .

This convenient fact allows us to find lower and upper bounds for λ and repeatedly bisect our interval until we are within ϵ of $k_{n,p}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

For this demonstration, we will take $n = 25$ and $p = 0.1$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

For this demonstration, we will take $n = 25$ and $p = 0.1$.

Since we're doing this by hand, we'll take our error margin e to be a modest 0.1.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 1: Find $k_{n,p}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 1: Find $k_{n,p}$.

Our value: $k_{n,p} = \frac{25 \cdot 0.1 \cdot 0.9}{(1 - 2 \cdot 0.1)^2} = 3.5156.$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 1: Find $k_{n,p}$.

Our value: $k_{n,p} = \frac{25 \cdot 0.1 \cdot 0.9}{(1 - 2 \cdot 0.1)^2} = 3.5156.$

Step 2: Find a and b such that $f(a) > k_{n,p} > f(b)$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 1: Find $k_{n,p}$.

Our value: $k_{n,p} = \frac{25 \cdot 0.1 \cdot 0.9}{(1 - 2 \cdot 0.1)^2} = 3.5156.$

Step 2: Find a and b such that $f(a) > k_{n,p} > f(b)$.

Our values: $a = 1, b = 3.$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

Calculate $c = \frac{a+b}{2}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

Calculate $c = \frac{a+b}{2}$.

- If $f(c) \leq k_{n,p} - 0.01$, we need a small value of c , so we take our new interval to be (a, c) .

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

Calculate $c = \frac{a + b}{2}$.

- ▶ If $f(c) \leq k_{n,p} - 0.01$, we need a small value of c , so we take our new interval to be (a, c) .
- ▶ If $f(c) \geq k_{n,p} + 0.01$, we need a larger value of c , so we take our new interval to be (c, b) .

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

Calculate $c = \frac{a+b}{2}$.

- ▶ If $f(c) \leq k_{n,p} - 0.01$, we need a small value of c , so we take our new interval to be (a, c) .
- ▶ If $f(c) \geq k_{n,p} + 0.01$, we need a larger value of c , so we take our new interval to be (c, b) .

Repeat this step until $f(c)$ is within e of $k_{n,p}$, or more precisely $k_{n,p} - 0.01 < f(c) < k_{n,p} + 0.01$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
-----------	-----	-----	-----	--------	----------------------------	----------------------------

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False
2	2.00	2.500	2.2500	3.7129	False	True

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False
2	2.00	2.500	2.2500	3.7129	False	True
3	2.25	2.500	2.3750	3.3252	True	False

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False
2	2.00	2.500	2.2500	3.7129	False	True
3	2.25	2.500	2.3750	3.3252	True	False
4	2.25	2.375	2.3125	3.5076	False	False

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False
2	2.00	2.500	2.2500	3.7129	False	True
3	2.25	2.500	2.3750	3.3252	True	False
4	2.25	2.375	2.3125	3.5076	False	False

We take the last value of c : 2.3125.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 5: Find the sign of $(1 - 2p)$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 5: Find the sign of $(1 - 2p)$.

Our $p = 0.1 \Rightarrow (1 - 2 \cdot 0.1) = 0.8 \Rightarrow$ positive.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 5: Find the sign of $(1 - 2p)$.

Our $p = 0.1 \Rightarrow (1 - 2 \cdot 0.1) = 0.8 \Rightarrow$ positive.

Step 6: Final answer: $\{\text{sign of } (1 - 2p)\}\lambda$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 5: Find the sign of $(1 - 2p)$.

Our $p = 0.1 \Rightarrow (1 - 2 \cdot 0.1) = 0.8 \Rightarrow$ positive.

Step 6: Final answer: $\{\text{sign of } (1 - 2p)\}\lambda$

Our final answer: $\lambda = 2.3125$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING σ

Once we have λ , we can easily find σ :

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING σ

Once we have λ , we can easily find σ :

$$\sigma = \sqrt{\frac{np(1-p)}{1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}}} = \sqrt{\frac{25 \cdot 0.1 \cdot 0.9}{1 - \frac{2}{\pi} \cdot \frac{2.3125^2}{1+2.3125^2}}} = 2.2029.$$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING μ

And with λ and σ , we can also find μ :

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING μ

And with λ and σ , we can also find μ :

$$\begin{aligned}\mu &= np - \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}} \\ &= 25 \cdot 0.1 - 2.2029 \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{2.3125}{\sqrt{1 + 2.3125^2}} \\ &= 0.8867.\end{aligned}$$

BIBLIOGRAPHY I

- B. C. Arnold and G. D. Lin. Characterizations of the skew-normal and generalized chi distributions. *Sankhy: The Indian Journal of Statistics*, 66(4):593–606, 2004.
- Adelchi Azzalini. A class of distributions which includes the normal ones. *Scandinavian Journal of Statistics*, 12: 171–178, 1985.
- Adelchi Azzalini. The skew-normal distribution and related multivariate families. *Scandinavian Journal of Statistics*, 32: 159–188, 2005.
- Lee J. Bain and Max Engelhardt. *Introduction to Probability and Mathematical Statistics*. Duxbury, 2nd edition, 1992.
- Ching-Hui Chang, Jyh-Jiuan Lin, Nabendu Pal, and Miao-Chen Chiang. A note on improved approximation of the binomial distribution by the skew-normal distribution. *American Statistical Association*, 62(2):167–170, May 2008.

BIBLIOGRAPHY II

- A. K. Gupta, T. T. Nguyen, and J. A. T. Sanqui. Characterization of the skew-normal distribution. *Annals of the Institute of Statistical Mathematics*, 56:351–360, 2004.
- A. O'Hagan and Tom Leonard. Bayes estimation subject to uncertainty about parameter constraints. *Biometrika*, 63: 201–203, 1976.
- Arthur Pewsey. Problems of inference for azzalini's skew-normal distribution. *Journal of Applied Statistics*, 27(7): 859–870, 2000.
- Martin Schader and Friedrich Schmid. Two rules of thumb for the approximation of the binomial distribution by the normal distribution. *American Statistical Association*, 43(1):23–24, February 1989.