

THE SKEW-NORMAL APPROXIMATION OF THE BINOMIAL DISTRIBUTION

Joyce Tipping

Truman State University

Spring 2011

INTRODUCTION

DEFINITION (BINOMIAL)

Let X represent the number of successes in a sequence of n independent Bernoulli trials with probability of success p on each trial.

INTRODUCTION

DEFINITION (BINOMIAL)

Let X represent the number of successes in a sequence of n independent Bernoulli trials with probability of success p on each trial.

X has discrete pdf

$$f_X(x) = \binom{n}{x} p^x q^{n-x}$$

INTRODUCTION

DEFINITION (BINOMIAL)

Let X represent the number of successes in a sequence of n independent Bernoulli trials with probability of success p on each trial.

X has discrete pdf

$$f_X(x) = \binom{n}{x} p^x q^{n-x}$$

and cdf

$$F_X(x) = P(X \leq x) = \sum_{k=0}^x f_X(k).$$

INTRODUCTION

The binomial cdf is easy to calculate for small n ...

INTRODUCTION

The binomial cdf is easy to calculate for small n ...

But as n gets larger, it becomes increasingly difficult.

INTRODUCTION

The binomial cdf is easy to calculate for small n ...

But as n gets larger, it becomes increasingly difficult.

For example ...

INTRODUCTION

When $n = 3$,

$$F(1) = \binom{3}{1} p^1 q^2 + \binom{3}{0} p^0 q^3$$

INTRODUCTION

When $n = 3$,

$$F(1) = \binom{3}{1} p^1 q^2 + \binom{3}{0} p^0 q^3$$

When $n = 25$,

$$\begin{aligned} F(12) = & \binom{25}{12} p^{12} q^{13} + \binom{25}{11} p^{11} q^{14} + \binom{25}{10} p^{10} q^{15} + \binom{25}{9} p^9 q^{16} \\ & + \binom{25}{8} p^8 q^{17} + \binom{25}{7} p^7 q^{18} + \binom{25}{6} p^6 q^{19} + \binom{25}{5} p^5 q^{20} \\ & + \binom{25}{4} p^4 q^{21} + \binom{25}{3} p^3 q^{22} + \binom{25}{2} p^2 q^{23} + \binom{25}{1} p^1 q^{24} \\ & + \binom{25}{0} p^0 q^{25} \end{aligned}$$

INTRODUCTION

A common technique is to use the normal distribution as an approximation:

$$F_X(x) \approx \Phi\left(\frac{x + 0.5 - \mu}{\sigma}\right),$$

where $\mu = np$, $\sigma = \sqrt{np(1-p)}$, and Φ is the standard normal cdf.

INTRODUCTION

A common technique is to use the normal distribution as an approximation:

$$F_X(x) \approx \Phi\left(\frac{x + 0.5 - \mu}{\sigma}\right),$$

where $\mu = np$, $\sigma = \sqrt{np(1-p)}$, and Φ is the standard normal cdf.

When does this work well? ...

INTRODUCTION

A common technique is to use the normal distribution as an approximation:


$$F_X(x) \approx \Phi\left(\frac{x + 0.5 - \mu}{\sigma}\right),$$

where $\mu = np$, $\sigma = \sqrt{np(1-p)}$, and Φ is the standard normal cdf.

When does this work well? ... In a nutshell, when the binomial is symmetric.


INTRODUCTION

The binomial is symmetric when $p = 0.5$



INTRODUCTION

The binomial is symmetric when $p = 0.5$ or n is very large.




INTRODUCTION

However, when n is medium and p is extreme ...

INTRODUCTION


However, when n is medium and p is extreme ...



the binomial is very skewed ...

INTRODUCTION


However, when n is medium and p is extreme ...



and the normal approximation doesn't work very well.

INTRODUCTION

However, when n is medium and p is extreme ...



Introducing ... the skew-normal distribution.

OUTLINE

Today's itinerary:

1. Skew-Normal distribution – basic properties

OUTLINE

Today's itinerary:

1. Skew-Normal distribution – basic properties
2. Method of Moments – derive an approximation

OUTLINE

Today's itinerary:

1. Skew-Normal distribution – basic properties
2. Method of Moments – derive an approximation
3. Accuracy – examine the accuracy of our approximation

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

DEFINITION (SKEW-NORMAL)

Let Y be a skew-normal distribution, with location parameter $\mu \in \mathbb{R}$, scale parameter $\sigma > 0$, and shape parameter $\lambda \in \mathbb{R}$. Then Y has pdf

$$f(x|\mu, \sigma, \lambda) = \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi\left(\frac{\lambda(x - \mu)}{\sigma}\right), \quad x \in \mathbb{R},$$

where ϕ is the standard normal pdf and Φ is the standard normal cdf.

We write $Y \sim SN(\mu, \sigma, \lambda)$.

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

DEFINITION (SKEW-NORMAL)

Let Y be a skew-normal distribution, with location parameter $\mu \in \mathbb{R}$, scale parameter $\sigma > 0$, and shape parameter $\lambda \in \mathbb{R}$. Then Y has pdf

$$f(x|\mu, \sigma, \lambda) = \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi\left(\frac{\lambda(x - \mu)}{\sigma}\right), \quad x \in \mathbb{R},$$

where ϕ is the standard normal pdf and Φ is the standard normal cdf.

We write $Y \sim SN(\mu, \sigma, \lambda)$.

Note: μ and σ are not intuitively related to the mean and variance.

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

So where did this funny-looking pdf come from? ...

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

So where did this funny-looking pdf come from? ...

LEMMA

If f_0 is a one-dimensional probability density function symmetric about 0, and G is a one-dimensional distribution function such that G' exists and is a density symmetric about 0, then

$$f(z) = 2 \cdot f_0(z) \cdot G\{w(z)\} \quad (-\infty < z < \infty)$$

is a density function for any odd function $w(\cdot)$. (Lemma 1, Azzalini, 2005)

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

So where did this funny-looking pdf come from? ...

LEMMA

If f_0 is a one-dimensional probability density function symmetric about 0, and G is a one-dimensional distribution function such that G' exists and is a density symmetric about 0, then

$$f(z) = 2 \cdot f_0(z) \cdot G\{w(z)\} \quad (-\infty < z < \infty)$$

is a density function for any odd function $w(\cdot)$. (Lemma 1, Azzalini, 2005)

- ▶ A pdf symmetric about 0 (kernel)

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

So where did this funny-looking pdf come from? ...

LEMMA

If f_0 is a one-dimensional probability density function symmetric about 0, and G is a one-dimensional distribution function such that G' exists and is a density symmetric about 0, then

$$f(z) = 2 \cdot f_0(z) \cdot G\{w(z)\} \quad (-\infty < z < \infty)$$

is a density function for any odd function $w(\cdot)$. (Lemma 1, Azzalini, 2005)

- ▶ A pdf symmetric about 0 (kernel)
- ▶ A cdf whose derivative is symmetric about 0

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

Basic properties:

$$E(Y) = \mu + b\delta\sigma$$

$$E(Y^2) = \mu^2 + 2b\delta\mu\sigma + \sigma^2$$

$$E(Y^3) = \mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3$$

$$\text{Var}(Y) = \sigma^2(1 - b^2\delta^2)$$

where $b = \sqrt{\frac{2}{\pi}}$ and $\delta = \frac{\lambda}{\sqrt{1 + \lambda^2}}$. (Pewsey, 2000)

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when $\lambda = 0$?

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when $\lambda = 0$?

$$f(x|\mu, \sigma, \lambda = 0) = \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0)$$

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when $\lambda = 0$?

$$\begin{aligned}f(x|\mu, \sigma, \lambda = 0) &= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0) \\&= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot 0.5\end{aligned}$$

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when $\lambda = 0$?

$$\begin{aligned}f(x|\mu, \sigma, \lambda = 0) &= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0) \\&= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot 0.5 \\&= \frac{1}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right)\end{aligned}$$

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when $\lambda = 0$?

$$\begin{aligned}f(x|\mu, \sigma, \lambda = 0) &= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0) \\&= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot 0.5 \\&= \frac{1}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \\&= \frac{1}{\sqrt{2\pi}\sigma} \cdot \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right),\end{aligned}$$

THE SKEW-NORMAL DISTRIBUTION: FOUNDATIONS

What happens when $\lambda = 0$?

$$\begin{aligned}f(x|\mu, \sigma, \lambda = 0) &= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot \Phi(0) \\&= \frac{2}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \cdot 0.5 \\&= \frac{1}{\sigma} \cdot \phi\left(\frac{x - \mu}{\sigma}\right) \\&= \frac{1}{\sqrt{2\pi}\sigma} \cdot \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right),\end{aligned}$$

which is the pdf of the normal distribution (μ, σ) .

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

DEFINITION (STANDARD SKEW-NORMAL)

The $SN(0, 1, \lambda)$ distribution is called the standard skew-normal and has pdf

$$f_Z(x|\lambda) = 2 \cdot \phi(x) \cdot \Phi(\lambda x), \quad x \in \mathbb{R}.$$

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

DEFINITION (STANDARD SKEW-NORMAL)

The $SN(0, 1, \lambda)$ distribution is called the standard skew-normal and has pdf

$$f_Z(x|\lambda) = 2 \cdot \phi(x) \cdot \Phi(\lambda x), \quad x \in \mathbb{R}.$$

Similar to the normal and standard normal, $Z = \frac{Y - \mu}{\sigma}$ and $Y = \sigma Z + \mu$.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (1)

If $Z \sim SN(0, 1, \lambda)$, then $(-Z) \sim SN(0, 1, -\lambda)$.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (1)

If $Z \sim SN(0, 1, \lambda)$, then $(-Z) \sim SN(0, 1, -\lambda)$.

Short proof:

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (1)

If $Z \sim SN(0, 1, \lambda)$, then $(-Z) \sim SN(0, 1, -\lambda)$.

Short proof:

$$f_{(-Z)}(z) = f_Z(-z)$$

Q.E.D.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (1)

If $Z \sim SN(0, 1, \lambda)$, then $(-Z) \sim SN(0, 1, -\lambda)$.

Short proof:

$$\begin{aligned}f_{(-Z)}(z) &= f_Z(-z) \\ &= 2 \cdot \phi(-z) \cdot \Phi(-\lambda z)\end{aligned}$$

Q.E.D.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (1)

If $Z \sim SN(0, 1, \lambda)$, then $(-Z) \sim SN(0, 1, -\lambda)$.

Short proof:

$$\begin{aligned}f_{(-Z)}(z) &= f_Z(-z) \\&= 2 \cdot \phi(-z) \cdot \Phi(-\lambda z) \\&= 2 \cdot \phi(z) \cdot \Phi(-\lambda z),\end{aligned}$$

Q.E.D.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (1)

If $Z \sim SN(0, 1, \lambda)$, then $(-Z) \sim SN(0, 1, -\lambda)$.

Short proof:


$$\begin{aligned}f_{(-Z)}(z) &= f_Z(-z) \\&= 2 \cdot \phi(-z) \cdot \Phi(-\lambda z) \\&= 2 \cdot \phi(z) \cdot \Phi(-\lambda z),\end{aligned}$$

which is the pdf of $SN(0, 1, -\lambda)$.

Q.E.D.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 1: $-SN(0, 1, \lambda) \sim SN(0, 1, -\lambda)$



THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (2)

If $Z \sim SN(0, 1, \lambda)$, then $Z^2 \sim \chi_1^2$ (chi-square with 1 degree of freedom).

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (2)

If $Z \sim SN(0, 1, \lambda)$, then $Z^2 \sim \chi_1^2$ (chi-square with 1 degree of freedom).

Short proof:

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (2)

If $Z \sim SN(0, 1, \lambda)$, then $Z^2 \sim \chi_1^2$ (chi-square with 1 degree of freedom).

Short proof:

Lemma 1 comes with a handy result, (Azzalini, 2005, page 161):

If $Y \sim f_0$ and $Z \sim f$, then $|Y| \stackrel{d}{=} |Z|$, where the notation $\stackrel{d}{=}$ denotes equality in distribution.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (2)

If $Z \sim SN(0, 1, \lambda)$, then $Z^2 \sim \chi_1^2$ (chi-square with 1 degree of freedom).

Short proof:

Lemma 1 comes with a handy result, (Azzalini, 2005, page 161):

If $Y \sim f_0$ and $Z \sim f$, then $|Y| \stackrel{d}{=} |Z|$, where the notation $\stackrel{d}{=}$ denotes equality in distribution.

Let $X \sim N(0, 1)$. Since $X^2 \sim \chi_1^2$ and $|X| \stackrel{d}{=} |Z|$, then $Z^2 \sim \chi_1^2$.
Q.E.D.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (3)

As $\lambda \rightarrow \pm\infty$, $SN(0, 1, \lambda)$ tends to the half normal distribution, $\pm|N(0, 1)|$.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (3)


As $\lambda \rightarrow \pm\infty$, $SN(0, 1, \lambda)$ tends to the half normal distribution, $\pm|N(0, 1)|$.

Let $X \sim |N(0, 1)|$. Then

$$f_X(x) = \begin{cases} 0 & \text{when } -\infty < x \leq 0 \\ 2\phi & \text{when } 0 < x < \infty \end{cases}.$$


THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$ as $\lambda \rightarrow \infty$:




THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$ as $\lambda \rightarrow \infty$:




THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$ as $\lambda \rightarrow \infty$:




THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$ as $\lambda \rightarrow \infty$:




THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow +|N(0, 1)|$ as $\lambda \rightarrow \infty$:




THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$ as $\lambda \rightarrow -\infty$:




THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$ as $\lambda \rightarrow -\infty$:




THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$ as $\lambda \rightarrow -\infty$:




THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$ as $\lambda \rightarrow -\infty$:



THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

Property 3: $SN(0, 1, \lambda) \rightarrow -|N(0, 1)|$ as $\lambda \rightarrow -\infty$:



THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (4)

The moment generating function of $SN(0, 1, \lambda)$ is

$$M(t|\lambda) = 2 \cdot \Phi(\delta t) \cdot e^{t^2/2},$$

where $\delta = \frac{\lambda}{\sqrt{1 + \lambda^2}}$ and $t \in (-\infty, \infty)$.

THE SKEW-NORMAL DISTRIBUTION: THE STANDARD SKEW-NORMAL

PROPERTY (4)

The moment generating function of $SN(0, 1, \lambda)$ is

$$M(t|\lambda) = 2 \cdot \Phi(\delta t) \cdot e^{t^2/2},$$

where $\delta = \frac{\lambda}{\sqrt{1 + \lambda^2}}$ and $t \in (-\infty, \infty)$.

According to Equation 5 in Azzalini (2005), the mgf of $SN(\mu, \sigma, \lambda)$ is

$$M(t) = E\{e^{tY}\} = 2 \cdot \exp\left(\mu t + \frac{\sigma^2 t^2}{2}\right) \cdot \Phi(\delta \sigma t).$$

Our result follows.

METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

1. Find the first three moments about the mean (central moments) of the binomial and the skew-normal.

METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

1. Find the first three moments about the mean (central moments) of the binomial and the skew-normal.

What are central moments?: $E(X)$, $E([X - E(X)]^2)$, $E([X - E(X)]^3)$.

METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

1. Find the first three moments about the mean (central moments) of the binomial and the skew-normal.

What are central moments?: $E(X)$, $E([X - E(X)]^2)$, $E([X - E(X)]^3)$.

2. Set them equal to each other.

METHOD OF MOMENTS: OVERVIEW

Now we're going to derive the approximation!

Game plan:

1. Find the first three moments about the mean (central moments) of the binomial and the skew-normal.

What are central moments?: $E(X)$, $E([X - E(X)]^2)$, $E([X - E(X)]^3)$.

2. Set them equal to each other.
3. Take n and p to be constants; solve for μ , σ , and λ .

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Let's start with the binomial ...

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Let $B \sim \text{Bin}(n, p)$.

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Let $B \sim \text{Bin}(n, p)$.

The first two central moments of B are just the mean and variance:

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Let $B \sim \text{Bin}(n, p)$.

The first two central moments of B are just the mean and variance:

$$E(B) = np$$

$$E([B - E(B)]^2) = \text{Var}(B) = np(1 - p)$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

The third one takes some elbow grease.

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

The third one takes some elbow grease.

First we'll need to find $E(B^2)$ and $E(B^3)$.

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Recall that $\text{Var}(B) = E(B^2) - [E(B)]^2$.

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Recall that $\text{Var}(B) = E(B^2) - [E(B)]^2$. Thus

$$E(B^2) = \text{Var}(B) + [E(B)]^2$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Recall that $\text{Var}(B) = E(B^2) - [E(B)]^2$. Thus

$$\begin{aligned} E(B^2) &= \text{Var}(B) + [E(B)]^2 \\ &= np(1-p) + n^2p^2 \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Recall that $\text{Var}(B) = E(B^2) - [E(B)]^2$. Thus

$$\begin{aligned} E(B^2) &= \text{Var}(B) + [E(B)]^2 \\ &= np(1-p) + n^2p^2 \\ &= np - np^2 + n^2p^2. \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

We will get $E(B^3)$ via the third factorial moment, $E[B(B-1)(B-2)]$.

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$E[B(B-1)(B-2)]$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} &E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \frac{n!}{x! (n-x)!} p^x q^{n-x} \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \frac{n!}{x! (n-x)!} p^x q^{n-x} \\ &= \sum_{x=3}^n \frac{n!}{(x-3)! (n-x)!} p^x q^{n-x} \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= \sum_{x=0}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \left\{ \binom{n}{x} p^x q^{n-x} \right\} \\ &= \sum_{x=3}^n x(x-1)(x-2) \cdot \frac{n!}{x! (n-x)!} p^x q^{n-x} \\ &= \sum_{x=3}^n \frac{n!}{(x-3)! (n-x)!} p^x q^{n-x} \\ &= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)! (n-x)!} p^{x-3} q^{n-x} \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let $y = x - 3$; then $x = y + 3$, and $x = 3, x = n \Rightarrow y = 0, y = n - 3$:

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let $y = x - 3$; then $x = y + 3$, and $x = 3, x = n \Rightarrow y = 0, y = n - 3$:

$$= n(n-1)(n-2)p^3 \cdot \sum_{y=0}^{n-3} \frac{(n-3)!}{y!(n-(y+3))!} p^y q^{n-(y+3)}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let $y = x - 3$; then $x = y + 3$, and $x = 3, x = n \Rightarrow y = 0, y = n - 3$:

$$= n(n-1)(n-2)p^3 \cdot \sum_{y=0}^{n-3} \frac{(n-3)!}{y!(n-(y+3))!} p^y q^{n-(y+3)}$$

$$= n(n-1)(n-2)p^3 \cdot \underbrace{\sum_{y=0}^{n-3} \frac{(n-3)!}{y!((n-3)-y)!} p^y q^{(n-3)-y}}_{\text{[pdf of } \textit{Bin}(n-3, p) \text{ summed over its domain]} = 1}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let $y = x - 3$; then $x = y + 3$, and $x = 3, x = n \Rightarrow y = 0, y = n - 3$:

$$= n(n-1)(n-2)p^3 \cdot \sum_{y=0}^{n-3} \frac{(n-3)!}{y!(n-(y+3))!} p^y q^{n-(y+3)}$$

$$= n(n-1)(n-2)p^3 \cdot \underbrace{\sum_{y=0}^{n-3} \frac{(n-3)!}{y!((n-3)-y)!} p^y q^{(n-3)-y}}_{\text{[pdf of } \textit{Bin}(n-3, p) \text{ summed over its domain]} = 1}$$

$$= n(n-1)(n-2)p^3$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

$$= \sum_{x=3}^n n(n-1)(n-2)p^3 \cdot \frac{(n-3)!}{(x-3)!(n-x)!} p^{x-3} q^{n-x}$$

Let $y = x - 3$; then $x = y + 3$, and $x = 3, x = n \Rightarrow y = 0, y = n - 3$:

$$= n(n-1)(n-2)p^3 \cdot \sum_{y=0}^{n-3} \frac{(n-3)!}{y!(n-(y+3))!} p^y q^{n-(y+3)}$$

$$= n(n-1)(n-2)p^3 \cdot \underbrace{\sum_{y=0}^{n-3} \frac{(n-3)!}{y!((n-3)-y)!} p^y q^{(n-3)-y}}_{\text{[pdf of } \textit{Bin}(n-3, p) \text{ summed over its domain]} = 1}$$

$$= n(n-1)(n-2)p^3$$

$$= n^3 p^3 - 3n^2 p^3 + 2np^3$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get $E(B^3)$, we expand the left side of the previous equation:

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get $E(B^3)$, we expand the left side of the previous equation:

$$E[B(B-1)(B-2)]$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get $E(B^3)$, we expand the left side of the previous equation:

$$\begin{aligned} E[B(B-1)(B-2)] \\ = E[B^3 - 3B^2 + 2B] \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get $E(B^3)$, we expand the left side of the previous equation:

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= E[B^3 - 3B^2 + 2B] \\ &= E(B^3) - 3E(B^2) + 2E(B) \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get $E(B^3)$, we expand the left side of the previous equation:

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= E[B^3 - 3B^2 + 2B] \\ &= E(B^3) - 3E(B^2) + 2E(B) \\ &= E(B^3) - 3(np - np^2 + n^2p^2) + 2np \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get $E(B^3)$, we expand the left side of the previous equation:

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= E[B^3 - 3B^2 + 2B] \\ &= E(B^3) - 3E(B^2) + 2E(B) \\ &= E(B^3) - 3(np - np^2 + n^2p^2) + 2np \\ &= E(B^3) - 3np + 3np^2 - 3n^2p^2 + 2np \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

To get $E(B^3)$, we expand the left side of the previous equation:

$$\begin{aligned} & E[B(B-1)(B-2)] \\ &= E[B^3 - 3B^2 + 2B] \\ &= E(B^3) - 3E(B^2) + 2E(B) \\ &= E(B^3) - 3(np - np^2 + n^2p^2) + 2np \\ &= E(B^3) - 3np + 3np^2 - 3n^2p^2 + 2np \\ &= E(B^3) + 3np^2 - 3n^2p^2 - np \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Left side: $E(B^3) + 3np^2 - 3n^2p^2 - np$

Right side: $n^3p^3 - 3n^2p^3 + 2np^3$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Left side: $E(B^3) + 3np^2 - 3n^2p^2 - np$

Right side: $n^3p^3 - 3n^2p^3 + 2np^3$

Set them equal and solve for $E(B^3)$:

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Left side: $E(B^3) + 3np^2 - 3n^2p^2 - np$

Right side: $n^3p^3 - 3n^2p^3 + 2np^3$

Set them equal and solve for $E(B^3)$:

$$E(B^3) + 3np^2 - 3n^2p^2 - np = n^3p^3 - 3n^2p^3 + 2np^3$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Left side: $E(B^3) + 3np^2 - 3n^2p^2 - np$

Right side: $n^3p^3 - 3n^2p^3 + 2np^3$

Set them equal and solve for $E(B^3)$:

$$E(B^3) + 3np^2 - 3n^2p^2 - np = n^3p^3 - 3n^2p^3 + 2np^3$$

$$\Rightarrow E(B^3) = n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$E\left([B - E(B)]^3\right)$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E \left([B - E(B)]^3 \right) \\ &= E \left(B^3 - 3B^2 E(B) + 3B[E(B)]^2 - [E(B)]^3 \right) \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \\ &= (n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np) \\ &\quad - 3(np - np^2 + n^2p^2)(np) + 2(np)^3 \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \\ &= (n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np) \\ &\quad - 3(np - np^2 + n^2p^2)(np) + 2(np)^3 \\ &= \cancel{n^3p^3} - \cancel{3n^2p^3} + 2np^3 - 3np^2 + \cancel{3n^2p^2} + np \\ &\quad - \cancel{3n^2p^2} + \cancel{3n^2p^3} - \cancel{3n^3p^3} + \cancel{2n^3p^3} \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \\ &= (n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np) \\ &\quad - 3(np - np^2 + n^2p^2)(np) + 2(np)^3 \\ &= \cancel{n^3p^3} - \cancel{3n^2p^3} + 2np^3 - 3np^2 + \cancel{3n^2p^2} + np \\ &\quad - \cancel{3n^2p^2} + \cancel{3n^2p^3} - \cancel{3n^3p^3} + \cancel{2n^3p^3} \\ &= 2np^3 - 3np^2 + np \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Now we can (finally!) compute the third central moment:

$$\begin{aligned} & E\left([B - E(B)]^3\right) \\ &= E\left(B^3 - 3B^2E(B) + 3B[E(B)]^2 - [E(B)]^3\right) \\ &= E(B^3) - 3E(B^2)E(B) + 3E(B)[E(B)]^2 - [E(B)]^3 \\ &= E(B^3) - 3E(B^2)E(B) + 2[E(B)]^3 \\ &= (n^3p^3 - 3n^2p^3 + 2np^3 - 3np^2 + 3n^2p^2 + np) \\ &\quad - 3(np - np^2 + n^2p^2)(np) + 2(np)^3 \\ &= \cancel{n^3p^3} - \cancel{3n^2p^3} + 2np^3 - 3np^2 + \cancel{3n^2p^2} + np \\ &\quad - \cancel{3n^2p^2} + \cancel{3n^2p^3} - \cancel{3n^3p^3} + \cancel{2n^3p^3} \\ &= 2np^3 - 3np^2 + np \\ &= np(p-1)(2p-1) \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE BINOMIAL

Let's restate our results:

$$E(B) = np,$$

$$E([B - E(B)]^2) = np(1 - p),$$

$$E([B - E(B)]^3) = np(p - 1)(2p - 1)$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Now lets move on to skew-normal ...

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Let $Y \sim SN(\mu, \sigma, \lambda)$.

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Let $Y \sim SN(\mu, \sigma, \lambda)$.

Again, the first and second central moments of Y are the mean and variance.

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Let $Y \sim SN(\mu, \sigma, \lambda)$.

Again, the first and second central moments of Y are the mean and variance.

$$\begin{aligned}E(Y) &= \mu + b\delta\sigma \\ \text{Var}(Y) &= \sigma^2(1 - b^2\delta^2)\end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$E([Y - E(Y)]^3)$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned} E([Y - E(Y)]^3) \\ = E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned} & E([Y - E(Y)]^3) \\ &= E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \\ &= (\mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3) \\ &\quad - 3(\mu^2 + 2b\delta\mu\sigma + \sigma^2)(\mu + b\delta\sigma) + 2(\mu + b\delta\sigma)^3 \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned} & E([Y - E(Y)]^3) \\ &= E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \\ &= (\mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3) \\ &\quad - 3(\mu^2 + 2b\delta\mu\sigma + \sigma^2)(\mu + b\delta\sigma) + 2(\mu + b\delta\sigma)^3 \\ &= \cancel{\mu^3} + \cancel{3b\delta\mu^2\sigma} + \cancel{3\mu\sigma^2} + \cancel{3b\delta\sigma^3} - b\delta^3\sigma^3 - \cancel{3\mu^3} - \cancel{3b\delta\mu^2\sigma} \\ &\quad - \cancel{6b\delta\mu^2\sigma} - \cancel{6b^2\delta^2\mu\sigma^2} - \cancel{3\mu\sigma^2} - \cancel{3b\delta\sigma^3} + 2\mu^3 + 6b\delta\mu^2\sigma \\ &\quad + \cancel{6b^2\delta^2\mu\sigma^2} + 2b^3\delta^3\sigma^3 \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned} & E([Y - E(Y)]^3) \\ &= E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \\ &= (\mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3) \\ &\quad - 3(\mu^2 + 2b\delta\mu\sigma + \sigma^2)(\mu + b\delta\sigma) + 2(\mu + b\delta\sigma)^3 \\ &= \cancel{\mu^3} + \cancel{3b\delta\mu^2\sigma} + \cancel{3\mu\sigma^2} + \cancel{3b\delta\sigma^3} - b\delta^3\sigma^3 - \cancel{3\mu^3} - \cancel{3b\delta\mu^2\sigma} \\ &\quad - \cancel{6b\delta\mu^2\sigma} - \cancel{6b^2\delta^2\mu\sigma^2} - \cancel{3\mu\sigma^2} - \cancel{3b\delta\sigma^3} + \cancel{2\mu^3} + \cancel{6b\delta\mu^2\sigma} \\ &\quad + \cancel{6b^2\delta^2\mu\sigma^2} + 2b^3\delta^3\sigma^3 \\ &= 2b^3\delta^3\sigma^3 - b\delta^3\sigma^3 \end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Again, the third one is a little harder:

$$\begin{aligned}E([Y - E(Y)]^3) &= E(Y^3) - 3E(Y^2)E(Y) + 2[E(Y)]^3 \\&= (\mu^3 + 3b\delta\mu^2\sigma + 3\mu\sigma^2 + 3b\delta\sigma^3 - b\delta^3\sigma^3) \\&\quad - 3(\mu^2 + 2b\delta\mu\sigma + \sigma^2)(\mu + b\delta\sigma) + 2(\mu + b\delta\sigma)^3 \\&= \cancel{\mu^3} + \cancel{3b\delta\mu^2\sigma} + \cancel{3\mu\sigma^2} + \cancel{3b\delta\sigma^3} - b\delta^3\sigma^3 - \cancel{3\mu^3} - \cancel{3b\delta\mu^2\sigma} \\&\quad - \cancel{6b\delta\mu^2\sigma} - \cancel{6b^2\delta^2\mu\sigma^2} - \cancel{3\mu\sigma^2} - \cancel{3b\delta\sigma^3} + 2\mu^3 + 6b\delta\mu^2\sigma \\&\quad + \cancel{6b^2\delta^2\mu\sigma^2} + 2b^3\delta^3\sigma^3 \\&= 2b^3\delta^3\sigma^3 - b\delta^3\sigma^3 \\&= b\delta^3\sigma^3(2b^2 - 1)\end{aligned}$$

METHOD OF MOMENTS: CENTRAL MOMENTS OF THE SKEW-NORMAL

Our results, restated:

$$E(Y) = \mu + b\delta\sigma = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}}$$

$$E([Y - E(Y)]^2) = \sigma^2(1 - b^2\delta^2) = \sigma^2 \left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2} \right)$$

$$E([Y - E(Y)]^3) = b\delta^3\sigma^3(2b^2 - 1) = \sigma^3 \sqrt{\frac{2}{\pi}} \left(\frac{\lambda}{\sqrt{1 + \lambda^2}} \right)^3 \left(\frac{4}{\pi} - 1 \right)$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We're finally ready to derive our approximation!

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We start by setting the central moments of the binomial equal to the central moments of the skew-normal.

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We start by setting the central moments of the binomial equal to the central moments of the skew-normal.

$$np = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}} \quad (1a)$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We start by setting the central moments of the binomial equal to the central moments of the skew-normal.

$$np = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}} \quad (1a)$$

$$np(1 - p) = \sigma^2 \left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2} \right) \quad (1b)$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

We start by setting the central moments of the binomial equal to the central moments of the skew-normal.

$$np = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}} \quad (1a)$$

$$np(1 - p) = \sigma^2 \left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2} \right) \quad (1b)$$

$$np(p - 1)(2p - 1) = \sigma^3 \sqrt{\frac{2}{\pi}} \left(\frac{\lambda}{\sqrt{1 + \lambda^2}} \right)^3 \left(\frac{4}{\pi} - 1 \right) \quad (1c)$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

To get λ , divide the cube of (1b) by the square of (1c):

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

To get λ , divide the cube of (1b) by the square of (1c):

$$\frac{\sigma^6 \left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}\right)^3}{\sigma^6 \cdot \frac{2}{\pi} \left(\frac{\lambda}{\sqrt{1+\lambda^2}}\right)^6 \left(\frac{4}{\pi} - 1\right)^2} = \frac{n^3 p^3 (1-p)^3}{n^2 p^2 (p-1)^2 (2p-1)^2}$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

To get λ , divide the cube of (1b) by the square of (1c):

$$\begin{aligned} \frac{\sigma^6 \left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}\right)^3}{\sigma^6 \cdot \frac{2}{\pi} \left(\frac{\lambda}{\sqrt{1+\lambda^2}}\right)^6 \left(\frac{4}{\pi} - 1\right)^2} &= \frac{n^3 p^3 (1-p)^3}{n^2 p^2 (p-1)^2 (2p-1)^2} \\ \Rightarrow \frac{\left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}\right)^3}{\frac{2}{\pi} \left(\frac{\lambda^2}{1+\lambda^2}\right)^3 \left(\frac{4}{\pi} - 1\right)^2} &= \frac{np(1-p)}{(1-2p)^2} \end{aligned} \quad (2)$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Equation (2) can be solved for λ^2 .

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Equation (2) can be solved for λ^2 .

Then take λ to be

$$\lambda = \{\text{sign of } (1 - 2p)\} \sqrt{\lambda^2}.$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Equation (2) can be solved for λ^2 .

Then take λ to be

$$\lambda = \{\text{sign of } (1 - 2p)\} \sqrt{\lambda^2}.$$

Why?

METHOD OF MOMENTS: DERIVING AN APPROXIMATION


Equation (2) can be solved for λ^2 .

Then take λ to be

$$\lambda = \{\text{sign of } (1 - 2p)\} \sqrt{\lambda^2}.$$


Why? Recall Property 3 ...

METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When $p < 0.5$:


METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When $p < 0.5$:

- ▶ The binomial skews right (weight shifts left) and approaches $+|N(0, 1)| \rightarrow \lambda$ is positive


METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When $p < 0.5$:


- ▶ The binomial skews right (weight shifts left) and approaches $+|N(0, 1)| \rightarrow \lambda$ is positive
- ▶ $(1 - 2p)$ is positive

METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When $p > 0.5$:


METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When $p > 0.5$:

- ▶ The binomial skews left (weight shifts right) and approaches $-|N(0, 1)| \rightarrow \lambda$ is negative

METHOD OF MOMENTS: DERIVING AN APPROXIMATION



When $p > 0.5$:

- ▶ The binomial skews left (weight shifts right) and approaches $-|N(0, 1)| \rightarrow \lambda$ is negative
- ▶ $(1 - 2p)$ is negative

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Once you have λ , solve for σ and then μ .

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Once you have λ , solve for σ and then μ .

$$np(1 - p) = \sigma^2 \left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2} \right) \Rightarrow \sigma = \sqrt{\frac{np(1 - p)}{1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1 + \lambda^2}}}$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Once you have λ , solve for σ and then μ .

$$np(1-p) = \sigma^2 \left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2} \right) \Rightarrow \sigma = \sqrt{\frac{np(1-p)}{1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}}}$$

$$np = \mu + \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1+\lambda^2}} \Rightarrow \mu = np - \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1+\lambda^2}}$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

When $p = 0.5$, the right hand side of (2)

$$\frac{\left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}\right)^3}{\frac{2}{\pi} \left(\frac{\lambda^2}{1+\lambda^2}\right)^3 \left(\frac{4}{\pi} - 1\right)^2} = \frac{np(1-p)}{(1-2p)^2}$$

is undefined.

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

When $p = 0.5$, the right hand side of (2)

$$\frac{\left(1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}\right)^3}{\frac{2}{\pi} \left(\frac{\lambda^2}{1+\lambda^2}\right)^3 \left(\frac{4}{\pi} - 1\right)^2} = \frac{np(1-p)}{(1-2p)^2}$$

is undefined.

Uh-oh.

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Fortunately, our intuition rescues us:

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Fortunately, our intuition rescues us:

When $p = 0.5$, the binomial is symmetric, so λ should equal 0.

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Fortunately, our intuition rescues us:

When $p = 0.5$, the binomial is symmetric, so λ should equal 0.

This takes us back to the usual normal approximation:

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Fortunately, our intuition rescues us:

When $p = 0.5$, the binomial is symmetric, so λ should equal 0.

This takes us back to the usual normal approximation:

$$\sigma = \sqrt{\frac{np(1-p)}{1 - \frac{2}{\pi} \cdot \frac{0^2}{1+0^2}}} = \sqrt{\frac{np(1-p)}{1}} = \sqrt{np(1-p)}$$

METHOD OF MOMENTS: DERIVING AN APPROXIMATION

Fortunately, our intuition rescues us:

When $p = 0.5$, the binomial is symmetric, so λ should equal 0.

This takes us back to the usual normal approximation:

$$\sigma = \sqrt{\frac{np(1-p)}{1 - \frac{2}{\pi} \cdot \frac{0^2}{1+0^2}}} = \sqrt{\frac{np(1-p)}{1}} = \sqrt{np(1-p)}$$

$$\mu = np - \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{0}{\sqrt{1+0^2}} = np - 0 = np$$

METHOD OF MOMENTS: RESTRICTIONS

Unfortunately, though better than the normal approximation, our skew-normal method isn't universal.

METHOD OF MOMENTS: RESTRICTIONS

To be able to solve for λ , we must restrict n and p by the following equation:

METHOD OF MOMENTS: RESTRICTIONS

To be able to solve for λ , we must restrict n and p by the following equation:

$$np(1 - p) \geq (1 - 2p)^2 \quad (3)$$

METHOD OF MOMENTS: RESTRICTIONS

To be able to solve for λ , we must restrict n and p by the following equation:

$$np(1 - p) \geq (1 - 2p)^2 \quad (3)$$

From (3), we can answer two questions:

METHOD OF MOMENTS: RESTRICTIONS

One: Given p , what is the least n necessary?

METHOD OF MOMENTS: RESTRICTIONS


One: Given p , what is the least n necessary?

$$n \geq \frac{(1 - 2p)^2}{p(1 - p)}$$

METHOD OF MOMENTS: RESTRICTIONS

One: Given p , what is the least n necessary?

$$n \geq \frac{(1 - 2p)^2}{p(1 - p)}$$



METHOD OF MOMENTS: RESTRICTIONS

Two: Given n , what is the range of possible p 's?

METHOD OF MOMENTS: RESTRICTIONS


Two: Given n , what is the range of possible p 's?

$$\frac{1}{2} - \frac{1}{2}\sqrt{\frac{n}{n+4}} \leq p \leq \frac{1}{2} + \frac{1}{2}\sqrt{\frac{n}{n+4}}$$

METHOD OF MOMENTS: RESTRICTIONS

Two: Given n , what is the range of possible p 's?

$$\frac{1}{2} - \frac{1}{2}\sqrt{\frac{n}{n+4}} \leq p \leq \frac{1}{2} + \frac{1}{2}\sqrt{\frac{n}{n+4}}$$



DEMONSTRATING IMPROVED ACCURACY

We have an approximation!!

DEMONSTRATING IMPROVED ACCURACY

We have an approximation!!

But is it more accurate? ...

DEMONSTRATING IMPROVED ACCURACY


We have an approximation!!

But is it more accurate? ... Answer: Yes!


DEMONSTRATING IMPROVED ACCURACY: VISUAL

The easiest way of gauging accuracy is by visual inspection.


DEMONSTRATING IMPROVED ACCURACY: VISUAL




DEMONSTRATING IMPROVED ACCURACY: VISUAL




DEMONSTRATING IMPROVED ACCURACY: VISUAL




DEMONSTRATING IMPROVED ACCURACY: VISUAL




DEMONSTRATING IMPROVED ACCURACY: VISUAL




DEMONSTRATING IMPROVED ACCURACY: VISUAL




DEMONSTRATING IMPROVED ACCURACY: VISUAL



DEMONSTRATING IMPROVED ACCURACY: VISUAL



DEMONSTRATING IMPROVED ACCURACY: VISUAL



DEMONSTRATING IMPROVED ACCURACY: MABS

A more numerical way of gauging accuracy is the maximal absolute error (*MABS*), defined as

$$\text{MABS}(n, p) = \max_{k \in \{0, 1, \dots, n\}} \left| F_{B(n, p)}(k) - F_{\text{appr}(n, p)}(k + 0.5) \right|.$$

DEMONSTRATING IMPROVED ACCURACY: MABS


A more numerical way of gauging accuracy is the maximal absolute error (*MABS*), defined as

$$\text{MABS}(n, p) = \max_{k \in \{0, 1, \dots, n\}} \left| F_{B(n, p)}(k) - F_{\text{appr}(n, p)}(k + 0.5) \right|.$$

(In English: The biggest vertical distance between the binomial cdf and the approximation curve.)


DEMONSTRATING IMPROVED ACCURACY: MABS

MABS as a function of p , with fixed $n = 25$:




DEMONSTRATING IMPROVED ACCURACY: MABS

MABS as a function of p , with fixed $n = 100$:




DEMONSTRATING IMPROVED ACCURACY: MABS

MABS as a function of n , with fixed $p = 0.05$:



DEMONSTRATING IMPROVED ACCURACY: MABS

MABS as a function of n , with fixed $p = 0.1$:



RESOURCES

A few resources you might find helpful ...

RESOURCES

Estimations of $SN(\mu, \sigma, \lambda)$ for $Bin(n, p)$

		n				
		25	50	100	250	500
p	0.05	(-0.11, 1.74, 4.56)	(0.79, 2.30, 2.54)	(2.85, 3.06, 1.86)	(9.58, 4.52, 1.38)	(21.32, 6.11, 1.15)
	0.10	(0.89, 2.20, 2.31)	(2.97, 2.94, 1.74)	(7.44, 3.94, 1.40)	(21.53, 5.88, 1.10)	(45.62, 8.01, 0.94)
	0.15	(2.02, 2.49, 1.79)	(5.32, 3.34, 1.43)	(12.25, 4.51, 1.19)	(33.77, 6.77, 0.96)	(70.30, 9.27, 0.82)
	0.20	(3.23, 2.67, 1.50)	(7.76, 3.61, 1.24)	(17.18, 4.89, 1.04)	(46.18, 7.39, 0.85)	(95.18, 10.16, 0.74)
	0.25	(4.49, 2.79, 1.29)	(10.28, 3.78, 1.09)	(22.20, 5.15, 0.93)	(58.71, 7.83, 0.76)	(120.22, 10.80, 0.67)
	0.30	(5.80, 2.85, 1.12)	(12.86, 3.88, 0.95)	(27.31, 5.32, 0.82)	(71.34, 8.12, 0.68)	(145.39, 11.24, 0.60)
	0.35	(7.17, 2.86, 0.96)	(15.50, 3.92, 0.83)	(32.49, 5.39, 0.72)	(84.09, 8.28, 0.60)	(170.70, 11.50, 0.53)
	0.40	(8.59, 2.83, 0.80)	(18.23, 3.89, 0.70)	(37.76, 5.39, 0.61)	(96.96, 8.32, 0.51)	(196.18, 11.60, 0.45)
	0.45	(10.12, 2.73, 0.61)	(21.08, 3.79, 0.53)	(43.21, 5.29, 0.47)	(110.07, 8.23, 0.40)	(221.93, 11.54, 0.35)
	0.50	(12.50, 2.50, 0.00)	(25.00, 3.54, 0.00)	(50.00, 5.00, 0.00)	(125.00, 7.91, 0.00)	(250.00, 11.18, 0.00)
	0.55	(14.88, 2.73, -0.61)	(28.92, 3.79, -0.53)	(56.79, 5.29, -0.47)	(139.93, 8.23, -0.40)	(278.07, 11.54, -0.35)
	0.60	(16.41, 2.83, -0.80)	(31.77, 3.89, -0.70)	(62.24, 5.39, -0.61)	(153.04, 8.32, -0.51)	(303.82, 11.60, -0.45)
	0.65	(17.83, 2.86, -0.96)	(34.50, 3.92, -0.83)	(67.51, 5.39, -0.72)	(165.91, 8.28, -0.60)	(329.30, 11.50, -0.53)
	0.70	(19.20, 2.85, -1.12)	(37.14, 3.88, -0.95)	(72.69, 5.32, -0.82)	(178.66, 8.12, -0.68)	(354.61, 11.24, -0.60)
	0.75	(20.51, 2.79, -1.29)	(39.72, 3.78, -1.09)	(77.80, 5.15, -0.93)	(191.29, 7.83, -0.76)	(379.78, 10.80, -0.67)
	0.80	(21.77, 2.67, -1.50)	(42.24, 3.61, -1.24)	(82.82, 4.89, -1.04)	(203.82, 7.39, -0.85)	(404.82, 10.16, -0.74)
	0.85	(22.98, 2.49, -1.79)	(44.68, 3.34, -1.43)	(87.75, 4.51, -1.19)	(216.23, 6.77, -0.96)	(429.70, 9.27, -0.82)
	0.90	(24.11, 2.20, -2.31)	(47.03, 2.94, -1.74)	(92.56, 3.94, -1.40)	(228.47, 5.88, -1.10)	(454.38, 8.01, -0.94)
	0.95	(25.11, 1.74, -4.56)	(49.21, 2.30, -2.54)	(97.15, 3.06, -1.86)	(240.42, 4.52, -1.38)	(478.68, 6.11, -1.15)

RESOURCES

All values in my project were computed using a Python library, which is freely available online:

<http://github.com/joycetipping/skew-normal-capstone/>

BIBLIOGRAPHY I

- B. C. Arnold and G. D. Lin. Characterizations of the skew-normal and generalized chi distributions. *Sankhy: The Indian Journal of Statistics*, 66(4):593–606, 2004.
- Adelchi Azzalini. A class of distributions which includes the normal ones. *Scandinavian Journal of Statistics*, 12: 171–178, 1985.
- Adelchi Azzalini. The skew-normal distribution and related multivariate families. *Scandinavian Journal of Statistics*, 32: 159–188, 2005.
- Lee J. Bain and Max Engelhardt. *Introduction to Probability and Mathematical Statistics*. Duxbury, 2nd edition, 1992.
- Ching-Hui Chang, Jyh-Jiuan Lin, Nabendu Pal, and Miao-Chen Chiang. A note on improved approximation of the binomial distribution by the skew-normal distribution. *American Statistical Association*, 62(2):167–170, May 2008.

BIBLIOGRAPHY II

- A. K. Gupta, T. T. Nguyen, and J. A. T. Sanqui. Characterization of the skew-normal distribution. *Annals of the Institute of Statistical Mathematics*, 56:351–360, 2004.
- A. O'Hagan and Tom Leonard. Bayes estimation subject to uncertainty about parameter constraints. *Biometrika*, 63: 201–203, 1976.
- Arthur Pewsey. Problems of inference for azzalini's skew-normal distribution. *Journal of Applied Statistics*, 27(7): 859–870, 2000.
- Martin Schader and Friedrich Schmid. Two rules of thumb for the approximation of the binomial distribution by the normal distribution. *American Statistical Association*, 43(1):23–24, February 1989.

Appendix

Calculating a skew-normal approximation

CALCULATING A SKEW-NORMAL APPROXIMATION

Although easier with a computer program, calculating estimates for μ , σ , and λ by hand is perfectly possible.

CALCULATING A SKEW-NORMAL APPROXIMATION

Although easier with a computer program, calculating estimates for μ , σ , and λ by hand is perfectly possible.

Here's an example!

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (2):

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (2):

$$\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi} \right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2} \right) = \frac{np(1 - p)}{(1 - 2p)^2} \quad (4)$$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (2):

$$\underbrace{\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right)}_{f(\lambda)} = \frac{np(1 - p)}{(1 - 2p)^2} \quad (4)$$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (2):

$$\underbrace{\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right)}_{f(\lambda)} = \underbrace{\frac{np(1 - p)}{(1 - 2p)^2}}_{k_{n,p}} \quad (4)$$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (2):

$$\underbrace{\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right)}_{f(\lambda)} = \underbrace{\frac{np(1 - p)}{(1 - 2p)^2}}_{k_{n,p}} \quad (4)$$

The closed form solution to (4) is pretty hideous, so we'll take a numerical approach.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

By far the biggest battle is λ .

We'll use a simplified version of equation (2):

$$\underbrace{\left(\frac{1 + \lambda^2}{\lambda^2} - \frac{2}{\pi}\right)^3 \left(\frac{\pi^3}{2(4 - \pi)^2}\right)}_{f(\lambda)} = \underbrace{\frac{np(1 - p)}{(1 - 2p)^2}}_{k_{n,p}} \quad (4)$$

The closed form solution to (4) is pretty hideous, so we'll take a numerical approach.

Our goal is to find λ such that $f(\lambda)$ is within a certain margin of error (e) of $k_{n,p}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Recall that the sign of λ is determined independently of the value. Thus, we can consider only $\lambda > 0$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Recall that the sign of λ is determined independently of the value. Thus, we can consider only $\lambda > 0$.

It is possible to show, by taking its derivative, that f is monotonically decreasing for positive λ .

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Recall that the sign of λ is determined independently of the value. Thus, we can consider only $\lambda > 0$.

It is possible to show, by taking its derivative, that f is monotonically decreasing for positive λ .

This convenient fact allows us to find lower and upper bounds for λ and repeatedly bisect our interval until we are within ϵ of $k_{n,p}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

For this demonstration, we will take $n = 25$ and $p = 0.1$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

For this demonstration, we will take $n = 25$ and $p = 0.1$.

Since we're doing this by hand, we'll take our error margin e to be a modest 0.01.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 1: Find $k_{n,p}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 1: Find $k_{n,p}$.

Our value: $k_{n,p} = \frac{25 \cdot 0.1 \cdot 0.9}{(1 - 2 \cdot 0.1)^2} = 3.5156.$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 1: Find $k_{n,p}$.

Our value: $k_{n,p} = \frac{25 \cdot 0.1 \cdot 0.9}{(1 - 2 \cdot 0.1)^2} = 3.5156.$

Step 2: Find a and b such that $f(a) > k_{n,p} > f(b)$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 1: Find $k_{n,p}$.

Our value: $k_{n,p} = \frac{25 \cdot 0.1 \cdot 0.9}{(1 - 2 \cdot 0.1)^2} = 3.5156.$

Step 2: Find a and b such that $f(a) > k_{n,p} > f(b)$.

Our values: $a = 1, b = 3.$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

Calculate $c = \frac{a+b}{2}$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

Calculate $c = \frac{a+b}{2}$.

- If $f(c) \leq k_{n,p} - 0.01$, we need a small value of c , so we take our new interval to be (a, c) .

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

Calculate $c = \frac{a + b}{2}$.

- ▶ If $f(c) \leq k_{n,p} - 0.01$, we need a small value of c , so we take our new interval to be (a, c) .
- ▶ If $f(c) \geq k_{n,p} + 0.01$, we need a larger value of c , so we take our new interval to be (c, b) .

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 3: Repeatedly bisect (a, b) until $f(c)$ is within e of $k_{n,p}$.

Calculate $c = \frac{a+b}{2}$.

- ▶ If $f(c) \leq k_{n,p} - 0.01$, we need a small value of c , so we take our new interval to be (a, c) .
- ▶ If $f(c) \geq k_{n,p} + 0.01$, we need a larger value of c , so we take our new interval to be (c, b) .

Repeat this step until $f(c)$ is within e of $k_{n,p}$, or more precisely $k_{n,p} - 0.01 < f(c) < k_{n,p} + 0.01$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
-----------	-----	-----	-----	--------	----------------------------	----------------------------

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False
2	2.00	2.500	2.2500	3.7129	False	True

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False
2	2.00	2.500	2.2500	3.7129	False	True
3	2.25	2.500	2.3750	3.3252	True	False

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False
2	2.00	2.500	2.2500	3.7129	False	True
3	2.25	2.500	2.3750	3.3252	True	False
4	2.25	2.375	2.3125	3.5076	False	False

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

(Step 3)

The following table shows our iterations:

Iteration	a	b	c	$f(c)$	$f(c) \leq k_{n,p} - 0.01$	$f(c) \geq k_{n,p} + 0.01$
1	2.00	3.000	2.5000	3.0164	True	False
2	2.00	2.500	2.2500	3.7129	False	True
3	2.25	2.500	2.3750	3.3252	True	False
4	2.25	2.375	2.3125	3.5076	False	False

We take the last value of c : 2.3125.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 5: Find the sign of $(1 - 2p)$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 5: Find the sign of $(1 - 2p)$.

Our $p = 0.1 \Rightarrow (1 - 2 \cdot 0.1) = 0.8 \Rightarrow$ positive.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 5: Find the sign of $(1 - 2p)$.

Our $p = 0.1 \Rightarrow (1 - 2 \cdot 0.1) = 0.8 \Rightarrow$ positive.

Step 6: Final answer: $\{\text{sign of } (1 - 2p)\}\lambda$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING λ

Step 5: Find the sign of $(1 - 2p)$.

Our $p = 0.1 \Rightarrow (1 - 2 \cdot 0.1) = 0.8 \Rightarrow$ positive.

Step 6: Final answer: $\{\text{sign of } (1 - 2p)\}\lambda$

Our final answer: $\lambda = 2.3125$.

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING σ

Once we have λ , we can easily find σ :

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING σ

Once we have λ , we can easily find σ :

$$\sigma = \sqrt{\frac{np(1-p)}{1 - \frac{2}{\pi} \cdot \frac{\lambda^2}{1+\lambda^2}}} = \sqrt{\frac{25 \cdot 0.1 \cdot 0.9}{1 - \frac{2}{\pi} \cdot \frac{2.3125^2}{1+2.3125^2}}} = 2.2029.$$

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING μ

And with λ and σ , we can also find μ :

CALCULATING A SKEW-NORMAL APPROXIMATION: FINDING μ

And with λ and σ , we can also find μ :

$$\begin{aligned}\mu &= np - \sigma \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{\lambda}{\sqrt{1 + \lambda^2}} \\ &= 25 \cdot 0.1 - 2.2029 \cdot \sqrt{\frac{2}{\pi}} \cdot \frac{2.3125}{\sqrt{1 + 2.3125^2}} \\ &= 0.8867.\end{aligned}$$

Here's the bibliography again ...

BIBLIOGRAPHY I

- B. C. Arnold and G. D. Lin. Characterizations of the skew-normal and generalized chi distributions. *Sankhy: The Indian Journal of Statistics*, 66(4):593–606, 2004.
- Adelchi Azzalini. A class of distributions which includes the normal ones. *Scandinavian Journal of Statistics*, 12: 171–178, 1985.
- Adelchi Azzalini. The skew-normal distribution and related multivariate families. *Scandinavian Journal of Statistics*, 32: 159–188, 2005.
- Lee J. Bain and Max Engelhardt. *Introduction to Probability and Mathematical Statistics*. Duxbury, 2nd edition, 1992.
- Ching-Hui Chang, Jyh-Jiuan Lin, Nabendu Pal, and Miao-Chen Chiang. A note on improved approximation of the binomial distribution by the skew-normal distribution. *American Statistical Association*, 62(2):167–170, May 2008.

BIBLIOGRAPHY II

- A. K. Gupta, T. T. Nguyen, and J. A. T. Sanqui. Characterization of the skew-normal distribution. *Annals of the Institute of Statistical Mathematics*, 56:351–360, 2004.
- A. O'Hagan and Tom Leonard. Bayes estimation subject to uncertainty about parameter constraints. *Biometrika*, 63: 201–203, 1976.
- Arthur Pewsey. Problems of inference for azzalini's skew-normal distribution. *Journal of Applied Statistics*, 27(7): 859–870, 2000.
- Martin Schader and Friedrich Schmid. Two rules of thumb for the approximation of the binomial distribution by the normal distribution. *American Statistical Association*, 43(1):23–24, February 1989.