


Calculus On Normed Vector Spaces

MTP REPORT

September 2017

Submitted by
Joydeep Medhi
Entry No. 2013MT60599

Supervisor
Dr. Amit Priyadarshi



Department of Mathematics
Indian Institute of Technology Delhi,
New Delhi, INDIA

1 Introduction

The aim of the project is to study the notion of derivatives on general normed vector spaces and do Calculus on them. We will study generalizations of several well-known theorems of Calculus. We will also look at some applications of these concepts.

2 Definitions

In this section, some basic definitions and elementary properties are discussed.

Norm: We will suppose that all vector spaces are real. Let E be a vector space. A mapping $\|\cdot\| : E \rightarrow \mathbb{R}$, is said to be a *norm* if, for all $x, y \in E$ and $\lambda \in \mathbb{R}$

- $\|x\| \geq 0$;
- $\|x\| = 0 \Leftrightarrow x = 0$;
- $\|\lambda x\| = |\lambda| \|x\|$;
- $\|x + y\| \leq \|x\| + \|y\|$.

The pair $(E, \|\cdot\|)$ is called a *normed vector space* and we say that $\|x\|$ is the norm of x .

Continuity: Suppose now that we have two normed vector spaces, $(E, \|\cdot\|_E)$ and $(F, \|\cdot\|_F)$. Let A be a subset of E , f a mapping of A into F and $a \in A$. We say that f is *continuous* at a if the following condition is satisfied:

for all $\epsilon > 0$, there exists $\delta > 0$ such that, if $x \in A$ and $\|x - a\|_E < \delta$, then $\|f(x) - f(a)\|_F < \epsilon$.
If f is *continuous* at every point $a \in A$, then we say that f is *continuous* on A .

Proposition 2.1. *The norm on a normed vector space is a continuous function.*

Proof. We have $\|x\| = \|x - y + y\| \leq \|x - y\| + \|y\| \Rightarrow \|x\| - \|y\| \leq \|x - y\|$

In the same way, $\|y\| - \|x\| \leq \|y - x\|$. As $\|y - x\| = \|x - y\|$, We have

$$|\|x\| - \|y\|| \leq \|x - y\|$$

And hence the continuity.

Proposition 2.2. *Let E and F be normed vector spaces, $A \subseteq E, a \in A$, f and g are mappings from E into F and $\lambda \in \mathbb{R}$.*

- *If f and g are continuous at a , then so is $f + g$.*
- *If f is continuous at a , then so is λf .*
- *If α is a real-valued function defined on E and both f and α are continuous at a , then so is αf .*

Proposition 2.3. Let $(E, \|\cdot\|_E)$ be a normed vector space

- The mapping $f : E \times E \longrightarrow E, (x, y) \mapsto x + y$ is continuous.
- The mapping $f : \mathbb{R} \times E \longrightarrow E, (\lambda, x) \mapsto \lambda x$ is continuous.

3 Differentiation

In this section we will be primarily concerned with extending the derivative defined for real-valued functions defined on an interval of \mathbb{R} . We will also consider minima and maxima of real-valued functions defined on a normed vector space.

3.1 Directional Derivatives

Let O be an open subset of a normed vector space E , f a real-valued function defined on O , $a \in O$ and u a nonzero element of E . The function $f_u : t \rightarrow f(a + tu)$ is defined on an open interval containing 0. If the derivative $\frac{df_u}{dt}(0)$ is defined, i.e., if the limit

$$\lim_{t \rightarrow 0} \frac{f(a + tu) - f(a)}{t}$$

exists, then it is called the *directional derivative* of f at a in the direction of u , i.e. $\partial_u f(a)$.

If $E = \mathbb{R}_n$ and e_i is its standard basis, then the directional derivative $e_i f(a)$ is called the i th partial derivative of f at a , or the derivative of f with respect to x_i at a .

$$\frac{\partial f}{\partial x_i} = \lim_{t \rightarrow 0} \frac{f(a_1, \dots, a_i + t, \dots, a_n) - f(a_1, \dots, a_n)}{t}$$

If for every point $x \in O$, the partial derivative $\frac{\partial f}{\partial x_i}(x)$ is defined, then we obtain the function i th partial derivative defined on O . If these functions are defined and continuous for all i , then we say that the function f is of class C^1 .

Example 3.1. If f is the function defined on \mathbb{R}^2 by $f(x, y) = xe^{xy}$, then the partial derivatives with respect to x and y are defined at all points $(x, y) \in \mathbb{R}^2$ and

$$\frac{\partial f}{\partial x}(x, y) = (1 + xy)e^{xy} \text{ and } \frac{\partial f}{\partial y}(x, y) = x^2 e^{xy}$$

As both are continuous, f is of class C^1 .

However, a function of two or more variables may have all its partial derivatives defined at a given point without being *continuous* there. Here is an example.

Example 3.2. Consider the function f defined on \mathbb{R}_2 by

$$f(x, y) = \begin{cases} \frac{x^6}{x^8 + (y - x^2)^2} & \text{if } (x, y) \neq (0, 0) \\ 0 & \text{otherwise} \end{cases}$$

We have (for x and y)

$$\lim_{t \rightarrow 0} \frac{t^6}{t^8 + t^4} / t = 0 \quad \text{and} \quad \lim_{t \rightarrow 0} \frac{0}{t^2} / t = 0$$

and so,

$$\frac{\partial f}{\partial x}(0, 0) = \frac{\partial f}{\partial y}(0, 0) = 0.$$

However, $\lim_{t \rightarrow 0} f(x, x^2) = \infty$, which indicates f is not continuous at 0.

Suppose now that O is an open subset of \mathbb{R}^n and f a mapping defined on O with image in \mathbb{R}^m . f has m coordinate mappings f_1, \dots, f_m . If $a \in O$ and the partial derivatives $\frac{\partial f_i}{\partial x_j}$ of a , for $1 \leq i \leq m$ and $1 \leq j \leq n$, are all defined, then the $m \times n$ matrix

$$J_f(a) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}$$

is called the *Jacobian Matrix* of f at a .

3.2 The Differential

Let E and F be normed vector spaces, O an open subset of E containing 0, and g a mapping from O into F such that $g(0) = 0$. If there exists a mapping ϵ , defined on a neighbourhood of $0 \in E$ and with image in F , such that $\lim_{h \rightarrow 0} \epsilon(h) = 0$ and

$$g(h) = \|h\|_E \epsilon(h),$$

then we write $g(h) = o(h)$ and say that g is "small o of h ".

The condition $g(h) = o(h)$ is independent of the norms we choose for two spaces E and F .

Let O be an open subset of a normed vector space E and f a mapping from O into a normed vector space F . If $a \in O$ and there is a continuous linear mapping $\phi : E \rightarrow F$ such that

$$f(a + h) = f(a) + \phi(h) + o(h)$$

when h is close to 0, then we say that f is *differentiable* at a .

Proposition 3.1. *If f is differentiable at a , then*

(a) *f is continuous at a ;*

(b) *ϕ is unique.*

Remark 3.1. *If E and F are normed vector spaces and $f : E \rightarrow F$ is constant, then $f'(a)$ is the zero mapping at any point $a \in E$. If $f : E \rightarrow F$ is linear and continuous, then $f'(a) = f$ at any point $a \in E$.*

Proposition 3.2. *Let f be a mapping defined on an open subset O of a normed vector space E with image in the cartesian product $F = F_1 \times \dots \times F_p$. Then f is differentiable at $a \in O$ if and only if the coordinate mappings f_i , for $i = 1, \dots, p$, are differentiable at a .*

$$f'(a) = (f'_1(a), \dots, f'_p(a))$$

Suppose that $\dim E = n < \infty$ and that e_i is a basis of E . If $x = \sum_{i=1}^n x_i e_i$, then

$$f'(a)x = \sum_{i=1}^n x_i f'(a)e_i = \sum_{i=1}^n \partial_{e_i} f(a) e_i^*(x),$$

where (e_i^*) is the dual basis of (e_i) . We thus obtain the expression. If $E = \mathbb{R}^n$ and (e_i) is its standard basis, then we usually write dx_i for e_i^* . This gives us the expression

$$f'(a)x = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(a) dx_i.$$

Differentiability at a given point: If we wish to determine whether a real-valued function f defined on an open subset of \mathbb{R}^n is differentiable at a given point a , then first we determine whether all its partial derivatives at a exist. If this is not the case, then f is not differentiable at a .

If all the partial derivatives exist, then we know that the only possibility for $f'(a)$ is the linear function $\phi = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(a) dx_i$. We consider the expression,

$$\frac{f(a + h) - f(a) - \phi(h)}{\|h\|} = \epsilon(h)$$

If $\lim_{h \rightarrow 0} \epsilon(h) = 0$, then f is differentiable at a , otherwise it is not.

3.3 Differentials of Compositions

Let E , F and G be normed vector spaces, O an open subset of E , U an open subset of F and $f : O \rightarrow F$, $g : U \rightarrow G$ be such that $f(O) \subset U$. Then the mapping $g \circ f$ is defined on O .

Theorem 3.1. *If f is differentiable at a and g is differentiable at $f(a)$, then $g \circ f$ is differentiable at a and*

$$(g \circ f)'(a) = g'(f(a)) \circ f'(a).$$

This expression is referred to as Chain Rule.

Corollary 3.1. *If in the above theorem the normed vector spaces are euclidian spaces, then*

$$J_{g \circ f}(a) = J_g(f(a)) \circ J_f(a).$$

Example 3.3. *Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ and $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by*

$$f(x, y, z) = (xy, e^{xz}) \quad g(u, v) = u^2v. \text{ Then,}$$

$$J_f(x, y, z) = \begin{bmatrix} y & x & 0 \\ ze^{xz} & 0 & xe^{xz} \end{bmatrix}$$

and

$$J_g(u, v) = [2uvu^2]$$

Multiplying the matrices $J_g(f(x, y, z))$ and $J_f(x, y, z)$, we obtain

$$J_{g \circ f}(x, y, z) = (2xy^2 + x^2y^2z)e^{xz}2x^2ye^{xz}x^3y^3e^{xz}.$$

3.4 Differentiability of the Norm

If E is a normed vector space with norm $\|\cdot\|$, then $\|\cdot\|$ is itself a mapping from E into \mathbb{R} and we can study its *differentiability*. We will write $Df(\|\cdot\|)(x)$ for differentiability of the norm at x (if exists).

Proposition 3.3. *Norm is not differentiable at the origin.*

Proof. Suppose $Df(\|\cdot\|)$ exists. Then for small non-zero values of h , we have

$$\|h\| = Df(\|\cdot\|)(0)h + o(h) \Rightarrow \lim_{h \rightarrow 0} \left(1 - Df(\|\cdot\|)\frac{h}{\|h\|}\right) = 0$$

And

$$\|h\| = \|-h\| = -Df(\|\cdot\|)(0)h + o(h) \Rightarrow \lim_{h \rightarrow 0} \left(1 + Df(\|\cdot\|)\frac{h}{\|h\|}\right) = 0$$

Summing the two limits we obtain $2 = 0$, which is a contradiction. Hence $Df(\|\cdot\|)(0)$ does not exist.

At points where the differential exists, we have the following interesting result:

Proposition 3.4. *Let E be a normed vector space and $\|\cdot\|$ its norm. If $\|\cdot\|$ is differentiable at $a \neq 0$ and $\lambda > 0$, then $\|\cdot\|$ is differentiable at λa and $Df(\|\cdot\|)(\lambda a) = Df(\|\cdot\|)(a)$. In addition, $|Df(\|\cdot\|)(a)|_{E^*} = 1$*

Proof. If $\|\cdot\|$ is differentiable at a , $\lambda > 0$ and $h \in E \setminus \{0\}$, then we have

$$\|\lambda a + h\| = \lambda \|a + \frac{h}{\lambda}\| = \lambda \left(\|a\| + Df(\|\cdot\|)\left(\frac{h}{\lambda}\right) + o\left(\frac{h}{\lambda}\right)\right) = \|\lambda a\| + Df(\|\cdot\|)(a)h + o(h)$$

It follows that $Df(\|\cdot\|)(\lambda a)$ exists and $Df(\|\cdot\|)(\lambda a) = Df(\|\cdot\|)(a)$.

For 2nd part, Consider the function, $f : \mathbb{R}_+^* \rightarrow \mathbb{R}, \lambda \mapsto \|\lambda a\|$

For a given $\lambda \in \mathbb{R}_+^*$ and $h \in \mathbb{R}$ sufficiently small, we have

$$\|(\lambda + h)a\| = (\lambda + h)\|a\|$$

and so

$$\lim_{h \rightarrow 0} \frac{\|(\lambda + h)a - \|\lambda a\|\|}{h} = \lim_{h \rightarrow 0} \frac{h\|a\|}{h} = \|a\|$$

Therefore $\dot{f}(\lambda) = \|a\|$ for all values of λ . On the other hand, $f = \|\cdot\| \circ \phi$, where $\phi(\lambda) = \lambda a$, and so

$$(f'(\lambda))s = Df(\|\cdot\|)(\lambda a)sa = a(Df(\|\cdot\|)(a))a$$

This implies, $\dot{f}(\lambda) = Df(\|\cdot\|)(a)a$ and hence $Df(\|\cdot\|)(a)a = \|a\|$. It follows that $|Df(\|\cdot\|)(a)|_{E^*} = 1$.

4 References

- [1] Avez, A.: Differential Calculus. J. Wiley and Sons Ltd, New York (1986)
- [2] Rodney Coleman: Calculus on Normed Vector Spaces (2012)
- [3] Dacorogna, B.: Introduction to the Calculus of Variations. Imperial College Press, London (2004)