

FAQ: How do I interpret odds ratios in logistic regression?

Introduction


When a binary outcome variable is modeled using logistic regression, it is assumed that the logit transformation of the outcome variable has a linear relationship with the predictor variables. This makes the interpretation of the regression coefficients somewhat tricky. In this page, we will walk through the concept of odds ratio and try to interpret the logistic regression results using the concept of odds ratio in a couple of examples.

From probability to odds to log of odds

Everything starts with the concept of probability. Let's say that the probability of success of some event is .8. Then the probability of failure is $1 - .8 = .2$. The odds of success are defined as the ratio of the probability of success over the probability of failure. In our example, the odds of success are $.8 / .2 = 4$. That is to say that the odds of success are 4 to 1. If the probability of success is .5, i.e., 50-50 percent chance, then the odds of success is 1 to 1.


The transformation from probability to odds is a monotonic transformation, meaning the odds increase as the probability increases or vice versa. Probability ranges from 0 and 1. Odds range from 0 and positive infinity. Below is a table of the transformation from probability to odds and we have also plotted for the range of p less than or equal to .9.

p	odds
.001	.001001
.01	.010101
.15	.1764706
.2	.25
.25	.3333333
.3	.4285714
.35	.5384616
.4	.6666667
.45	.8181818
.5	1
.55	1.222222
.6	1.5
.65	1.857143
.7	2.333333
.75	3
.8	4
.85	5.666667
.9	9
.999	999
.9999	9999



The transformation from odds to log of odds is the log transformation. Again this is a monotonic transformation. That is to say, the greater the odds, the greater the log of odds and vice versa. The table below shows the relationship among the probability, odds and log of odds. We have also shown the plot of log odds against odds.

p	odds	logodds
.001	.001001	-6.906755
.01	.010101	-4.59512
.15	.1764706	-1.734601
.2	.25	-1.386294
.25	.3333333	-1.098612
.3	.4285714	-.8472978
.35	.5384616	-.6190392
.4	.6666667	-.4054651
.45	.8181818	-.2006707
.5	1	0
.55	1.222222	.2006707
.6	1.5	.4054651
.65	1.857143	.6190392
.7	2.333333	.8472978
.75	3	1.098612
.8	4	1.386294
.85	5.666667	1.734601
.9	9	2.197225
.999	999	6.906755
.9999	9999	9.21024



Why do we take all the trouble doing the transformation from probability to log odds? One reason is that it is usually difficult to model a variable which has restricted range, such as probability. This transformation is an attempt to get around the restricted range problem. It maps probability ranging between 0 and 1 to log odds ranging from negative infinity to positive infinity. Another reason is that among all of the infinitely many choices of transformation, the log of odds is one of the easiest to understand and interpret. This transformation is called logit transformation. The other common choice is the probit transformation, which will not be covered here.

A logistic regression model allows us to establish a relationship between a binary outcome variable and a group of predictor variables. It models the logit-transformed probability as a linear relationship with the predictor variables. More formally, let y be the binary outcome variable indicating failure/success with 0/1 and p be the probability of y to be 1, $p = \text{prob}(y=1)$. Let x_1, \dots, x_k be a set of predictor variables. Then the logistic regression of y on x_1, \dots, x_k estimates parameter values for $\beta_0, \beta_1, \dots, \beta_k$ via maximum likelihood method of the following equation.

$$\text{logit}(p) = \log(p/(1-p)) = \beta_0 + \beta_1 x_1 + \dots + \beta_k x_k$$

In terms of probabilities, the equation above is translated into

$$p = \exp(\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k) / (1 + \exp(\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k))$$

We are now ready for a few examples of logistic regressions. We will use a sample dataset, [sample.csv](#), for the purpose of illustration. The data set has 200 observations and the outcome variable used will be **hon**, indicating if a student is in an honors class or not. So our $p = \text{prob}(\text{hon}=1)$. We will purposely ignore all the significance tests and focus on the meaning of the regression coefficients. The output on this page was created using Stata with some editing.

Logistic regression with no predictor variables

Let's start with the simplest logistic regression, a model without any predictor variables. In an equation, we are modeling

$$\text{logit}(p) = \beta_0$$

Logistic regression	Number of obs	=	200
	LR chi2(0)	=	0.00
	Prob > chi2	=	.
Log likelihood = -111.35502	Pseudo R2	=	0.0000

hon	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]
intercept	-1.12546	.1644101	-6.85	0.000	-1.447697 - .8032217

This means $\log(p/(1-p)) = -1.12546$. What is p here? It turns out that p is the overall probability of being in honors class ($\text{hon} = 1$). Let's take a look at the frequency table for **hon**.

hon	Freq.	Percent	Cum.
0	151	75.50	75.50
1	49	24.50	100.00
Total	200	100.00	

So $p = 49/200 = .245$. The odds are $.245/(1-.245) = .3245$ and the log of the odds (logit) is $\log(.3245) = -1.12546$. In other words, the intercept from the model with no predictor variables is the estimated log odds of being in honors class for the whole population of interest. We can also transform the log of the odds back to a probability: $p = \exp(-1.12546) / (1 + \exp(-1.12546)) = .245$, if we like.

Logistic regression with a single dichotomous predictor variables

Now let's go one step further by adding a binary predictor variable, **female**, to the model. Writing it in an equation, the model describes the following linear relationship.

$$\text{logit}(p) = \beta_0 + \beta_1 \text{female}$$

Logistic regression	Number of obs	=	200
	LR chi2(1)	=	3.10
	Prob > chi2	=	0.0781
Log likelihood = -109.80312	Pseudo R2	=	0.0139

hon	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]
female	.5927822	.3414294	1.74	0.083	-.0764072 1.261972
intercept	-1.470852	.2689555	-5.47	0.000	-1.997995 -.9437087

Before trying to interpret the two parameters estimated above, let's take a look at the crosstab of the variable **hon** with **female**.

hon	female		Total
	male	female	
0	74	77	151
1	17	32	49
Total	91	109	200

In our dataset, what are the odds of a male being in the honors class and what are the odds of a female being in the honors class? We can manually calculate these odds from the table: for males, the odds of being in the honors class are $(17/91)/(74/91) = 17/74 = .23$; and for females, the odds of being in the honors class are $(32/109)/(77/109) = 32/77 = .42$. The ratio of the odds for female to the odds for male is $(32/77)/(17/74) = (32*74)/(77*17) = 1.809$. So the odds for males are 17 to 74, the odds for females are 32 to 77, and the odds for female are about 81% higher than the odds for males.

Now we can relate the odds for males and females and the output from the logistic regression. The intercept of -1.471 is the log odds for males since male is the reference group (**female** = 0). Using the odds we calculated above for males, we can confirm this: $\log(.23) = -1.47$. The coefficient for **female** is the log of odds ratio between the female group and male group: $\log(1.809) = .593$. So we can get the odds ratio by exponentiating the coefficient for female. Most statistical packages display both the raw regression coefficients and the exponentiated coefficients for logistic regression models. The table below is created by Stata.

Logistic regression				Number of obs	=	200
				LR chi2(1)	=	3.10
				Prob > chi2	=	0.0781
				Pseudo R2	=	0.0139
Log likelihood = -109.80312						
hon	Odds Ratio	Std. Err.	z	P> z	[95% Conf. Interval]	
female	1.809015	.6176508	1.74	0.083	.9264389	3.532379

Logistic regression with a single continuous predictor variable

Another simple example is a model with a single continuous predictor variable such as the model below. It describes the relationship between students' math scores and the log odds of being in an honors class.

$$\text{logit}(p) = \beta_0 + \beta_1 \cdot \text{math}$$

Logistic regression				Number of obs	=	200
				LR chi2(1)	=	55.64
				Prob > chi2	=	0.0000
				Pseudo R2	=	0.2498
Log likelihood = -83.536619						
hon	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
math	.1563404	.0256095	6.10	0.000	.1061467	.206534
intercept	-9.793942	1.481745	-6.61	0.000	-12.69811	-6.889775

In this case, the estimated coefficient for the intercept is the log odds of a student with a math score of zero being in an honors class. In other words, the odds of being in an honors class when the math score is zero is $\exp(-9.793942) = .00005579$. These odds are very low, but if we look at the distribution of the variable **math**, we will see that no one in the sample has math score lower than 30. In fact, all the test scores in the data set were standardized around mean of 50 and standard deviation of 10. So the intercept in this model corresponds to the log odds of being in an honors class when **math** is at the *hypothetical* value of zero.

How do we interpret the coefficient for math? The coefficient and intercept estimates give us the following equation:

$$\log(p/(1-p)) = \logit(p) = -9.793942 + .1563404 \cdot \text{math}$$

Let's fix **math** at some value. We will use 54. Then the conditional logit of being in an honors class when the math score is held at 54 is

$$\log(p/(1-p))(\text{math}=54) = -9.793942 + .1563404 \cdot 54.$$

We can examine the effect of a one-unit increase in math score. When the math score is held at 55, the conditional logit of being in an honors class is

$$\log(p/(1-p))(\text{math}=55) = -9.793942 + .1563404 \cdot 55.$$

Taking the difference of the two equations, we have the following:

$$\log(p/(1-p))(\text{math}=55) - \log(p/(1-p))(\text{math}=54) = .1563404.$$

We can say now that the coefficient for **math** is the difference in the log odds. In other words, for a one-unit increase in the math score, the expected change in log odds is .1563404.

Can we translate this change in log odds to the change in odds? Indeed, we can. Recall that logarithm converts multiplication and division to addition and subtraction. Its inverse, the exponentiation converts addition and subtraction back to multiplication and division. If we exponentiate both sides of our last equation, we have the following:

$$\exp[\log(p/(1-p))(\text{math}=55) - \log(p/(1-p))(\text{math}=54)] = \exp(\log(p/(1-p))(\text{math}=55)) / \exp(\log(p/(1-p))(\text{math}=54)) = \text{odds}(\text{math}=55) / \text{odds}(\text{math}=54) = \exp(.1563404) = 1.1692241.$$

So we can say for a one-unit increase in math score, we expect to see about 17% increase in the odds of being in an honors class. This 17% of increase does not depend on the value that math is held at.

Logistic regression with multiple predictor variables and no interaction terms

In general, we can have multiple predictor variables in a logistic regression model.

$$\text{logit}(p) = \log(p/(1-p)) = \beta_0 + \beta_1 \cdot x_1 + \dots + \beta_k \cdot x_k$$

Applying such a model to our example dataset, each estimated coefficient is the expected change in the log odds of being in an honors class for a unit increase in the corresponding predictor variable holding the other predictor variables constant at certain value. Each exponentiated coefficient is the ratio of two odds, or the change in odds in the multiplicative scale for a unit increase in the corresponding predictor variable holding other variables at certain value. Here is an example.

$$\text{logit}(p) = \log(p/(1-p)) = \beta_0 + \beta_1 \cdot \text{math} + \beta_2 \cdot \text{female} + \beta_3 \cdot \text{read}$$

Logistic regression				Number of obs	=	200
				LR chi2(3)	=	66.54
				Prob > chi2	=	0.0000
				Pseudo R2	=	0.2988
Log likelihood = -78.084776						

hon	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
math	.1229589	.0312756	3.93	0.000	.0616599	.1842578
female	.979948	.4216264	2.32	0.020	.1535755	1.80632
read	.0590632	.0265528	2.22	0.026	.0070207	.1111058
intercept	-11.77025	1.710679	-6.88	0.000	-15.12311	-8.417376

This fitted model says that, holding **math** and **reading** at a fixed value, the odds of getting into an honors class for females (**female** = 1) over the odds of getting into an honors class for males (**female** = 0) is $\exp(.979948) = 2.66$. In terms of percent change, we can say that the odds for females are 166% higher than the odds for males. The coefficient for **math** says that, holding **female** and **reading** at a fixed value, we will see 13% increase in the odds of getting into an honors class for a one-unit increase in math score since $\exp(.1229589) = 1.13$.

Logistic regression with an interaction term of two predictor variables

FAQ: How do I interpret odds ratios in logistic

9/26/16, 9:12 AM

In all the previous examples, we have said that the regression coefficient of a variable corresponds to the change in log odds and its exponentiated form corresponds to the odds ratio. This is only true when our model does not have any interaction terms. When a model has interaction term(s) of two predictor variables, it attempts to describe how the effect of a predictor variable depends on the level/value of another predictor variable. The interpretation of the regression coefficients become more involved.

Let's take a simple example.

$$\text{logit}(p) = \log(p/(1-p)) = \beta_0 + \beta_1 \text{female} + \beta_2 \text{math} + \beta_3 \text{female} \text{math}$$

Logistic regression

Number of obs = 200
LR chi2(3) = 62.94
Prob > chi2 = 0.0000
Pseudo R2 = 0.2826

Log likelihood = -79.883301

hon	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]
female	-2.899863	3.094186	-0.94	0.349	-8.964357 3.164631
math	.1293781	.0358834	3.61	0.000	.0590479 .1997082
femalemath	.0669951	.05346	1.25	0.210	-.0377846 .1717749
intercept	-8.745841	2.12913	-4.11	0.000	-12.91886 -4.572823

In the presence of interaction term of **female** by **math**, we can no longer talk about the effect of **female**, holding all other variables at certain value, since it does not make sense to fix **math** and **femalemath** at certain value and still allow female change from 0 to 1!

In this simple example where we examine the interaction of a binary variable and a continuous variable, we can think that we actually have two equations: one for males and one for females. For males (**female**=0), the equation is simply

$$\text{logit}(p) = \log(p/(1-p)) = \beta_0 + \beta_2 \text{math}.$$

For females, the equation is

$$\text{logit}(p) = \log(p/(1-p)) = (\beta_0 + \beta_1) + (\beta_2 + \beta_3) \text{math}.$$


Now we can map the logistic regression output to these two equations. So we can say that the coefficient for **math** is the effect of **math** when **female** = 0. More explicitly, we can say that for male students, a one-unit increase in **math** score yields a change in log odds of (.13 + .067) = 0.197. In terms of odds ratios, we can say that for male students, the odds ratio is $\exp(.13) = 1.14$ for a one-unit increase in **math** score and the odds ratio for female students is $\exp(.197) = 1.22$ for a one-unit increase in **math** score. The ratio of these two odds ratios (female over male) turns out to be the exponentiated coefficient for the interaction term of **female** by **math**: $1.22/1.14 = \exp(.067) = 1.07$.


[How to cite this page](#)


[Report an error on this page or leave a comment](#)

The content of this web site should not be construed as an endorsement of any particular web site, book, or software product by the University of California.

IDRE RESEARCH TECHNOLOGY GROUP

 High Performance Computing

 Statistical Computing

 GIS and Visualization

High Performance Computing

GIS

Statistical Computing

Hoffman2 Cluster

Mapshare

Classes

Hoffman2 Account Application

Visualization

Conferences

Hoffman2 Usage Statistics

3D Modeling

Reading Materials

UC Grid Portal

Technology Sandbox

IDRE Listserv

UCLA Grid Portal

Tech Sandbox Access

IDRE Resources

Shared Cluster & Storage

Data Centers

Social Sciences Data Archive

About IDRE

UCLA

ABOUT CONTACT NEWS EVENTS OUR EXPERTS

© 2016 UC Regents Terms of Use & Privacy Policy

