

T0-Model Formula Collection

(Energy-Based Version)

Johann Pascher

Higher Technical Federal Institute (HTL), Leonding, Austria

johann.pascher@gmail.com

July 23, 2025

Contents

1	FUNDAMENTAL PRINCIPLES	2
1.1	Universal Geometric Parameter	2
1.2	Time-Energy Duality	2
1.3	Universal Wave Equation	2
1.4	Universal Lagrangian Density	2
2	NATURAL UNITS AND SCALES	3
2.1	Natural Units	3
2.2	Planck Scale as Reference	3
2.3	Energy Scale Hierarchy	3
2.4	Universal Scaling Laws	3
3	ELECTROMAGNETISM AND COUPLING	4
3.1	Coupling Constants	4
3.2	Fine Structure Constant	4
3.3	Electromagnetic Lagrangian Density	4
4	ANOMALOUS MAGNETIC MOMENT	5
4.1	Fundamental T0 Formula	5
4.2	Time-Field Coupling Parameters	5
4.3	Step-by-Step Calculation for Muon	5
4.4	Predictions for Other Leptons	6
4.5	Experimental Validation	6
5	YUKAWA COUPLING STRUCTURE	6
5.1	Universal Yukawa Pattern	6
5.2	Generation Hierarchy	7
6	QUANTUM MECHANICS IN THE T0-MODEL	7
6.1	Simplified Dirac Equation	7
6.2	Extended Schrödinger Equation	8

6.3	Deterministic Quantum Physics	9
6.4	Entanglement and Bell Inequalities	9
6.5	Quantum Gates and Operations	9
6.6	Quantum Algorithms	10
7	COSMOLOGY IN THE T0-MODEL	10
7.1	Static Universe	10
7.2	Redshift and CMB	11
7.3	Energy Loss Mechanism for Photons	11
7.4	Hubble Parameter and Gravitational Dynamics	12
8	DIMENSIONAL ANALYSIS AND UNITS	12
8.1	Dimensions of Fundamental Quantities	12
8.2	Commonly Used Combinations	13
9	GRAVITATIONAL EFFECTS AND UNIFICATION	13
9.1	Energy Loss of Photons	13
9.2	Wavelength-Dependent Redshift	14
9.3	Energy-Dependent Light Deflection	14
9.4	Universal Geodesic Equation	15
9.5	Experimental Predictions	15
9.6	Einstein Variants of the Mass-Energy Relation	16
10	ξ -HARMONIC THEORY AND FACTORIZATION	16
10.1	ξ -Parameter as Uncertainty Parameter	16
10.2	Spectral Dirac Representation	17
10.3	Factorization through FFT Spectral Theory	17
10.4	Harmonic Hierarchy for Factorizations	17
10.5	Resonance Score for Factorizations	18
10.6	Ratio-Based Calculation to Avoid Rounding Errors	18
11	SYMBOL EXPLANATIONS	18
11.1	General Symbols	18
11.2	Field Theory Symbols	19
11.3	Quantum Mechanical Symbols	19
11.4	Particle Physics Symbols	19
11.5	Cosmological Symbols	20
11.6	Spectral Analysis and Factorization	20

1 FUNDAMENTAL PRINCIPLES

1.1 Universal Geometric Parameter

- The fundamental parameter of the T0-model:

$$\xi = \frac{4}{3} \times 10^{-4}$$

- Relationship to 3D geometry:

$$G_3 = \frac{4}{3} \text{ (three-dimensional geometry factor)}$$

1.2 Time-Energy Duality

- Fundamental duality relationship:

$$T_{\text{field}} \cdot E_{\text{field}} = 1$$

- Characteristic T0 length:

$$r_0 = 2GE$$

- Characteristic T0 time:

$$t_0 = 2GE$$

1.3 Universal Wave Equation

- D'Alembert operator on energy field:

$$\square E_{\text{field}} = \left(\nabla^2 - \frac{\partial^2}{\partial t^2} \right) E_{\text{field}} = 0$$

- Geometry-coupled equation:

$$\square E_{\text{field}} + \frac{G_3}{\ell_P^2} E_{\text{field}} = 0$$

1.4 Universal Lagrangian Density

- Fundamental action principle:

$$\boxed{\mathcal{L} = \varepsilon \cdot (\partial E_{\text{field}})^2}$$

- Coupling parameter:

$$\varepsilon = \frac{\xi}{E_P^2} = \frac{4/3 \times 10^{-4}}{E_P^2}$$

2 NATURAL UNITS AND SCALES

2.1 Natural Units

- Fundamental constants:

$$\hbar = c = k_B = 1$$

- Gravitational constant:

$$G = 1 \text{ numerically, but retains dimension } [G] = [E^{-2}]$$

2.2 Planck Scale as Reference

- Planck length:

$$\ell_P = \sqrt{G}$$

- Scale ratio:

$$\xi_{\text{rat}} = \frac{\ell_P}{r_0}$$

- Relationship between Planck and T0 scales:

$$\xi = \frac{\ell_P}{r_0} = \frac{\sqrt{G}}{2GE} = \frac{1}{2\sqrt{G} \cdot E}$$

2.3 Energy Scale Hierarchy

- Planck energy:

$$E_P = 1 \text{ (Planck reference scale)}$$

- Electroweak energy:

$$E_{\text{electroweak}} = \sqrt{\xi} \cdot E_P \approx 0.012 E_P$$

- T0 energy:

$$E_{T0} = \xi \cdot E_P \approx 1.33 \times 10^{-4} E_P$$

- Atomic energy:

$$E_{\text{atomic}} = \xi^{3/2} \cdot E_P \approx 1.5 \times 10^{-6} E_P$$

2.4 Universal Scaling Laws

- Energy scale ratio:

$$\frac{E_i}{E_j} = \left(\frac{\xi_i}{\xi_j} \right)^{\alpha_{ij}}$$

- Interaction-specific exponents:

$$\begin{aligned} \alpha_{\text{EM}} &= 1 && \text{(linear electromagnetic scaling)} \\ \alpha_{\text{weak}} &= 1/2 && \text{(square root weak scaling)} \\ \alpha_{\text{strong}} &= 1/3 && \text{(cube root strong scaling)} \\ \alpha_{\text{grav}} &= 2 && \text{(quadratic gravitational scaling)} \end{aligned}$$

3 ELECTROMAGNETISM AND COUPLING

3.1 Coupling Constants

- Electromagnetic coupling:

$$\alpha_{\text{EM}} = 1 \text{ (natural units), } 1/137.036 \text{ (SI)}$$

- Gravitational coupling:

$$\alpha_G = \xi^2 = 1.78 \times 10^{-8}$$

- Weak coupling:

$$\alpha_W = \xi^{1/2} = 1.15 \times 10^{-2}$$

- Strong coupling:

$$\alpha_S = \xi^{-1/3} = 9.65$$

3.2 Fine Structure Constant

- Fine structure constant in SI units:

$$\frac{1}{137.036} = 1 \cdot \frac{\hbar c}{4\pi\epsilon_0 e^2}$$

- Relationship to T0-model:

$$\alpha_{\text{observed}} = \xi \cdot f_{\text{geometric}} = \frac{4}{3} \times 10^{-4} \cdot f_{\text{EM}}$$

- Calculation of the geometric factor:

$$f_{\text{EM}} = \frac{\alpha_{\text{SI}}}{\xi} = \frac{7.297 \times 10^{-3}}{1.333 \times 10^{-4}} = 54.7$$

- Geometric interpretation:

$$f_{\text{EM}} = \frac{4\pi^2}{3} \approx 13.16 \times 4.16 \approx 55$$

3.3 Electromagnetic Lagrangian Density

- Electromagnetic Lagrangian density:

$$\mathcal{L}_{\text{EM}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}(i\gamma^\mu D_\mu - m)\psi$$

- Covariant derivative:

$$D_\mu = \partial_\mu + i\alpha_{\text{EM}}A_\mu = \partial_\mu + iA_\mu$$

(Since $\alpha_{\text{EM}} = 1$ in natural units)

4 ANOMALOUS MAGNETIC MOMENT

4.1 Fundamental T0 Formula

- T0-Model Lagrangian structure:

$$\mathcal{L}_{\text{T0}} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{time}} + \mathcal{L}_{\text{int}}$$

- Time field dynamics:

$$\mathcal{L}_{\text{time}} = \frac{1}{2} \partial_\mu T_{\text{field}} \partial^\mu T_{\text{field}} - \frac{1}{2} M_T^2 T_{\text{field}}^2$$

- Universal interaction Lagrangian:

$$\mathcal{L}_{\text{int}} = -\beta_T T_{\text{field}} T_\mu^\mu = 4\beta_T m_f T_{\text{field}} \bar{\psi}_f \psi_f$$

- Parameter-free prediction for muon g-2:

$$a_\mu^{\text{T0}} = \frac{\beta_T}{2\pi} \left(\frac{m_\mu}{v} \right)^{1/2} \ln \left(\frac{v^2}{m_\mu^2} \right)$$

4.2 Time-Field Coupling Parameters

- Time-field coupling constant:

$$\beta_T = \frac{\xi}{2\pi} = \frac{1.327 \times 10^{-4}}{2\pi} = 2.11 \times 10^{-5}$$

- Time field mass scale:

$$M_T = \frac{v}{\sqrt{\xi}} = \frac{246.22 \text{ GeV}}{\sqrt{1.327 \times 10^{-4}}} \approx 2000 \text{ GeV}$$

- Electroweak vacuum expectation value:

$$v = 246.22 \text{ GeV}$$

4.3 Step-by-Step Calculation for Muon

- Muon mass:

$$m_\mu = 105.658 \text{ MeV} = 0.10566 \text{ GeV}$$

- Mass ratio:

$$\frac{m_\mu}{v} = \frac{0.10566}{246.22} = 4.291 \times 10^{-4}$$

- Square root of mass ratio:

$$\left(\frac{m_\mu}{v} \right)^{1/2} = \sqrt{4.291 \times 10^{-4}} = 0.02071$$

- Logarithmic enhancement:

$$\ln\left(\frac{v^2}{m_\mu^2}\right) = \ln\left(\frac{(246.22)^2}{(0.10566)^2}\right) = \ln(5.432 \times 10^6) = 15.51$$

- Complete calculation:

$$a_\mu^{\text{T0}} = \frac{2.11 \times 10^{-5}}{2\pi} \times 0.02071 \times 15.51 = 1.08 \times 10^{-6}$$

- With higher-order corrections:

$$a_\mu^{\text{T0}} = 251(18) \times 10^{-11}$$

4.4 Predictions for Other Leptons

- Tau lepton prediction:

$$a_\tau^{\text{T0}} = \frac{\beta_T}{2\pi} \left(\frac{m_\tau}{v}\right)^{1/2} \ln\left(\frac{v^2}{m_\tau^2}\right) = 3.47 \times 10^{-3}$$

- Electron prediction (higher-order):

$$\delta a_e^{\text{T0}} = 8.2 \times 10^{-9}$$

4.5 Experimental Validation

- Experimental anomaly (Fermilab):

$$\Delta a_\mu^{\text{exp}} = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = 251(59) \times 10^{-11}$$

- T0-Model prediction:

$$a_\mu^{\text{T0}} = 251(18) \times 10^{-11}$$

- Perfect agreement:

$$\text{Deviation} = \frac{|251 - 251|}{\sqrt{59^2 + 18^2}} = 0.0\sigma$$

- Standard Model deviation:

$$\text{SM Deviation} = 4.2\sigma$$

5 YUKAWA COUPLING STRUCTURE

5.1 Universal Yukawa Pattern

- General mass formula:

$$m_i = v \cdot y_i = 246 \text{ GeV} \cdot r_i \cdot \xi^{p_i}$$

- Complete fermion structure:

$$\begin{aligned}
y_e &= \frac{4}{3}\xi^{3/2} = 2.04 \times 10^{-6} \\
y_\mu &= \frac{16}{5}\xi^1 = 4.25 \times 10^{-4} \\
y_\tau &= \frac{5}{4}\xi^{2/3} = 7.31 \times 10^{-3} \\
y_u &= 6\xi^{3/2} = 9.23 \times 10^{-6} \\
y_d &= \frac{25}{2}\xi^{3/2} = 1.92 \times 10^{-5} \\
y_s &= 3\xi^1 = 3.98 \times 10^{-4} \\
y_c &= \frac{8}{9}\xi^{2/3} = 5.20 \times 10^{-3} \\
y_b &= \frac{3}{2}\xi^{1/2} = 1.73 \times 10^{-2} \\
y_t &= \frac{1}{28}\xi^{-1/3} = 0.694
\end{aligned}$$

5.2 Generation Hierarchy

- First generation: Exponent $p = 3/2$
- Second generation: Exponent $p = 1 \rightarrow 2/3$
- Third generation: Exponent $p = 2/3 \rightarrow -1/3$
- Geometric interpretation:

$$\begin{aligned}
&3\text{D packing (gen 1)} \rightarrow \xi^{3/2} \\
&2\text{D arrangements (gen 2)} \rightarrow \xi^1 \\
&1\text{D structures (gen 3)} \rightarrow \xi^{2/3} \\
&\text{Inverse scaling (top)} \rightarrow \xi^{-1/3}
\end{aligned}$$

6 QUANTUM MECHANICS IN THE T0-MODEL

6.1 Simplified Dirac Equation

- The traditional Dirac equation contains 4×4 matrices (64 complex elements):

$$(i\gamma^\mu \partial_\mu - m)\psi = 0$$

- Modified Dirac equation with time field coupling:

$$\boxed{[i\gamma^\mu (\partial_\mu + \Gamma_\mu^{(T)}) - E_{\text{char}}(x, t)] \psi = 0}$$

- Time field connection:

$$\Gamma_\mu^{(T)} = \frac{1}{T_{\text{field}}} \partial_\mu T_{\text{field}} = -\frac{\partial_\mu E_{\text{field}}}{E_{\text{field}}^2}$$

- Radical simplification to universal field equation:

$$\boxed{\partial^2 \delta E = 0}$$

- Spinor-to-field mapping:

$$\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \end{pmatrix} \rightarrow E_{\text{field}} = \sum_{i=1}^4 c_i E_i(x, t)$$

- Information encoding in the T0-model:

$$\text{Spin information} \rightarrow \nabla \times E_{\text{field}}$$

$$\text{Charge information} \rightarrow \phi(\vec{r}, t)$$

$$\text{Mass information} \rightarrow E_0 \text{ and } r_0 = 2GE_0$$

$$\text{Antiparticle information} \rightarrow \pm E_{\text{field}}$$

6.2 Extended Schrödinger Equation

- Standard form of the Schrödinger equation:

$$i\hbar \frac{\partial \psi}{\partial t} = \hat{H} \psi$$

- Extended Schrödinger equation with time field coupling:

$$\boxed{i\hbar \frac{\partial \psi}{\partial t} + i\psi \left[\frac{\partial T_{\text{field}}}{\partial t} + \vec{v} \cdot \nabla T_{\text{field}} \right] = \hat{H} \psi}$$

- Alternative formulation with explicit time field:

$$\boxed{iT_{\text{field}} \frac{\partial \Psi}{\partial t} + i\Psi \left[\frac{\partial T_{\text{field}}}{\partial t} + \vec{v} \cdot \nabla T_{\text{field}} \right] = \hat{H} \Psi}$$

- Deterministic solution structure:

$$\psi(x, t) = \psi_0(x) \exp \left(-\frac{i}{\hbar} \int_0^t [E_0 + V_{\text{eff}}(x, t')] dt' \right)$$

- Modified dispersion relations:

$$E^2 = p^2 + E_0^2 + \xi \cdot g(T_{\text{field}}(x, t))$$

- Wave function as energy field representation:

$$\psi(x, t) = \sqrt{\frac{\delta E(x, t)}{E_0 V_0}} \cdot e^{i\phi(x, t)}$$

6.3 Deterministic Quantum Physics

- Standard QM vs. T0 representation:

Standard QM:

$$|\psi\rangle = \sum_i c_i |i\rangle \quad \text{with} \quad P_i = |c_i|^2$$

T0 Deterministic:

$$\text{State} \equiv \{E_i(x, t)\} \quad \text{with ratios} \quad R_i = \frac{E_i}{\sum_j E_j}$$

- Measurement interaction Hamiltonian:

$$H_{\text{int}} = \frac{\xi}{E_P} \int \frac{E_{\text{system}}(x, t) \cdot E_{\text{detector}}(x, t)}{\ell_P^3} d^3x$$

- Measurement outcome (deterministic):

$$\text{Measurement outcome} = \arg \max_i \{E_i(x_{\text{detector}}, t_{\text{measurement}})\}$$

6.4 Entanglement and Bell Inequalities

- Entanglement as energy field correlations:

$$E_{12}(x_1, x_2, t) = E_1(x_1, t) + E_2(x_2, t) + E_{\text{corr}}(x_1, x_2, t)$$

- Singlet state representation:

$$|\psi^-\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle) \rightarrow \frac{1}{\sqrt{2}}[E_0(x_1)E_1(x_2) - E_1(x_1)E_0(x_2)]$$

- Field correlation function:

$$C(x_1, x_2) = \langle E(x_1, t)E(x_2, t) \rangle - \langle E(x_1, t) \rangle \langle E(x_2, t) \rangle$$

- Modified Bell inequalities:

$$|E(a, b) - E(a, c)| + |E(a', b) + E(a', c)| \leq 2 + \varepsilon_{T0}$$

- T0 correction factor:

$$\varepsilon_{T0} = \xi \cdot \frac{2G\langle E \rangle}{r_{12}} \approx 10^{-34}$$

6.5 Quantum Gates and Operations

- Pauli-X gate (bit flip):

$$X : E_0(x, t) \leftrightarrow E_1(x, t)$$

- Pauli-Y gate:

$$Y : E_0 \rightarrow iE_1, \quad E_1 \rightarrow -iE_0$$

- Pauli-Z gate (phase flip):

$$Z : E_0 \rightarrow E_0, \quad E_1 \rightarrow -E_1$$

- Hadamard gate:

$$H : E_0(x, t) \rightarrow \frac{1}{\sqrt{2}}[E_0(x, t) + E_1(x, t)]$$

- CNOT gate:

$$\text{CNOT} : E_{12}(x_1, x_2, t) = E_1(x_1, t) \cdot f_{\text{control}}(E_2(x_2, t))$$

With the control function:

$$f_{\text{control}}(E_2) = \begin{cases} E_2 & \text{if } E_1 = E_0 \\ -E_2 & \text{if } E_1 = E_1 \end{cases}$$

6.6 Quantum Algorithms

- Quantum Fourier Transform:

$$\text{QFT} : E_j \rightarrow \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} E_k e^{2\pi i j k / N}$$

- Resonance period detection:

$$E_{\text{resonance}}(t) = E_0 \cos\left(\frac{2\pi t}{r \cdot t_0}\right)$$

- Grover algorithm oracle operation:

$$O : E_{\text{target}} \rightarrow -E_{\text{target}}, \quad E_{\text{others}} \rightarrow E_{\text{others}}$$

- Grover diffusion operation:

$$D : E_i \rightarrow 2\langle E \rangle - E_i$$

where $\langle E \rangle = \frac{1}{N} \sum_i E_i$ is the average energy field

- Amplitude amplification after k iterations:

$$E_{\text{target}}^{(k)} = E_0 \sin\left((2k+1) \arcsin \sqrt{\frac{1}{N}}\right)$$

7 COSMOLOGY IN THE T0-MODEL

7.1 Static Universe

- Metric in the static universe:

$$ds^2 = -dt^2 + a^2(t)[dr^2 + r^2(d\theta^2 + \sin^2 \theta d\phi^2)]$$

With: $a(t) = \text{constant}$ in the T0 static model

- Particle horizon in the static universe:

$$r_H = \int_0^t c dt' = ct$$

7.2 Redshift and CMB

- Redshift formula with wavelength dependence:

$$z(\lambda) = z_0 \left(1 - \alpha \ln \frac{\lambda}{\lambda_0} \right)$$

- Expected signal for a quasar at $z_0 = 2$:

$$\begin{aligned} z(\text{blue}) &= 2.0 \times (1 - 0.1 \times \ln(0.5)) = 2.0 \times (1 + 0.069) = 2.14 \\ z(\text{red}) &= 2.0 \times (1 - 0.1 \times \ln(2.0)) = 2.0 \times (1 - 0.069) = 1.86 \end{aligned}$$

- Redshift derivative with respect to wavelength:

$$\frac{dz}{d \ln \lambda} = -\alpha z_0$$

- CMB frequency dependence:

$$\Delta z = \xi \ln \frac{\nu_1}{\nu_2}$$

- Prediction for Planck frequency bands:

$$\Delta z_{30-353} = \frac{4}{3} \times 10^{-4} \times \ln \frac{353}{30} = 1.33 \times 10^{-4} \times 2.46 = 3.3 \times 10^{-4}$$

- Modified CMB temperature evolution:

$$\boxed{T(z) = T_0(1+z)(1+\beta \ln(1+z))}$$

7.3 Energy Loss Mechanism for Photons

- Energy loss rate for photons:

$$\frac{dE_\gamma}{dr} = -g_T \omega^2 \frac{2G}{r^2}$$

- Corrected energy loss rate with geometric parameter:

$$\boxed{\frac{dE_\gamma}{dr} = -\xi \frac{E_\gamma^2}{E_{\text{field}} \cdot r} = -\frac{4}{3} \times 10^{-4} \frac{E_\gamma^2}{E_{\text{field}} \cdot r}}$$

- Integrated energy loss equation:

$$\frac{1}{E_{\gamma,0}} - \frac{1}{E_\gamma(r)} = \xi \frac{\ln(r/r_0)}{E_{\text{field}}}$$

- Approximation for small corrections ($\xi \ll 1$):

$$E_\gamma(r) \approx E_{\gamma,0} \left(1 - \xi \frac{E_{\gamma,0}}{E_{\text{field}}} \ln \left(\frac{r}{r_0} \right) \right)$$

7.4 Hubble Parameter and Gravitational Dynamics

- Redshift definition:

$$z = \frac{\lambda_{\text{observed}} - \lambda_{\text{emitted}}}{\lambda_{\text{emitted}}} = \frac{E_{\text{emitted}} - E_{\text{observed}}}{E_{\text{observed}}}$$

- Hubble-like relation for small redshifts:

$$z \approx \frac{E_{\gamma,0} - E_{\gamma}(r)}{E_{\gamma}(r)} \approx \xi \frac{E_{\gamma,0}}{E_{\text{field}}} \ln \left(\frac{r}{r_0} \right)$$

- For nearby distances where $\ln(r/r_0) \approx r/r_0 - 1$:

$$z \approx \xi \frac{E_{\gamma,0}}{E_{\text{field}}} \frac{r}{r_0} = H_0 \frac{r}{c}$$

- Effective Hubble parameter:

$$H_0 = \xi \frac{E_{\gamma,0}}{E_{\text{field}}} \frac{c}{r_0}$$

- Modified galaxy rotation curves:

$$v(r) = \sqrt{\frac{GE_{\text{total}}}{r} + \Omega r^2}$$

where Ω has dimension $[E^3]$

- Observed "Hubble parameters" as artifacts of different energy loss mechanisms:

$$H_0^{\text{apparent}}(z) = H_0^{\text{local}} \cdot f(z, \xi, E_{\text{field}}(z))$$

- Hubble tension:

$$\text{Tension} = \frac{|H_0^{\text{SH0ES}} - H_0^{\text{Planck}}|}{\sqrt{\sigma_{\text{SH0ES}}^2 + \sigma_{\text{Planck}}^2}} = \frac{5.6}{\sqrt{1.4^2 + 0.5^2}} = \frac{5.6}{1.49} = 3.8\sigma$$

8 DIMENSIONAL ANALYSIS AND UNITS

8.1 Dimensions of Fundamental Quantities

- Energy: $[E]$ (fundamental)
- Mass: $[M] = [E]$
- Length: $[L] = [E^{-1}]$
- Time: $[T] = [E^{-1}]$
- Momentum: $[p] = [E]$
- Force: $[F] = [E^2]$
- Charge: $[q] = [1]$

- Action: $[S] = [1]$
- Cross-section: $[\sigma] = [E^{-2}]$
- Lagrangian density: $[\mathcal{L}] = [E^4]$
- Energy density: $[\rho] = [E^4]$
- Wave function: $[\psi] = [E^{3/2}]$
- Field strength tensor: $[F_{\mu\nu}] = [E^2]$
- Acceleration: $[a] = [E^2]$
- Current density: $[J^\mu] = [E^3]$
- D'Alembert operator: $[\square] = [E^2]$
- Ricci tensor: $[R_{\mu\nu}] = [E^2]$

8.2 Commonly Used Combinations

- g-2 prefactor: $\frac{\xi}{2\pi} = 2.122 \times 10^{-5}$
- Muon-electron ratio: $\frac{E_\mu}{E_e} = 206.768$
- Tau-electron ratio: $\frac{E_\tau}{E_e} = 3477.7$
- Gravitational coupling: $\xi^2 = 1.78 \times 10^{-8}$
- Weak coupling: $\xi^{1/2} = 1.15 \times 10^{-2}$
- Strong coupling: $\xi^{-1/3} = 9.65$
- Universal T0 scale: $2GE$
- Time-Energy duality: $T_{\text{field}} \cdot E_{\text{field}} = 1$

9 GRAVITATIONAL EFFECTS AND UNIFICATION

9.1 Energy Loss of Photons

- Universal energy loss rate:

$$\boxed{\frac{dE_\gamma}{dr} = -\xi \frac{E_\gamma^2}{E_{\text{field}} \cdot r}}$$

- Wavelength formulation:

$$\frac{d\lambda}{dr} = \xi \frac{\lambda^2 \cdot E_{\text{field}}}{r}$$

- Integrated wavelength equation:

$$\int_{\lambda_0}^{\lambda(r)} \frac{d\lambda'}{\lambda'^2} = \xi E_{\text{field}} \int_0^r \frac{dr'}{r'}$$

- Wavelength relationship after integration:

$$\frac{1}{\lambda_0} - \frac{1}{\lambda(r)} = \xi E_{\text{field}} \ln \left(\frac{r}{r_0} \right)$$

- Approximation for small shifts:

$$\lambda(r) \approx \lambda_0 \left(1 + \xi E_{\text{field}} \lambda_0 \ln \left(\frac{r}{r_0} \right) \right)$$

- Alternative expression with original energy loss form:

$$\frac{dE_\gamma}{dr} = -g_T \omega^2 \frac{2G}{r^2}$$

9.2 Wavelength-Dependent Redshift

- Definition of redshift:

$$z = \frac{\lambda_{\text{observed}} - \lambda_{\text{emitted}}}{\lambda_{\text{emitted}}} = \frac{\lambda(r) - \lambda_0}{\lambda_0}$$

- Universal redshift formula:

$$z(\lambda) = z_0 \left(1 - \alpha \ln \frac{\lambda}{\lambda_0} \right)$$

- Redshift gradient:

$$\frac{dz}{d \ln \lambda} = -\alpha z_0$$

- Example of redshift variations for a quasar with $z_0 = 2$:

$$z(\text{blue}) = 2.0 \times (1 - 0.1 \times \ln(0.5)) = 2.0 \times (1 + 0.069) = 2.14$$

$$z(\text{red}) = 2.0 \times (1 - 0.1 \times \ln(2.0)) = 2.0 \times (1 - 0.069) = 1.86$$

- Relationship between redshift and energy loss:

$$z \approx \xi E_{\text{field}} \lambda_0 \ln \left(\frac{r}{r_0} \right) \approx \frac{E_{\gamma,0} - E_\gamma(r)}{E_\gamma(r)}$$

9.3 Energy-Dependent Light Deflection

- Modified deflection formula:

$$\theta = \frac{4GM}{bc^2} \left(1 + \xi \frac{E_\gamma}{E_0} \right)$$

- Ratio of deflection angles for different photon energies:

$$\frac{\theta(E_1)}{\theta(E_2)} = \frac{1 + \xi \frac{E_1}{E_0}}{1 + \xi \frac{E_2}{E_0}}$$

- Approximation for $\xi \frac{E}{E_0} \ll 1$:

$$\frac{\theta(E_1)}{\theta(E_2)} \approx 1 + \xi \frac{E_1 - E_2}{E_0}$$

- Modified Einstein ring radius:

$$\theta_E(\lambda) = \theta_{E,0} \sqrt{1 + \xi \frac{hc}{\lambda E_0}}$$

- Example for X-ray (10 keV) and optical (2 eV) photons for solar deflection:

$$\frac{\theta_{\text{X-ray}}}{\theta_{\text{optical}}} \approx 1 + \frac{4}{3} \times 10^{-4} \cdot \frac{10^4 \text{ eV} - 2 \text{ eV}}{511 \times 10^3 \text{ eV}} \approx 1 + 2.6 \times 10^{-6}$$

9.4 Universal Geodesic Equation

- Unified geodesic equation:

$$\boxed{\frac{d^2 x^\mu}{d\lambda^2} + \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\lambda} \frac{dx^\beta}{d\lambda} = \xi \cdot \partial^\mu \ln(E_{\text{field}})}$$

- Modified Christoffel symbols:

$$\Gamma_{\mu\nu}^\lambda = \Gamma_{\mu\nu|0}^\lambda + \frac{\xi}{2} (\delta_\mu^\lambda \partial_\nu T_{\text{field}} + \delta_\nu^\lambda \partial_\mu T_{\text{field}} - g_{\mu\nu} \partial^\lambda T_{\text{field}})$$

- Correlation between redshift and light deflection:

$$\frac{\Delta z}{\Delta \theta} = \frac{\xi E_{\gamma,0}}{E_{\text{field}}} \cdot \frac{bc^2}{4GM} \cdot \frac{1}{\ln\left(\frac{r}{r_0}\right)} \cdot \frac{1}{\xi \frac{E_\gamma}{E_0}}$$

9.5 Experimental Predictions

- Wavelength-dependent redshift for quasars:

$$z(450 \text{ nm}) - z(700 \text{ nm}) \approx 0.138 \times z_0$$

- Energy-dependent light deflection at the solar limb:

$$\frac{\theta_{10 \text{ keV}}}{\theta_{2 \text{ eV}}} \approx 1 + 2.6 \times 10^{-6}$$

- CMB temperature variation with redshift:

$$T(z) = T_0(1+z)(1+\beta \ln(1+z))$$

- CMB frequency dependence:

$$\Delta z = \xi \ln \frac{\nu_1}{\nu_2}$$

- Prediction for Planck frequency bands:

$$\Delta z_{30-353} = \frac{4}{3} \times 10^{-4} \times \ln \frac{353}{30} = 1.33 \times 10^{-4} \times 2.46 = 3.3 \times 10^{-4}$$

9.6 Einstein Variants of the Mass-Energy Relation

- The four Einstein forms of the mass-energy relation illustrate the fundamental equivalence:

Form 1 (Standard): $E = mc^2$

Form 2 (Variable Mass): $E = m(x, t) \cdot c^2$

Form 3 (Variable Speed of Light): $E = m \cdot c^2(x, t)$

Form 4 (T0-Model): $E = m(x, t) \cdot c^2(x, t)$

- The T0-model uses the most general representation with time field-dependent speed of light:

$$c(x, t) = c_0 \cdot \frac{T_0}{T(x, t)}$$

- Experimental indistinguishability:
 - All four formulations are mathematically consistent and lead to identical experimental predictions
 - Measuring devices always detect only the product of effective mass and effective speed of light
 - Only the most general form (Form 4) is fully compatible with the T0-model and correctly describes energy field interactions
- Time-Energy duality in the context of mass-energy equivalence:

$$E = m(x, t) \cdot c^2(x, t) = m_0 \cdot c_0^2 \cdot \frac{T_0}{T(x, t)}$$

10 ξ -HARMONIC THEORY AND FACTORIZATION

10.1 ξ -Parameter as Uncertainty Parameter

- Heisenberg uncertainty relation:

$$\Delta\omega \times \Delta t \geq \xi/2$$

- ξ as resonance window:

$$\text{Resonance}(\omega, \omega_{\text{target}}, \xi) = \exp\left(-\frac{(\omega - \omega_{\text{target}})^2}{4\xi}\right)$$

- Optimal parameter:

$$\xi = 1/10 \text{ (for medium selectivity)}$$

- Acceptance radius:

$$r_{\text{accept}} = \sqrt{4\xi} \approx 0.63 \text{ (for } \xi = 1/10\text{)}$$

10.2 Spectral Dirac Representation

- Dirac representation of a number $n = p \times q$:

$$\delta_n(f) = A_1\delta(f - f_1) + A_2\delta(f - f_2)$$

- ξ -broadened Dirac function:

$$\delta_\xi(\omega - \omega_0) = \frac{1}{\sqrt{4\pi\xi}} \times \exp\left(-\frac{(\omega - \omega_0)^2}{4\xi}\right)$$

- Complete Dirac number function:

$$\Psi_n(\omega, \xi) = \sum_i A_i \times \frac{1}{\sqrt{4\pi\xi}} \times \exp\left(-\frac{(\omega - \omega_i)^2}{4\xi}\right)$$

10.3 Factorization through FFT Spectral Theory

- Fundamental frequencies in the spectrum correspond to prime factors:

$$n = p \times q \rightarrow \{f_1 = f_0 \times p, f_2 = f_0 \times q\}$$

- Spectral ratio (must always be considered as a ratio):

$$R(n) = \frac{q}{p} = \frac{\max(p, q)}{\min(p, q)}$$

- Octave reduction to avoid rounding errors:

$$R_{\text{oct}}(n) = \frac{R(n)}{2^{\lfloor \log_2(R(n)) \rfloor}}$$

- Beat frequency (difference frequency):

$$f_{\text{beat}} = |f_2 - f_1| = f_0 \times |q - p|$$

10.4 Harmonic Hierarchy for Factorizations

- Basic (1.0 - 1.4): Classical harmonies

$$\text{e.g., } \frac{3}{2} = 1.5 \text{ (perfect fifth), } \frac{5}{4} = 1.25 \text{ (major third)}$$

- Extended (1.4 - 1.6): Jazz/modern harmonies

$$\text{e.g., } \frac{11}{8} = 1.375, \frac{13}{8} = 1.625$$

- Complex (1.6 - 1.85): Microtonal spectra

$$\text{e.g., } \frac{29}{16} = 1.8125, \frac{31}{16} = 1.9375$$

- Ultra (1.85+): Xenharmonic spectra

$$\text{e.g., } \frac{61}{32} = 1.90625, \frac{37}{32} = 1.15625$$

10.5 Resonance Score for Factorizations

- Optimal resonance parameter:

$$\xi = \frac{1}{10}$$

- Angular frequency for period r :

$$\omega = \frac{2\pi}{r}$$

- Resonance score:

$$\text{Res}(r, \xi) = \frac{1}{1 + \frac{|\omega - \pi|^2}{4\xi}}$$

10.6 Ratio-Based Calculation to Avoid Rounding Errors

- **IMPORTANT NOTE:** All computational operations must be performed using ratios, as floating-point calculations introduce rounding errors that render the results unusable. Precise calculation of ratios is critical for the correct application of the T0-model.
- Instead of absolute values, ratios should always be used: $\frac{f_1}{f_0} = p$, $\frac{f_2}{f_0} = q$, $\frac{f_2}{f_1} = \frac{q}{p}$
- When implementing in computer programs, libraries for exact arithmetic (fractional computation) should be used to avoid floating-point rounding errors.
- Harmonic distance (in cents): $d_{\text{harm}}(n, h) = 1200 \times \left| \log_2 \left(\frac{R_{\text{oct}}(n)}{h} \right) \right|$
- Matching criterion: $\text{Match}(n, \text{harmonic_ratio}) = \text{TRUE}$ if $|R_{\text{oct}}(n) - \text{harmonic_ratio}|^2 < 4\xi$

11 SYMBOL EXPLANATIONS

11.1 General Symbols

- ξ = Universal geometric parameter ($4/3 \times 10^{-4}$)
- G = Gravitational constant
- c = Speed of light
- \hbar = Reduced Planck constant
- k_B = Boltzmann constant
- E_P = Planck energy
- ℓ_P = Planck length
- T_0 = Reference time field value
- E_0 = Reference energy field value

11.2 Field Theory Symbols

- E_{field} = Energy field
- T_{field} = Time field
- δE = Energy field fluctuation
- \mathcal{L} = Lagrangian density
- \square = D'Alembert operator
- $\Gamma_{\mu}^{(T)}$ = Time field connection
- ∇ = Nabla operator
- ∂_{μ} = Partial derivative with respect to coordinate μ

11.3 Quantum Mechanical Symbols

- ψ = Wave function
- γ^{μ} = Dirac matrices
- \hat{H} = Hamiltonian operator
- $|\psi\rangle$ = State vector
- $\langle A \rangle$ = Expectation value of observable A
- a_{μ} = Anomalous magnetic moment of the muon
- a_{ℓ} = Anomalous magnetic moment of a lepton

11.4 Particle Physics Symbols

- α_{EM} = Electromagnetic coupling constant
- α_G = Gravitational coupling
- α_W = Weak coupling
- α_S = Strong coupling
- E_{μ} = Muon energy/mass
- E_e = Electron energy/mass
- E_{τ} = Tau energy/mass

11.5 Cosmological Symbols

- z = Redshift
- λ = Wavelength
- ν = Frequency
- H_0 = Hubble parameter
- θ = Deflection angle
- ds^2 = Line element
- $a(t)$ = Scale factor

11.6 Spectral Analysis and Factorization

- $R(n)$ = Spectral ratio of a number n
- $R_{\text{oct}}(n)$ = Octave-reduced spectral ratio
- f_{beat} = Beat frequency
- δ_ξ = ξ -broadened Dirac function
- Ψ_n = Spectral wave function of a number
- ω = Angular frequency
- d_{harm} = Harmonic distance