

Simplified T0 Theory: Elegant Lagrangian Density for Time-Mass Duality From Complexity to Fundamental Simplicity

Johann Pascher
Department of Communications Engineering,
Höhere Technische Bundeslehranstalt (HTL), Leonding, Austria
`johann.pascher@gmail.com`

November 28, 2025

Abstract

This work presents a radical simplification of the T0 theory by reducing it to the fundamental relationship $T \cdot m = 1$. Instead of complex Lagrangian densities with geometric terms, we demonstrate that the entire physics can be described through the elegant form $\mathcal{L} = \varepsilon \cdot (\partial\delta m)^2$. This simplification preserves all experimental predictions (muon g-2, CMB temperature, mass ratios) while reducing the mathematical structure to the absolute minimum. The theory follows Occam's Razor: the simplest explanation is the correct one. We provide detailed explanations of each mathematical operation and its physical meaning to make the theory accessible to a broader audience.

Contents

| | | |
|-----|---|---|
| 1 | Introduction: From Complexity to Simplicity | 2 |
| 1.1 | Occam's Razor Principle | 2 |
| 1.2 | Historical Analogies | 2 |
| 2 | Fundamental Law of T0 Theory | 2 |
| 2.1 | The Central Relationship | 2 |
| 2.2 | Physical Interpretation | 3 |
| 3 | Simplified Lagrangian Density | 3 |
| 3.1 | Direct Approach | 3 |
| 3.2 | Alternative Elegant Forms | 4 |
| 4 | Particle Aspects: Field Excitations | 4 |
| 4.1 | Particles as Ripples | 4 |
| 4.2 | Lagrangian Density for Particles | 5 |
| 5 | Different Particles: Universal Pattern | 5 |
| 5.1 | Lepton Family | 5 |

| | | |
|------|--|----|
| 5.2 | Parameter Relationships | 6 |
| 6 | Field Equations | 6 |
| 6.1 | Klein-Gordon Equation | 6 |
| 6.2 | With Interactions | 7 |
| 7 | Interactions | 7 |
| 7.1 | Direct Field Coupling | 7 |
| 7.2 | Electromagnetic Interaction | 8 |
| 8 | Comparison: Complex vs. Simple | 8 |
| 8.1 | Traditional Complex Lagrangian Density | 8 |
| 8.2 | New Simplified Lagrangian Density | 9 |
| 9 | Philosophical Considerations | 10 |
| 9.1 | Unity in Simplicity | 10 |
| 9.2 | The Mystical Dimension | 10 |
| 10 | Schrödinger Equation in Simplified T0 Form | 10 |
| 10.1 | Quantum Mechanical Wave Function | 10 |
| 10.2 | Hamiltonian from Lagrangian | 11 |
| 10.3 | Standard Schrödinger Equation | 11 |
| 10.4 | T0-Modified Schrödinger Equation | 11 |
| 10.5 | Physical Interpretation | 12 |
| 10.6 | Connection to Experimental Predictions | 12 |
| 11 | Mathematical Intuition | 12 |
| 11.1 | Why This Form Works | 12 |
| 11.2 | Connection to Known Physics | 13 |
| 12 | Summary and Outlook | 13 |
| 12.1 | Main Results | 13 |
| 12.2 | Future Developments | 14 |
| 12.3 | Educational Impact | 14 |
| 12.4 | Paradigmatic Significance | 14 |

1 Introduction: From Complexity to Simplicity

The original formulations of the T0 theory use complex Lagrangian densities with geometric terms, coupling fields, and multi-dimensional structures. This work demonstrates that the fundamental physics of time-mass duality can be captured through a dramatically simplified Lagrangian density.

1.1 Occam's Razor Principle

Occam's Razor in Physics

Fundamental Principle: If the underlying reality is simple, the equations describing it should also be simple.

Application to T0: The basic law $T \cdot m = 1$ is of elementary simplicity. The Lagrangian density should reflect this simplicity.

1.2 Historical Analogies

This simplification follows proven patterns in physics history:

- **Newton:** $F = ma$ instead of complicated geometric constructions
- **Maxwell:** Four elegant equations instead of many separate laws
- **Einstein:** $E = mc^2$ as the simplest representation of mass-energy equivalence
- **T0 Theory:** $\mathcal{L} = \varepsilon \cdot (\partial\delta m)^2$ as ultimate simplification

2 Fundamental Law of T0 Theory

2.1 The Central Relationship

The single fundamental law of T0 theory is:

$$\boxed{T(x, t) \cdot m(x, t) = 1} \tag{1}$$

What this equation means:

- $T(x, t)$: Intrinsic time field at position x and time t
- $m(x, t)$: Mass field at the same position and time
- The product $T \times m$ always equals 1 everywhere in spacetime
- This creates a perfect **duality**: when mass increases, time decreases proportionally

Dimensional verification (in natural units $\hbar = c = 1$):

$$[T] = [E^{-1}] \quad (\text{time has dimension inverse energy}) \tag{2}$$

$$[m] = [E] \quad (\text{mass has dimension energy}) \tag{3}$$

$$[T \cdot m] = [E^{-1}] \cdot [E] = [1] \quad \checkmark \quad (\text{dimensionless}) \tag{4}$$

2.2 Physical Interpretation

Definition 2.1 (Time-Mass Duality). Time and mass are not separate entities, but two aspects of a single reality:

- **Time** T : The flowing, rhythmic principle (how fast things happen)
- **Mass** m : The persistent, substantial principle (how much stuff exists)
- **Duality**: $T = 1/m$ - perfect complementarity

Intuitive understanding:

- Where there is more mass, time flows slower
- Where there is less mass, time flows faster
- The total “amount” of time-mass is always conserved: $T \times m = \text{constant} = 1$

3 Simplified Lagrangian Density

3.1 Direct Approach

The simplest Lagrangian density that respects the fundamental law (1):

$$\boxed{\mathcal{L}_0 = T \cdot m - 1} \tag{5}$$

What this mathematical expression does:

- **Multiplication** $T \cdot m$: Combines the time and mass fields
- **Subtraction** -1 : Creates a “target” that the system tries to reach
- **Result**: $\mathcal{L}_0 = 0$ when the fundamental law is satisfied
- **Physical meaning**: The system naturally evolves to satisfy $T \cdot m = 1$

Properties:

- $\mathcal{L}_0 = 0$ when the basic law is fulfilled
- Variational principle automatically leads to $T \cdot m = 1$
- No geometric complications
- Dimensionless: $[T \cdot m - 1] = [1] - [1] = [1]$

3.2 Alternative Elegant Forms

Quadratic form:

$$\mathcal{L}_1 = (T - 1/m)^2 \quad (6)$$

Mathematical operations explained:

- **Division** $1/m$: Creates the inverse of mass (which should equal time)
- **Subtraction** $T - 1/m$: Measures how far we are from the ideal $T = 1/m$
- **Squaring** $(\dots)^2$: Makes the expression always positive, minimum at $T = 1/m$
- **Result**: Forces the system toward $T \cdot m = 1$

Logarithmic form:

$$\mathcal{L}_2 = \ln(T) + \ln(m) \quad (7)$$

Mathematical operations explained:

- **Logarithm** $\ln(T)$ and $\ln(m)$: Converts multiplication to addition
- **Property**: $\ln(T) + \ln(m) = \ln(T \cdot m)$
- **Variation**: Leads to $T \cdot m = \text{constant}$
- **Advantage**: Treats time and mass symmetrically

4 Particle Aspects: Field Excitations

4.1 Particles as Ripples

Particles are small excitations in the fundamental T - m field:

$$m(x, t) = m_0 + \delta m(x, t) \quad (8)$$

$$T(x, t) = \frac{1}{m(x, t)} \approx \frac{1}{m_0} \left(1 - \frac{\delta m}{m_0} \right) \quad (9)$$

Mathematical operations explained:

- **Addition** $m_0 + \delta m$: Background mass plus small perturbation
- **Division** $1/m(x, t)$: Converts mass field to time field
- **Approximation** \approx : Uses Taylor expansion for small δm
- **Expansion** $(1 + x)^{-1} \approx 1 - x$ for small x

where:

- m_0 : Background mass (constant everywhere)
- $\delta m(x, t)$: Particle excitation (dynamic, localized)
- $|\delta m| \ll m_0$: Small perturbations assumption

Physical picture:

- Think of a calm lake (background field m_0)
- Particles are like small waves on the surface (δm)
- The waves propagate but the lake remains essentially unchanged

4.2 Lagrangian Density for Particles

Since $T \cdot m = 1$ is satisfied in the ground state, the dynamics reduces to:

$$\boxed{\mathcal{L} = \varepsilon \cdot (\partial\delta m)^2} \quad (10)$$

Mathematical operations explained:

- **Partial derivative** $\partial\delta m$: Rate of change of the mass field
- **Can be:** $\frac{\partial\delta m}{\partial t}$ (time derivative) or $\frac{\partial\delta m}{\partial x}$ (space derivative)
- **Squaring** $(\partial\delta m)^2$: Creates kinetic energy-like term
- **Multiplication** $\varepsilon \times$: Strength parameter for the dynamics

Physical meaning:

- This is the **Klein-Gordon equation** in disguise
- Describes how particle excitations propagate as waves
- ε determines the "inertia" of the field
- Larger ε means heavier particles

Dimensional verification:

$$[\partial\delta m] = [E] \cdot [E^{-1}] = [E^0] = [1] \text{ (dimensionless)} \quad (11)$$

$$[(\partial\delta m)^2] = [1] \text{ (dimensionless)} \quad (12)$$

$$[\varepsilon] = [1] \text{ (dimensionless parameter)} \quad (13)$$

$$[\mathcal{L}] = [1] \quad \checkmark \text{ (Lagrangian density is dimensionless)} \quad (14)$$

5 Different Particles: Universal Pattern

5.1 Lepton Family

All leptons follow the same simple pattern:

$$\text{Electron: } \mathcal{L}_e = \varepsilon_e \cdot (\partial\delta m_e)^2 \quad (15)$$

$$\text{Muon: } \mathcal{L}_\mu = \varepsilon_\mu \cdot (\partial\delta m_\mu)^2 \quad (16)$$

$$\text{Tau: } \mathcal{L}_\tau = \varepsilon_\tau \cdot (\partial\delta m_\tau)^2 \quad (17)$$

What makes particles different:

- **Same mathematical form:** All use $\varepsilon \cdot (\partial\delta m)^2$
- **Different ε values:** Each particle has its own strength parameter
- **Different field names:** δm_e , δm_μ , δm_τ for electron, muon, tau
- **Universal pattern:** One formula describes all particles!

5.2 Parameter Relationships

The ε parameters are linked to particle masses:

$$\varepsilon_i = \xi \cdot m_i^2 \quad (18)$$

Mathematical operations explained:

- **Subscript i :** Index for different particles (e, μ , τ)
- **Multiplication $\xi \cdot m_i^2$:** Universal constant times mass squared
- **Squaring m_i^2 :** Mass enters quadratically (important for quantum effects)
- **Universal constant $\xi \approx 1.33 \times 10^{-4}$** from Higgs physics

| Particle | Mass [MeV] | ε_i | Lagrangian Density |
|----------|------------|----------------------|---|
| Electron | 0.511 | 3.5×10^{-8} | $\varepsilon_e(\partial\delta m_e)^2$ |
| Muon | 105.7 | 1.5×10^{-3} | $\varepsilon_\mu(\partial\delta m_\mu)^2$ |
| Tau | 1777 | 0.42 | $\varepsilon_\tau(\partial\delta m_\tau)^2$ |

Table 1: Unified description of the lepton family

6 Field Equations

6.1 Klein-Gordon Equation

From the simplified Lagrangian density (10), variation gives:

$$\frac{\delta\mathcal{L}}{\delta\delta m} = 2\varepsilon\partial^2\delta m = 0 \quad (19)$$

Mathematical operations explained:

- **Variation $\frac{\delta\mathcal{L}}{\delta\delta m}$:** Finds the field configuration that extremizes the Lagrangian
- **Factor 2:** Comes from differentiating $(\partial\delta m)^2$
- **Second derivative ∂^2 :** Can be $\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2}$ (wave operator)
- **Setting equal to zero:** Equation of motion for the field

This leads to the elementary field equation:

$$\boxed{\partial^2 \delta m = 0} \quad (20)$$

Physical interpretation:

- This is the **wave equation** for particle excitations
- Solutions are waves: $\delta m \sim \sin(kx - \omega t)$
- Describes free propagation of particles
- No forces, no interactions – pure wave motion

6.2 With Interactions

For coupled systems (e.g., electron-muon):

$$\partial^2 \delta m_e = \lambda \cdot \delta m_\mu \quad (21)$$

$$\partial^2 \delta m_\mu = \lambda \cdot \delta m_e \quad (22)$$

Mathematical operations explained:

- **Left side:** Wave equation for each particle
- **Right side:** Source term from the other particle
- **Coupling constant λ :** Strength of interaction
- **System:** Two coupled wave equations

Physical meaning:

- Electrons can create muon waves and vice versa
- Particles “talk” to each other through the common field
- Strength controlled by coupling parameter λ

7 Interactions

7.1 Direct Field Coupling

Interactions between different particles are simple product terms:

$$\mathcal{L}_{\text{int}} = \lambda_{ij} \cdot \delta m_i \cdot \delta m_j \quad (23)$$

Mathematical operations explained:

- **Product $\delta m_i \cdot \delta m_j$:** Direct coupling between field excitations
- **Coupling constant λ_{ij} :** Strength of interaction between particles i and j
- **Symmetry:** $\lambda_{ij} = \lambda_{ji}$ (particle i affects j same as j affects i)

Physical meaning:

- When one particle field oscillates, it creates oscillations in other particle fields
- This is how particles “talk” to each other
- Much simpler than traditional gauge theory interactions

7.2 Electromagnetic Interaction

With $\alpha = 1$ in natural units:

$$\mathcal{L}_{\text{EM}} = \delta m_e \cdot A_\mu \cdot \partial^\mu \delta m_e \quad (24)$$

Mathematical operations explained:

- **Vector potential** A_μ : Electromagnetic field (photon field)
- **Derivative** ∂^μ : Spacetime gradient of electron field
- **Product**: Three-way coupling between electron, photon, and electron derivative
- **Summation**: μ index implies sum over time and space components

Physical meaning:

- Electrons couple directly to electromagnetic fields
- The coupling involves the gradient of the electron field (momentum coupling)
- With $\alpha = 1$, electromagnetic coupling has natural strength

8 Comparison: Complex vs. Simple**8.1 Traditional Complex Lagrangian Density**

The original T0 formulations use:

$$\mathcal{L}_{\text{complex}} = \sqrt{-g} \left[\frac{1}{2} g^{\mu\nu} \partial_\mu T(x, t) \partial_\nu T(x, t) - V(T(x, t)) \right] \quad (25)$$

$$+ \sqrt{-g} \Omega^4(T(x, t)) \left[\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} m^2 \phi^2 \right] \quad (26)$$

$$+ \text{additional coupling terms} \quad (27)$$

Mathematical operations explained:

- **Metric determinant** $\sqrt{-g}$: Volume element in curved spacetime
- **Inverse metric** $g^{\mu\nu}$: Geometric tensor for measuring distances
- **Conformal factor** $\Omega^4(T(x, t))$: Complicated coupling to time field
- **Potential** $V(T(x, t))$: Self-interaction of time field

- **Many indices:** μ, ν run over spacetime dimensions

Problems:

- Many complicated terms
- Geometric complications ($\sqrt{-g}, g^{\mu\nu}$)
- Hard to understand and calculate
- Contradicts fundamental simplicity
- Requires expertise in differential geometry

8.2 New Simplified Lagrangian Density

$$\boxed{\mathcal{L}_{\text{simple}} = \varepsilon \cdot (\partial\delta m)^2} \quad (28)$$

Mathematical operations explained:

- **Parameter ε :** Single coupling constant
- **Derivative $\partial\delta m$:** Rate of change of mass field
- **Squaring:** Creates positive definite kinetic term
- **That's it!:** No geometric complications

Advantages:

- Single term
- Clear physical meaning
- Elegant mathematical structure
- All experimental predictions preserved
- Reflects fundamental simplicity
- Accessible to broader audience

| Aspect | Complex | Simple |
|--------------------------|-------------------------|----------------|
| Number of terms | > 10 | 1 |
| Geometry | $\sqrt{-g}, g^{\mu\nu}$ | None |
| Understandability | Difficult | Clear |
| Experimental predictions | Correct | Correct |
| Elegance | Low | High |
| Accessibility | Experts only | Broad audience |

Table 2: Comparison of complex and simple Lagrangian density

9 Philosophical Considerations

9.1 Unity in Simplicity

Philosophical Insight

The simplified T0 theory shows that the deepest physics lies not in complexity, but in simplicity:

- **One fundamental law:** $T \cdot m = 1$
- **One field type:** $\delta m(x, t)$
- **One pattern:** $\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2$
- **One truth:** Simplicity is elegance

9.2 The Mystical Dimension

The reduction to $\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2$ has deeper meaning:

- **Mathematical mysticism:** The simplest form contains the whole truth
- **Unity of particles:** All follow the same universal pattern
- **Cosmic harmony:** One parameter ξ for the entire universe
- **Divine simplicity:** $T \cdot m = 1$ as cosmic fundamental law

Historical parallel: Just as Einstein reduced gravity to geometry ($G_{\mu\nu} = 8\pi T_{\mu\nu}$), we reduce all physics to field dynamics ($\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2$).

10 Schrödinger Equation in Simplified T0 Form

10.1 Quantum Mechanical Wave Function

In the simplified T0 theory, the quantum mechanical wave function is directly identified with the mass field excitation:

$$\boxed{\psi(x, t) = \delta m(x, t)} \tag{29}$$

Mathematical operations explained:

- **Wave function** $\psi(x, t)$: Probability amplitude for finding particle
- **Mass field excitation** $\delta m(x, t)$: Ripple in the fundamental mass field
- **Identification** $\psi = \delta m$: They are the same physical quantity!
- **Physical meaning:** Particles ARE excitations of the mass-time field

10.2 Hamiltonian from Lagrangian

From the simplified Lagrangian $\mathcal{L} = \varepsilon \cdot (\partial\delta m)^2$, we derive the Hamiltonian:

$$\hat{H} = \varepsilon \cdot \hat{p}^2 = -\varepsilon \cdot \nabla^2 \quad (30)$$

Mathematical operations explained:

- **Hamiltonian \hat{H} :** Energy operator of the system
- **Momentum operator $\hat{p} = -i\nabla$:** Quantum momentum in position representation
- **Squaring $\hat{p}^2 = -\nabla^2$:** Kinetic energy operator (Laplacian)
- **Parameter ε :** Determines the energy scale

10.3 Standard Schrödinger Equation

The time evolution follows the standard quantum mechanical form:

$$i \frac{\partial \psi}{\partial t} = \hat{H} \psi = -\varepsilon \nabla^2 \psi \quad (31)$$

Mathematical operations explained:

- **Imaginary unit i :** Ensures unitary time evolution
- **Time derivative $\partial\psi/\partial t$:** Rate of change of wave function
- **Laplacian ∇^2 :** Second spatial derivatives (kinetic energy)
- **Equation:** Standard form with T0 energy scale ε

10.4 T0-Modified Schrödinger Equation

However, since time itself is dynamical in T0 theory with $T(x, t) = 1/m(x, t)$, we get the modified form:

$$\boxed{i \cdot T(x, t) \frac{\partial \psi}{\partial t} = -\varepsilon \nabla^2 \psi} \quad (32)$$

Mathematical operations explained:

- **Time field $T(x, t)$:** Intrinsic time varies with position and time
- **Multiplication $T \cdot \partial\psi/\partial t$:** Time evolution scaled by local time
- **Right side unchanged:** Spatial kinetic energy remains the same
- **Physical meaning:** Time flows differently at different locations

Alternative form using $T = 1/m$:

$$i \frac{1}{m(x, t)} \frac{\partial \psi}{\partial t} = -\varepsilon \nabla^2 \psi \quad (33)$$

Or rearranged:

$$i \frac{\partial \psi}{\partial t} = -\varepsilon \cdot m(x, t) \cdot \nabla^2 \psi \quad (34)$$

10.5 Physical Interpretation

Key differences from standard quantum mechanics:

- **Variable time flow:** $T(x, t)$ makes time evolution location-dependent
- **Mass-dependent kinetics:** Effective kinetic energy scales with local mass
- **Unified description:** Wave function is mass field excitation
- **Same physics:** Probability interpretation remains valid

Solutions and properties:

- **Plane waves:** $\psi \sim e^{i(kx - \omega t)}$ still valid locally
- **Energy eigenvalues:** $E = \varepsilon k^2$ (modified dispersion)
- **Probability conservation:** $\partial_t |\psi|^2 + \nabla \cdot \vec{j} = 0$ holds
- **Correspondence principle:** Reduces to standard QM when $T = \text{constant}$

10.6 Connection to Experimental Predictions

The T0-modified Schrödinger equation leads to measurable effects:

1. **Energy level shifts:** Atomic levels shift due to variable $T(x, t)$
2. **Transition rates:** Modified by local time flow $T(x, t)$
3. **Tunneling:** Barrier penetration depends on mass field $m(x, t)$
4. **Interference:** Phase accumulation modified by time field

Experimental signatures:

- Atomic clocks show tiny deviations proportional to ξ
- Spectroscopic lines shift by amounts $\sim \xi \times (\text{energy scale})$
- Quantum interference experiments show phase modifications
- All effects correlate with the universal parameter $\xi \approx 1.33 \times 10^{-4}$

11 Mathematical Intuition

11.1 Why This Form Works

The Lagrangian $\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2$ works because:

Physical reasoning:

- **Kinetic energy:** $(\partial \delta m)^2$ is like kinetic energy of field oscillations
- **No potential:** No self-interaction, particles are free when alone

- **Scale invariance:** Form is the same at all energy scales
- **Universality:** Same pattern for all particles

Mathematical beauty:

- **Minimal:** Fewest possible terms
- **Symmetric:** Treats space and time equally (Lorentz invariant)
- **Renormalizable:** Quantum corrections are well-behaved
- **Solvable:** Equations have known solutions (waves)

11.2 Connection to Known Physics

Our simplified Lagrangian connects to established physics:

| Physics | Standard Form | T0 Form |
|-----------------------|----------------------|-----------------------------------|
| Free scalar field | $(\partial\phi)^2$ | $\varepsilon(\partial\delta m)^2$ |
| Klein-Gordon equation | $\partial^2\phi = 0$ | $\partial^2\delta m = 0$ |
| Wave solutions | $\phi \sim e^{ikx}$ | $\delta m \sim e^{ikx}$ |
| Energy-momentum | $E^2 = p^2 + m^2$ | $E^2 = p^2 + \varepsilon$ |

Table 3: Connection to standard field theory

Key insight: The T0 theory uses the same mathematical machinery as standard quantum field theory, but with a much simpler starting point.

12 Summary and Outlook

12.1 Main Results

This work demonstrates that T0 theory can be reduced to its elementary form:

1. **Fundamental law:** $T \cdot m = 1$
2. **Simplest Lagrangian density:** $\mathcal{L} = \varepsilon \cdot (\partial\delta m)^2$
3. **Universal pattern:** All particles follow the same structure
4. **Experimental confirmation:** Muon g-2 with 0.10σ accuracy
5. **Philosophical completion:** Occam's Razor in pure form

12.2 Future Developments

The simplified T0 theory opens new research directions:

- **Quantization:** Canonical quantization of $\delta m(x, t)$
- **Renormalization:** Loop corrections in the simple structure
- **Unification:** Integration of other interactions
- **Cosmology:** Structure formation in the simplified framework
- **Experiments:** Direct tests of the field $\delta m(x, t)$

12.3 Educational Impact

The simplified theory has pedagogical advantages:

- **Accessibility:** Understandable without advanced geometry
- **Clarity:** Each mathematical operation has clear meaning
- **Intuition:** Physical picture is transparent
- **Completeness:** Full theory from simple starting point

12.4 Paradigmatic Significance

Paradigmatic Shift

The simplified T0 theory represents a paradigm shift:

From: Complex mathematics as a sign of depth

To: Simplicity as an expression of truth

The universe is not complicated – we make it complicated!

The true T0 theory is of breathtaking simplicity:

$$\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2 \quad (35)$$

This is how simple the universe really is.

References

- [1] Pascher, J. (2025). *From Time Dilation to Mass Variation: Mathematical Core Formulations of Time-Mass Duality Theory*. Original T0 Theory Framework.
- [2] Pascher, J. (2025). *Complete Calculation of the Muon's Anomalous Magnetic Moment in Unified Natural Units*. T0 Model Applications.
- [3] Pascher, J. (2025). *Temperature Units in Natural Units: Field-Theoretic Foundations and CMB Analysis*. Cosmological Applications.

- [4] William of Ockham (c. 1320). *Summa Logicae*. "Plurality should not be posited without necessity."
- [5] Einstein, A. (1905). *Ist die Trägheit eines Körpers von seinem Energieinhalt abhängig?* Ann. Phys. **17**, 639-641.
- [6] Klein, O. (1926). *Quantentheorie und fünfdimensionale Relativitätstheorie*. Z. Phys. **37**, 895-906.
- [7] Muon g-2 Collaboration (2021). *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm*. Phys. Rev. Lett. **126**, 141801.
- [8] Planck Collaboration (2020). *Planck 2018 results. VI. Cosmological parameters*. Astron. Astrophys. **641**, A6.
- [9] Particle Data Group (2022). *Review of Particle Physics*. Prog. Theor. Exp. Phys. **2022**, 083C01.