

The T0 Energy Field Model:

Mathematical Formulation

January 2025

Abstract

The T0 model describes physical phenomena through a universal energy field $E_{\text{field}}(x, t)$ with the parameter $\xi = \frac{4}{3} \times 10^{-4}$. The field equation is $\square E_{\text{field}} = 0$, the Lagrangian density $\mathcal{L} = \xi(\partial E)^2$. The model uses standard natural units with $\hbar = c = 1$.

Fundamental quantities:

- Characteristic energy: $E_0 = \sqrt{m_e \cdot m_\mu} = 7.348 \text{ MeV}$
- Fine structure constant: $\alpha = \xi(E_0/1 \text{ MeV})^2 \approx 1/137$
- Gravitational constant: $G = \xi^2/(4m_e) \times \text{factors}$

Predictions: Lepton masses with 2% accuracy, anomalous magnetic moments $a_\ell = \frac{\xi}{2\pi}(E_\ell/E_e)^2$, fine structure constant with 0.03% agreement.

Detailed derivations: See Document 011 (fine structure), 012 (gravitation), 018 (g-2 geometric), 019 (Lagrangian).

Contents

1	Units Convention	4
1.1	T0 natural units (Heaviside-Lorentz)	4
1.2	Dimensions in natural units	4
2	Time-Energy Duality	4
2.1	Fundamental relation	4
2.2	Intrinsic time field	4
3	Universal Field Equation	5
3.1	Wave equation	5
3.2	With sources	5
4	Lagrangian Density	5

4.1	Universal Lagrangian density	5
4.2	Euler-Lagrange equation	5
5	Characteristic Energy	5
5.1	Definition	5
5.2	Numerical values	6
5.3	Usage	6
6	The Parameter ξ	6
6.1	Definition	6
6.2	Geometric components	6
7	Fine Structure Constant	6
7.1	In T0 units	6
7.2	Reconstruction of SI value	7
7.3	Numerical calculation	7
7.4	Dimension check	7
8	Gravitational Constant	7
8.1	T0 formula	7
8.2	Fundamental relation	8
9	Characteristic Lengths	8
9.1	T0 characteristic length	8
9.2	Derivation	8
9.3	Time scale	8
10	Scale Hierarchy	8
10.1	Planck length as reference	8
10.2	Scale ratio	9
11	Particles as Field Excitations	9
11.1	Classification by energy	9
11.2	Antiparticles	9
12	Lepton Masses	9
13	Anomalous Magnetic Moments	9
13.1	Definition	9
13.2	T0 prediction formula	10
13.3	Muon	10
13.4	Electron	10
13.5	Tau	10

14	Three Field Geometries	10
14.1	Type 1: Localized spherical	10
14.2	Type 2: Localized non-spherical	11
14.3	Type 3: Extended homogeneous	11
15	Mathematical Identities	11
15.1	Energy field normalization	11
15.2	Duality consistency	11
16	Dimensional Analysis Verifications	11
16.1	Field equation	11
16.2	Characteristic length	12
16.3	Lagrangian density	12
16.4	Anomalous magnetic moment	12
17	Formula Reference	12
17.1	Fundamental equations	12
17.2	Derived constants	12
17.3	Characteristic scales	12
17.4	Prediction formulas	13
18	Numerical Values	13
18.1	Fundamental constants (in natural units)	13
18.2	T0 parameters	13
18.3	Lepton energies	13
18.4	Energy ratios	14
19	Calculation Examples	14
19.1	Muon g-2	14
19.2	Fine structure constant	14
19.3	Characteristic length (electron)	15
A	Symbol Reference	15
B	Unit Conversions	16
B.1	Natural \rightarrow SI	16
B.2	Standard natural units	16
C	Relations to Other Documents	16

1 Units Convention

1.1 T0 natural units (Heaviside-Lorentz)

T0 uses Heaviside-Lorentz natural units:

$$\hbar = c = 4\pi\epsilon_0 = 1 \quad (1)$$

In this system:

- Fine structure constant: $\alpha = e^2 = 1$ (by convention in T0)
- Energy = Mass: $E = m$
- Length = Time = Energy⁻¹: $[L] = [T] = [E^{-1}]$

Note: In SI units, $\alpha \approx 1/137$. The choice $\alpha = 1$ in T0 is a unit convention that simplifies formulas.

1.2 Dimensions in natural units

$$[E] = E \quad (2)$$

$$[m] = E \quad (3)$$

$$[t] = E^{-1} \quad (4)$$

$$[L] = E^{-1} \quad (5)$$

$$[G] = E^{-2} \quad (6)$$

$$[\partial_\mu] = E \quad (7)$$

2 Time-Energy Duality

2.1 Fundamental relation

$$T_{\text{field}}(x, t) \cdot E_{\text{field}}(x, t) = 1 \quad (8)$$

with $[T_{\text{field}}] = E^{-1}$ and $[E_{\text{field}}] = E$.

2.2 Intrinsic time field

$$T_{\text{field}}(x, t) = \frac{1}{E_{\text{field}}(x, t)} \quad (9)$$

3 Universal Field Equation

3.1 Wave equation

$$\square E_{\text{field}} = 0 \quad (10)$$

with d'Alembert operator:

$$\square = \nabla^2 - \frac{\partial^2}{\partial t^2} \quad (11)$$

3.2 With sources

$$\nabla^2 E_{\text{field}} = 4\pi G\rho \cdot E_{\text{field}} \quad (12)$$

Dimension check: $[E^3] = [E^{-2}][E^4][E] = [E^3] \checkmark$

4 Lagrangian Density

4.1 Universal Lagrangian density

$$\mathcal{L} = \xi \cdot (\partial_\mu E_{\text{field}})(\partial^\mu E_{\text{field}}) \quad (13)$$

with $\xi = \frac{4}{3} \times 10^{-4}$.

4.2 Euler-Lagrange equation

$$\frac{\partial \mathcal{L}}{\partial E} - \partial_\mu \frac{\partial \mathcal{L}}{\partial (\partial_\mu E)} = 0 \quad (14)$$

yields:

$$\square E_{\text{field}} = 0 \quad (15)$$

5 Characteristic Energy

5.1 Definition

The characteristic energy E_0 is the geometric mean of electron and muon mass (derivation in Document 011):

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (16)$$

5.2 Numerical values

From experimental masses:

$$E_0 = \sqrt{0.511 \times 105.66} \quad (17)$$

$$= \sqrt{53.99} \quad (18)$$

$$= 7.348 \text{ MeV} \quad (19)$$

Theoretical T0 value:

$$E_0^{\text{T0}} = 7.398 \text{ MeV} \quad (20)$$

Deviation: 0.7% (within geometric corrections)

5.3 Usage

E_0 serves as energy scale for:

- Fine structure constant: $\alpha = \xi(E_0/1 \text{ MeV})^2$
- Normalization of electromagnetic effects
- Scaling of anomalous magnetic moments

6 The Parameter ξ

6.1 Definition

$$\xi = \frac{4}{3} \times 10^{-4} = 1.3333 \times 10^{-4} \quad (21)$$

Dimensionless: $[\xi] = 1$.

6.2 Geometric components

$$\xi = G_3 \times S_{\text{ratio}} \quad (22)$$

where:

- $G_3 = \frac{4}{3}$: Geometric factor (sphere-cube ratio)
- $S_{\text{ratio}} = 10^{-4}$: Scale ratio

7 Fine Structure Constant

7.1 In T0 units

In T0 natural units: $\alpha = 1$ (by convention)

7.2 Reconstruction of SI value

Important: The SI value can be reconstructed from T0 parameters!

$$\alpha_{\text{SI}} = \xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2 \quad (23)$$

7.3 Numerical calculation

With $\xi = \frac{4}{3} \times 10^{-4}$ and $E_0 = 7.398 \text{ MeV}$:

$$\alpha_{\text{SI}} = 1.3333 \times 10^{-4} \times (7.398)^2 \quad (24)$$

$$= 1.3333 \times 10^{-4} \times 54.73 \quad (25)$$

$$= 7.297 \times 10^{-3} \quad (26)$$

$$= \frac{1}{137.04} \quad (27)$$

Experimental: $\alpha_{\text{exp}} = \frac{1}{137.036}$

Agreement: 0.03%

7.4 Dimension check

$$[\alpha_{\text{SI}}] = [\xi] \times \left[\frac{E}{E} \right]^2 = 1 \times 1 = 1 \quad \checkmark \quad (28)$$

8 Gravitational Constant

8.1 T0 formula

The gravitational constant is derived from ξ and m_e (derivation in Document 012):

$$G = \frac{\xi^2}{4m_e} \times C_{\text{dim}} \times C_{\text{conv}} \quad (29)$$

where:

- C_{dim} : Dimension correction
- C_{conv} : SI conversion factor

8.2 Fundamental relation

In natural units:

$$\xi = 2\sqrt{G \cdot m_e} \quad (30)$$

Solved for G :

$$G_{\text{nat}} = \frac{\xi^2}{4m_e} \quad (31)$$

Dimension: $[G] = [E^{-2}]$ in natural units.

9 Characteristic Lengths

9.1 T0 characteristic length

$$r_0 = 2GE \quad (32)$$

Dimension: $[r_0] = [E^{-2}][E] = [E^{-1}] = [L] \checkmark$

9.2 Derivation

For spherically symmetric point source $\rho(r) = E_0 \delta^3(\vec{r})$:

Solution of $\nabla^2 E = 4\pi G \rho E$:

$$E(r) = E_0 \left(1 - \frac{r_0}{r}\right) \quad (33)$$

with $r_0 = 2GE_0$.

9.3 Time scale

$$t_0 = \frac{r_0}{c} = r_0 = 2GE \quad (34)$$

(since $c = 1$)

10 Scale Hierarchy

10.1 Planck length as reference

$$\ell_P = \sqrt{G} = 1 \quad (\text{in nat. units}) \quad (35)$$

10.2 Scale ratio

$$\xi_{\text{ratio}} = \frac{\ell_P}{r_0} = \frac{\sqrt{G}}{2GE} = \frac{1}{2\sqrt{G} \cdot E} \quad (36)$$

For $E \sim 1$ GeV:

$$\frac{r_0}{\ell_P} \sim 10^7 \quad (\text{sub-Planck}) \quad (37)$$

11 Particles as Field Excitations

11.1 Classification by energy

Particle	Energy [MeV]
Electron	0.511
Muon	105.658
Tau	1776.86

11.2 Antiparticles

Negative field excitations: $E_{\text{field}} < 0$

12 Lepton Masses

The T0 model predicts lepton masses (derivation in Document 003):

Lepton	T0 [MeV]	Exp [MeV]	Δ [%]
Electron	0.507	0.511	0.87
Muon	103.5	105.7	2.09
Tau	1815	1777	2.16

13 Anomalous Magnetic Moments

13.1 Definition

Magnetic moment:

$$\mu = g \cdot \frac{e}{2m} \cdot \frac{\hbar}{2} \quad (38)$$

Anomalous magnetic moment:

$$a = \frac{g - 2}{2} \quad (39)$$

13.2 T0 prediction formula

$$a_\ell = \frac{\xi}{2\pi} \left(\frac{E_\ell}{E_e} \right)^2 \quad (40)$$

13.3 Muon

$$\frac{E_\mu}{E_e} = \frac{105.658}{0.511} = 206.768 \quad (41)$$

$$a_\mu = \frac{1.3333 \times 10^{-4}}{2\pi} \times (206.768)^2 \quad (42)$$

$$= 2.122 \times 10^{-5} \times 42,753 \quad (43)$$

$$= 1.166 \times 10^{-3} \quad (44)$$

13.4 Electron

$$a_e = \frac{\xi}{2\pi} = 2.122 \times 10^{-5} \quad (45)$$

13.5 Tau

$$a_\tau = \frac{\xi}{2\pi} \left(\frac{1776.86}{0.511} \right)^2 = 1.28 \times 10^{-3} \quad (46)$$

14 Three Field Geometries

14.1 Type 1: Localized spherical

$$E(r) = E_0 \left(1 - \frac{\beta}{r} \right), \quad \beta = r_0 \quad (47)$$

Application: Individual particles (electron, muon, tau)

14.2 Type 2: Localized non-spherical

$$E(\vec{r}) = E_0 \left(1 - \frac{\beta_{ij} r_i r_j}{r^3} \right) \quad (48)$$

Application: Bound systems

14.3 Type 3: Extended homogeneous

Effective parameter:

$$\xi_{\text{eff}} = \frac{\xi}{2} = \frac{2}{3} \times 10^{-4} \quad (49)$$

Application: Cosmology (see Document 026)

15 Mathematical Identities

15.1 Energy field normalization

$$E_{\text{field}}(\vec{r}, t) = E_0 \cdot f(\vec{r}, t) \cdot e^{i\phi(\vec{r}, t)} \quad (50)$$

with:

- E_0 : Characteristic energy
- $f(\vec{r}, t)$: Normalized profile
- $\phi(\vec{r}, t)$: Phase

15.2 Duality consistency

Time-mass (Document 003): $T \cdot m = 1$

Time-energy (this document): $T \cdot E = 1$

In natural units ($c = 1$):

$$E = mc^2 = m \quad \Rightarrow \quad T \cdot m = T \cdot E \quad (51)$$

16 Dimensional Analysis Verifications

16.1 Field equation

$$[\nabla^2 E] = [L^{-2}][E] = [E^2][E] = [E^3] \quad (52)$$

$$[4\pi G \rho E] = [E^{-2}][E^4][E] = [E^3] \quad \checkmark \quad (53)$$

16.2 Characteristic length

$$[r_0] = [2GE] = [E^{-2}][E] = [E^{-1}] = [L] \quad \checkmark \quad (54)$$

16.3 Lagrangian density

$$[\mathcal{L}] = [\xi][(\partial E)^2] = [1][E^2] = [E^2] \quad (\text{correct for Lagrangian density}) \quad (55)$$

16.4 Anomalous magnetic moment

$$[a_\ell] = [\xi] \left[\frac{E^2}{E^2} \right] = [1][1] = [1] \quad \checkmark \quad (56)$$

17 Formula Reference

17.1 Fundamental equations

$$\text{Duality: } T_{\text{field}} \cdot E_{\text{field}} = 1 \quad (57)$$

$$\text{Wave equation: } \square E_{\text{field}} = 0 \quad (58)$$

$$\text{With sources: } \nabla^2 E = 4\pi G \rho E \quad (59)$$

$$\text{Lagrangian density: } \mathcal{L} = \xi(\partial E)^2 \quad (60)$$

17.2 Derived constants

$$\text{Characteristic energy: } E_0 = \sqrt{m_e \cdot m_\mu} = 7.348 \text{ MeV} \quad (61)$$

$$\text{Fine structure constant: } \alpha = \xi(E_0/1 \text{ MeV})^2 \approx 1/137 \quad (62)$$

$$\text{Gravitational constant: } G = \frac{\xi^2}{4m_e} \times \text{factors} \quad (63)$$

17.3 Characteristic scales

$$\text{T0 length: } r_0 = 2GE \quad (64)$$

$$\text{T0 time: } t_0 = 2GE \quad (65)$$

$$\text{Planck length: } \ell_P = \sqrt{G} = 1 \quad (66)$$

$$\text{Scale ratio: } \xi_{\text{ratio}} = \frac{1}{2\sqrt{GE}} \quad (67)$$

17.4 Prediction formulas

$$\text{g-2 formula: } a_\ell = \frac{\xi}{2\pi} \left(\frac{E_\ell}{E_e} \right)^2 \quad (68)$$

$$\text{Parameter: } \xi = \frac{4}{3} \times 10^{-4} \quad (69)$$

$$\text{Effective parameter: } \xi_{\text{eff}} = \frac{\xi}{2} \quad (70)$$

18 Numerical Values

18.1 Fundamental constants (in natural units)

$$\hbar = 1 \quad (71)$$

$$c = 1 \quad (72)$$

$$\alpha = \frac{1}{137.036} \approx 7.297 \times 10^{-3} \quad (73)$$

$$G = 1 \text{ (numerically, dimension } [E^{-2}]) \quad (74)$$

18.2 T0 parameters

$$\xi = \frac{4}{3} \times 10^{-4} = 1.3333 \times 10^{-4} \quad (75)$$

$$\xi^2 = 1.7778 \times 10^{-8} \quad (76)$$

$$\frac{\xi}{2\pi} = 2.1221 \times 10^{-5} \quad (77)$$

$$\xi_{\text{eff}} = 6.6667 \times 10^{-5} \quad (78)$$

$$E_0 = 7.348 \text{ MeV (from exp. masses)} \quad (79)$$

$$E_0^{\text{T0}} = 7.398 \text{ MeV (theoretical)} \quad (80)$$

18.3 Lepton energies

$$E_e = 0.511 \text{ MeV} \quad (81)$$

$$E_\mu = 105.658 \text{ MeV} \quad (82)$$

$$E_\tau = 1776.86 \text{ MeV} \quad (83)$$

18.4 Energy ratios

$$\frac{E_\mu}{E_e} = 206.768 \quad (84)$$

$$\frac{E_\tau}{E_e} = 3477.2 \quad (85)$$

$$\frac{E_\tau}{E_\mu} = 16.817 \quad (86)$$

19 Calculation Examples

19.1 Muon g-2

Given:

- $\xi = 1.3333 \times 10^{-4}$
- $E_\mu = 105.658 \text{ MeV}$
- $E_e = 0.511 \text{ MeV}$

Calculation:

$$\frac{E_\mu}{E_e} = \frac{105.658}{0.511} = 206.768 \quad (87)$$

$$\left(\frac{E_\mu}{E_e}\right)^2 = 42,753.3 \quad (88)$$

$$\frac{\xi}{2\pi} = \frac{1.3333 \times 10^{-4}}{6.2832} = 2.1221 \times 10^{-5} \quad (89)$$

$$a_\mu = 2.1221 \times 10^{-5} \times 42,753.3 \quad (90)$$

$$= 1.1659 \times 10^{-3} \quad (91)$$

19.2 Fine structure constant

Given:

- $\xi = 1.3333 \times 10^{-4}$
- $E_0 = 7.398 \text{ MeV}$

Calculation:

$$\left(\frac{E_0}{1 \text{ MeV}}\right)^2 = (7.398)^2 = 54.73 \quad (92)$$

$$\alpha = 1.3333 \times 10^{-4} \times 54.73 \quad (93)$$

$$= 7.297 \times 10^{-3} \quad (94)$$

$$= \frac{1}{137.04} \quad (95)$$

Experimental: $\alpha_{\text{exp}} = \frac{1}{137.036}$
 Deviation: 0.03%

19.3 Characteristic length (electron)

Given:

- $E_e = 0.511 \text{ MeV} = 0.511 \times 1.6 \times 10^{-13} \text{ J} = 8.2 \times 10^{-14} \text{ J}$
- $G = 6.674 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$
- $c = 3 \times 10^8 \text{ m/s}$

Conversion to natural units:

$$r_0 = 2GE \approx 10^{-28} \text{ m} \quad (96)$$

Planck comparison:

$$\frac{r_0}{\ell_P} = \frac{10^{-28}}{1.6 \times 10^{-35}} \approx 10^7 \quad (97)$$

A Symbol Reference

Symbol	Meaning	Dimension
ξ	Fundamental parameter	1
E_0	Characteristic energy	E
E_{field}	Universal energy field	E
T_{field}	Intrinsic time field	E^{-1}
r_0	T0 characteristic length	$L = E^{-1}$
t_0	T0 characteristic time	$T = E^{-1}$
ℓ_P	Planck length	$L = E^{-1}$
G	Gravitational constant	E^{-2}
α	Fine structure constant	1
a_ℓ	Anomalous magnetic moment	1
E_e, E_μ, E_τ	Lepton energies	E

m_e, m_μ, m_τ	Lepton masses (= E in nat. units)	E
\mathcal{L}	Lagrangian density	E^4
\square	d'Alembert operator	E^2
ξ_{eff}	Effective parameter ($\xi/2$)	1

B Unit Conversions

B.1 Natural \rightarrow SI

$$1 \text{ (Energy)} = 1 \text{ GeV} = 1.6 \times 10^{-10} \text{ J} \quad (98)$$

$$1 \text{ (Length)} = \frac{\hbar c}{1 \text{ GeV}} = 0.197 \text{ fm} \quad (99)$$

$$1 \text{ (Time)} = \frac{\hbar}{1 \text{ GeV}} = 6.58 \times 10^{-25} \text{ s} \quad (100)$$

B.2 Standard natural units

In standard convention ($\hbar = c = 1$):

- $\alpha = \frac{e^2}{4\pi\epsilon_0} \approx \frac{1}{137}$ (dimensionless)
- All quantities in powers of energy
- Physical predictions identical to other conventions

C Relations to Other Documents

- **Document 003:** Time-mass duality, foundations, origin of ξ
- **Document 011:** Fine structure constant (geometric derivation)
- **Document 012:** Gravitational constant (systematic derivation)
- **Document 018:** Geometric g-2 formulation (fractal geometry)
- **Document 019:** Lagrangian formulation (quantum field theory)
- **Document 026:** Cosmology ($\xi_{\text{eff}} = \xi/2$)

All formulations are based on $\xi = \frac{4}{3} \times 10^{-4}$.