

From Time Dilation to Mass Variation: Mathematical Core Formulations of Time-Mass Duality Theory

Johann Pascher

March 29, 2025

Zusammenfassung

This work presents the essential mathematical formulations of time-mass duality theory, focusing on the fundamental equations and their physical interpretations. The theory establishes a duality between two complementary descriptions of reality: the standard view with time dilation and constant rest mass, and the T0 model with absolute time and variable mass. Central to this framework is the intrinsic time $T(x) = \frac{\hbar}{\max(mc^2, \omega)}$, which enables a unified treatment of massive particles and photons. The mathematical formulations include modified Lagrangian densities that emphasize emergent gravitation and energy-loss redshift in a static universe.

Inhaltsverzeichnis

1	Introduction to Time-Mass Duality	2
1.1	Relationship to the Standard Model	2
2	Emergent Gravitation from the Intrinsic Time Field	2
3	Mathematical Foundations: Intrinsic Time	2
4	Modified Derivative Operators	3
5	Modified Field Equations	3
6	Modified Lagrangian Density for the Higgs Field	3
7	Modified Lagrangian Density for Fermions	3
8	Modified Lagrangian Density for Gauge Bosons	3
9	Complete Total Lagrangian Density	3
10	Cosmological Implications	3
11	Derivation of β_T in the T0 Model	4

1 Introduction to Time-Mass Duality

The time-mass duality theory proposes an alternative framework:

1. Standard View: $t' = \gamma_{\text{Lorentz}} t$, $m_0 = \text{const.}$
2. T0 Model: $T_0 = \text{const.}$, $m = \gamma_{\text{Lorentz}} m_0$

1.1 Relationship to the Standard Model

The T0 model extends the Standard Model with:

1. Intrinsic Time Field: $T(x) = \frac{\hbar}{\max(mc^2, \omega)}$
2. Higgs Field: Φ with dynamic mass coupling
3. Fermion Fields: ψ with Yukawa coupling
4. Gauge Boson Fields: A_μ with $T(x)$ interaction

2 Emergent Gravitation from the Intrinsic Time Field

Theorem 2.1 (Emergence of Gravitation). *Gravitation arises from gradients of the intrinsic time field:*

$$\nabla T(x) = -\frac{\hbar}{m^2 c^2} \nabla m \quad (1)$$

with the modified potential:

$$\Phi(r) = -\frac{GM}{r} + \kappa r, \quad \kappa \approx 4.8 \times 10^{-11} \text{ m/s}^2 \quad (2)$$

Beweis. From $T(x) = \frac{\hbar}{mc^2}$ for massive particles:

$$\nabla T(x) = -\frac{\hbar}{m^2 c^2} \nabla m \quad (3)$$

With $m(\vec{r}) = m_0(1 + \frac{\Phi_g}{c^2})$:

$$\nabla m = \frac{m_0}{c^2} \nabla \Phi_g \quad (4)$$

Thus:

$$\nabla T(x) \approx -\frac{\hbar}{m_0 c^4} \nabla \Phi_g \quad (5)$$

□

3 Mathematical Foundations: Intrinsic Time

Theorem 3.1 (Intrinsic Time).

$$T(x) = \frac{\hbar}{\max(mc^2, \omega)} \quad (6)$$

4 Modified Derivative Operators

Definition 4.1 (Modified Derivative). The modified covariant derivative in the T0 model is:

$$\partial_\mu \Psi + \Psi \partial_\mu T(x) = \partial_\mu \Psi + \Psi \partial_\mu T(x) \quad (7)$$

5 Modified Field Equations

Theorem 5.1 (Modified Schrödinger Equation).

$$i\hbar T(x) \frac{\partial}{\partial t} \Psi + i\hbar \Psi \frac{\partial T(x)}{\partial t} = \hat{H} \Psi \quad (8)$$

6 Modified Lagrangian Density for the Higgs Field

Theorem 6.1 (Higgs Lagrangian Density). *The Lagrangian density of the Higgs field with coupling to $T(x)$ is:*

$$\begin{aligned} \mathcal{L}_{Higgs-T} = & |T(x)(\partial_\mu + igA_\mu)\Phi + \Phi \partial_\mu T(x)|^2 + \frac{1}{2} \partial_\mu T(x) \partial^\mu T(x) - V(T(x), \Phi), \\ & T(x)(\partial_\mu + igA_\mu)\Phi + \Phi \partial_\mu T(x) = T(x)(\partial_\mu + igA_\mu)\Phi + \Phi \partial_\mu T(x) \end{aligned} \quad (9)$$

7 Modified Lagrangian Density for Fermions

Theorem 7.1 (Fermion Lagrangian Density).

$$\mathcal{L}_{Fermion} = \bar{\psi} i \gamma^\mu (\partial_\mu \psi + \psi \partial_\mu T(x)) - y \bar{\psi} \Phi \psi \quad (10)$$

8 Modified Lagrangian Density for Gauge Bosons

Theorem 8.1 (Gauge Boson Lagrangian Density).

$$\mathcal{L}_{Boson} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \partial_\mu T(x) \partial^\mu T(x) \quad (11)$$

9 Complete Total Lagrangian Density

Theorem 9.1 (Total Lagrangian Density).

$$\mathcal{L}_{Total} = \mathcal{L}_{Boson} + \mathcal{L}_{Fermion} + \mathcal{L}_{Higgs-T} + \mathcal{L}_{intrinsic}, \quad \mathcal{L}_{intrinsic} = \frac{1}{2} \partial_\mu T(x) \partial^\mu T(x) - V(T(x)) \quad (12)$$

10 Cosmological Implications

The T0 model has the following implications:

- Modified Gravitational Potential: $\Phi(r) = -\frac{GM}{r} + \kappa r$, $\kappa \approx 4.8 \times 10^{-11} \text{ m/s}^2$
- Cosmic Redshift: $1 + z = e^{\alpha d}$, $\alpha \approx 2.3 \times 10^{-28} \text{ m}^{-1}$
- Wavelength Dependence: $z(\lambda) = z_0(1 + \beta_T \ln(\lambda/\lambda_0))$, $\beta_T \approx 0.008$ (SI units)

11 Derivation of β_T in the T0 Model

The parameter β_T describes the coupling of the intrinsic time field $T(x)$ to physical phenomena such as wavelength-dependent redshift. In the T0 model, β_T is precisely derived as:

$$\beta_T = \frac{\lambda_h^2 v^2}{16\pi^3} \cdot \frac{1}{m_h^2} \cdot \frac{1}{\xi} \quad (13)$$

where λ_h is the Higgs self-coupling, v is the Higgs vacuum expectation value, m_h is the Higgs mass, and $\xi \approx 1.33 \times 10^{-4}$ is a dimensionless parameter defining the characteristic length scale $r_0 = \xi \cdot l_P$ (l_P : Planck length). In natural units, $\beta_T = 1$ holds, representing an exact theoretical prediction derived directly from the model parameters, as detailed in [11]. A comprehensive derivation and discussion of this parameter can be found in [11].

Literatur

- [1] Pascher, J. (2025). [Time as an Emergent Property in Quantum Mechanics: A Connection Between Relativity, Fine-Structure Constant, and Quantum Dynamics](#). March 23, 2025.
- [2] Pascher, J. (2025). [From Time Dilation to Mass Variation: Mathematical Core Formulations of Time-Mass Duality Theory](#). March 29, 2025.
- [3] Pascher, J. (2025). [Dynamic Mass of Photons and Its Implications for Nonlocality in the T0 Model](#).
- [4] Pascher, J. (2025). [The Necessity of Extending Standard Quantum Mechanics and Quantum Field Theory](#). March 27, 2025.
- [5] Pascher, J. (2025). [Mass Variation in Galaxies: An Analysis in the T0 Model with Emergent Gravitation](#). March 30, 2025.
- [6] Pascher, J. (2025). [Mathematical Formulation of the Higgs Mechanism in Time-Mass Duality](#). March 28, 2025.
- [7] Pascher, J. (2025). [Field Theory and Quantum Correlations: A New Perspective on Instantaneity](#). March 28, 2025.
- [8] Pascher, J. (2025). [Compensatory and Additive Effects: An Analysis of Measurement Differences Between the T0 Model and the \$\Lambda\$ CDM Standard Model](#). April 2, 2025.
- [9] Pascher, J. (2025). [Real Consequences of Reformulating Time and Mass in Physics: Beyond the Planck Scale](#). March 24, 2025.
- [10] Pascher, J. (2025). [Energy as a Fundamental Unit: Natural Units with \$\alpha_{EM} = 1\$ in the T0 Model](#). March 26, 2025.
- [11] Pascher, J. (2025). [Unified Unit System in the T0 Model: The Consistency of \$\alpha = 1\$ and \$\beta = 1\$](#) . April 5, 2025.
- [12] Pascher, J. (2025). [Adjustment of Temperature Units in Natural Units and CMB Measurements](#). April 2, 2025.
- [13] Pascher, J. (2025). [Time-Mass Duality Theory \(T0 Model\): Derivation of Parameters \$\kappa\$, \$\alpha\$, and \$\beta\$](#) . April 4, 2025.
- [14] Pascher, J. (2025). [Emergent Gravitation in the T0 Model: A Comprehensive Derivation](#). April 1, 2025.