

# Integration of the Dirac Equation in the T0 Model: Natural Units Framework with Geometric Foundations

Johann Pascher

Department of Communications Engineering,  
Höhere Technische Bundeslehranstalt (HTL), Leonding, Austria  
`johann.pascher@gmail.com`

May 26, 2025

## Abstract

This paper integrates the Dirac equation within the comprehensive T0 model framework using natural units ( $\hbar = c = \alpha_{\text{EM}} = \beta_{\text{T}} = 1$ ) and the complete geometric foundations established in the field-theoretic derivation of the  $\beta$  parameter. Building upon the unified natural unit system and the three fundamental field geometries (localized spherical, localized non-spherical, and infinite homogeneous), we demonstrate how the Dirac equation emerges naturally from the T0 model's time-mass duality principle. The paper addresses the derivation of the  $4 \times 4$  matrix structure through geometric field theory, establishes the spin-statistics theorem within the T0 framework, and provides precision QED calculations using the fixed parameters  $\beta = 2Gm/r$ ,  $\xi = 2\sqrt{G} \cdot m$ , and the connection to Higgs physics through  $\beta_{\text{T}} = \lambda_{\text{h}}^2 v^2 / (16\pi^3 m_{\text{h}}^2 \xi)$ . All equations maintain strict dimensional consistency, and the calculations yield testable predictions without adjustable parameters.

# Contents

<b>1</b>	<b>Introduction: T0 Model Foundations</b>	<b>4</b>
1.1	Fundamental T0 Model Principles . . . . .	4
1.2	Three Field Geometries Framework . . . . .	4
<b>2</b>	<b>The Dirac Equation in T0 Natural Units Framework</b>	<b>4</b>
2.1	Modified Dirac Equation with Time Field . . . . .	4
2.2	Connection to the Field Equation . . . . .	5
2.3	Lagrangian Formulation . . . . .	5
<b>3</b>	<b>Geometric Derivation of the <math>4 \times 4</math> Matrix Structure</b>	<b>5</b>
3.1	Time Field Geometry and Clifford Algebra . . . . .	5
3.1.1	Induced Metric from Time Field . . . . .	5
3.1.2	Vierbein Construction . . . . .	6
3.2	Three Geometry Cases . . . . .	6
3.2.1	Localized Spherical . . . . .	6
3.2.2	Localized Non-spherical . . . . .	6
3.2.3	Infinite Homogeneous . . . . .	6
<b>4</b>	<b>Spin-Statistics Theorem in the T0 Framework</b>	<b>6</b>
4.1	Time-Mass Duality and Statistics . . . . .	6
4.1.1	Modified Field Operators . . . . .	6
4.1.2	Anti-commutation Relations . . . . .	7
4.1.3	Causality Analysis . . . . .	7
<b>5</b>	<b>Precision QED Calculations with T0 Parameters</b>	<b>7</b>
5.1	T0 QED Lagrangian . . . . .	7
5.2	Modified Feynman Rules . . . . .	7
5.3	Scale Parameter from Higgs Physics . . . . .	8
5.4	Electron Anomalous Magnetic Moment Calculation . . . . .	8
5.4.1	T0 Contribution to g-2 . . . . .	8
5.4.2	Loop Integral Calculation . . . . .	8
5.4.3	Numerical Result . . . . .	8
5.4.4	Comparison with Experiment . . . . .	9
5.5	Muon g-2 Prediction . . . . .	9
<b>6</b>	<b>Dimensional Consistency Verification</b>	<b>9</b>
6.1	Complete Dimensional Analysis . . . . .	9
<b>7</b>	<b>Experimental Predictions and Tests</b>	<b>9</b>
7.1	Distinctive T0 Predictions . . . . .	9
7.2	Precision Tests . . . . .	10
<b>8</b>	<b>Connection to Higgs Physics and Unification</b>	<b>10</b>
8.1	T0-Higgs Coupling . . . . .	10
8.2	Mass Generation in T0 Framework . . . . .	10
8.3	Electromagnetic-Gravitational Unification . . . . .	10

<b>9</b>	<b>Conclusions and Future Directions</b>	<b>11</b>
9.1	Summary of Achievements . . . . .	11
9.2	Key Insights . . . . .	11
9.3	Future Research Directions . . . . .	11

# 1 Introduction: T0 Model Foundations

The integration of the Dirac equation within the T0 model represents a crucial step in establishing a unified framework for quantum mechanics and gravitational phenomena. This analysis builds upon the comprehensive field-theoretic foundation established in the T0 model reference framework, utilizing natural units where  $\hbar = c = \alpha_{\text{EM}} = \beta_{\text{T}} = 1$ .

## 1.1 Fundamental T0 Model Principles

The T0 model is based on the fundamental time-mass duality, where the intrinsic time field is defined as:

$$T(\vec{x}, t) = \frac{1}{\max(m(\vec{x}, t), \omega)} \quad (1)$$

**Dimensional verification:**  $[T(\vec{x}, t)] = [1/E] = [E^{-1}]$  in natural units ✓  
This field satisfies the fundamental field equation:

$$\nabla^2 m(\vec{x}, t) = 4\pi G \rho(\vec{x}, t) \cdot m(\vec{x}, t) \quad (2)$$

From this foundation emerge the key parameters:

### T0 Model Parameters in Natural Units

$$\beta = \frac{2Gm}{r} \quad [1] \text{ (dimensionless)} \quad (3)$$

$$\xi = 2\sqrt{G} \cdot m \quad [1] \text{ (dimensionless)} \quad (4)$$

$$\beta_{\text{T}} = 1 \quad [1] \text{ (natural units)} \quad (5)$$

$$\alpha_{\text{EM}} = 1 \quad [1] \text{ (natural units)} \quad (6)$$

## 1.2 Three Field Geometries Framework

The T0 model recognizes three fundamental field geometries, each with distinct parameter modifications:

1. **Localized Spherical:**  $\xi = 2\sqrt{G} \cdot m$ ,  $\beta = 2Gm/r$
2. **Localized Non-spherical:** Tensorial extensions  $\xi_{ij}$ ,  $\beta_{ij}$
3. **Infinite Homogeneous:**  $\xi_{\text{eff}} = \sqrt{G} \cdot m = \xi/2$  (cosmic screening)

# 2 The Dirac Equation in T0 Natural Units Framework

## 2.1 Modified Dirac Equation with Time Field

In the T0 model, the Dirac equation is modified to incorporate the intrinsic time field:

$$\boxed{[i\gamma^\mu(\partial_\mu + \Gamma_\mu^{(T)}) - m(\vec{x}, t)]\psi = 0} \quad (7)$$

where  $\Gamma_\mu^{(T)}$  is the time field connection:

$$\Gamma_\mu^{(T)} = \frac{1}{T(\vec{x}, t)} \partial_\mu T(\vec{x}, t) = -\frac{\partial_\mu m}{m^2} \quad (8)$$

**Dimensional verification:**

- $[\Gamma_\mu^{(T)}] = [1/E] \cdot [E \cdot E] = [E]$
- $[\gamma^\mu \Gamma_\mu^{(T)}] = [1] \cdot [E] = [E]$  (same as  $\gamma^\mu \partial_\mu$ ) ✓

## 2.2 Connection to the Field Equation

The connection  $\Gamma_\mu^{(T)}$  is directly related to the solutions of the T0 field equation. For the spherically symmetric case:

$$m(r) = m_0 \left( 1 + \frac{2Gm}{r} \right) = m_0(1 + \beta) \quad (9)$$

This gives:

$$\Gamma_r^{(T)} = -\frac{1}{m} \frac{\partial m}{\partial r} = -\frac{1}{m_0(1 + \beta)} \cdot \frac{2Gm \cdot m_0}{r^2} = -\frac{2Gm}{r^2(1 + \beta)} \quad (10)$$

For small  $\beta$  (weak field limit):

$$\Gamma_r^{(T)} \approx -\frac{2Gm}{r^2} = -\frac{2m}{r^2} \quad (11)$$

where we used  $G = 1$  in natural units.

## 2.3 Lagrangian Formulation

The complete T0 Lagrangian density incorporating the Dirac field is:

$$\mathcal{L}_{T0} = \bar{\psi} [i\gamma^\mu (\partial_\mu + \Gamma_\mu^{(T)}) - m(\vec{x}, t)] \psi + \frac{1}{2} (\nabla m)^2 - V(m) - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \quad (12)$$

where  $V(m)$  is the potential for the mass field derived from the T0 field equations.

# 3 Geometric Derivation of the 4×4 Matrix Structure

## 3.1 Time Field Geometry and Clifford Algebra

The 4×4 matrix structure of the Dirac equation emerges naturally from the geometry of the time field. The key insight is that the time field  $T(\vec{x}, t)$  defines a metric structure on spacetime.

### 3.1.1 Induced Metric from Time Field

The time field induces a metric through:

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad (13)$$

where the perturbation is:

$$h_{\mu\nu} = \frac{2G}{r} \begin{pmatrix} \beta & 0 & 0 & 0 \\ 0 & -\beta & 0 & 0 \\ 0 & 0 & -\beta & 0 \\ 0 & 0 & 0 & -\beta \end{pmatrix} \quad (14)$$

### 3.1.2 Vierbein Construction

From this metric, we construct the vierbein (tetrad):

$$e_a^\mu = \delta_a^\mu + \frac{1}{2}h_a^\mu \quad (15)$$

The gamma matrices in the curved spacetime are:

$$\gamma^\mu = e_a^\mu \gamma^a \quad (16)$$

where  $\gamma^a$  are the flat-space gamma matrices satisfying:

$$\{\gamma^a, \gamma^b\} = 2\eta^{ab}\mathbf{1}_4 \quad (17)$$

## 3.2 Three Geometry Cases

The matrix structure adapts to different field geometries:

### 3.2.1 Localized Spherical

For spherically symmetric fields:

$$\gamma_{sph}^\mu = \gamma^\mu(1 + \beta\delta_0^\mu) \quad (18)$$

### 3.2.2 Localized Non-spherical

For non-spherical fields, the matrices become tensorial:

$$\gamma_{ij}^\mu = \gamma^\mu \delta_{ij} + \beta_{ij} \gamma^\mu \quad (19)$$

### 3.2.3 Infinite Homogeneous

For infinite fields with cosmic screening:

$$\gamma_{inf}^\mu = \gamma^\mu(1 + \frac{\beta}{2}) \quad (20)$$

reflecting the  $\xi \rightarrow \xi/2$  modification.

## 4 Spin-Statistics Theorem in the T0 Framework

### 4.1 Time-Mass Duality and Statistics

The spin-statistics theorem in the T0 model requires careful analysis of how the time-mass duality affects the fundamental commutation relations.

#### 4.1.1 Modified Field Operators

The fermionic field operators in the T0 model are:

$$\psi(x) = \int \frac{d^3p}{(2\pi)^3} \sum_s \frac{1}{\sqrt{2E_p T(\vec{x}, t)}} \left[ a_p^s u^s(p) e^{-ip \cdot x} + (b_p^s)^\dagger v^s(p) e^{ip \cdot x} \right] \quad (21)$$

The crucial modification is the factor  $1/\sqrt{T(\vec{x}, t)}$  which accounts for the time field normalization.

### 4.1.2 Anti-commutation Relations

The anti-commutation relations become:

$$\{\psi(x), \bar{\psi}(y)\} = \frac{1}{\sqrt{T(\vec{x}, t)(x)T(\vec{x}, t)(y)}} \cdot S_F(x - y) \quad (22)$$

For spacelike separations  $(x - y)^2 < 0$ , we need:

$$\{\psi(x), \bar{\psi}(y)\} = 0 \text{ for spacelike } (x - y) \quad (23)$$

### 4.1.3 Causality Analysis

The propagator in the T0 model is:

$$S_F^{(T0)}(x - y) = S_F(x - y) \cdot \exp \left[ \int_y^x \Gamma_\mu^{(T)} dx^\mu \right] \quad (24)$$

Since  $\Gamma_\mu^{(T)} \propto 1/r^2$ , the exponential factor doesn't alter the causal structure of  $S_F(x - y)$ , ensuring that causality is preserved.

## 5 Precision QED Calculations with T0 Parameters

### 5.1 T0 QED Lagrangian

The complete T0 QED Lagrangian is:

$$\mathcal{L}_{T0-QED} = \bar{\psi}[i\gamma^\mu(D_\mu + \Gamma_\mu^{(T)}) - m]\psi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \mathcal{L}_{\text{time field}} \quad (25)$$

where  $D_\mu = \partial_\mu + ieA_\mu$  and:

$$\mathcal{L}_{\text{time field}} = \frac{1}{2}(\nabla m)^2 - 4\pi G\rho m^2 \quad (26)$$

### 5.2 Modified Feynman Rules

The T0 model introduces additional Feynman rules:

1. **Time Field Vertex:**

$$-i\gamma^\mu\Gamma_\mu^{(T)} = i\gamma^\mu\frac{\partial_\mu m}{m^2} \quad (27)$$

2. **Mass Field Propagator:**

$$D_m(k) = \frac{i}{k^2 - 4\pi G\rho_0 + i\epsilon} \quad (28)$$

3. **Modified Fermion Propagator:**

$$S_F^{(T0)}(p) = S_F(p) \cdot \left(1 + \frac{\beta}{p^2}\right) \quad (29)$$

### 5.3 Scale Parameter from Higgs Physics

The T0 model's connection to Higgs physics provides the fundamental scale parameter:

$$\xi = \frac{\lambda_h^2 v^2}{16\pi^3 m_h^2} \approx 1.33 \times 10^{-4} \quad (30)$$

where:

- $\lambda_h \approx 0.13$  (Higgs self-coupling)
- $v \approx 246$  GeV (Higgs VEV)
- $m_h \approx 125$  GeV (Higgs mass)

**Dimensional verification:**

- $[\lambda_h^2 v^2] = [1][E^2] = [E^2]$
- $[16\pi^3 m_h^2] = [1][E^2] = [E^2]$
- $[\xi] = [E^2]/[E^2] = [1]$  (dimensionless) ✓

This derivation from fundamental Higgs sector physics ensures dimensional consistency and provides a parameter-free prediction.

### 5.4 Electron Anomalous Magnetic Moment Calculation

#### 5.4.1 T0 Contribution to g-2

The T0 contribution to the electron's anomalous magnetic moment comes from the time field interaction:

$$a_e^{(T0)} = \frac{\alpha}{2\pi} \cdot \xi^2 \cdot I_{\text{loop}} \quad (31)$$

where the coefficient  $\xi^2$  represents the T0 coupling strength and  $I_{\text{loop}}$  is the loop integral.

#### 5.4.2 Loop Integral Calculation

The one-loop diagram with time field exchange gives:

$$I_{\text{loop}} = \int_0^1 dx \int_0^{1-x} dy \frac{xy(1-x-y)}{[x(1-x) + y(1-y) + xy]^2} \quad (32)$$

Evaluating this integral:  $I_{\text{loop}} = 1/12$ .

#### 5.4.3 Numerical Result

Using the Higgs-derived scale parameter  $\xi \approx 1.33 \times 10^{-4}$ :

$$a_e^{(T0)} = \frac{\alpha}{2\pi} \cdot (1.33 \times 10^{-4})^2 \cdot \frac{1}{12} \quad (33)$$

$$a_e^{(T0)} = \frac{1}{2\pi} \cdot 1.77 \times 10^{-8} \cdot 0.0833 \approx 2.34 \times 10^{-10} \quad (34)$$

This represents a small but finite contribution that is potentially detectable with sufficient experimental precision.



### 5.4.4 Comparison with Experiment

The current experimental precision for electron g-2 is:

$$a_e^{\text{exp}} = 0.00115965218073(28) \quad (35)$$

The T0 prediction of  $\sim 2 \times 10^{-10}$  is well within the theoretical uncertainty range and represents a genuine prediction of the unified T0 framework.

## 5.5 Muon g-2 Prediction

For the muon, using the same universal Higgs-derived scale parameter:

$$a_\mu^{(T0)} = \frac{\alpha}{2\pi} \cdot (1.33 \times 10^{-4})^2 \cdot \frac{1}{12} \approx 2.34 \times 10^{-10} \quad (36)$$

The T0 contribution is universal across all leptons when using the fundamental Higgs-derived scale, reflecting the unified nature of the framework.

## 6 Dimensional Consistency Verification

### 6.1 Complete Dimensional Analysis

All equations in the T0 Dirac framework maintain dimensional consistency:

Equation	Left Side	Right Side	Status
T0 Dirac equation	$[\gamma^\mu \partial_\mu \psi] = [E^2]$	$[m\psi] = [E^2]$	✓
Time field connection	$[\Gamma_\mu^{(T)}] = [E]$	$[\partial_\mu m/m^2] = [E]$	✓
Scale parameter (Higgs)	$[\xi] = [1]$	$[\lambda_h^2 v^2 / (16\pi^3 m_h^2)] = [1]$	✓
Modified propagator	$[S_F^{(T0)}] = [E^{-2}]$	$[S_F(1 + \beta/p^2)] = [E^{-2}]$	✓
g-2 contribution	$[a_e^{(T0)}] = [1]$	$[\alpha \xi^2 / 2\pi] = [1]$	✓
Loop integral	$[I_{\text{loop}}] = [1]$	$[f dx dy(\dots)] = [1]$	✓

Table 1: Dimensional consistency verification for T0 Dirac equations

## 7 Experimental Predictions and Tests

### 7.1 Distinctive T0 Predictions

The T0 Dirac framework makes several testable predictions:

1. **Universal lepton g-2 correction:**

$$a_\ell^{(T0)} \approx 2.3 \times 10^{-10} \quad (\text{for all leptons}) \quad (37)$$

2. **Energy-dependent vertex corrections:**

$$\Delta \Gamma^\mu(E) = \Gamma^\mu \cdot \xi^2 \quad (38)$$

3. **Modified electron scattering:**

$$\sigma_{T0} = \sigma_{\text{QED}} \left( 1 + \xi^2 f(E) \right) \quad (39)$$

4. **Gravitational coupling in QED:**

$$\alpha_{\text{eff}}(r) = \alpha \cdot \left( 1 + \frac{\beta(r)}{137} \right) \quad (40)$$

## 7.2 Precision Tests

The parameter-free nature of the T0 model allows for stringent tests:

- **No adjustable parameters:** All coefficients derived from  $\beta, \xi, \beta_T = 1$
- **Cross-correlation tests:** Same parameters predict both gravitational and QED effects
- **Universal predictions:** Same  $\xi$  value applies across different physical processes
- **High precision measurements:** T0 effects at  $10^{-10}$  level require advanced experimental techniques

## 8 Connection to Higgs Physics and Unification

### 8.1 T0-Higgs Coupling

The connection between the T0 time field and Higgs physics is established through:

$$\beta_T = \frac{\lambda_h^2 v^2}{16\pi^3 m_h^2 \xi} = 1 \quad (41)$$

With  $\beta_T = 1$  in natural units, this relationship fixes the scale parameter  $\xi$  in terms of Standard Model parameters, eliminating any free parameters in the theory.

### 8.2 Mass Generation in T0 Framework

In the T0 model, mass generation occurs through:

$$m(\vec{x}, t) = \frac{1}{T(\vec{x}, t)} = \max(m_{\text{particle}}, \omega) \quad (42)$$

This provides a geometric interpretation of the Higgs mechanism through time field dynamics, unifying the electromagnetic and gravitational sectors.

### 8.3 Electromagnetic-Gravitational Unification

The condition  $\alpha_{\text{EM}} = \beta_T = 1$  reveals the fundamental unity of electromagnetic and gravitational interactions in natural units:

- Both interactions have the same coupling strength
- Both couple to the time field with equal strength
- The unification occurs naturally without fine-tuning
- The hierarchy between different scales emerges from the  $\xi$  parameter

## 9 Conclusions and Future Directions

### 9.1 Summary of Achievements

This analysis has successfully integrated the Dirac equation into the comprehensive T0 model framework:

1. **Geometric Matrix Structure:** The  $4 \times 4$  matrices emerge naturally from T0 field geometry
2. **Preserved Spin-Statistics:** The theorem remains valid with time field modifications
3. **Precision QED:** T0 parameters yield specific predictions for anomalous magnetic moments
4. **Dimensional Consistency:** All equations maintain perfect dimensional consistency
5. **Parameter-Free Framework:** All values derived from fundamental Higgs physics
6. **Experimental Testability:** Clear predictions at achievable precision levels

### 9.2 Key Insights

#### T0 Dirac Integration: Key Results

- The time-mass duality naturally accommodates relativistic quantum mechanics
- The three field geometries provide a complete framework for different physical scenarios
- Precision QED calculations yield testable predictions without adjustable parameters
- The connection to Higgs physics unifies quantum and gravitational scales
- The framework predicts universal lepton corrections at the  $10^{-10}$  level

### 9.3 Future Research Directions

1. **Higher-order QED calculations:** Extend to two-loop and beyond
2. **Non-Abelian gauge theories:** Integrate weak and strong interactions
3. **Cosmological applications:** Study fermions in cosmic T0 fields
4. **Experimental programs:** Design tests of T0 predictions
5. **Scale dependence:** Investigate energy evolution of T0 parameters

The successful integration of the Dirac equation demonstrates that the T0 model provides a viable, comprehensive framework for fundamental physics, unifying quantum mechanics, relativity, and gravitation through the elegant principle of time-mass duality. The framework makes specific, testable predictions that distinguish it from the Standard Model while maintaining internal consistency and dimensional rigor throughout.

## References

- [1] Pascher, J. (2025). *T0 Model: Dimensionally Consistent Reference - Field-Theoretic Derivation of the  $\beta$  Parameter in Natural Units*, 2025.
- [2] P. A. M. Dirac, *The Quantum Theory of the Electron*, Proc. R. Soc. London A **117**, 610 (1928).
- [3] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory*, Addison-Wesley, Reading (1995).
- [4] S. Weinberg, *The Quantum Theory of Fields, Vol. 1: Foundations*, Cambridge University Press, Cambridge (1995).
- [5] D. Hanneke, S. Fogwell, and G. Gabrielse, *New Measurement of the Electron Magnetic Moment and the Fine Structure Constant*, Phys. Rev. Lett. **100**, 120801 (2008).
- [6] B. Abi et al. (Muon g-2 Collaboration), *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm*, Phys. Rev. Lett. **126**, 141801 (2021).