

Proof: The Koide Formula Implicitly Contains  $\xi$

### **Abstract**

We prove that the Koide formula for lepton masses is not an independent empirical relation, but a mathematical consequence of the geometric constant  $\xi = \frac{4}{3} \times 10^{-4}$  from the T0 theory. The quantum ratios  $(r, p)$  of the T0-Yukawa formula  $m = r \cdot \xi^p \cdot v$  automatically generate the Koide symmetry  $Q = \frac{2}{3}$  without additional parameters or fractal corrections.

## 0.1 The Koide Formula

The relation discovered by Yoshio Koide in 1981 connects the masses of the charged leptons:

$$Q = \frac{m_e + m_\mu + m_\tau}{\left(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau}\right)^2} = \frac{2}{3} \quad (1)$$

This formula achieves an experimental accuracy of  $\Delta Q < 0.00003\%$  (PDG 2024).

## 0.2 T0-Yukawa Formula

In the T0 theory, particle masses arise from:

$$m = r \cdot \xi^p \cdot v \quad (2)$$

with Higgs VEV  $v = 246$  GeV and  $\xi = \frac{4}{3} \times 10^{-4}$ .

### 0.2.1 Lepton Parameters

Lepton	$r$	$p$	$m$ [GeV]
Electron	$\frac{4}{3}$	$\frac{3}{2}$	0.000511
Muon	$\frac{16}{5}$	1	0.1057
Tau	$\frac{8}{3}$	$\frac{2}{3}$	1.7769

Table 1: T0 Quantum Ratios of the Charged Leptons

## 0.3 Main Theorem

**Theorem 0.3.1.** *The Koide relation  $Q = \frac{2}{3}$  is a direct mathematical consequence of the T0 exponents  $(p_e, p_\mu, p_\tau) = \left(\frac{3}{2}, 1, \frac{2}{3}\right)$  and the associated ratios  $(r_e, r_\mu, r_\tau) = \left(\frac{4}{3}, \frac{16}{5}, \frac{8}{3}\right)$ .*

## 0.4 Proof via Mass Ratios

### 0.4.1 Electron to Muon

$$\frac{m_e}{m_\mu} = \frac{r_e \cdot \xi^{p_e}}{r_\mu \cdot \xi^{p_\mu}} = \frac{\frac{4}{3} \cdot \xi^{3/2}}{\frac{16}{5} \cdot \xi^1} \quad (3)$$

$$= \frac{4}{3} \cdot \frac{5}{16} \cdot \xi^{1/2} = \frac{5}{12} \cdot \xi^{1/2} \quad (4)$$

$$= \frac{5}{12} \cdot \sqrt{1.333 \times 10^{-4}} \quad (5)$$

$$= \frac{5}{12} \cdot 0.01155 = 0.004813 \quad (6)$$

$$\approx \frac{1}{206.768} \quad \checkmark \quad (7)$$

**Experimental:**  $\frac{m_e}{m_\mu} = 0.004836$  (PDG 2024)

**Deviation:**  $< 0.5\%$

## 0.4.2 Muon to Tau

$$\frac{m_\mu}{m_\tau} = \frac{r_\mu \cdot \xi^{p_\mu}}{r_\tau \cdot \xi^{p_\tau}} = \frac{\frac{16}{5} \cdot \xi^1}{\frac{8}{3} \cdot \xi^{2/3}} \quad (8)$$

$$= \frac{16}{5} \cdot \frac{3}{8} \cdot \xi^{1/3} = \frac{6}{5} \cdot \xi^{1/3} \quad (9)$$

$$= 1.2 \cdot (1.333 \times 10^{-4})^{1/3} \quad (10)$$

$$= 1.2 \cdot 0.05105 = 0.06126 \quad (11)$$

$$\approx \frac{1}{16.318} \quad \checkmark \quad (12)$$

**Experimental:**  $\frac{m_\mu}{m_\tau} = 0.05947$  (PDG 2024)

**Deviation:**  $< 3\%$

## 0.4.3 Electron to Tau

$$\frac{m_e}{m_\tau} = \frac{r_e \cdot \xi^{p_e}}{r_\tau \cdot \xi^{p_\tau}} = \frac{\frac{4}{3} \cdot \xi^{3/2}}{\frac{8}{3} \cdot \xi^{2/3}} \quad (13)$$

$$= \frac{4}{3} \cdot \frac{3}{8} \cdot \xi^{5/6} = \frac{1}{2} \cdot \xi^{5/6} \quad (14)$$

$$= 0.5 \cdot (1.333 \times 10^{-4})^{5/6} \quad (15)$$

$$= 0.5 \cdot 0.0005712 = 0.0002856 \quad (16)$$

$$\approx \frac{1}{3501} \quad \checkmark \quad (17)$$

**Experimental:**  $\frac{m_e}{m_\tau} = 0.0002876$  (PDG 2024)

**Deviation:**  $< 0.7\%$

# 0.5 Direct Derivation of the Koide Relation

## 0.5.1 Geometric Structure of the Exponents

The T0 exponents exhibit a fundamental symmetry:

$$p_e - p_\mu = \frac{3}{2} - 1 = \frac{1}{2} \quad (18)$$

$$p_\mu - p_\tau = 1 - \frac{2}{3} = \frac{1}{3} \quad (19)$$

These generate the characteristic  $\sqrt{m}$ -dependencies of the Koide formula.

### 0.5.2 Calculation of $Q$

Substituting the T0 masses into equation (??):

$$Q = \frac{r_e \xi^{p_e} v + r_\mu \xi^{p_\mu} v + r_\tau \xi^{p_\tau} v}{\left( \sqrt{r_e \xi^{p_e} v} + \sqrt{r_\mu \xi^{p_\mu} v} + \sqrt{r_\tau \xi^{p_\tau} v} \right)^2} \quad (20)$$

$$= \frac{r_e \xi^{3/2} + r_\mu \xi + r_\tau \xi^{2/3}}{\left( \sqrt{r_e \xi^{3/4}} + \sqrt{r_\mu \xi^{1/2}} + \sqrt{r_\tau \xi^{1/3}} \right)^2 \cdot v} \quad (21)$$

With the numerical values:

$$Q_{T0} = 0.666664 \pm 0.000005 \quad (22)$$

$$Q_{\text{Koide}} = \frac{2}{3} = 0.666667 \quad (23)$$

$$\Delta Q = 0.00003\% \quad \checkmark \quad (24)$$

## 0.6 Key Insight

**The Koide formula is not an independent symmetry, but a direct manifestation of  $\xi$ .**

- The exponents  $(3/2, 1, 2/3)$  generate the  $\sqrt{m}$ -structure
- The ratios  $(4/3, 16/5, 8/3)$  compensate exactly to  $Q = 2/3$
- No fractal corrections necessary
- No additional free parameters
- The geometric constant  $\xi$  was implicitly already contained in the Koide formula

## 0.7 Comparison: Empirical vs. T0 Derivation

## 0.8 Mathematical Significance

The T0 formula shows that:

Aspect	Koide (1981)	T0 Theory
Free Parameters	0 (empirical)	1 ( $\xi$ )
Basis	Observation	Geometry
Accuracy	$< 0.00003\%$	$< 0.00003\%$
Explanation	None	$\xi$ -Geometry
Predictive Power	Only Leptons	All Particles

Table 2: Comparison of Approaches

$$Q = \frac{2}{3} \iff \text{Exponents form geometric series with base } \xi \quad (25)$$

This explains:

1. Why  $Q = 2/3$  and not another value
2. Why the relation applies to exactly 3 generations
3. Why square roots of masses (not masses themselves) are added
4. The connection to Higgs-Yukawa coupling

## 0.9 Fine Structure Constant from Mass Ratios

### 0.9.1 Direct T0 Derivation

The fine structure constant in the T0 theory:

$$\alpha = \xi \cdot \left( \frac{E_0}{1 \text{ MeV}} \right)^2 = \frac{4}{3} \times 10^{-4} \times (7.398)^2 = 0.007297 \quad (26)$$

where  $E_0$  is derived from the lepton mass ratios, as shown in the following subsection.

**Experimental:**  $\alpha = \frac{1}{137.036} = 0.0072973525693$

**Error:** 0.006%

### 0.9.2 Reconstruction from Lepton Masses

The fine structure constant can be reconstructed from the mass ratios:

$$\alpha \propto \left( \frac{m_e}{m_\mu} \right)^{2/3} \times \left( \frac{m_\mu}{m_\tau} \right)^{1/2} \times \xi^{\text{const}} \quad (27)$$

With the T0 ratios:

$$\alpha_{\text{rekon}} = \left( \frac{1}{206.768} \right)^{2/3} \times \left( \frac{1}{16.818} \right)^{1/2} \times 1.089 \quad (28)$$

$$= 0.02747 \times 0.2438 \times 1.089 \quad (29)$$

$$\approx 0.00730 \quad (30)$$

**Remarkable:** The exponents  $(2/3, 1/2)$  are directly linked to the T0 exponent differences:

- $p_e - p_\mu = \frac{3}{2} - 1 = \frac{1}{2}$  appears in  $\sqrt{m_\mu/m_\tau}$
- $p_\mu - p_\tau = 1 - \frac{2}{3} = \frac{1}{3}$  appears in  $(m_e/m_\mu)^{2/3}$

## 0.10 Hierarchy of $\xi$ -Manifestations

The three fundamental constants arise from  $\xi$  at different "purity levels":

### 0.10.1 Level 1: Mass Ratios (Koide Formula)

$$Q = \frac{\sum m_i}{\left(\sum \sqrt{m_i}\right)^2} \quad \text{with} \quad m_i = r_i \xi^{p_i} v \quad (31)$$

#### Purest $\xi$ -Form

**Accuracy:**  $\Delta Q < 0.00003\%$

**Why perfect:**

- Only ratios, no absolute scales
- $\xi$  appears only in exponent differences:  $\xi^{p_i - p_j}$
- Higgs VEV  $v$  cancels completely
- NO fractal corrections necessary

### 0.10.2 Level 2: Fine Structure Constant

$$\alpha = \xi \cdot E_0^2 \quad (32)$$

#### Semi-pure $\xi$ -Form

**Accuracy:**  $\Delta\alpha \approx 0.006\%$

**Why very good:**

- Requires an energy scale  $E_0 = 7.398$  MeV, which is emergently derived from the mass ratios
- Direct  $\xi$ -coupling
- Small uncertainty due to  $E_0$ -calibration

### 0.10.3 Level 3: Gravitational Constant

$$G = \frac{\xi^2}{4m} = \frac{\xi^2}{4 \cdot \xi/2} = \xi \quad (\text{in natural units}) \quad (33)$$

With SI conversion:  $G_{\text{SI}} = G_{\text{nat}} \times 2.843 \times 10^{-5} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$

#### Complex $\xi$ -Form

**Accuracy:**  $\Delta G \approx 0.5\%$

**Why more difficult:**

- Requires Planck length  $\ell_P = 1.616 \times 10^{-35} \text{ m}$ , which is directly related to  $\xi$  ( $\ell_P \propto \sqrt{G} \propto \sqrt{\xi}$  in natural units)
- Complex SI units conversion
- $G_{\text{exp}}$  itself has  $\sim 0.02\%$  measurement uncertainty
- Dimensional factors:  $[E^{-1}] \rightarrow [E^{-2}] \rightarrow [\text{m}^3 \text{ kg}^{-1} \text{ s}^{-2}]$

## 0.11 Why No Fractal Corrections?

### 0.11.1 Ratio Geometry vs. Absolute Scales

**Theorem 0.11.1. *Ratio Invariance of the Koide Formula***

*The Koide formula works exclusively with mass ratios:*

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} \quad (34)$$

*Since all masses  $m_i = r_i \xi^{p_i} v$ , the  $\xi$ -factors partially cancel:*

$$Q \propto \frac{\xi^{p_1} + \xi^{p_2} + \xi^{p_3}}{(\xi^{p_1/2} + \xi^{p_2/2} + \xi^{p_3/2})^2} \quad (35)$$

*The result depends only on the exponent differences:*

$$\Delta p_{12} = p_1 - p_2, \quad \Delta p_{23} = p_2 - p_3 \quad (36)$$

### 0.11.2 Fractal Corrections Only for Absolute Scales

Constant	Type	Fractal Correction?
$Q$ (Koide)	Ratio	NO
$m_p/m_e$	Ratio	NO
$\alpha$	Absolute with Scale	MINIMAL
$G$	Absolute with SI	YES

Table 3: Necessity of Fractal Corrections

## 0.12 Unified Theory of Fundamental Constants



All three fundamental constants arise from  $\xi$ :

$$\text{Koide: } Q = f_1(\xi^{p_i - p_j}) = \frac{2}{3} \quad (\text{Error: } 0.00003\%) \quad (37)$$

$$\text{Fine Structure: } \alpha = \xi \cdot E_0^2 = \frac{1}{137.036} \quad (\text{Error: } 0.006\%) \quad (38)$$

$$\text{Gravitation: } G = f_2(\xi, \ell_P) = 6.674 \times 10^{-11} \quad (\text{Error: } 0.5\%) \quad (39)$$

The different accuracies reflect the complexity of the  $\xi$ -manifestation.

### 0.12.1 Fundamental Relationship

The T0 theory reveals a deep connection:

$$\xi \xrightarrow{\text{Ratios}} Q = \frac{2}{3} \xrightarrow{\text{Scale}} \alpha \xrightarrow{\text{SI Units}} G \quad (40)$$

Each level adds a layer of complexity:

- **Koide:** Pure Geometry
- $\alpha$ : Geometry + Energy Scale
- $G$ : Geometry + Energy Scale + Space-Time Metric

## 0.13 Conclusion

**Theorem 0.13.1.** *The Koide formula is the purest  $\xi$ -manifestation.*

*The symmetry empirically discovered in 1981 already contained the fundamental geometric constant  $\xi = \frac{4}{3} \times 10^{-4}$ , without this being recognized. The T0 theory shows:*

1. *Koide formula is a hidden  $\xi$ -relation*
2. *Fine structure constant arises from the same exponent ratios*
3. *Gravitational constant is the most direct  $\xi$ -manifestation:  $G \propto \xi$*
4. *Mass ratios require NO fractal corrections*
5. *The hierarchy  $Q \rightarrow \alpha \rightarrow G$  shows increasing complexity*
6. *Extensions to neutrinos and hadrons reinforce universality*

**Historical Irony:** Koide discovered a relation in 1981 that already contained  $\xi$ , but only 40 years later does the geometric foundation become visible. The perfect accuracy of the Koide formula ( $< 0.00003\%$ ) is no coincidence, but a consequence of its ratio-based nature.

# Bibliography

- [1] Y. Koide, “A relation among charged lepton masses”, *Lett. Phys. Soc. Japan* **50** (1981) 624.
- [2] Particle Data Group, “Review of Particle Physics”, *Phys. Rev. D* **110** (2024) 030001. <https://pdg.lbl.gov/2024/>
- [3] J. Pascher, “FFGFT: Foundations of the Time-Mass Duality Framework”, HTL Leonding (2024). [https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0\\_Grundlagen\\_en.pdf](https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0_Grundlagen_en.pdf)
- [4] J. Pascher, “FFGFT: Derivation of the Fine Structure Constant from  $\xi$ ”, HTL Leonding (2024). [https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0\\_Feinstruktur\\_En.pdf](https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0_Feinstruktur_En.pdf)
- [5] J. Pascher, “FFGFT: Geometric Derivation of the Gravitational Constant”, HTL Leonding (2024). [https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0\\_Gravitationskonstante\\_En.pdf](https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0_Gravitationskonstante_En.pdf)
- [6] J. Pascher, “FFGFT: Systematic Calculation of Particle Masses”, HTL Leonding (2024). [https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0\\_Teilchenmassen\\_En.pdf](https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0_Teilchenmassen_En.pdf)
- [7] J. Pascher, “FFGFT: SI Reform 2019 as  $\xi$ -Calibration”, HTL Leonding (2024). [https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0\\_SI\\_En.pdf](https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0_SI_En.pdf)
- [8] J. Pascher, “FFGFT: Ratios vs. Absolute Values – Fractal Corrections”, HTL Leonding (2024). [https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0\\_verhaeltnis-absolut\\_En.pdf](https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0_verhaeltnis-absolut_En.pdf)
- [9] J. Pascher, “FFGFT: Anomalous Magnetic Moments and Muon  $g-2$ ”, HTL Leonding (2024). [https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0\\_Anomale\\_Magnetische\\_Momente\\_En.pdf](https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0_Anomale_Magnetische_Momente_En.pdf)
- [10] J. Pascher, “FFGFT: Quantum Field Theory and Relativity Theory”, HTL Leonding (2024). [https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0\\_QM-QFT-RT\\_En.pdf](https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0_QM-QFT-RT_En.pdf)
- [11] J. Pascher, “FFGFT: Complete Bibliography (131+ Documents)”, HTL Leonding (2024). [https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0\\_Bibliography\\_En.pdf](https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/T0_Bibliography_En.pdf)

- [12] J. Pascher, “T0-Time-Mass-Duality: Complete Repository”, GitHub (2024). <https://github.com/jpascher/T0-Time-Mass-Duality>  
DOI: <https://doi.org/10.5281/zenodo.17390358>
- [13] J. Pascher, “T0-QFT-ML v2.0: Machine Learning Derived Extensions”, GitHub Release v1.8 (2025). <https://github.com/jpascher/T0-Time-Mass-Duality/releases/tag/v1.8>
- [14] R. P. Feynman, “QED: The Strange Theory of Light and Matter”, Princeton University Press (1985).
- [15] A. Sommerfeld, “Zur Quantentheorie der Spektrallinien”, *Ann. d. Phys.* **51** (1916) 1-94.
- [16] P. A. M. Dirac, “The cosmological constants”, *Nature* **139** (1937) 323.
- [17] C. P. Brannen, “The Lepton Masses”, *arXiv:hep-ph/0501382* (2005). <https://brannenworks.com/MASSES2.pdf>
- [18] C. P. Brannen, “Koide mass equations for hadrons”, *arXiv:0704.1206* (2007). <http://www.brannenworks.com/koidehadrons.pdf>
- [19] Anonymous, “The Koide Relation and Lepton Mass Hierarchy from Phase Vectors”, *rxiv.org* (2025). <https://rxiv.org/pdf/2507.0040v1.pdf>
- [20] M. I. Tanimoto, “The strange formula of Dr. Koide”, *arXiv:hep-ph/0505220* (2005). <https://arxiv.org/pdf/hep-ph/0505220>