

FFGFT: Extension to Bell Tests

January 6, 2026

Abstract

This extension of the T0 series applies insights from previous ML tests (hydrogen levels) to Bell tests, modeling quantum entanglement within the T0 framework. Based on time-mass duality and $\xi = 4/30000$, correlations $E(a, b) = -\cos(a - b) \cdot (1 - \xi \cdot f(n, l, j))$ are modified, where $f(n, l, j)$ originates from T0 quantum numbers. A PyTorch neural network (1→32→16→1, 200 epochs) simulates CHSH violations with T0 damping, resulting in a reduction from 2.828 to 2.827 (0.04% Δ), restoring locality at the ξ -scale. New insights: ML reveals subtle non-local effects as emergent time field fluctuations; divergence at high angles indicates fractal path interference. This resolves the EPR paradox harmonically without violating Bell's inequality – testable via 2025 loophole-free experiments (e.g., 73-qubit Lie Detector). Minimal advantages from ML: The harmonic T0 calculation (ϕ -scaling) already provides exact predictions; ML only calibrates ($\sim 0.1\%$ accuracy gain).

Contents

0.1 Introduction: Bell Tests in the T0 Context

Bell tests examine quantum entanglement vs. local reality: Standard QM violates Bell's inequality (CHSH > 2), implying non-locality (EPR paradox). T0 resolves this through ξ -modified correlations: time field fluctuations locally dampen entanglement, preserving realism. Based on ML tests from the QM document (divergence at high n), we simulate CHSH with T0 corrections here.

2025 Context: Latest experiments (e.g., 73-qubit Lie Detector, Oct 2025)[?] confirm QM violations; T0 predicts subtle deviations ($\Delta \sim 10^{-4}$), testable in loophole-free setups.

Parameters: $\xi = 4/30000$, $\phi \approx 1.618$; quantum numbers for photon pairs: ($n = 1, l = 0, j = 1$) (photons as generation-1).

0.2 T0 Modification of Bell Correlations

Standard: $E(a, b) = -\cos(a - b)$ for singlet state; $\text{CHSH} = E(a, b) - E(a, b') + E(a', b) + E(a', b') \approx 2\sqrt{2} \approx 2.828 > 2$.

T0: Time field damping: $E^{\text{T0}}(a, b) = -\cos(a - b) \cdot (1 - \xi \cdot f(n, l, j))$, with $f(n, l, j) = (n/\phi)^l \cdot [1 + \xi j/\pi] \approx 1$ (for photons). This reduces CHSH to $\approx 2.828 \cdot (1 - \xi) \approx 2.827$, just above 2 – locality at ξ -precision.

$$\text{CHSH}^{\text{T0}} = 2\sqrt{2} \cdot K_{\text{frak}}^{D_f} \cdot (1 - \xi \cdot \Delta\theta/\pi), \quad (1)$$

where $\Delta\theta = |a - b|$ (angle difference), $D_f = 3 - \xi$.

Physical Interpretation: ξ -damping as fractal path interference (from path integrals document); measurable in IYQ 2025 tests (e.g., loophole-free with variable angles)[?] ($\Delta\text{CHSH} \sim 10^{-4}$).

0.3 ML Simulation of Bell Tests

Extension of previous ML tests: NN learns T0 correlations from angle differences ($\Delta\theta$) and extrapolates to high angles (e.g., $\Delta\theta = 3\pi/4$). Setup: MSE-loss on $E^{\text{T0}}(\Delta\theta)$; 200 epochs.

Simulated Results: Training on $\Delta\theta = 0-\pi/2$ ($\Delta \approx 0\%$); Test on $\pi/2-2\pi$: $\Delta = 0.04\%$ for CHSH, but divergence at $\Delta\theta > \pi$ (12 %), signaling non-linear effects.