

The Complete Field Theory of FFGFT

Johann Pascher

2026

Contents

I Foundations and Motivation	4
1 What is FFGFT?	4
2 The Fundamental Scale of the Torus	4
3 The Fractal Dimension	5
II The Torus Modes and Their Quantum Numbers	5
4 Quantum Numbers of the Modes	5
5 Explicit Mode Functions	6
III The Non-Closure Theorem	7
6 Algebraic Structure of the Modes	7
7 The Theorem	8

IV The Lagrangian and the Bridge to the Mode Operator	8
8 The Free Lagrangian	9
9 The Torus Correction Term	9
10The Mode Operator	10
11The Bridge Formula	10
V Mass Generation as a Binding Phenomenon	10
12The Binding Picture	11
13Not Fine-Tuning — a Binding Phenomenon	11
14Connection to the Lagrangian Extension	12
15The Verification Condition	12
VI The Field-Theoretic Formulation	13
16Hilbert Space and Recursion Operator	13
17Algebraic Structure	13
18The Renormalisation Group	13
VII Open Mathematical Tasks	14
19Four Precisely Formulated Problems	14
VIII Summary	14

20 Overview of Results

15

Part I

Foundations and Motivation

1 What is FFGFT?

FFGFT (Fundamental Fractal Geometric Field Theory), also called T0 Time-Mass Duality, describes all known elementary particles as bound oscillation modes of a **universal fractal torus**. The central idea: observed masses and interactions follow directly from geometry, without arbitrary parameters.

The theory requires only two fundamental inputs:

- **The geometric constant** $\xi_0 = \frac{4}{3} \times 10^{-4}$ — not postulated but derived from the tetrahedral symmetry of the 4D torus and the self-consistency of the electron-proton-muon system (Doc. 180).
- **The Planck length** ℓ_P — the natural length scale of quantum gravity.

All physical quantities — particle masses, coupling constants, cosmological parameters — follow from these two quantities.

2 The Fundamental Scale of the Torus

The characteristic length scale of the universal torus:

$$L_0 = \frac{\ell_P}{\xi_0} \approx 7500 \cdot \ell_P \approx 6.14 \times 10^{-16} \text{ GeV}^{-1} \quad (1)$$

This is the **fundamental period** of the universal torus. All particles are modes on this torus — like overtones of a vibrating string, but in three spatial dimensions. The Compton wavelength $L_i = 2\pi/m_i$ is the wave-

length of the respective mode, not the size of a separate torus.

3 The Fractal Dimension

An ordinary torus has dimension 3. The FFGFT torus has a **fractal correction**:

$$D_f = 3 - \xi_0 \approx 2.9998667 \quad (2)$$

This minimal deviation is the origin of all particle masses. It generates a **non-linear dispersion relation** — analogous to an anharmonic crystal lattice. On a simple torus ($D_f = 3$) masses would be proportional to wave number: $m_n \propto n$, ratios integer. In reality:

$$\frac{m_\mu}{m_e} \approx 206.8, \quad \frac{m_\tau}{m_e} \approx 3477$$

These non-integer ratios require the fractal dispersion.

Part II

The Torus Modes and Their Quantum Numbers

4 Quantum Numbers of the Modes

Each mode on the torus is characterised by three quantum numbers: n (principal), l (orbital angular momentum, transverse component) and j (spin, $j = l \pm \frac{1}{2}$ for fermions).

The wave vector of a mode is quantised:

$$\mathbf{k}_{n,l} = \frac{2\pi}{L_0}(k_x, k_y, k_z), \quad k_x, k_y, k_z \in \mathbb{Z} \quad (3)$$

Connection to the quantum numbers:

$$n = k_x^2 + k_y^2 + k_z^2 \quad (4)$$

(principal = squared total length)

$$l = \sqrt{k_x^2 + k_y^2} \quad (5)$$

(orbital = transverse component)

The r_i prefactors and p_i exponents are discrete winding ratios of the 4D torus — not continuous parameters. For the leptons:

Particle	n	l	(k_x, k_y, k_z)	r_i	p_i
Electron	1	0	(1, 0, 0)	4/3	3/2
Muon	2	1	(1, 1, 0)	16/5	1
Tau	3	2	(1, 1, 1)	25/9	2/3

These values are the complete enumeration of discrete torus modes — analogous to the integers in hydrogen, where one does not ask for a formula for $n = 1, 2, 3$. The table (Doc. 006) is the theory.

5 Explicit Mode Functions

On the torus $T^3 \times \mathbb{R}$, the eigenfunctions of the d'Alembertian $\square = \partial_t^2 - \nabla^2$ are:

$$\phi_{n,l,j}(t, \mathbf{x}) = \frac{1}{\sqrt{V}} e^{i\mathbf{k}_{n,l} \cdot \mathbf{x}} \cdot Y_{l,j}(\theta, \varphi) \cdot e^{-im_{n,l,j} t} \quad (6)$$

with $V = L_0^3$ (torus volume) and spherical harmonics $Y_{l,j}(\theta, \varphi)$.

These functions are **orthonormal**:

$$\int_{T^3} \phi_{n,l,j}^*(\mathbf{x}) \phi_{n',l',j'}(\mathbf{x}) d^3x = \delta_{nn'} \delta_{ll'} \delta_{jj'} \quad (7)$$

and satisfy the **Klein-Gordon equation**:

$$\square \phi_{n,l,j} = -m_{n,l,j}^2 \phi_{n,l,j} \quad (8)$$

The spin structure follows from the winding numbers of the torus (Doc. 051): spin- s corresponds to torus winding (n_θ, n_ϕ) with $n_\phi/n_\theta = 2s$. For spin- $\frac{1}{2}$: $(1,1)$; the 720° symmetry follows directly from the topology.

Part III

The Non-Closure Theorem

6 Algebraic Structure of the Modes

The particle masses follow the geometric mass formula:

$$m_{n,l,j} = r_{n,l,j} \cdot \xi_0^{p_{n,l,j}} \cdot v, \quad v = 246 \text{ GeV} \quad (9)$$

The prefactors r_i are rational numbers (labels of torus modes); the exponents p_i are rational scale parameters.

Restriction: Numerical substitution is permitted. Symbolic cross-multiplication is forbidden.

7 The Theorem

Non-Closure Theorem of FFGFT Torus Modes

For all pairs of different torus modes $i \neq j$:

$$\forall i \neq j: (r_i \cdot r_j, p_i + p_j) \notin \mathcal{P}$$

where $\mathcal{P}$ is the set of all admissible pairs (r, p) in FFGFT. The set $\mathcal{P}$ is **not closed** under cross-operations between different torus modes.

Proof of violation (example):

$$r_e \cdot r_\mu = \frac{4}{3} \cdot \frac{16}{5} = \frac{64}{15} \approx 4.267$$

$$p_e + p_\mu = \frac{3}{2} + 1 = \frac{5}{2}$$

In the complete mode table (Doc. 006), no particle has $r = 64/15$ and $p = 5/2$. The pair $(64/15, 5/2) \notin \mathcal{P}$.

Why does the theorem hold? The cause lies in the orthogonality of the modes. In Hilbert space, the product of two different basis vectors is structurally meaningless. Likewise, $r_e \cdot r_\mu$ is computable as a number but belongs to no torus mode.

Practical rule:

Permitted	Forbidden
Numerical: $m_e \cdot m_\mu \approx 53.9 \text{ MeV}^2$	Symbolic: $r_e \cdot r_\mu = 64/15$ as mass formula
$m_i = r_i \cdot \xi_0^{p_i} \cdot v$ for fixed i	$(r_i \cdot r_j) \cdot \xi_0^{p_i + p_j} \cdot v$ for $i \neq j$
Mass ratios m_i/m_j numerically	Cross-multiplication of different r_i, r_j

Part IV

The Lagrangian and the Bridge to the Mode Operator

8 The Free Lagrangian

Starting point is the free Klein-Gordon Lagrangian:

$$\mathcal{L}_0 = \frac{1}{2}(\partial\delta m)^2 \quad (10)$$

The scalar field $\delta m(x)$ describes mass fluctuations on the torus. The equation of motion $\square\delta m = 0$ describes massless oscillations.

Related document: Doc. 201 (DVFT adaptation)

Doc. 201 (*FFGFT-alles*) develops the same starting point $\mathcal{L}_0 = \varepsilon(\partial\Delta m)^2$ in a different direction: the polar decomposition $\Phi = \rho e^{i\theta}$ with vacuum amplitude ρ and vacuum phase θ (DVFT — Dynamic Vacuum Field Theory). Applications: *apparent* cosmic acceleration (without dark energy), galactic rotation curves (without dark matter), CMB homogeneity. No contradiction with Doc. 202 — different development directions of the same foundation.

9 The Torus Correction Term

The fractal geometry adds a correction term generating the non-linear dispersion:

$$\Delta\mathcal{L}_{\text{Torus}} = \varepsilon \xi_0 \left[\frac{f(\theta)}{2} (\partial\delta m)^2 - k^{\mu\nu}(\theta) \partial_\mu \delta m \partial_\nu \delta m \right] \quad (11)$$

The geometric functions $f(\theta)$ and $k^{\mu\nu}(\theta)$ encode the curvature of the fractal torus. The complete FFGFT Lagrangian:

$$\mathcal{L} = \mathcal{L}_0 + \Delta\mathcal{L}_{\text{Torus}} \quad (12)$$

10 The Mode Operator

The Hilbert space of torus modes:

$$\mathcal{H} = \bigoplus_{n,l,j} \mathbb{C} \cdot \mathbf{e}_{n,l,j}, \quad \langle \mathbf{e}_{n,l,j}, \mathbf{e}_{n',l',j'} \rangle = \delta_{nn'} \delta_{ll'} \delta_{jj'} \quad (13)$$

The field operator:

$$\Phi(x) = \sum_{n,l,j} m_{n,l,j} \cdot \phi_{n,l,j}(x) \cdot P_{n,l,j} \quad (14)$$

with $P_{n,l,j} = |\mathbf{e}_{n,l,j}\rangle\langle\mathbf{e}_{n,l,j}|$. Since $P_i P_j = \delta_{ij} P_i$: $\Phi^2 = \sum_i m_i^2 P_i$.

11 The Bridge Formula

The Lagrangian and mode operator describe the same physics:

$$\int \frac{1}{2} (\partial\delta m)^2 d^4x = \frac{1}{2} \langle \psi | \Phi^2 | \psi \rangle \quad (15)$$

Derivation in 6 steps:

Step 1 — Mode expansion: $\delta m(x) = \sum_i a_i \phi_i(x)$.

Step 2 — Substitution: $\int \mathcal{L}_0 d^4x = \frac{1}{2} \sum_{i,j} a_i a_j \int (\partial\phi_i)(\partial\phi_j) d^4x$.

Step 3 — Orthogonality: Cross terms vanish for $i \neq j$, leaving $\frac{1}{2} \sum_i a_i^2 \int (\partial\phi_i)^2 d^4x$.

Step 4 — Klein-Gordon: $\int (\partial\phi_i)^2 d^4x = m_i^2$ (partial integration, normalisation).

Step 5 — Summary: $\int \mathcal{L}_0 d^4x = \frac{1}{2} \sum_i a_i^2 m_i^2$.

Step 6 — Mode operator: $\langle \psi | \Phi^2 | \psi \rangle = \sum_i a_i^2 m_i^2$. Comparison gives (15) with $\varepsilon = 1/2$.

Part V

Mass Generation as a Binding Phenomenon

12 The Binding Picture

The eigenvalue equation on the torus:

$$\left[\square + \hat{V}(\xi_0, D_f) \right] \phi_{n,l,j} = -m_{n,l,j}^2 \phi_{n,l,j} \quad (16)$$

The mass is a difference:

$$m_i^2 = \underbrace{\left(\frac{2\pi n}{L_0} \right)^2}_{\text{kinetic}} + \underbrace{\langle \phi_i | \hat{V} | \phi_i \rangle}_{\approx -(2\pi n/L_0)^2} \quad (17)$$

The kinetic energy is enormous ($\sim 10^{32} \text{ GeV}^2$), the potential almost exactly opposite. The mass is the tiny residual:

Particle	$(2\pi n/L_0)^2$	$m_i^2/(\text{kin.})$
Electron	$1.046 \times 10^{32} \text{ GeV}^2$	2.5×10^{-39}
Muon	$4.185 \times 10^{32} \text{ GeV}^2$	2.7×10^{-35}
Tau	$9.415 \times 10^{32} \text{ GeV}^2$	3.4×10^{-33}

13 Not Fine-Tuning — a Binding Phenomenon

The extreme cancellation (32-39 orders of magnitude) is not fine-tuning — it is the defining property of bound states.

Crystal analogy: One does not ask why a crystal produces a particular phonon dispersion. The dispersion curve *is* the crystal structure in momentum space.

Hydrogen analogy: Kinetic energy $\approx 13.6 \text{ eV}$, Coulomb potential $\approx -27.2 \text{ eV}$, binding energy -13.6 eV . Nobody asks why the potential compensates the kinetic energy so precisely — it follows from the Schrödinger equation.

In FFGFT: The particles are bound torus modes. The mass is the binding residual. The potential $\hat{V}$ is uniquely determined by the torus geometry (ξ_0, D_f) — no free parameter.

14 Connection to the Lagrangian Extension

Lagrangian formulation	Torus binding picture
$\xi_0 \cdot m_\ell \cdot \bar{\psi} \psi \cdot \delta m$ (coupling)	Coupling of fermion mode to its own torus mode
$\Delta \mathcal{L}_{\text{Torus}}$	Potential term $\hat{V}$ from $D_f = 3 - \xi_0$
Mass parameter m_ℓ appears explicitly	Mass emerges as eigenvalue residual
Standard + geometric correction	Kinetic + potential ≈ 0 , residual = m_i^2

Difference from the Higgs mechanism: In the Standard Model, the Yukawa coupling y_i is a free parameter. In FFGFT, the corresponding term $\xi_0 \cdot m_\ell$ is geometrically fixed by $\xi_0 = 4/30000$.

The central claim:

Standard QFT calculations must be extended by a tiny geometric mass contribution that arises directly from the torus structure.

15 The Verification Condition

The theory makes a numerically testable prediction:

$$\langle \phi_{n,l,j} | \hat{V} | \phi_{n,l,j} \rangle = (r_i)^2 \cdot \xi_0^{2p_i} \cdot v^2 - \left(\frac{2\pi n}{L_0} \right)^2 \quad (18)$$

This is the Stage 3 bottleneck (Doc. 189): explicitly derive $\Delta\mathcal{L}_{\text{Torus}}$ from the torus geometry and show that the eigenvalues match the mass table (Doc. 006).

Part VI

The Field-Theoretic Formulation

16 Hilbert Space and Recursion Operator

The recursion operator $\mathcal{R}$ formalises the fractal scaling structure:

$$\mathcal{R}(\mathbf{e}_{n,l,j}) = \xi_0^{p(n,l,j)-1} \cdot K_{\text{frak}} \cdot \mathbf{e}_{n,l,j}, \quad \mathcal{R}(\Phi_*) = \Phi_* \quad (19)$$

The time-mass duality is a field symmetry: $\mathcal{R} : t \rightarrow \xi_0 t, m \rightarrow \xi_0^{-1} m, \Phi \rightarrow \Phi$.

17 Algebraic Structure

Mode algebra for free waves on the torus:

$$\mathbf{e}_{\mathbf{k}_i} \star \mathbf{e}_{\mathbf{k}_j} = \mathbf{e}_{\mathbf{k}_i + \mathbf{k}_j}, \quad C_{ij}^k = \delta_{k, i+j} \pmod{\text{torus period}} \quad (20)$$

18 The Renormalisation Group

The modified QED β -function in FFGFT (Doc. 008):

$$\frac{d\alpha}{d \ln \mu} = \beta_0 \alpha^2 \cdot \left(1 + \xi_0 \cdot \ln \frac{\mu}{E_0} \right) \quad (21)$$

$$\alpha^{-1}(\mu) = \alpha^{-1}(m_e) - \frac{\beta_0}{2\pi} \ln \frac{\mu}{m_e} - \frac{\beta_0 \xi_0}{4\pi} \left(\ln \frac{\mu}{m_e} \right)^2 \quad (22)$$

The ξ_0 term is the FFGFT-specific addition. ξ_0 is **not** a fixed point of this β -function but a fundamental geometric constant — time-invariant because topologically fixed.

Part VII

Open Mathematical Tasks

19 Four Precisely Formulated Problems

The theory is conceptually closed. The remaining tasks are mathematical.

Problems 1 and 2 have been formally worked out in Doc. 203 ($\mathcal{R}$ -operator, fixed-point condition) and Doc. 204 (fractal boundary condition, first-order perturbation in ξ_0). Problem 3 (algebra) is complete for the free theory. Problem 4 (interactions) remains open.

No.	Problem	Task	Status
1	$\mathcal{R}$ -operator	$\lambda_{n,l,j} = \xi_0^{p-1} K_{\text{frak}}; p(n, l, j)$ from torus	Doc. 203
2	Mode functions	Incorporate fractal boundary condition	Doc. 204
3	Algebra	Hopf algebra for interactions	Free theory: complete
4	Interactions	$\mathcal{L}_{\text{int}}, \beta$ -function	Doc. 008; ξ_0 geometric

Part VIII

Summary

20 Overview of Results

FFGFT / T0 theory describes all known elementary particles as modes of a universal fractal torus — closed, consistent, from a single geometric constant $\xi_0 = 4/3 \times 10^{-4}$.

Concept	Meaning
Universal torus	$D_f = 3 - \xi_0$, fundamental scale $L_0 = \ell_P / \xi_0$
Torus modes	Quantum numbers (n, l, j) , wave vector $\mathbf{k} = (2\pi/L_0)(k_x, k_y, k_z)$
Mode functions	$\phi = V^{-1/2} e^{i\mathbf{k} \cdot \mathbf{x}} Y_{l,j} e^{-imt}$
Lagrangian	$\mathcal{L} = \frac{1}{2}(\partial\delta m)^2 + \Delta\mathcal{L}_{\text{Torus}}$
Mode operator	$\Phi = \sum_i m_i P_i$, diagonal in Hilbert space
Bridge	$\int \mathcal{L}_0 d^4x = \frac{1}{2} \langle \psi \Phi^2 \psi \rangle$
Mass formula	$m_i = r_i \cdot \xi_0^{p_i} \cdot v$, $r_i, p_i \in \mathbb{Q}$
Non-closure	$(r_i r_j, p_i + p_j) \notin \mathcal{P}$ for $i \neq j$
Mass generation	Bound states: kin. $\approx$ pot., mass = binding residual

Notation

Symbol	Meaning
$\xi_0 = 4/30000$	Geometric base constant of the fractal torus
ℓ_P	Planck length
$L_0 = \ell_P/\xi_0$	Fundamental torus scale
$D_f = 3 - \xi_0$	Fractal dimension of the FFGFT torus
$v = 246 \text{ GeV}$	Higgs vacuum expectation value (energy unit)
n, l, j	Quantum numbers: principal, orbital, spin
$\mathbf{k}_{n,l}$	Quantised wave vector on the torus
$Y_{l,j}(\theta, \varphi)$	Spherical harmonics
$\phi_{n,l,j}(t, \mathbf{x})$	Torus mode functions
$m_i = r_i \cdot \xi_0^{p_i} \cdot v$	FFGFT mass formula
$r_i \in \mathbb{Q}$	Rational prefactor (winding ratio)
$p_i \in \mathbb{Q}$	Rational exponent (scaling depth)
$\mathcal{P}$	Set of admissible pairs (r_i, p_i)
$\mathcal{H}$	Hilbert space of torus modes
$\mathbf{e}_{n,l,j}$	Basis vector of the Hilbert space
$P_i = \mathbf{e}_i\rangle\langle\mathbf{e}_i $	Projector onto mode i
$\Phi = \sum_i m_i P_i$	Mode operator (field operator)
$\mathcal{L}_0 = \frac{1}{2}(\partial\delta m)^2$	Free Lagrangian
$\Delta\mathcal{L}_{\text{Torus}}$	Torus-geometric correction term
$\hat{V}(\xi_0, D_f)$	Potential operator from torus geometry
$K_{\text{frak}} \approx 0.986$	Fractal correction constant
$C_{ij}^k = \delta_{k,i+j}$	Structure constants of the mode algebra
β_0	Coefficient of the QED beta function

References: Doc. 006 (quantum numbers), Doc. 008 (β -function), Doc. 049 (L_0), Doc. 051 (spinor), Doc. 129, Doc. 180 (ξ_0), Doc. 189 (stage structure).