

Factorisation Classes in $G(6)$ and the Harmonics of the Primes:

Entry rule $k = \text{ord}_p(3)$, chords as partitions, and the boundary of the Galois layer

Johann Pascher
ORCID 0009-0000-6518-4064

2 September 2026

Contents

Factorisation Classes in $G(6)$ and the Harmonics of the Primes	2
.1 Starting point: algebraic equals geometric	1
.2 Theorem A: Factorisation classes in $G(6)$	1
Irreducible cubic polynomials over $GF(3)$	1
The five classes	1
.3 Theorem B: Entry rule of the harmonic primes	2
.4 Theorem C: Chords as partitions of 8	3
.5 Theorem D: Discriminating power and the boundary of the Galois layer	3
Density of Galois products	3
The threshold: Galois grid versus fractal correction	4
Methodological rule	4
.6 Negative result: tau sector	4
.7 Summary	4

Factorisation Classes in $G(6)$ and the Harmonics of the Primes

Correction notes

Corrections from the calculation review (1 Oct. 2026). The following passages are corrected or annotated in the text; each carries a dated note with the previous value:

- Table of factorisation classes, order-13 type $g_i^2(x+1)^2$: order 13 $\rightarrow$ 26.
- Entry rule: "from $k = 7$ on, only primes ≥ 41 " $\rightarrow$ from $k = 7$ on no new prime ≤ 13 enters (later ones: 17, 19, 23, 29, 31, 37, ...).
- Overtone series: below 5 cents "at $n \approx 250$ " $\rightarrow$ from $n \approx 350$ (at $n = 250$: 6.9 cents).

Abstract

Doc. 341 shows that $\text{GF}(27)$ enters $G(6)$ as an order structure through the characteristic polynomial $\chi_X = (x^3 + 2x + 1)^2(x + 1)^2$. This document asks which further factorisations χ admits in $G(6) \cong M_8(\text{GF}(9))$, and shows that the same factorisation rule organises the harmonic patterns of the primes (Doc. 057, Doc. 159).

Four results: **(A)** Over $\text{GF}(3)$ there are eight irreducible cubic polynomials, four of them primitive (order 26) and four of order 13; they generate five factorisation classes of χ in $G(6)$ **[B]**. **(B)** A prime p enters $|\text{GF}(3^k)^*|$ exactly at $k = \text{ord}_p(3)$; the harmonic primes enter in the order 13 ($k=3$), 5 ($k=4$), 11 ($k=5$), 7 ($k=6$); from $k = 7$ on, no new prime ≤ 13 enters (later ones: 17, 19, 23, 29, 31, 37, ...) **[B]**. *(Corrected on 1 Oct. 2026: previously "after which only large primes appear" – e.g. $\text{ord}_{23}(3) = 11$, $\text{ord}_{17}(3) = 16$; see entry rule below.)* **(C)** Element orders in $G(6)$ are partitions of 8 with an lcm rule — the algebraic form of a chord of overtones **[B]**. **(D)** From $k \geq 4$ on, the Galois grid is finer than the fractal-recursive correction (1.05 %, Doc. 338); Galois products then become statistically indistinguishable from higher torus modes. This is not a contradiction but the algebraic form of the mode convergence of Doc. 159 **[B]**.

Negative result: the tau sector is not a Galois product at 0.05 % **[S]**. Check scripts: `pruef_342_faktorisierung_primzahlen.py`, `pruef_342b_hoehere_harmonien.py`, `pruef_342c_dichte_hoehere_harmonien.py`.

.1 Starting point: algebraic equals geometric

The following status markers are used throughout:

- [K]** *Confirmed* — algebraically or numerically fully proved; all check-script assertions passed; derived directly from ξ or a corpus-established derivation.
- [B]** *Established* — numerically secured by sampling or explicit construction; analytical proof present or trivial.
- [S]** *Speculative* — indication or conjecture, not yet fully proved.

Docs. 338 and 341 showed that mass ratios from torus winding numbers (r_i, p_i) and from Galois group orders yield the same number: $(m_\mu/m_e)^2 = 43200 = |\text{GF}(9)^*|^2 \cdot 5^2 \cdot |\text{GF}(27)|$. The question whether the geometric formulation (vectors on T^4) is “more precise” than the algebraic one is ill-posed: winding numbers on S^1 and elements of $\text{GF}(3^k)$ are both integer, exact objects. The bridge is the Frobenius automorphism $g \mapsto g^3$ on $\text{GF}(3^k)$, which represents the $\mathbb{Z}_3$ rotation of the orbifold $T^4/\mathbb{Z}_3$ (Doc. 314, Doc. 336).

The algebraic route has one methodological advantage: polynomial factorisation can be enumerated systematically, geometric search cannot. This document exploits that.

.2 Theorem A: Factorisation classes in $G(6)$

Irreducible cubic polynomials over $\text{GF}(3)$

A monic cubic over $\text{GF}(3)$ is irreducible iff it has no root in $\{0, 1, 2\}$. There are eight such polynomials; their companion matrices have order 26 or 13 in $\text{GL}_3(\text{GF}(3))$ **[B]**:

Table 1: Irreducible cubics over $\text{GF}(3)$ and their order

Label	Polynomial	Order
f_1	$x^3 + 2x + 1$	26
f_2	$x^3 + 2x^2 + 1$	26
f_3	$x^3 + 2x^2 + x + 1$	26
f_4	$x^3 + x^2 + 2x + 1$	26
g_1	$x^3 + 2x + 2$	13
g_2	$x^3 + x^2 + 2$	13
g_3	$x^3 + x^2 + x + 2$	13
g_4	$x^3 + 2x^2 + 2x + 2$	13

The number four is forced: $\text{GF}(27)^*$ has $\varphi(26) = 12$ primitive elements in Frobenius orbits of length 3, hence $12/3 = 4$ minimal polynomials. Doc. 341 fixes only order 26 for the leptons, not a class; the assignment of charged leptons to the Dirac layer $\{f_3, f_4\}$ and neutrinos to the Majorana layer $\{f_1, f_2\}$ follows in Doc. 343, Theorem D''' **[S]**.

The five classes

In $G(6) \cong M_8(\text{GF}(9))$, χ splits over $\text{GF}(3)$ into factors whose degrees sum to 8. With cubic and linear factors this gives five classes **[B]**:

Table 2: χ factorisation classes with GF(27) content in G(6)

Type	$\chi(x)$	Count	Order
Doc.-341 type	$f_i^2 (x+1)^2$	4	26
Mixed primitive	$f_i f_j (x+1)^2, i \neq j$	6	26
Order-13 type	$g_i^2 (x+1)^2$	4	26 (sign partners of the f_i , Doc. 343)
Primitive $\times$ 13	$f_i g_j (x+1)^2$	16	26
Thin	$f_i (x+1)^5$	4	26

(Corrected on 1 Oct. 2026: previously order 13 for the order-13 type – the roots of g_i have order 13, the factor $(x+1)^2$ contributes the eigenvalue -1 (order 2); in the semisimple case the order is $\text{lcm}(13, 2) = 26$, with a Jordan block 78. Order 13 would arise only with $(x-1)^2 = (x+2)^2$; the name of the type refers to g_i .)

The four primitive polynomials are the four Frobenius conjugacy classes $f_1 = \{g, g^3, g^9\}$, $f_2 = \{g^{17}, g^{23}, g^{25}\}$, $f_3 = \{g^5, g^{15}, g^{19}\}$, $f_4 = \{g^7, g^{11}, g^{21}\}$ of a generator g of $\text{GF}(27)^*$ (root of f_1). Doc. 341 fixes only order 26 for the leptons, not a class; the assignment of charged leptons to the Dirac layer $\{f_3, f_4\}$ and neutrinos to the Majorana layer $\{f_1, f_2\}$ follows in Doc. 343, Theorem D''' [S].

.3 Theorem B: Entry rule of the harmonic primes

Doc. 057 describes primes as elementary relational steps: 2 octave, 3 fifth, 5 major third, 7 natural seventh, 11 eleventh, 13 thirteenth. A p -limit system admits primes up to p as interval building blocks.

Table 3: Order of $\text{GF}(3^k)^*$ and entry of new primes

k	$ \text{GF}(3^k)^* $	prime factors	new	harmonics
1	2	2	2	octave
2	8	2^3	—	GF(9), Matzke's $\mathbb{Z}_3\mathbb{C}$
3	26	$2 \cdot 13$	13	thirteenth; leptons (Doc. 338)
4	80	$2^4 \cdot 5$	5	major third; Doug's G(3) orders
5	242	$2 \cdot 11^2$	11	eleventh; neutrinos (Doc. 340)
6	728	$2^3 \cdot 7 \cdot 13$	7	seventh; with 13 on one carrier
7	2186	$2 \cdot 1093$	1093	no small harmonics left
8	6560	$2^5 \cdot 5 \cdot 41$	41	—
12	531440	$2^4 \cdot 5 \cdot 7 \cdot 13 \cdot 73$	73	all four united

Entry rule [B]: A prime $p \neq 3$ divides $3^k - 1$ iff $\text{ord}_p(3) \mid k$; it therefore first enters at $k = \text{ord}_p(3)$. This yields the order 13, 5, 11, 7 for $k = 3, 4, 5, 6$. All four harmonic primes above 3 enter between $k = 3$ and $k = 6$; from $k = 7$ on, no new prime ≤ 13 enters (the smallest later ones are 17, 19, 23, 29, 31, 37, ...; $k = 7$ brings 1093, a Wieferich prime, $k = 8$ brings 41). (Corrected on 1 Oct. 2026: previously “from $k = 7$ on, only primes ≥ 41 appear” – by the entry rule itself: $\text{ord}_{23}(3) = 11$ ($3^{11} - 1 = 2 \cdot 23 \cdot 3851$), $\text{ord}_{17}(3) = 16$, $\text{ord}_{19}(3) = \text{ord}_{37}(3) = 18$, $\text{ord}_{29}(3) = 28$, $\text{ord}_{31}(3) = 30$.)

This is the algebraic reason why practically relevant harmonics end at the 13-limit, and why the FFGFT corpus has so far needed only the primes $\{2, 3, 5, 7, 11, 13\}$. The neutrino eleven of Doc. 340 ($\Delta m_{\text{atm}}^2 = (11^2 - 1) m_\nu^2$) sits at $k = 5$; this had not been located in the corpus before.

$\mathbb{Z}_3$ compatibility: for $p \equiv 1 \pmod{3}$, $\mathbb{Z}_3 \subset \text{GF}(p)^*$. This holds for 7 and 13, not for 2, 5, 11. The two $\mathbb{Z}_3$ -compatible primes are exactly those that appear as structural numbers in Doc. 289 (top quark) and Docs. 338/341 (leptons) [B].

.4 Theorem C: Chords as partitions of 8

For semisimple $X \in M_8(\text{GF}(9))$ with χ split over $\text{GF}(3)$ into factors of degrees $d_1 + \dots + d_r = 8$, the eigenvalues lie in $\text{GF}(3^{d_i})$ and $\text{ord}(X)$ divides $\text{lcm}(3^{d_i} - 1)$ [B]. A partition of 8 is thus the algebraic form of a chord of overtones $k \in \{d_i\}$; the order is the common period.

Table 4: Partitions of 8 and maximal semisimple order

Partition	$\text{lcm}(3^{d_i}-1)$	reading
(3, 3, 1, 1)	26	pure $\text{GF}(27)$ chord (Doc. 341)
(3, 3, 2)	104	$= 2^3 \cdot 13$: $\text{GF}(9)$ with $\text{GF}(27)$
(4, 3, 1)	1040	$= 2^4 \cdot 5 \cdot 13$: third with thirteenth
(6, 2), (6, 1, 1)	728	$= 2^3 \cdot 7 \cdot 13$: seventh with thirteenth
(5, 3)	3146	$= 2 \cdot 11^2 \cdot 13$: eleventh with thirteenth
(8)	6560	$= 2^5 \cdot 5 \cdot 41$: maximal

The number of irreducible resp. primitive polynomials of degree k over $\text{GF}(3)$ grows quickly (Möbius formula): degree 3: 8/4, degree 4: 18/8, degree 6: 116/48, degree 8: 810/320. The space of possible chords is therefore large — which leads directly to the question of discriminating power.

.5 Theorem D: Discriminating power and the boundary of the Galois layer

Density of Galois products

Forming all products of up to three building blocks $\{3^k, 3^k - 1, \text{prime divisors}\}$ for $k \leq k_{\max}$ with exponents $\{-1, 1, 2\}$, the mean logarithmic spacing of neighbouring values in the window $10 < x < 10^4$ drops as follows [B]:

Table 5: Mean log spacing of Galois products

$k_{\max}$	values in window	mean spacing	expected chance hits at 0.5 %
2	54	12.8 %	0.08
3	218	3.17 %	0.32
4	650	1.06 %	0.94
5	1423	0.49 %	2.06
6	2503	0.28 %	3.62
8	5311	0.13 %	7.69

From $k_{\max} \approx 5$ on, the product set is denser than 0.5 %: every number is “hit”. A control test confirms this: at $k_{\max} = 6$, ten physical mass ratios have 2.9 hits on average, two hundred random numbers 3.7. The hit rate is at chance level [B].

The threshold: Galois grid versus fractal correction

Doc. 338 quantifies the fractal-recursive higher-winding correction between bare value and measurement at 1.05 % for $(m_\mu/m_e)^2$. The Galois grid is coarser than twice this correction up to $k = 3$ (3.17 % > 2.10 %) and hence discriminating; from $k = 4$ on it lies below (1.06 % < 2.10 %) **[B]**.

Consequence. The exact Galois layer ends structurally where the fractal correction begins. Up to GF(27) the algebra is exact and the 1 % is a higher-mode correction; above that, the algebra itself is finer than the correction and dissolves into the higher winding modes. Galois products with $k \geq 4$ are statistically indistinguishable from higher torus modes.

This is not a contradiction. It is the same mechanism as in the overtone series: the interval $(n+1) : n$ shrinks from 1200 cents ($n = 1$) via 105 cents ($n = 16$) to below 7 cents at $n \approx 250$ and below 5 cents from $n \approx 350$; the series becomes quasi-continuous. (*Corrected on 1 Oct. 2026: previously "below 5 cents at $n \approx 250$ " – $1200 \log_2(251/250) = 6.9$ cents; below 5 cents only from $n = 346$.)* Doc. 159 describes exactly this mode convergence on the fractal torus; here it appears in algebraic form.

Methodological rule

- $k \leq 3$ (GF(3), GF(9), GF(27)): identity layer. A hit with few building blocks is significant and may carry **[K]/[B]** if grounded geometrically in ξ (as 43200 in Doc. 338).
- $k = 4$ (GF(81), 5-limit): transition region; only with additional justification.
- $k \geq 5$: no status from product search. A relation holds here only if derived geometrically from ξ .

.6 Negative result: tau sector

At a tolerance of 0.05 %, none of the ratios m_τ/m_μ , m_τ/m_e or their squares is a product of up to three Galois building blocks with $k \leq 6$ **[S]**. Hits at 0.5 % exist (e.g. $7 \cdot 11^2/3$ for $(m_\tau/m_\mu)^2$ at 0.17 %) but lie at chance level by Theorem D. The tau ratio remains grounded in the corpus through winding numbers (Doc. 258, Doc. 310), not through a Galois identity.

.7 Summary

Table 6: Results and status

Result	Content	Status
A	8 irreducible cubics, 4 primitive; 5 χ classes	[B]
B	entry rule $k = \text{ord}_p(3)$; order 13, 5, 11, 7	[B]
B'	7, 13 are $\mathbb{Z}_3$ -compatible ($p \equiv 1 \pmod{3}$)	[B]
C	chords = partitions of 8, order = lcm	[B]
D	Galois grid < 2× fractal correction from $k = 4$	[B]
D'	hit rate physical = random for $k \geq 5$	[B]
—	tau sector no Galois product at 0.05 %	[S]
—	physics of the classes f_2, f_3, f_4	[S]

Check scripts

Directory 2/python/Dok342_Skripte/, wrapper run_all_342.sh:

- pruef_342_faktorisierung_primzahlen.py — Theorem A, p -limit assignment
- pruef_342b_hoehere_harmonien.py — Theorems B, C; product search with look-elsewhere note; Koide
- pruef_342c_dichte_hoehere_harmonien.py — Theorem D; density, random control, overtone analogue, threshold

All assertions passed.

Note on the use of AI assistance

This document was prepared with the assistance of Claude (Anthropic). All counts and order determinations are verified by Python check scripts with exact arithmetic. The physical interpretations and conclusions are those of Johann Pascher (ORCID 0009-0000-6518-4064).

Bibliography

- [1] J. Pascher, *Doc. 057: Relational number system*, FFGFT corpus.
- [2] J. Pascher, *Doc. 159: Harmonic structure of the torus*, FFGFT corpus.
- [3] J. Pascher, *Doc. 338: Mass calculations and Galois comparison*, FFGFT corpus, 28 August 2026.
- [4] J. Pascher, *Doc. 340: Neutrino mass hierarchy from $GF(27)^*$* , FFGFT corpus, 30 August 2026.
- [5] J. Pascher, *Doc. 341: $GF(27)$ in GALG and FFGFT*, FFGFT corpus, 1 September 2026.