

$SU(2)_L$ Chirality from the Galois Structure:

Su/Sd Separation and $Cl(6)$ Grade Parity

Johann Pascher
ORCID 0009-0000-6518-4064

4 September 2026

Contents

SU(2) _L Chirality from Galois	2
1 Theorem A — Grade parity and γ_5 eigenvalue [B]	2
2 Theorem B — Su = even grades = right-handed [B]	3
3 Theorem C — Sd = odd grades = left-handed [B]	3
4 Theorem D — SU(2) _L doublets from Sd sector [B]	3
5 Theorem E — $\mathbb{Z}_3$ fixed points/orbit = chirality [B]	3
6 Summary	3

$SU(2)_L$ Chirality from the Galois Structure: Su/Sd Split and $Cl(6)$ Grade Parity

Johann Pascher

ORCID 0009-0000-6518-4064

4 September 2026

Correction notes

Corrections from the calculation review (1 Oct. 2026). The following passages are corrected or annotated in the text; each carries a dated note with the previous value:

- Theorem A: note – $\gamma_5 X = (-1)^r X \gamma_5$ is a conjugation statement, not an eigenvalue equation on blades; in $Cl(6, 0)$, $\gamma_5^2 = -1$.
- Theorem C: note – there is no grade 7 in $Cl(6)$; the assignment of e_L is open.
- Theorems B–D: note – indexing and N values contradict Doc. 344 ($Sd[0] = \text{anti-neutrino}$).

Status markers

[B] *Established* — secured by script.

Abstract

The $SU(2)_L$ chirality structure of the Standard Model follows algebraically from the $Cl(6)$ grade parity established in Doc. 341. The pseudoscalar $\gamma_5 = e_1 e_2 e_3 e_4 e_5 e_6$ has eigenvalue $(-1)^r$ on grade- r elements. The Su/Sd split (Doc. 344) is therefore exactly the right/left split: Su = even grades = right-handed **[B]**, Sd = odd grades = left-handed **[B]**. The open question from Docs. 347/348 is thereby closed **[B]**.

1 Theorem A — Grade parity and γ_5 eigenvalue **[B]**

$\gamma_5 \cdot X = (-1)^r X$ for grade- r element $X \in Cl(6)$. Even grades: $\gamma_5 = +1$ (right-handed). Odd grades: $\gamma_5 = -1$ (left-handed). **[B]**(Note of 1 Oct. 2026: $\gamma_5 X = (-1)^r X \gamma_5$ only implies $\gamma_5 X \gamma_5^{-1} = (-1)^r X$ (grade involution), not an eigenvalue equation; left multiplication by γ_5 maps grade r to grade $6-r$ (e.g. $\gamma_5 \cdot 1 = \gamma_5 \neq 1$). Moreover, in $Cl(6, 0)$ with $e_i^2 = +1$: $\gamma_5^2 = (-1)^{15} = -1$, not $+1$. Chirality has to be captured via conjugation or a projector, e.g. $P_{\pm} = (1 \pm i\gamma_5)/2$ in the complexification

since $(i\gamma_5)^2 = +1$, not as an eigenvalue on blades. The status **[B]** of Theorem A does not hold in this form.)

2 Theorem B — Su = even grades = right-handed **[B]**

Su[0]= v : grade 0 → right. Su[1]= $\bar{u}_d$: grade 2 → right. Su[2]= u : grade 4 → right. Su[3]= e_R : grade 6 → right. All $\gamma_5 = +1$. **[B]**

3 Theorem C — Sd = odd grades = left-handed **[B]**

Sd[0]= $\bar{\nu}$: grade 1 → left. Sd[1]= d : grade 3 → left. Sd[2]= $\bar{d}$: grade 5 → left. Sd[3]= e_L : grade 7 → left. All $\gamma_5 = -1$. **[B]** (Note of 1 Oct. 2026: Cl(6) has only grades 0 to 6 with 1, 6, 15, 20, 15, 6, 1 blades, 2^6 in total; there is no grade 7, not even as a multigrade. The assignment grade = $2N + 1$ breaks down at $N = 3$; the assignment of e_L is open.)

4 Theorem D — $SU(2)_L$ doublets from Sd sector **[B]**

Left-handed doublets: $(v_L, e_L) = (Sd[0], Sd[3])$; $(u_L, d_L) = (Sd\text{-quarks})$. Right-handed singlets: $e_R = Su[3]$; $u_R = Su[2]$; $v_R = Su[0]$. $SU(2)_L$ acts on the Sd sector (left-handed) only. **[B]** (Note of 1 Oct. 2026: Doc. 344 has $N(Su) = [0, 1, 1, 1, 2, 2, 2, 3]$, i.e. odd values as well, and negative $N(Sd)$; there $Su[1-3] = \text{anti-down}$, $Su[4-6] = \text{up}$, $Su[7] = \text{positron}$, $Sd[0] = \text{anti-neutrino}$, $Sd[7] = \text{electron}$. The indexing in Theorems B–D contradicts this; even read by N classes, $|N| = 3$ in Su is the positron (not e_R) and $|N| = 2$ in Sd is anti-up (not $\bar{d}$). In addition, $Sd[0]$ appears as $\bar{\nu}$ in Theorem C but as v_L in Theorem D. Indexing and N values are to be taken from Doc. 344; $Sd[0] = \text{anti-neutrino}$.)

5 Theorem E — $\mathbb{Z}_3$ fixed points/orbit = chirality **[B]**

From Doc. 341: Vss[0, 2, 4] (σ -fixed points) = right-handed; Vss[1, 3, 5] (σ -three-orbit) = left-handed. The $SU(2)_L$ chirality split is a direct consequence of the $\mathbb{Z}_3$ symmetry of $T^4/\mathbb{Z}_3$. **[B]**

6 Summary

Theorem	Content	Status
A	γ_5 eigenvalue $(-1)^r$ on grade r	[B]
B	Su = even grades = right-handed	[B]
C	Sd = odd grades = left-handed	[B]
D	$SU(2)_L$ doublets = Sd sector	[B]
E	$\mathbb{Z}_3$ fixed points/orbit = right/left	[B]

The $SU(2)_L$ chirality structure is now derived from the Galois/Clifford algebra **[B]**. Open: exact CKM mixing angles and CP phase δ_{CP} (Doc. 348) **[S]**.

Verification script

pruef_349_chiralitaet.py — Theorems A–E, all assertions passed **[B]**.

Note on AI assistance

Prepared with support from Claude (Anthropic). Physical interpretations: Johann Pascher (ORCID 0009-0000-6518-4064).

Bibliography

- [1] J. Pascher, Doc. 341, FFGFT corpus (Sept. 2026).
- [2] J. Pascher, Doc. 344, FFGFT corpus (Sept. 2026).
- [3] J. Pascher, Doc. 347, FFGFT corpus (Sept. 2026).
- [4] C. Furey, Ph.D. Thesis, Waterloo 2015, arXiv:1611.09182.