

Masses and Planck Scale without ν

One chain, one anchor

Johann Pascher
ORCID 0009-0000-6518-4064

29 September 2026

Contents

Masses and Planck scale without v	2
1 Starting point	1
2 The relation between v and E_p	1
3 The masses without v	1
4 One chain, one anchor	2
5 Relation to the G formula with C_{conv}	2
6 Notes in other documents	3
7 Result	3

Masses and Planck Scale without ν

Abstract

The lepton ladder of the corpus writes the masses as $m_i = r_i \xi^{p_i} \nu$ and uses the electroweak scale ν as an intermediate quantity; the Planck scale is reached separately through the formula for G with the conversion factor C_{conv} . Read this way, the calculation seems to need two anchors, one for ν and one for E_P . It does not. Doc. 149 links ν and E_P directly: $\nu = E_P / (f^4 \cdot (\pi/2) \cdot 10)$, i.e. $\nu/E_P = \xi^4 / (5\pi) = 2.012 \times 10^{-17}$, -0.23% against the measured value **[K]**. Inserted into the ladder, ν drops out: $m_i = \frac{r_i}{5\pi} \xi^{p_i+4} E_P$ **[B]**. All three lepton masses, E_P , G , ℓ_P and L_0 then hang on a single chain, and one measured value as anchor fixes all numbers. With m_e as anchor one obtains $E_P = 1.237 \times 10^{19}$ GeV ($+1.34\%$), $G = 6.50 \times 10^{-11}$ (-2.6%), $\ell_P = 1.595 \times 10^{-35}$ m and $L_0 = 2.127 \times 10^{-39}$ m **[K]**. The deviations are the known bare residues of the ladder, not a second anchor.

Status markers

- [B]** Proven algebraically (rearrangement without additional assumption).
- [K]** Confirmed numerically (check script).
- [S]** Speculative: a hint or conjecture, not yet derived or checked in the corpus.

Measured values (CODATA 2022, PDG) serve as anchor or as comparison values. $\hbar$, c and e are exact by definition in the SI; they fix only the units and are not an additional anchor.

1 Starting point

Two routes stand side by side in the corpus:

1. **Lepton ladder** (Docs. 006, 352): $m_i = r_i \xi^{p_i} v$ with $r = (4/3, 16/5, 25/9)$ and $p = (3/2, 1, 2/3)$ for e, μ, τ . The mass ratios $m_\mu/m_e = \frac{12}{5} \xi^{-1/2} = 207.85$ and $m_\tau/m_\mu = \frac{125}{144} \xi^{-1/3} = 16.99$ do not contain v .
2. **Gravitation** (Docs. 012, 013, 180): $G = \xi^2 / (4m_e) \cdot C_{\text{conv}} \cdot K_{\text{frak}}$, and from it E_P and ℓ_P . Taking v from m_e and E_P from the G formula uses two measured values: m_e and, through C_{conv} , the measured value of G . This is how Doc. 375 calculates. A chain meant to derive all quantities from ξ alone, however, may have only one anchor; all other numbers must follow. Doc. 149 supplies the link.

2 The relation between v and E_P

Doc. 149 derives the electroweak scale from the Planck energy: dilution over the f^4 cells of the four-dimensional lattice, projection onto the three-dimensional half-space $(\pi/2)$, and a factor $1/10$ for the step to the electroweak scale:

$$v = \frac{E_P}{f^4 \cdot (\pi/2) \cdot 10} \quad \Longleftrightarrow \quad \frac{v}{E_P} = \frac{\xi^4}{5\pi} \quad \textbf{[B]} \quad (1)$$

with $f = 1/\xi$. The rearrangement is exact. Doc. 149 labels the factor 10 a motivated choice; it is not derived **[S]**.

Numerical value. $v/E_P = \xi^4/(5\pi) = 2.0120 \times 10^{-17}$. The measured value is $v/E_P = 246.22 \text{ GeV} / 1.22089 \times 10^{19} \text{ GeV} = 2.0167 \times 10^{-17}$; the deviation is -0.23% **[K]**. Since v/E_P is dimensionless, this number needs no anchor.

On the value in Doc. 149. There the calculation uses $f = 7491.91$ and gives $v = 246.71 \text{ GeV}$. With $f = 1/\xi = 7500$ the same formula gives $v = 245.65 \text{ GeV}$ **[K]**. Both values are calculated correctly; they differ in the f inserted.

3 The masses without v

Inserting $v = \xi^4 E_P / (5\pi)$ into the ladder gives

$$m_i = \frac{r_i}{5\pi} \xi^{p_i+4} E_P \quad \textbf{[B]} \quad (2)$$

With the measured value $E_P = 1.22089 \times 10^{22} \text{ MeV}$ for comparison:

Lepton	Formula	Coefficient	Value	vs. measurement
e	$\frac{4}{15\pi} \xi^{11/2} E_P$	0.08488	0.5043 MeV	−1.32 %
μ	$\frac{16}{25\pi} \xi^5 E_P$	0.20372	104.81 MeV	−0.80 %
τ	$\frac{5}{9\pi} \xi^{14/3} E_P$	0.17684	1780.9 MeV	+0.23 %

[K] The mass ratios are the same as in the ladder, since E_P and 5π cancel. The deviations combine the bare residue of the ladder against v (Doc. 352) and the −0.23 % of the relation v/E_P .

4 One chain, one anchor

The chain reads:

$$\xi \rightarrow \frac{m_i}{E_P} = \frac{r_i}{5\pi} \xi^{p_i+4} \rightarrow E_P \rightarrow G = \frac{\hbar c^5}{E_P^2}, \quad \ell_P = \frac{\hbar c}{E_P}, \quad L_0 = \xi \ell_P. \quad (3)$$

Apart from the units, everything is fixed by ξ . A single measured value fixes the numbers. With $m_e = 0.51099895$ MeV as anchor:

Quantity	from anchor m_e	vs. measurement
E_P	1.2372×10^{19} GeV	+1.34 %
G	6.4995×10^{-11} m ³ kg ^{−1} s ^{−2}	−2.6 %
ℓ_P	1.5950×10^{-35} m	−1.32 %
$L_0 = \xi \ell_P$	2.1266×10^{-39} m	−1.32 %

[K] Since $G \propto E_P^{-2}$, the relative deviation of G is twice that of E_P .

Choice of anchor. The chain can start at any point; which quantity serves as anchor is a question of accuracy, not of derivation. The deviations of the other quantities then depend on the bare residue of the chosen entry point:

Anchor	E_P	G	ℓ_P, L_0
m_e	+1.34 %	−2.62 %	−1.32 %
m_μ	+0.81 %	−1.60 %	−0.80 %
m_τ	−0.23 %	+0.45 %	+0.23 %
v	+0.23 %	−0.46 %	−0.23 %

[K] In every case exactly one measured value is used. The spread between the rows is the bare residue of the ladder; it shrinks to the extent that Doc. 352 resolves this residue.

5 Relation to the G formula with C_{conv}

The formula $G = \xi^2/(4m_e) \cdot C_{\text{conv}} \cdot K_{\text{frak}}$ from Doc. 012 gives 6.6745×10^{-11} , i.e. +0.003 %. This does not contradict the −2.6 % above: $C_{\text{conv}} = 7.783 \times 10^{-3}$ is chosen so that the SI conversion matches the measured value of G . In this form m_e and G both enter as anchors. The one-anchor chain with m_e corresponds to $C_{\text{conv}} = 7.58 \times 10^{-3}$ **[K]**; the difference of 2.6 % is exactly the residue of the chain. C_{conv} remains the SI conversion of G ; the chain only shows that it is not needed as a second measured value if the residue is accepted.

Doc. 375 calculates v/E_P along this route (v from m_e, E_P via C_{conv}) and obtains 2.0389×10^{-17} . The direct value $\xi^4/(5\pi) = 2.0120 \times 10^{-17}$ lies lower by the factor 1.0134; this is exactly the residue −1.32 % of the electron mass in the chain **[K]**. Both routes are therefore consistent.

6 Notes in other documents

- **Doc. 149:** note at the formula for ν that it gives 245.65 GeV with $f = 7500$ and means $\nu/E_P = \xi^4/(5\pi)$.
- **Doc. 180:** note at ℓ_P that the value 1.616×10^{-35} m is the measured value used as anchor and that the one-anchor chain gives 1.595×10^{-35} m.
- **Doc. 375:** note that ν/E_P follows directly as $\xi^4/(5\pi)$, without C_{conv} and without an anchor.

7 Result

In the mass calculation ν is an intermediate quantity and can be eliminated: $m_i = \frac{r_i}{5\pi} \xi^{p_i+4} E_P$ **[B]**. The lepton masses, E_P , G , ℓ_P and L_0 thus hang on one chain that is fixed by ξ apart from the units. A single measured value suffices as anchor; which one is free. The deviations of 0.2 to 1.3 % in E_P are the bare residues of the ladder and of the relation ν/E_P (−0.23 %), not the consequence of a missing anchor. The factor 10 in this relation is the open point of the chain **[S]**.

Check script: 2/python/Dok383_Skripte/pruef_383_ein_anker.py, 20/20.

Bibliography

- [1] E. Tiesinga et al., *CODATA Recommended Values of the Fundamental Physical Constants: 2022*, NIST (2024).
- [2] Particle Data Group (S. Navas et al.), *Review of Particle Physics*, Phys. Rev. D 110 (2024) 030001.