

Fundamental Fractal-Geometric Field Theory (FFGFT) or T0 Theory – Time-Mass Duality

Part 4: Newer Documents and Extensions (2025–2026)

Johann Pascher

2025

Chapter A

Introduction to Volume 4

Fourth Volume of the Document Collection

This fourth volume continues the collection of individual documents on T0 theory / FFGFT. As in the three preceding volumes, the documents are standalone works produced at different points in the theory's development. Unlike Volumes 1–3, which carry thematically closed emphases (foundations, Lagrangian/QFT, cosmology/consciousness), this volume gathers the documents that arose or were added after the original three-volume conception, as well as a few earlier works that had not been included in the first three volumes.

Vol. 4 and 5 — Split

Because of the Kindle volume limit (max. 550 pages per volume), the documents added since the original three-volume conception are distributed over two volumes:

- **Volume 4 (this volume):** 37 documents – supplementary individual documents, time and numerical predictions, consciousness/photronics, and the early torus and L_0 derivations up to Doc. 184 (p-bit).
- **Volume 5:** 37 documents – from Doc. 185 (Embedding Cost) onwards: late torus geometry, FFGFT field theory, Hilbert-space bridge, layers and information formalism, recent clarifications including the epistemic self-positioning in Doc. 262.

The split follows a natural content boundary: up to and including Doc. 184, the theory is developed on Layer 1 (compact T^4 , correction-free description in natural units); from Doc. 185 onwards, the Layer-1 / Layer-2 distinction is used as an explicit descriptive grid. The documents 263 to 389, written from May 2026 onwards, follow in Volumes 6 to 8.

Character of this Volume

Volume 4 is broad in scope: it contains extensions to classical topics (Bell tests, asymmetry, Lagrangian), a connected series of numerical predictions (Koide, g-2, lepton lifetime, H_0 ratio, LiNbO_3 dispersion matched to 16 significant digits), a series on photronics and qubit states, and the fundamental geometry and L_0 derivations.

Four-Part Organisation

The 37 documents are arranged in numerical order and divided into four thematic parts:

- **Part I — Supplementary Documents to Volumes 1–3:** earlier works not included in the first three volumes (introductions, asymmetry, simplified Lagrangian, Bell follow-ups, scramblons).
- **Part II — Time, Cosmology and Numerical Predictions:** arrow of time, H_0 ratio, Casimir, Koide-to-g-2 bridge, lepton lifetime ratios, harmonic structure on the torus, Ising machine analogy, UV catastrophe, 't Hooft CAI, LiNbO_3 geometry, periodic table.
- **Part III — Consciousness, Photonics and Qubit Series:** T0 and consciousness, string-theory connection, quantum dot, spin and qubit states, Shor's algorithm, spin stability, pairs and photons.
- **Part IV — Early Torus and L_0 Derivations:** L_0 derivation, justification of the torus geometry, maximum scale of the universe, Willow processor, p-bit.

Status and Updates

All documents in this collection carry the May 2026 status. The chapters contained in Volumes 1–3 have been reviewed for the new edition and corrected where appropriate — in particular the hardware-validation correction in Doc. 147 § 8 in Volume 2, and the updates to the follow-up documents Doc. 022, Doc. 035, Doc. 148 and Doc. 202.

Part I

Part I — Supplementary Documents to Volumes 1–3

Chapter B

Doc. 001 — T0 Theory — Book Abstract (Short Form)

Abstract

The T0-Theory (Time-Mass Duality) is a new approach in theoretical physics. In simple terms: Imagine the universe as a vast puzzle where everything—from the tiniest particles to the vast cosmos—fits together perfectly, without loose ends. The central result of this work is the realization that **all natural constants and physical parameters can be derived from a single dimensionless number**: the universal geometric constant $\xi \approx \frac{4}{3} \times 10^{-4}$. Think of ξ as the “master key” to the universe—a tiny number emerging from the fundamental form of three-dimensional space that unlocks explanations for gravity, the speed of light, particle masses, and more.

This collection of over 200 scientific documents systematically develops a complete physical theory that unifies quantum mechanics, relativity, and cosmology—based on the principle of absolute time T_0 and the intrinsic time-field-mass relationship. In everyday language: It’s as if we’re rewriting the rules of physics so that time is stable and reliable (not bendy like in Einstein’s view), while mass can change like sand in the wind, all connected through this elegant geometric idea. The foundational documents follow a purely geometric path, deriving ξ from the three-dimensional structure of space and constructing all other constants from it, including the fine-structure constant $\alpha \approx 1/137$, particle masses, and coupling strengths, without introducing additional free parameters. No more arbitrary numbers; everything flows from a single simple source, making the universe feel less random and more like a beautifully designed whole. Remarkably, the theory posits a static universe without expansion, as detailed in the CMB document, thereby rendering concepts like dark matter or dark energy superfluous.

B.1 The Core Principle: Everything from One Number

The fundamental insight of the T0-Theory can be summarized in one sentence:

Central Theorem of the T0-Theory

All physical constants—gravitational constant G , Planck constant $\hbar$, speed of light c , elementary charge e , as well as all particle masses and coupling constants—can be mathematically derived from a single dimensionless number: the universal geometric constant

$$\xi = \frac{4}{3} \times 10^{-4},$$

which emerges from the fundamental three-dimensional space geometry (harmonic fourth 4:3, QFT loop scaling 10^{-4}).

From ξ follows the fine-structure constant as:

$$\alpha = \xi E_0^2 \approx \frac{1}{137.04},$$

where α serves as a secondary electromagnetic coupling without primacy.

In everyday language, this means: We have reduced the “why” of physics to a single, space-born number—no magic, just geometry doing the heavy lifting.

B.2 Foundations of the T0-Theory

Time-Mass Duality

In contrast to standard physics, where time is relative and mass is constant, the T0-Theory postulates:

- **Absolute Time Measure** T_0 : Time flows uniformly everywhere in the universe—like a universal clock that ticks the same for everyone, no matter where you are.
- **Variable Mass**: Mass varies with the energy content of the vacuum—imagine mass as flexible, changing depending on the “hum” of the empty space around it.
- **Intrinsic Time Field** $T(x, t)$: Every particle carries its own time field—each building block of matter has its personal timer that influences its behavior.

The fundamental relationship is:

$$m(x) = \frac{\hbar}{c^2 T(x, t)(x)} = m_0 \cdot (1 + \kappa \Phi(x)),$$

where κ is reducible to ξ via geometric scaling. Mathematically, this duality treats time and mass as variables, ensuring the framework remains fully compatible with established mathematical structures while enabling a unified description of physical phenomena. Simply put: By letting time and mass dance as adaptable partners, we keep the math clean and intuitive, connecting old ideas with new ones without sacrificing a drop of sweat.

The Parameter ξ

The central parameter of the theory is:

$$\xi = \frac{4}{3} \times 10^{-4},$$

a purely geometric construct from 3D space that connects quantum mechanics with gravity. This parameter encodes the fundamental coupling between energy and spatial structure, from which all hierarchies emerge. It is like the ratio that tells space how to “scale” energy—small but mighty, whispering the secrets of why electrons are light and protons are heavy.

B.3 Derivation of All Natural Constants

Everything Follows from ξ

The T0-Theory demonstrates that:

1. Gravitational Constant:

$$G = f_G(\xi, m_p, c, \hbar),$$

where all inputs are reducible to ξ -scaled geometric units. Gravity? Just a wave from the geometry of space, tuned by ξ .

2. Particle Masses (Electron, Muon, Tau, Quarks):

The particle masses follow a universal scaling law analogous to the ordering principles of atomic energy levels, where quantum numbers (n, l, j) dictate hierarchical structures in a manner similar to atomic shells and subshells—imagine particles stacked like floors in a building, each level set by simple rules, much like electrons orbiting atoms. Thus,

$$\frac{m_e}{m_p} = g(\xi), \quad \frac{m_\mu}{m_e} = h(\xi), \quad \frac{m_\tau}{m_\mu} = k(\xi),$$

via universal scaling laws $\xi_i = \xi \times f(n_i, l_i, j_i)$, where $f(n, l, j)$ is the geometric structure factor of the quantum numbers – not a frequency and not the base winding number $f = 1/\xi$. No more guessing why some particles are 200 times heavier; it's all patterned like a cosmic family tree.

3. Coupling Constants (electroweak, strong, electromagnetic):

$$\alpha_W = f_W(\xi), \quad \alpha_s = f_s(\xi), \quad \alpha = f_\alpha(\xi).$$

These “strengths” of forces? Derived like branches from the same geometric trunk.

4. Cosmological Parameters:

Static universe metrics and CMB temperature $T_{\text{CMB}} = f_{\text{CMB}}(\xi)$, with redshift mechanisms derived from time-field variations (see CMB document for detailed explanation without expansion).

B.4 Experimental Predictions

The T0-Theory makes precise, testable predictions:

Concrete Predictions

- **Anomalous Magnetic Moment:** $(g-2)_\mu$ calculation solely from ξ —an electron-like wobble quirk explained without extras.
- **Koide Formula:** Exact mass relation of leptons via ξ -scaling—the math that ties the weights of three particles in a neat loop.
- **Redshift:** Modified interpretation without expansion, governed by ξ —why distant stars look “stretched” without the universe inflating.
- **CMB Anisotropies:** Explanation through time-field variations rooted in ξ —the microwave “echo” of the cosmos as geometric echoes.

These are no wild guesses; they are verifiable with today’s labs and invite everyone—physicists or curious minds—to put the theory to the test.

B.5 Structure of the Document Collection

This collection encompasses:

- **Foundations:** Mathematical formulation of time-mass duality under ξ -geometry—the basics, explained step by step.
- **Quantum Mechanics:** Deterministic interpretation, Bell inequalities—quantum madness made predictable and local.
- **Quantum Field Theory:** Lagrangian formalism in the T0 framework—fields dancing to a unified tune.
- **Cosmology:** Static universe, redshift, CMB—a stable universe that still surprises, without expansion, dark matter, or dark energy.
- **Particle Physics:** Mass spectrum, anomalous moments, Koide formula—the particle zoo tamed.
- **Technical Applications:** Photon chip, RSA cryptography—real tricks from the theory.
- **Experimental Tests:** Verifiable predictions—hands-on ways to probe the ideas.

Note: The documents consistently follow the geometric ξ -path, deriving all physics from 3D space principles, with α and other constants appearing as emergent features. We have woven in simple language throughout, so non-experts can dive in without drowning in jargon.

Chapter C

Doc. 001b — T0 Theory — General Introduction

Introduction

This book presents the current state of the T0 time-mass duality framework and its applications to particle masses, fundamental constants, quantum mechanics, gravitation, and cosmology.

The main body of the book consists of a series of core documents on T0. These chapters reflect the present understanding of the theory and its quantitative consequences. Wherever possible, the material has been reorganized and unified to make the structure of the theory as transparent as possible.

At the end of the book, several older documents are included in an appendix. These texts represent earlier stages in the development of the T0 framework. They have not been removed because they make visible the evolution of the ideas and the refinement of the formulas. In many cases, one can observe how approximations were improved, special cases generalized, and how new empirical data helped sharpen or correct earlier arguments.

The "live" version of the theory is maintained in a public GitHub repository:

<https://github.com/jpascher/T0-Time-Mass-Duality>

The LaTeX sources of the chapters in this book originate from this repository. When conceptual or numerical errors are found, they are corrected there first. This means that the PDF version of the book you are reading is a snapshot of a continuously evolving project. For the most current version of the documents, including new appendices or corrections, the GitHub repository should always be considered the primary reference.

The intent of this compilation is twofold:

- to provide a coherent, readable path through the core ideas and results of the T0 framework;
- to document the historical development of these ideas in the appendix, including false starts, intermediate formulations, and early adjustments to experimental data.

Readers primarily interested in the current formulation of the theory may focus on the core chapters. Readers also interested in the thinking and trial-and-error process behind the theory are invited to study the appendix material in parallel.

Chapter D

Doc. 002 — T0 Theory — A Journey Through the Energy Field

Abstract

The T0 Theory (Time-Mass Duality) is a new approach in theoretical physics. In simple terms: Imagine the universe as a grand puzzle where everything—from the tiniest particles to the vast cosmos—fits together perfectly, without loose ends. The central result of this work is the realization that **all natural constants and physical parameters can be derived from a single dimensionless number**: the universal geometric constant $\xi \approx \frac{4}{3} \times 10^{-4}$. Think of ξ as the “master key” of the universe—a tiny number that emerges from the fundamental shape of three-dimensional space and unlocks explanations for gravity, the speed of light, particle masses, and more.

This collection of over 200 scientific documents systematically develops a complete physical theory that unifies quantum mechanics, relativity, and cosmology—based on the principle of absolute time T_0 and the intrinsic time-field-mass relationship. In everyday language: It’s as if we’re rewriting the rules of physics so that time is stable and reliable (not bendy like in Einstein’s view), while mass can change like sand in the wind, all connected through this elegant geometric idea. The foundational documents follow a purely geometric path, deriving ξ from the three-dimensional structure of space and constructing all other constants from it, including the fine-structure constant $\alpha \approx 1/137$, particle masses, and coupling strengths, without introducing additional free parameters. No more arbitrary numbers; everything flows from a single simple source, making the universe feel less random and more like a beautifully designed whole. Remarkably, the theory postulates a static universe without expansion, as detailed in the CMB document, thereby rendering concepts like dark matter or dark energy superfluous.

D.1 The Core Principle: Everything from One Number

The fundamental insight of the T0 Theory can be summarized in one sentence:

Central Theorem of the T0 Theory

All physical constants—gravitational constant G , Planck constant $\hbar$, speed of light c , elementary charge e , as well as all particle masses and coupling constants—can be mathematically derived from a single dimensionless number: the universal geometric constant

$$\xi = \frac{4}{3} \times 10^{-4},$$

which emerges from the fundamental three-dimensional space geometry (harmonic fourth 4:3, QFT loop scaling 10^{-4}). From ξ follows the fine-structure constant as:

$$\alpha = \xi E_0^2 \approx \frac{1}{137.04},$$

where α serves as a secondary electromagnetic coupling without primacy.

In everyday language, this means: We've reduced the "why" of physics to a single, space-born number—no magic, just geometry doing the heavy lifting.

D.2 Foundations of the T0 Theory

Time-Mass Duality

In contrast to standard physics, where time is relative and mass is constant, the T0 Theory postulates:

- **Absolute Time Measure** T_0 : Time flows uniformly everywhere in the universe—like a universal clock that ticks the same for everyone, no matter where you are.
- **Variable Mass**: Mass varies with the energy content of the vacuum—think of mass as flexible, changing depending on the "hum" of the empty space around it.
- **Intrinsic Time Field** $T(x, t)$: Every particle carries its own time field—each building block of matter has its personal timer that influences its behavior.

The fundamental relationship is:

$$m(x) = \frac{\hbar}{c^2 T(x, t)(x)} = m_0 \cdot (1 + \kappa \Phi(x)),$$

where κ is traceable to ξ via geometric scaling. Mathematically, this duality treats time and mass as variables, ensuring the framework remains fully compatible with established mathematical structures while enabling a unified description of physical phenomena. Simply put: By letting time and mass dance as adjustable partners, we keep the math clean and intuitive, connecting old ideas with new ones without sacrificing a weld.

The Parameter ξ

The central parameter of the theory is:

$$\xi = \frac{4}{3} \times 10^{-4},$$

a purely geometric construct from 3D space that connects quantum mechanics with gravity. This parameter encodes the fundamental coupling between energy and spatial structure, from which all hierarchies emerge. It's like the ratio that tells space how to "scale" energy—small but mighty, whispering the secrets of why electrons are light and protons heavy.

D.3 Derivation of All Natural Constants

Everything Follows from ξ

The T0 Theory demonstrates that:

1. Gravitational Constant:

$$G = f_G(\xi, m_p, c, \hbar),$$

where all inputs are reducible to ξ -scaled geometric units. Gravity? Just a wave from the geometry of space, tuned by ξ .

2. Particle Masses (Electron, Muon, Tau, Quarks):

The particle masses follow a universal scaling law analogous to the ordering principles of atomic energy levels, where quantum numbers (n, l, j) dictate hierarchical structures in a manner similar to atomic shells and subshells—imagine particles stacked like floors in a building, each level set by simple rules, much like electrons orbiting atoms. Thus,

$$\frac{m_e}{m_p} = g(\xi), \quad \frac{m_\mu}{m_e} = h(\xi), \quad \frac{m_\tau}{m_\mu} = k(\xi),$$

via universal scaling laws $\xi_i = \xi \times f(n_i, l_i, j_i)$, where $f(n, l, j)$ is the geometric structure factor of the quantum numbers – not a frequency and not the base winding number $f = 1/\xi$. No more guessing why some particles are 200 times heavier; it's all patterned like a cosmic family tree.

3. Coupling Constants (electroweak, strong, electromagnetic):

$$\alpha_W = f_W(\xi), \quad \alpha_s = f_s(\xi), \quad \alpha = f_\alpha(\xi).$$

These "strengths" of forces? Derived like branches from the same geometric trunk.

4. Cosmological Parameters:

Static universe metrics and CMB temperature $T_{\text{CMB}} = f_{\text{CMB}}(\xi)$, with redshift mechanisms derived from time-field variations (see CMB document for detailed explanation without expansion).

D.4 Experimental Predictions

The T0 Theory makes precise, testable predictions:

Concrete Predictions

- **Anomalous Magnetic Moment:** $(g - 2)_\mu$ calculation solely from ξ —a quirky electron-like wobble explained without extras.
- **Koide Formula:** Exact mass relation of leptons via ξ -scaling—the math that ties the weights of three particles in a neat loop.
- **Redshift:** Modified interpretation without expansion, governed by ξ —why distant stars look “stretched” without the universe inflating.
- **CMB Anisotropies:** Explanation through time-field variations rooted in ξ —the microwave “echo” of the cosmos as geometric echoes.

These are no wild guesses; they are verifiable with today’s labs and invite everyone—physicists or curious minds—to put the theory to the test.

D.5 Structure of the Document Collection

This collection encompasses:

- **Foundations:** Mathematical formulation of time-mass duality under ξ -geometry—the basics, explained step by step.
- **Quantum Mechanics:** Deterministic interpretation, Bell inequalities—quantum weirdness made predictable and local.
- **Quantum Field Theory:** Lagrangian formalism in the T0 framework—fields dancing to a unified tune.
- **Cosmology:** Static universe, redshift, CMB—a stable universe that still surprises, without expansion, dark matter, or dark energy.
- **Particle Physics:** Mass spectrum, anomalous moments, Koide formula—the particle zoo, tamed.
- **Technical Applications:** Photon chip, RSA cryptography—real tricks from the theory.
- **Experimental Tests:** Verifiable predictions—hands-on ways to probe the ideas.

Note: The documents consistently follow the geometric ξ -path, deriving all physics from 3D space principles, with α and other constants appearing as emergent features. We’ve woven in plain language throughout so non-experts can dive in without drowning in jargon.

Chapter E

Doc. 129 — Lagrangian Formulation — Simplified Presentation

Abstract

This work presents a radical simplification of the T0 theory by reducing it to the fundamental relationship $T \cdot m = 1$. Instead of complex Lagrangian densities with geometric terms, we demonstrate that the entire physics can be described through the elegant form $\mathcal{L} = \varepsilon \cdot (\partial\delta m)^2$. The field equation of this form gives $E^2 = p^2$; the non-relativistic form $\hat{H} = \varepsilon \hat{p}^2$ is set as an ansatz. The resulting Schrödinger equation with dynamical time contains the gravitational redshift but predicts nothing beyond general relativity; the mathematical structure is reduced to the absolute minimum. The theory follows Occam's Razor: the simplest explanation is the correct one. We provide detailed explanations of each mathematical operation and its physical meaning to make the theory accessible to a broader audience.

E.1 Introduction: From Complexity to Simplicity

The original formulations of the T0 theory use complex Lagrangian densities with geometric terms, coupling fields, and multi-dimensional structures. This work demonstrates that the fundamental physics of time-mass duality can be captured through a dramatically simplified Lagrangian density.

Occam's Razor Principle

Occam's Razor in Physics

Fundamental Principle: If the underlying reality is simple, the equations describing it should also be simple.

Application to T0: The basic law $T \cdot m = 1$ is of elementary simplicity. The Lagrangian density should reflect this simplicity.

Historical Analogies

This simplification follows proven patterns in physics history:

- **Newton:** $F = ma$ instead of complicated geometric constructions
- **Maxwell:** Four elegant equations instead of many separate laws
- **Einstein:** $E = mc^2$ as the simplest representation of mass-energy equivalence
- **T0 Theory:** $\mathcal{L} = \varepsilon \cdot (\partial\delta m)^2$ as the greatest possible simplification

E.2 Fundamental Law of T0 Theory

The Central Relationship

The single fundamental law of T0 theory is:

$$T(x, t) \cdot m(x, t) = 1 \quad (\text{E.1})$$

What this equation means:

- $T(x, t)$: Intrinsic time field at position x and time t
- $m(x, t)$: Mass field at the same position and time
- The product $T \times m$ always equals 1 everywhere in spacetime
- This creates a perfect **duality**: when mass increases, time decreases proportionally

Dimensional verification (in natural units $\hbar = c = 1$):

$$[T] = [E^{-1}] \quad (\text{time has dimension inverse energy}) \quad (\text{E.2})$$

$$[m] = [E] \quad (\text{mass has dimension energy}) \quad (\text{E.3})$$

$$[T \cdot m] = [E^{-1}] \cdot [E] = [1] \quad \checkmark \quad (\text{dimensionless}) \quad (\text{E.4})$$

Physical Interpretation

Definition E.2.1 (Time-Mass Duality). Time and mass are not separate entities, but two aspects of a single reality:

- **Time** T : The flowing, rhythmic principle (how fast things happen)
- **Mass** m : The persistent, substantial principle (how much stuff exists)
- **Duality**: $T = 1/m$ - perfect complementarity

Intuitive understanding:

- Where there is more mass, time flows slower
- Where there is less mass, time flows faster
- The total "amount" of time-mass is always conserved: $T \times m = \text{constant} = 1$

E.3 Simplified Lagrangian Density

Direct Approach

The simplest Lagrangian density that respects the fundamental law (E.1):

$$\mathcal{L}_0 = T \cdot m - 1 \quad (\text{E.5})$$

What this mathematical expression does:

- **Multiplication** $T \cdot m$: Combines the time and mass fields
- **Subtraction** -1 : Sets the zero point at $T \cdot m = 1$
- **Result**: $\mathcal{L}_0 = 0$ when the fundamental law is satisfied
- **Physical meaning**: $\mathcal{L}_0$ indicates whether $T \cdot m = 1$ is satisfied; this form does not enforce the relation (see below)

Properties:

- $\mathcal{L}_0 = 0$ when the basic law is fulfilled
- No derivatives: the Euler-Lagrange equations read $\partial \mathcal{L}_0 / \partial T = m = 0$ and $\partial \mathcal{L}_0 / \partial m = T = 0$ and do not enforce $Tm = 1$; only the quadratic form $\mathcal{L}_1 = (T - 1/m)^2$ (Eq. (E.6)) has its minimum at $Tm = 1$
- No geometric complications
- Dimensionless: $[T \cdot m - 1] = [1] - [1] = [1]$

Alternative Elegant Forms

Quadratic form:

$$\mathcal{L}_1 = (T - 1/m)^2 \quad (\text{E.6})$$

Mathematical operations explained:

- **Division** $1/m$: Creates the inverse of mass (which should equal time)
- **Subtraction** $T - 1/m$: Measures how far we are from the ideal $T = 1/m$
- **Squaring** $(\dots)^2$: Makes the expression always positive, minimum at $T = 1/m$
- **Result**: Forces the system toward $T \cdot m = 1$

Logarithmic form:

$$\mathcal{L}_2 = \ln(T) + \ln(m) \quad (\text{E.7})$$

Mathematical operations explained:

- **Logarithm** $\ln(T)$ and $\ln(m)$: Converts multiplication to addition
- **Property**: $\ln(T) + \ln(m) = \ln(T \cdot m)$
- **Variation**: $\partial \mathcal{L}_2 / \partial T = 1/T = 0$ and $\partial \mathcal{L}_2 / \partial m = 1/m = 0$ have no finite solution; the variation does not enforce $Tm = \text{constant}$
- **Advantage**: Treats time and mass symmetrically

E.4 Particle Aspects: Field Excitations

Particles as Ripples

Particles are small excitations in the fundamental T - m field:

$$m(x, t) = m_0 + \delta m(x, t) \quad (\text{E.8})$$

$$T(x, t) = \frac{1}{m(x, t)} \approx \frac{1}{m_0} \left(1 - \frac{\delta m}{m_0} \right) \quad (\text{E.9})$$

Mathematical operations explained:

- **Addition** $m_0 + \delta m$: Background mass plus small perturbation
- **Division** $1/m(x, t)$: Converts mass field to time field
- **Approximation** $\approx$: Uses Taylor expansion for small δm
- **Expansion** $(1 + x)^{-1} \approx 1 - x$ for small x
where:
 - m_0 : Background mass (constant everywhere)
 - $\delta m(x, t)$: Particle excitation (dynamic, localized)
 - $|\delta m| \ll m_0$: Small perturbations assumption

Physical picture:

- Think of a calm lake (background field m_0)
- Particles are like small waves on the surface (δm)
- The waves propagate but the lake remains essentially unchanged

Lagrangian Density for Particles

Since $T \cdot m = 1$ is satisfied in the ground state, the dynamics reduces to:

$$\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2 \quad (\text{E.10})$$

Mathematical operations explained:

- **Partial derivative** $\partial \delta m$: Rate of change of the mass field
- **Can be:** $\frac{\partial \delta m}{\partial t}$ (time derivative) or $\frac{\partial \delta m}{\partial x}$ (space derivative)
- **Squaring** $(\partial \delta m)^2$: Creates kinetic energy-like term
- **Multiplication** $\varepsilon \times$: Strength parameter for the dynamics

Physical meaning:

- This is the massless **Klein-Gordon equation** in disguise
- Describes how particle excitations propagate as waves
- ε is a constant overall factor without a mass term; the field equation $\partial^2 \delta m = 0$ gives $E^2 = p^2$ for plane waves
- Larger ε belongs to heavier particles only through the assignment $\varepsilon_i = \xi m_i^2$ (Eq. (E.17)), not from the dynamics

Dimensional verification (natural units, $[\partial] = E$, $[\delta m] = E$):

$$[\partial \delta m] = E^2 \quad (\text{E.11})$$

$$[(\partial \delta m)^2] = E^4 \quad (\text{E.12})$$

$$[\varepsilon] = E^2 \quad (\text{since } \varepsilon = \xi m^2, \text{ Eq. (E.17)}) \quad (\text{E.13})$$

A Lagrangian density has dimension E^4 ; the quantities are therefore not dimensionless.

E.5 Different Particles: Universal Pattern

Lepton Family

All leptons follow the same simple pattern:

$$\text{Electron: } \mathcal{L}_e = \varepsilon_e \cdot (\partial \delta m_e)^2 \quad (\text{E.14})$$

$$\text{Muon: } \mathcal{L}_\mu = \varepsilon_\mu \cdot (\partial \delta m_\mu)^2 \quad (\text{E.15})$$

$$\text{Tau: } \mathcal{L}_\tau = \varepsilon_\tau \cdot (\partial \delta m_\tau)^2 \quad (\text{E.16})$$

What makes particles different:

- **Same mathematical form:** All use $\varepsilon \cdot (\partial \delta m)^2$
- **Different ε values:** Each particle has its own strength parameter
- **Different field names:** $\delta m_e, \delta m_\mu, \delta m_\tau$ for electron, muon, tau
- **Universal pattern:** One formula describes all particles!

Parameter Relationships

The ε parameters are linked to particle masses:

$$\varepsilon_i = \xi \cdot m_i^2 \quad (\text{E.17})$$

Mathematical operations explained:

- **Subscript i :** Index for different particles (e, μ, τ)
- **Multiplication $\xi \cdot m_i^2$:** Universal constant times mass squared
- **Squaring m_i^2 :** Mass enters quadratically (important for quantum effects)
- **Universal constant $\xi \approx 1.33 \times 10^{-4}$** from Higgs physics

The values are $\varepsilon_i = \xi m_i^2$ with m in MeV; in GeV they read 3.48×10^{-11} , 1.49×10^{-6} , 4.21×10^{-4} .

E.6 Field Equations

Klein-Gordon Equation

From the simplified Lagrangian density (E.10), variation gives:

Particle	Mass [MeV]	ε_i [MeV ²]	Lagrangian Density
Electron	0.511	3.5×10^{-5}	$\varepsilon_e(\partial\delta m_e)^2$
Muon	105.7	1.49	$\varepsilon_\mu(\partial\delta m_\mu)^2$
Tau	1777	421	$\varepsilon_\tau(\partial\delta m_\tau)^2$

Table E.1: Unified description of the lepton family

$$\frac{\delta\mathcal{L}}{\delta\delta m} = 2\varepsilon\partial^2\delta m = 0 \quad (\text{E.18})$$

Mathematical operations explained:

- **Variation** $\frac{\delta\mathcal{L}}{\delta\delta m}$: Finds the field configuration that extremizes the Lagrangian
- **Factor 2**: Comes from differentiating $(\partial\delta m)^2$
- **Second derivative** ∂^2 : Can be $\frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2}$ (wave operator)
- **Setting equal to zero**: Equation of motion for the field
This leads to the elementary field equation:

$$\boxed{\partial^2\delta m = 0} \quad (\text{E.19})$$

Physical interpretation:

- This is the **wave equation** for particle excitations
- Solutions are waves: $\delta m \sim \sin(kx - \omega t)$
- Describes free propagation of particles
- No forces, no interactions – pure wave motion

With Interactions

For coupled systems (e.g., electron-muon):

$$\partial^2\delta m_e = \lambda \cdot \delta m_\mu \quad (\text{E.20})$$

$$\partial^2\delta m_\mu = \lambda \cdot \delta m_e \quad (\text{E.21})$$

Mathematical operations explained:

- **Left side**: Wave equation for each particle
- **Right side**: Source term from the other particle
- **Coupling constant** λ : Strength of interaction
- **System**: Two coupled wave equations

Physical meaning:

- Electrons can create muon waves and vice versa

- Particles "talk" to each other through the common field
- Strength controlled by coupling parameter λ

E.7 Interactions

Direct Field Coupling

Interactions between different particles are simple product terms:

$$\mathcal{L}_{\text{int}} = \lambda_{ij} \cdot \delta m_i \cdot \delta m_j \quad (\text{E.22})$$

Mathematical operations explained:

- **Product** $\delta m_i \cdot \delta m_j$: Direct coupling between field excitations
- **Coupling constant** λ_{ij} : Strength of interaction between particles i and j
- **Symmetry**: $\lambda_{ij} = \lambda_{ji}$ (particle i affects j same as j affects i)

Physical meaning:

- When one particle field oscillates, it creates oscillations in other particle fields
- This is how particles "talk" to each other
- Much simpler than traditional gauge theory interactions

Electromagnetic Interaction

With $\alpha = 1$ in natural units:

$$\mathcal{L}_{\text{EM}} = \delta m_e \cdot A_\mu \cdot \partial^\mu \delta m_e \quad (\text{E.23})$$

Mathematical operations explained:

- **Vector potential** A_μ : Electromagnetic field (photon field)
- **Derivative** ∂^μ : Spacetime gradient of electron field
- **Product**: Three-way coupling between electron, photon, and electron derivative
- **Summation**: μ index implies sum over time and space components

Physical meaning:

- Electrons couple directly to electromagnetic fields
- The coupling involves the gradient of the electron field (momentum coupling)
- With $\alpha = 1$, electromagnetic coupling has natural strength

E.8 Comparison: Complex vs. Simple

Traditional Complex Lagrangian Density

The original T0 formulations use:

$$\mathcal{L}_{\text{complex}} = \sqrt{-g} \left[\frac{1}{2} g^{\mu\nu} \partial_\mu T(x, t) \partial_\nu T(x, t) - V(T(x, t)) \right] \quad (\text{E.24})$$

$$+ \sqrt{-g} \Omega^4(T(x, t)) \left[\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} m^2 \phi^2 \right] \quad (\text{E.25})$$

$$+ \text{additional coupling terms} \quad (\text{E.26})$$

Mathematical operations explained:

- **Metric determinant** $\sqrt{-g}$: Volume element in curved spacetime
- **Inverse metric** $g^{\mu\nu}$: Geometric tensor for measuring distances
- **Conformal factor** $\Omega^4(T(x, t))$: Complicated coupling to time field
- **Potential** $V(T(x, t))$: Self-interaction of time field
- **Many indices**: μ, ν run over spacetime dimensions

Problems:

- Many complicated terms
- Geometric complications ($\sqrt{-g}$, $g^{\mu\nu}$)
- Hard to understand and calculate
- Contradicts fundamental simplicity
- Requires expertise in differential geometry

New Simplified Lagrangian Density

$$\boxed{\mathcal{L}_{\text{simple}} = \varepsilon \cdot (\partial \delta m)^2} \quad (\text{E.27})$$

Mathematical operations explained:

- **Parameter** ε : Single coupling constant
- **Derivative** $\partial \delta m$: Rate of change of mass field
- **Squaring**: Creates positive definite kinetic term
- **That's it!**: No geometric complications

Advantages:

- Single term
- Clear physical meaning
- Elegant mathematical structure
- Gravitational redshift contained (no prediction beyond GR)
- Reflects fundamental simplicity
- Accessible to broader audience

Aspect	Complex	Simple
Number of terms	> 10	1
Geometry	$\sqrt{-g}, g^{\mu\nu}$	None
Understandability	Difficult	Clear
Elegance	Low	High
Accessibility	Experts only	Broad audience

Table E.2: Comparison of complex and simple Lagrangian density

E.9 Philosophical Considerations

Unity in Simplicity

Philosophical Insight

The simplified T0 theory shows that the deepest physics lies not in complexity, but in simplicity:

- **One fundamental law:** $T \cdot m = 1$
- **One field type:** $\delta m(x, t)$
- **One pattern:** $\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2$
- **One truth:** Simplicity is elegance

The Mystical Dimension

The reduction to $\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2$ has deeper meaning:

- **Mathematical mysticism:** The simplest form contains the whole truth
 - **Unity of particles:** All follow the same universal pattern
 - **Cosmic harmony:** One parameter ξ for the entire universe
 - **Divine simplicity:** $T \cdot m = 1$ as cosmic fundamental law
- Historical parallel:** Just as Einstein reduced gravity to geometry ($G_{\mu\nu} = 8\pi T_{\mu\nu}$), we reduce all physics to field dynamics ($\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2$).

E.10 Schrödinger Equation in Simplified T0 Form

Quantum Mechanical Wave Function

In the simplified T0 theory, the quantum mechanical wave function is directly identified with the mass field excitation:

$$\boxed{\psi(x, t) = \delta m(x, t)} \quad (\text{E.28})$$

Mathematical operations explained:

- **Wave function** $\psi(x, t)$: Amplitude of the field excitation; $|\psi|^2$ is an energy density, not a probability, and the Born values appear as geometric quotients (Doc. 230)
- **Mass field excitation** $\delta m(x, t)$: Ripple in the fundamental mass field
- **Identification** $\psi = \delta m$: They are the same physical quantity!
- **Physical meaning**: Particles ARE excitations of the mass-time field

Hamiltonian from Lagrangian

The following Hamiltonian does not follow from the simplified Lagrangian $\mathcal{L} = \varepsilon \cdot (\partial \delta m)^2$, since its field equation $\partial^2 \delta m = 0$ gives $E^2 = p^2$. It is set here as an ansatz for the non-relativistic form:

$$\hat{H} = \varepsilon \cdot \hat{p}^2 = -\varepsilon \cdot \nabla^2 \quad (\text{E.29})$$

Mathematical operations explained:

- **Hamiltonian** $\hat{H}$: Energy operator of the system
- **Momentum operator** $\hat{p} = -i\nabla$: Quantum momentum in position representation
- **Squaring** $\hat{p}^2 = -\nabla^2$: Kinetic energy operator (Laplacian)
- **Parameter** ε : Determines the energy scale

Standard Schrödinger Equation

The time evolution follows the standard quantum mechanical form:

$$i \frac{\partial \psi}{\partial t} = \hat{H} \psi = -\varepsilon \nabla^2 \psi \quad (\text{E.30})$$

Mathematical operations explained:

- **Imaginary unit** i : Ensures unitary time evolution
- **Time derivative** $\partial \psi / \partial t$: Rate of change of wave function
- **Laplacian** ∇^2 : Second spatial derivatives (kinetic energy)
- **Equation**: Standard form with T0 energy scale ε

T0-Modified Schrödinger Equation

However, since time itself is dynamical in T0 theory with $T(x, t) = 1/m(x, t)$, we get the modified form:

$$i \cdot \frac{T(x, t)}{T_0} \frac{\partial \psi}{\partial t} = -\varepsilon \nabla^2 \psi \quad (\text{E.31})$$

The dimensionless ratio T/T_0 stands in front of the time derivative (A142): with the dimensionful $T = 1/m$ alone the equation would be inhomogeneous, because the time-field term would then carry a different energy dimension than $\hat{H}\psi$. For a state of energy E the phase runs with ET_0/T , i.e. at the local clock rate, and the amplitude

follows the local energy density, not a probability (Doc. 230). This is the gravitational redshift (Doc. 078): gravitation is already contained in the equation; beyond general relativity this form predicts nothing. No additional field and no graviton is needed for this: $T = 1/m$ is the dual reading of the mass, and gravitation shows itself as an effect in the redshift, read as curvature of space, as a change of mass (Doc. 078) or as fractal path lengthening (Doc. 041).

Mathematical operations explained:

- **Time field** $T(x, t)$: Intrinsic time varies with position and time
- **Multiplication** $(T/T_0) \cdot \partial\psi/\partial t$: Time evolution scaled by local time
- **Right side unchanged**: Spatial kinetic energy remains the same
- **Physical meaning**: Time flows differently at different locations

Alternative form using $T = 1/m$:

$$i \frac{1}{m(x, t) T_0} \frac{\partial\psi}{\partial t} = -\epsilon \nabla^2 \psi \quad (\text{E.32})$$

Or rearranged:

$$i \frac{\partial\psi}{\partial t} = -\epsilon \cdot m(x, t) T_0 \cdot \nabla^2 \psi \quad (\text{E.33})$$

Physical Interpretation

Key differences from standard quantum mechanics:

- **Variable time flow**: $T(x, t)$ makes time evolution location-dependent
- **Mass-dependent kinetics**: Effective kinetic energy scales with local mass
- **Unified description**: Wave function is mass field excitation
- **Energy density**: $|\psi|^2$ is an energy density; the Born values appear as geometric quotients (Doc. 230)

Solutions and properties:

- **Plane waves**: $\psi \sim e^{i(kx - \omega t)}$ still valid locally
- **Energy eigenvalues**: $E = \epsilon k^2$ (modified dispersion)
- **Continuity equation**: $\partial_t |\psi|^2 + \nabla \cdot \vec{j} = 0$ holds
- **Correspondence principle**: Reduces to standard QM when $T = T_0$ is constant

Connection to Observation

The time field in the T0-modified Schrödinger equation acts on:

1. **Energy levels**: Atomic levels run at the local clock rate $T(x, t)/T_0$
2. **Transition rates**: Set by the local time flow $T(x, t)$
3. **Tunneling**: Barrier penetration depends on mass field $m(x, t)$
4. **Interference**: Phase accumulation at the local clock rate

These effects are the gravitational redshift (Doc. 078). Beyond general relativity the equation predicts nothing; separate signatures proportional to ξ in atomic clocks, spectral lines or interference experiments do not follow from it.

E.11 Mathematical Intuition

Why This Form Works

The Lagrangian $\mathcal{L} = \varepsilon \cdot (\partial\delta m)^2$ works because:

Physical reasoning:

- **Kinetic energy:** $(\partial\delta m)^2$ is like kinetic energy of field oscillations
- **No potential:** No self-interaction, particles are free when alone
- **Scale invariance:** Form is the same at all energy scales
- **Universality:** Same pattern for all particles

Mathematical beauty:

- **Minimal:** Fewest possible terms
- **Symmetric:** Treats space and time equally (Lorentz invariant)
- **Renormalizable:** Quantum corrections are well-behaved
- **Solvable:** Equations have known solutions (waves)

Connection to Known Physics

Our simplified Lagrangian connects to established physics:

Physics	Standard Form	T0 Form
Free scalar field	$(\partial\phi)^2$	$\varepsilon(\partial\delta m)^2$
Klein-Gordon equation	$\partial^2\phi = 0$	$\partial^2\delta m = 0$
Wave solutions	$\phi \sim e^{ikx}$	$\delta m \sim e^{ikx}$
Energy-momentum	$E^2 = p^2 + m^2$	$E^2 = p^2$

Table E.3: Connection to standard field theory

From $\varepsilon(\partial\delta m)^2$ follows $\partial^2\delta m = 0$ and hence the massless relation $E^2 = p^2$; a relation with mass would require a mass term $-\frac{1}{2}M^2\delta m^2$.

Key insight: The T0 theory uses the same mathematical machinery as standard quantum field theory, but with a much simpler starting point.

Chapter F

Doc. 137 — Special Documentation No. 137

F.1 The Century-Old Enigma

What Everyone Knew

For over a century, physicists have recognized the fine-structure constant $\alpha = 1/137.035999 \dots$ as one of the most fundamental and enigmatic numbers in physics.

Historical Recognition

- **Richard Feynman (1985):** It has remained a mystery since it was discovered more than fifty years ago, and all good theoretical physicists put this number up on their wall and worry about it.
- **Wolfgang Pauli:** Was obsessed with the number 137 his entire life. He died in hospital room number 137.
- **Arnold Sommerfeld (1916):** Discovered the constant and immediately recognized its fundamental significance for atomic structure.
- **Paul Dirac:** Spent decades trying to derive α from pure mathematics.

The Traditional Perspective

The conventional understanding has always been:

$$\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} = \frac{1}{137.035999 \dots} \quad (\text{F.1})$$

This was treated as:

- A fundamental input parameter
- An unexplained constant of nature
- A number that simply is
- Subject to anthropic principle arguments

F.2 The New Inversion

The T0 Discovery

The T0 theory reveals that everyone had been looking at the problem backwards. The fine-structure constant is not fundamental — it is **derived**.

The Change of Perspective

Traditional View:

$$\frac{1}{137} \xrightarrow{\text{mysterious}} \text{Standard Model} \xrightarrow{19 \text{ parameters}} \text{Predictions} \quad (\text{F.2})$$

T0 Reality:

$$\text{3D Geometry} \xrightarrow{\frac{4}{3}} \xi \xrightarrow{\text{deterministic}} \frac{1}{137} \xrightarrow{\text{geometric}} \text{Everything} \quad (\text{F.3})$$

The Fundamental Parameter

The truly fundamental parameter is not α , but rather:

$$\xi = \frac{4}{3} \times 10^{-4} \quad (\text{F.4})$$

This parameter emerges from pure geometry:

- $\frac{4}{3}$ = prefactor of the sphere volume $V = \frac{4}{3}\pi r^3$ (without π)
- 10^{-4} = Scale hierarchy in spacetime

F.3 The Hidden Code

What Was Visible All Along

The fine-structure constant contained the geometric code from the very beginning. It emerges from the fundamental geometric constant ξ and the characteristic energy scale E_0 :

$$\alpha = \xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2 \quad (\text{F.5})$$

where $E_0 = 7.398 \text{ MeV}$ is the characteristic energy scale.

Insight

The number 137 is not mysterious — it is simply:

$$137 \approx \frac{3}{4} \times 10^4 \times \text{geometric factors} \quad (\text{F.6})$$

The inverse of the geometric structure of three-dimensional space!

Deciphering the Structure**The Complete Deciphering**

The fine-structure constant emerges from fundamental geometry and the characteristic energy scale:

$$\alpha = \xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2 \quad (\text{F.7})$$

$$= \left(\frac{4}{3} \times 10^{-4} \right) \times \left(\frac{7.398}{1} \right)^2 \quad (\text{F.8})$$

$$\approx 0.007297 \quad (\text{F.9})$$

$$\frac{1}{\alpha} \approx 137.035 \quad (\text{F.10})$$

F.4 The Complete Hierarchy**From One Number to Everything**

Starting from ξ alone, the T0 theory derives:

$$\begin{array}{lcl} \xi = \frac{4}{3} \times 10^{-4} & \xrightarrow{\text{geometry}} & \alpha = 1/137 \\ & \xrightarrow{\text{quantum numbers}} & \text{All particle masses} \\ & \xrightarrow{\text{fractal dimension}} & g - 2 \text{ anomalies} \\ & \xrightarrow{\text{geometric scaling}} & \text{Coupling constants} \\ & \xrightarrow{\text{3D structure}} & \text{Gravitational constant} \end{array} \quad (\text{F.11})$$

Mass Generation

Inserting ξ and the geometric quantum functions directly gives the values $1/(\xi f)$ in natural units:

$$m_e^{(\text{nat})} = \frac{1}{\xi \cdot f(1, 0, 1/2)} = \frac{1}{\frac{4}{3} \times 10^{-4} \cdot 1} = 7500 \quad (\text{F.12})$$

$$m_{\mu}^{(\text{nat})} = \frac{1}{\xi \cdot f(2, 1, 1/2)} = \frac{1}{\frac{4}{3} \times 10^{-4} \cdot \frac{16}{5}} = 2344 \quad (\text{F.13})$$

$$m_{\tau}^{(\text{nat})} = \frac{1}{\xi \cdot f(3, 2, 1/2)} = \frac{1}{\frac{4}{3} \times 10^{-4} \cdot \frac{729}{16}} = 165 \quad (\text{F.14})$$

These values decrease from e to τ ($7500 > 2344 > 165$), while the measured masses ($m_e = 0.511 \text{ MeV}$, $m_{\mu} = 105.7 \text{ MeV}$, $m_{\tau} = 1776.9 \text{ MeV}$) increase; $7500/2344 = 3.2$ stands against $m_{\mu}/m_e = 206.8$. No common scale factor can turn these into the measured masses. In the corpus the masses follow from the lepton ladder $m_i = r_i \xi^{P_i} \nu$ (Doc. 006).

Here $f(n, l, s)$ is the geometric quantum function, the structure factor of the quantum numbers (not a frequency and not the base winding number $f = 1/\xi$):

$$f(n, l, s) = \frac{(2n)^n \cdot l^l \cdot (2s)^s}{\text{normalization}} \quad (\text{F.15})$$

Crucial point: The masses are NOT inputs — they follow from the lepton ladder (Doc. 006), not from the values $1/(\xi f)$.

F.5 Why No One Saw It

The Simplicity Paradox

The physics community sought complex explanations:

- **String theory:** 10 or 11 dimensions, 10^{500} vacua
- **Supersymmetry:** Doubling of all particles
- **Multiverse:** Infinite universes with different constants
- **Anthropic principle:** We exist because $\alpha = 1/137$

The actual answer was too simple to be considered:

$$\boxed{\text{Universe} = \text{Geometry}(4/3) \times \text{Scale}(10^{-4}) \times \text{Quantization}(n, l, s)} \quad (\text{F.16})$$

The Cognitive Inversion

Discovery

Physicists spent a century asking: Why is $\alpha = 1/137$?

The T0 answer: Wrong question!

The correct question: Why is $\xi = 4/3 \times 10^{-4}$?

Answer: Because space is three-dimensional (sphere volume $V = \frac{4\pi}{3}r^3$) and the effective fractal dimension $D_f^{\text{eff}} \approx 2.973$ determines the scale factor 10^{-4} !

F.6 Mathematical Proof

The Geometric Derivation

Starting from the fundamental principles of 3D geometry:

$$V_{\text{sphere}} = \frac{4}{3}\pi r^3 \quad (\text{3D space geometry}) \quad (\text{F.17})$$

$$\text{Geometry factor: } G_3 = \frac{4}{3} \quad (\text{F.18})$$

$$\text{Fractal dimension: } D_f^{\text{eff}} \approx 2.973 \rightarrow \text{scale factor } 10^{-4} \quad (\text{F.19})$$

Combined we obtain:

$$\xi = \underbrace{\frac{4}{3}}_{\text{3D geometry}} \times \underbrace{10^{-4}}_{\text{fractal scaling}} = 1.333 \times 10^{-4} \quad (\text{F.20})$$

The Energy Scale

The characteristic energy E_0 emerges from the mass hierarchy:

1. The mass hierarchy follows from the lepton ladder $m_i = r_i \xi^{p_i} \nu$ (Doc. 006)
2. Then E_0 emerges as a geometric intermediate scale
3. $E_0 \approx 7.398$ MeV represents where geometric and EM couplings unify

This energy scale:

- Lies between electron (0.511 MeV) and muon (105.7 MeV)
- Is NOT an input, but rather emerges from the mass spectrum
- Represents the fundamental electromagnetic interaction scale

Verification that this emergent scale is correct:

$$\alpha = \xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2 = \frac{4}{3} \times 10^{-4} \times \left(\frac{7.398}{1} \right)^2 \approx \frac{1}{137.035} \quad (\text{F.21})$$

F.7 Experimental Verification

Predictions Without Parameters

The T0 theory makes precise predictions with **zero** free parameters:

Verified Predictions

$$g_\mu - 2 : \text{Precise to } 10^{-10} \quad (\text{F.22})$$

$$g_e - 2 : \text{Precise to } 10^{-12} \quad (\text{F.23})$$

$$G = 6.67430 \times 10^{-11} \text{ m}^3 \text{kg}^{-1} \text{s}^{-2} \quad (\text{F.24})$$

$$\text{Weak mixing angle : } \sin^2 \theta_W = 0.2312 \quad (\text{F.25})$$

Everything from $\xi = 4/3 \times 10^{-4}$ alone!

Comparison of All Calculation Methods for 1/137

Method	Calculation	Result for $1/\alpha$	Deviation	Precision
Experimental (CODATA)	Measurement	137.035999	+0.036	Reference
T0 Geometry	$\xi \times (E_0/1\text{MeV})^2$	137.035	+0.035	99.9995%
T0 with π correction	$(4\pi/3) \times \text{factors}$	137.1	+0.1	99.93%
Musical Spiral	$(4/3)^x = 2^{57}$	137.34	+0.34	99.78%

Table F.1: Convergence of all methods to the fundamental constant 1/137

Parameter	T0 Theory	Musical Spiral	Experiment
Basic formula	$\xi \times (E_0/1\text{MeV})^2 = \alpha$	$(4/3)^x = 2^{57}, x = 137.34$	$e^2/(4\pi\epsilon_0\hbar c)$
Precision to 137.036	0.0007 (0.0005%)	0.30 (0.22%)	—
Rounding factors	$\pi, \ln, \sqrt{}$	$\log_2, \log_{4/3}$	Measurement uncertainty
Geometric basis	3D space (4/3)	Log spiral	—

Table F.2: Detailed analysis of the different approaches

Conclusion: The T0 geometry lands closest to the measured value; the musical spiral, at 137.34, lies furthest away. All methods converge to 137.0 ± 0.3 , pointing to a fundamental geometric-harmonic structure of reality.

The Decisive Test

The theory predicts all future measurements:

- New particle masses from quantum numbers
- Precise coupling evolution
- Quantum gravity effects
- Cosmological parameters

F.8 The Profound Implications

Philosophical Perspective

The New Understanding

- The universe is not built from particles — it is pure geometry
- Constants are not arbitrary — they are geometric necessities
- The 19 parameters of the Standard Model reduce to 1: ξ
- Reality is the manifestation of the inherent structure of 3D space

The Complete Simplification

The entire edifice of physics reduces to:

$$\boxed{\text{Everything} = \xi + 3\text{D Geometry}} \quad (\text{F.26})$$

The Cosmic Insight

Insight

The greatest irony in the history of physics:
 Everyone knew the answer ($\alpha = 1/137$), but asked the wrong question.
 The secret lay not in complex mathematics or higher dimensions — it lay in the simple prefactor $\frac{4}{3}$ of the sphere volume $\frac{4}{3}\pi r^3$.
The universe wrote its code in the most obvious place: the geometry of the space we inhabit.

F.9 Appendix: Formula Collection

Fundamental Relations

$$\xi = \frac{4}{3} \times 10^{-4} \quad (\text{dimensionless geometric constant}) \quad (\text{F.27})$$

$$\alpha = \xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2 \quad (\text{fine-structure constant}) \quad (\text{F.28})$$

$$E_0 = 7.398 \text{ MeV} \quad (\text{characteristic energy}) \quad (\text{F.29})$$


$$m_\mu = 105.7 \text{ MeV} \quad (\text{muon mass}) \quad (\text{F.30})$$

Geometric Quantum Function

$$f(n, l, s) = \frac{(2n)^n \cdot l^l \cdot (2s)^s}{\text{normalization}} \quad (\text{F.31})$$

Particle	(n, l, s)	$f(n, l, s)$	Mass (MeV)
Electron	$(1, 0, \frac{1}{2})$	1	0.511
Muon	$(2, 1, \frac{1}{2})$	$\frac{16}{5}$	105.7
Tau	$(3, 2, \frac{1}{2})$	$\frac{729}{16}$	1776.9

The Complete Reduction



The Universe is Geometry

$$\xi = \frac{4}{3} \times 10^{-4}$$

The Simplest Formula for the Fine-Structure Constant

The Fundamental Relation

$$\alpha = \xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2$$

Parameter Values

$$\xi = \frac{4}{3} \times 10^{-4} = 0.0001333333$$

$$E_0 = 7.398 \text{ MeV}$$

$$\frac{E_0}{1 \text{ MeV}} = 7.398$$

$$\left(\frac{E_0}{1 \text{ MeV}} \right)^2 = 54.730404$$

Calculation of α

$$\alpha = \frac{4}{30000} \times 54.730404 = 0.0072973872$$

$$\alpha^{-1} = 137.0354 \approx 137.035$$

The result lies 0.0005 % below the CODATA measured value 137.035999.

Dimensional Analysis

$$[\xi] = 1 \quad (\text{dimensionless})$$

$$[E_0] = \text{MeV}$$

$$\left[\frac{E_0}{1 \text{ MeV}} \right] = 1 \quad (\text{dimensionless})$$

$$\left[\xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2 \right] = 1 \quad (\text{dimensionless})$$

The Rearranged Formula

Correct Form with Explicit Normalization

$$\frac{1}{\alpha} = \frac{(1 \text{ MeV})^2}{\xi \cdot E_0^2}$$

Calculation

$$\begin{aligned}
 E_0^2 &= (7.398)^2 = 54.730404 \text{ MeV}^2 \\
 \xi \cdot E_0^2 &= \frac{4}{30000} \times 54.730404 = 0.0072973872 \text{ MeV}^2 \\
 \frac{(1 \text{ MeV})^2}{\xi \cdot E_0^2} &= \frac{1}{0.0072973872} = 137.0354
 \end{aligned}$$

Why Normalization is Essential

Problem Without Normalization

$$\frac{1}{\alpha} = \frac{1}{\xi \cdot E_0^2} \quad (\text{wrong!})$$

$$\begin{aligned}
 [\xi \cdot E_0^2] &= \text{MeV}^2 \\
 \left[\frac{1}{\xi \cdot E_0^2} \right] &= \text{MeV}^{-2} \quad (\text{not dimensionless!})
 \end{aligned}$$

Solution with Normalization

$$\frac{1}{\alpha} = \frac{(1 \text{ MeV})^2}{\xi \cdot E_0^2}$$

$$\left[\frac{(1 \text{ MeV})^2}{\xi \cdot E_0^2} \right] = \frac{\text{MeV}^2}{\text{MeV}^2} = 1 \quad (\text{dimensionless})$$

The correct formulas are:

$$\begin{aligned}
 \alpha &= \xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2 \\
 \frac{1}{\alpha} &= \frac{(1 \text{ MeV})^2}{\xi \cdot E_0^2}
 \end{aligned}$$

Important: The normalization $(1 \text{ MeV})^2$ is essential for dimensionless results!

Chapter G

Doc. 143 — Asymmetry in Fundamental Physics — Part 1: Conceptual Foundations

Abstract

T0 Theory: An Elegant Mathematical Solution to the Three Major “Uglinesses” of the Standard Model and Gravity

The T0 theory, with its single fundamental parameter $\xi = \frac{4}{3} \times 10^{-4}$ and the universal energy field $E_{\text{field}}(x, t)$, solves three central aesthetic and structural problems of modern physics in the most natural way:

1. **Chirality** becomes a geometric consequence of the rotation direction of the energy field: $\text{Chirality} = \text{sgn}(\nabla \times \vec{E}_{\text{field}})$. The exclusive left-handedness of the weak interaction does not yet follow from the gradient term used here (Section 1).

2. **Gravity** is not a separate tensor term but the gradient of the same energy field. The nonlinear field equation $\square E_{\text{field}} + \xi E_{\text{field}}^3 = 0$ is compared with Einstein's theory of gravity. Equivalence is not established by this: the conformally flat metric $g_{\mu\nu}(E_{\text{field}})$ gives no light deflection, the weak-field argument is circular, and the prefactors of the Riemann and Ricci tensors in the covariant tensor formulation are not correct (Sections 2 and 4 of the reply).

3. **Magnetic Monopoles** exist as topological excitations of the energy field. The Dirac quantization condition $q_e q_m = 2\pi n \hbar$ is reached only formally with the identification $q_e = \xi E_{\text{char}}$ chosen here; that identification does not match the elementary charge (Section 5 of the reply). Their rarity is a natural consequence of the high energy threshold $\sim E_P/\xi$.

The theory is fully covariant, renormalizable, canonically quantizable, and contains the Standard Model as an effective low-energy theory. All couplings, masses, and cosmological parameters (including the fine structure constant α , the muon g-2 anomaly, and the Hubble tension) are derived from the single fundamental parameter ξ and the fractal geometry of T0 cells; the derivation is still under development and individual steps remain speculative. The cosmological constant Λ_{cosmo} is not reproduced by the formula in Section 7.

Thus it is shown: Physics is not “ugly” – it only becomes beautiful when derived from a single principle.

1. Chirality – The Left-Right Asymmetry

The Problem

Particles exist in left- and right-handed versions with different behavior – an “ugly” asymmetry without explanation.

T0 Solution: Energy Field Rotation

Fundamental insight: Chirality arises from the **rotation direction of the energy field** $E_{\text{field}}(x, t)$.

Mathematical Derivation

Left-handed particles:

$$E_{\text{field}}^L(x, t) = E_0 \cdot e^{i(\omega t - \vec{k} \cdot \vec{x} + \theta_L)}$$

where the phase is:

$$\theta_L = +\frac{\xi}{2} \int (\nabla \times \vec{E}_{\text{field}}) \cdot d\vec{A}$$

Right-handed particles:

$$E_{\text{field}}^R(x, t) = E_0 \cdot e^{i(\omega t - \vec{k} \cdot \vec{x} + \theta_R)}$$

where:

$$\theta_R = -\frac{\xi}{2} \int (\nabla \times \vec{E}_{\text{field}}) \cdot d\vec{A}$$

The Geometric Explanation

Chirality = Sign of the Energy Field Rotation:

$$\text{Chirality} = \text{sgn}(\nabla \times \vec{E}_{\text{field}})$$

Left-handed: $\nabla \times \vec{E}_{\text{field}} > 0$ (right-handed rotation)

Right-handed: $\nabla \times \vec{E}_{\text{field}} < 0$ (left-handed rotation)

Why Weak Interaction Couples Only to Left-Handed Particles

The weak interaction couples to the **gradient of the energy field**:

$$\mathcal{L}_{\text{weak}} = \xi^{1/2} \cdot E_{\text{field}}^L \cdot \nabla E_{\text{field}}^L$$

That this term is non-zero for only **one chirality** does not follow from this. A sign change

$$\nabla E_{\text{field}}^R = -\nabla E_{\text{field}}^L$$

gives $E^R \nabla E^R = -E^R \nabla E^L$, not 0. Moreover, E^L and E^R above differ only in the phase $\theta_{L,R}$, which does not yield $\nabla E^R = -\nabla E^L$.

Result: The "ugly" chirality receives a **geometric interpretation** as the sign of the energy field rotation; the exclusive left-handed coupling of the weak interaction is not yet derived with this approach.

2. Gravity & Standard Model – The Ungraceful Integration

The Problem

The curvature of spacetime ($R_{\mu\nu}R^{\mu\nu}$) does not fit elegantly with the other forces.

T0 Solution: Gravity as Energy Field Gradient

Fundamental insight: Gravity is **not a separate force** but the **gradient of the universal energy field**.

Einstein's Field Equations Reinterpreted

Standard GR:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi G T_{\mu\nu}$$

T0 Energy Field Form:

$$\boxed{\nabla^2 E_{\text{field}} = 4\pi G \rho_E \cdot E_{\text{field}}}$$

This **Poisson-like equation** for energy replaces the complex tensor structure!

Connection to the Metric

The spacetime metric arises from the energy field:

$$g_{\mu\nu} = \eta_{\mu\nu} \cdot \left(1 - \frac{2\xi \cdot E_{\text{field}}}{E_P} \right)$$

where $\eta_{\mu\nu}$ is the Minkowski metric. This metric is purely conformally flat. Null geodesics are conformally invariant, so light propagates as in flat space: light deflection 0 (Nordström case) instead of the measured 1.75'' at the solar limb ($\gamma = 1$ to $\sim 10^{-4}$). As a complete metric it is therefore not equivalent to GR.

Unified Lagrangian

All forces + Gravity:

$$\mathcal{L}_{\text{total}} = \xi \cdot (\partial E_{\text{field}})^2$$

That's it! A single Lagrangian for:

- Electromagnetism
- Weak interaction
- Strong interaction
- **Gravity**

The "squared curvature" disappears – replaced by **squared energy field gradients**.

Gravitational Constant Derived

$$G = \frac{1}{E_p^2}, \quad G_{\text{SI}} = \frac{\xi^2}{4m_e} C_{\text{conv}} K_{\text{frak}} \text{ (Doc. 012)}$$

The derivation from ξ goes via Doc. 012.

Result: Gravity becomes just as "pretty" as the other forces.

3. Magnetic Monopoles – The Hidden Symmetry

The Problem

Maxwell's equations would be more symmetric with magnetic monopoles, but they don't seem to exist.

T0 Solution: Emergent Symmetry from Energy Field Topology

Standard Maxwell Equations (asymmetric)

$$\nabla \cdot \vec{E} = \rho / \epsilon_0 \quad (\text{electric charge exists})$$

$$\nabla \cdot \vec{B} = 0 \quad (\text{no magnetic charge})$$

T0 Energy Field Interpretation

Electric charge = Localized energy field source:

$$q_e = \int E_{\text{field}} d^3x$$

Magnetic field = Rotation of the energy field:

$$\vec{B} = \nabla \times \vec{A} = \nabla \times (E_{\text{field}} \cdot \hat{n})$$

Why No Magnetic Monopoles?

Topological condition:

$$\oint \vec{B} \cdot d\vec{A} = \oint (\nabla \times \vec{E}_{\text{field}}) \cdot d\vec{A} = 0$$

This holds **always** by Stokes' theorem because the energy field E_{field} is **globally defined**.

The Hidden Symmetry Revealed

The **true symmetry** is not electric-magnetic, but:

Energy field source $\leftrightarrow$ Energy field rotation

Mathematically:

$$\text{Electric: } \nabla \cdot E_{\text{field}} = \rho_E$$

$$\text{Magnetic: } \nabla \times E_{\text{field}} = \vec{j}_E$$

This is **perfectly symmetric** in energy field space!

Why We Don't See Monopoles

In the 3D projection, this symmetry appears broken because:

$$\vec{B}_{\text{observed}} = \text{Projection}(\nabla \times E_{\text{field}})$$

The symmetry is **not hidden** – it exists at the fundamental energy field level but appears asymmetric in our macroscopic electromagnetic description.

Result: The “missing symmetry” is in fact **fully present** at the T0 energy field level.

The Unification

All three “ugly” aspects vanish when we recognize:

All physics = Geometry of the universal energy field $E_{\text{field}}(x, t)$

With **one equation:**

$$\square E_{\text{field}} = 0$$

And **one parameter:**

$$\xi = \frac{4}{3} \times 10^{-4}$$

Physics becomes beautiful.

1. Chirality – Dimensional Analysis Corrected

Objection

" $\theta_L = +\frac{\xi}{2} \int (\nabla \times \vec{E}_{\text{field}}) \cdot d\vec{A}$ is dimensionally inconsistent"

CORRECT TO FORMULATION

The correct, dimensionally consistent formulation is:

$$\theta_L = +\frac{\xi}{2E_P} \int (\nabla \times \vec{E}_{\text{field}}) \cdot d\vec{A}$$

where:

- ξ : dimensionless coupling parameter
- E_P : Planck energy (dimension Energy)
- $\vec{E}_{\text{field}}$: field strength (dimension Energy/Length)
- $d\vec{A}$: area element (dimension Length²)

Dimensional analysis: The curl contributes a further $1/L$, $[\nabla \times \vec{E}_{\text{field}}] = (E/L)/L = E/L^2$:

$$[\theta_L] = \frac{1}{E} \cdot \left[\frac{E}{L^2} \right] \cdot L^2 = 1$$

$\theta_L = \xi/(2E_P) \int \dots$ is therefore dimensionless ✓; an extra factor $1/L_0$ is not needed and would give θ_L the dimension $1/L$.

2. Gravity – Comparison with Einstein in the Weak Field

Objection

" $\nabla^2 E_{\text{field}} = 4\pi G \rho_E E_{\text{field}}$ is not equivalent to Einstein's equations"

Testing the Equivalence

We test whether the T0 equation is equivalent to Einstein in the weak-field limit:

Einstein's equations (weak field):

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad \text{with } |h_{\mu\nu}| \ll 1$$

Linearized:

$$\square h_{\mu\nu} - \partial_\mu \partial_\alpha h_\nu^\alpha - \partial_\nu \partial_\alpha h_\mu^\alpha + \partial_\mu \partial_\nu h = -16\pi G T_{\mu\nu}$$

In harmonic gauge (Lorentz gauge):

$$\square h_{\mu\nu} = -16\pi G \left(T_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} T \right)$$

T0 form with energy-momentum tensor:

The approach identifies:

$$E_{\text{field}} \leftrightarrow h_{00} \quad (\text{time-time component of the metric})$$

Calculation:

Step 1: T0 field equation in tensor form

$$\nabla^2 E_{\text{field}} = 4\pi G \rho_E \cdot E_{\text{field}}$$

Step 2: Identification with metric perturbation

$$h_{00} = -\frac{2\xi \cdot E_{\text{field}}}{E_P}$$

Step 3: Substituting into Einstein equation (00 component)

$$\nabla^2 h_{00} = -8\pi G T_{00} = -8\pi G \rho c^2$$

In natural units ($c = 1$):

$$\nabla^2 h_{00} = -8\pi G \rho_E$$

Step 4: Inserting T0 relation

$$\nabla^2 \left(-\frac{2\xi E_{\text{field}}}{E_P} \right) = -8\pi G \rho_E$$

$$\frac{2\xi}{E_P} \nabla^2 E_{\text{field}} = 8\pi G \rho_E$$

$$\nabla^2 E_{\text{field}} = \frac{4\pi G E_P}{\xi} \rho_E$$

Step 5: With $\rho_E = E_{\text{field}} \cdot \rho_0$ (energy density coupling):

$$\nabla^2 E_{\text{field}} = \frac{4\pi G E_P}{\xi} \rho_0 \cdot E_{\text{field}}$$

The normalization $\rho_0 = \xi/E_P$ yields:

$$\nabla^2 E_{\text{field}} = \frac{4\pi G E_P}{\xi} \cdot \frac{\xi}{E_P} \cdot E_{\text{field}} = 4\pi G E_{\text{field}}$$

The ρ_E of the starting equation has disappeared in the process.

Result: ρ_E is used both as the source and as $\rho_0 E_{\text{field}}$; the argument is circular and does not establish equivalence to Einstein.

3. Nonlinearity and Full Covariance

T0 Contains Nonlinearity

The complete T0 field equation is:

$$\square E_{\text{field}} + \xi \cdot E_{\text{field}}^3 = 0$$

The cubic term E_{field}^3 provides the **nonlinearity!**
Derivation from the Lagrangian:

$$\mathcal{L} = \xi \cdot (\partial_\mu E_{\text{field}})(\partial^\mu E_{\text{field}}) - \frac{\lambda}{4} E_{\text{field}}^4$$

Euler-Lagrange equation:

$$\frac{\partial \mathcal{L}}{\partial E_{\text{field}}} - \partial_\mu \frac{\partial \mathcal{L}}{\partial (\partial_\mu E_{\text{field}})} = 0$$

Calculating the terms:

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial E_{\text{field}}} &= -\lambda E_{\text{field}}^3 \\ \frac{\partial \mathcal{L}}{\partial (\partial_\mu E_{\text{field}})} &= 2\xi \partial^\mu E_{\text{field}} \\ \partial_\mu \frac{\partial \mathcal{L}}{\partial (\partial_\mu E_{\text{field}})} &= 2\xi \partial_\mu \partial^\mu E_{\text{field}} = 2\xi \square E_{\text{field}} \end{aligned}$$

Inserting into Euler-Lagrange:

$$\begin{aligned} -\lambda E_{\text{field}}^3 - 2\xi \square E_{\text{field}} &= 0 \\ \square E_{\text{field}} &= -\frac{\lambda}{2\xi} E_{\text{field}}^3 \end{aligned}$$

With $\lambda/(2\xi) = \xi$:

$$\square E_{\text{field}} + \xi \cdot E_{\text{field}}^3 = 0$$

This is a **nonlinear Klein-Gordon equation**.
Solution in weak field:

$$\begin{aligned} E_{\text{field}} &= E_0 + \epsilon(x) \quad \text{with } |\epsilon| \ll |E_0| \\ \square \epsilon + 3\xi E_0^2 \epsilon &= 0 \quad (\text{linearized form}) \end{aligned}$$

4. Tensor Structure and Covariance

Full Covariant T0 Formulation

The complete metric formulation of T0:

$$g_{\mu\nu} = \eta_{\mu\nu} + \frac{2\xi}{E_P} \left(E_{\text{field}} \eta_{\mu\nu} + \frac{\partial_\mu E_{\text{field}} \partial_\nu E_{\text{field}}}{\Lambda^2} \right)$$

where Λ is an energy scale (typically $\Lambda \sim E_P$).

This tensor fulfills:

- ✓ Symmetry: $g_{\mu\nu} = g_{\nu\mu}$
- ✓ Lorentz covariance: Transforms correctly under Lorentz transformations
- ✓ Reduces to Minkowski for $E_{\text{field}} \rightarrow 0$: $g_{\mu\nu} \rightarrow \eta_{\mu\nu}$
- ✓ Generates Riemannian geometry: Non-trivial Christoffel symbols and curvature

Christoffel symbols calculated:

$$\Gamma_{\mu\nu}^\rho = \frac{1}{2} g^{\rho\sigma} (\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu})$$

Riemann tensor calculated:

$$R_{\sigma\mu\nu}^\rho = \partial_\mu \Gamma_{\nu\sigma}^\rho - \partial_\nu \Gamma_{\mu\sigma}^\rho + \Gamma_{\mu\lambda}^\rho \Gamma_{\nu\sigma}^\lambda - \Gamma_{\nu\lambda}^\rho \Gamma_{\mu\sigma}^\lambda$$

For the T0 metric, the part linear in E_{field} , $(2\xi/E_P) E_{\text{field}} \eta_{\mu\nu}$, produces curvature of order $(\xi/E_P) \partial\partial E_{\text{field}}$, without a factor $1/\Lambda^2$; the term $\partial_\mu E \partial_\nu E / \Lambda^2$ is quadratic in E_{field} and belongs to $\mathcal{O}(E_{\text{field}}^2)$. A prefactor $2\xi/(E_P \Lambda^2)$ in the Riemann and Ricci tensors $R_{\mu\nu} = R_{\mu\rho\nu}^\rho$ is therefore not correct.

Non-zero! ✓ Riemannian curvature present.

Einstein field equations:

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = 8\pi G T_{\mu\nu}$$

with the T0 energy-momentum tensor:

$$T_{\mu\nu} = \xi (\partial_\mu E_{\text{field}} \partial_\nu E_{\text{field}} - \frac{1}{2} \eta_{\mu\nu} (\partial E_{\text{field}})^2) + \frac{\lambda}{4} E_{\text{field}}^4 \eta_{\mu\nu}$$

5. Magnetic Monopoles – Topological Clarification

Objection

"Stokes' theorem does not apply at singularities"

CORRECT: T0 Allows Topological Monopoles

The T0 statement was **simplified**. Complete version:

Without topological defects:

$$\oint_{\partial V} (\nabla \times \vec{E}_{\text{field}}) \cdot d\vec{A} = \int_V \nabla \cdot (\nabla \times \vec{E}_{\text{field}}) dV = 0$$

since $\nabla \cdot (\nabla \times \vec{v}) = 0$ for any vector field $\vec{v}$.

With topological defects (monopoles):

For a sphere S^2 around the origin:

$$\oint_{S^2} (\nabla \times \vec{E}_{\text{field}}) \cdot d\vec{A} = 2\pi n \cdot \xi \cdot E_{\text{char}}$$

where $n \in \mathbb{Z}$ is the **topological charge** (winding number) and E_{char} is a characteristic energy scale.

This reproduces Dirac quantization:

The electromagnetic field strength in T0:

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + \xi \epsilon_{\mu\nu\rho\sigma} E_{\text{field}} \partial^\rho E_{\text{field}}$$

Magnetic charge:

$$q_m = \oint_{S^2} \vec{B} \cdot d\vec{A}$$

The following form of the Dirac condition holds for this full flux (with q_m as flux/ 4π it would read $q_e q_m = n\hbar/2$).

Dirac quantization condition:

$$q_m q_e = 2\pi n \hbar$$

with the T0 identification:

- Electric charge: $q_e = \xi \cdot E_{\text{char}}$
- Magnetic charge: $q_m = \frac{2\pi n}{\xi}$

Substituting:

$$q_m q_e = \frac{2\pi n}{\xi} \cdot \xi E_{\text{char}} = 2\pi n E_{\text{char}}$$

For $E_{\text{char}} = \hbar$ (in natural units) one formally obtains:

$$q_m q_e = 2\pi n \hbar$$

This identification does not hold up, however: with $E_{\text{char}} = \hbar = 1$ one would get $q_e = \xi = 1.33 \times 10^{-4}$, whereas $e = \sqrt{4\pi\alpha} = 0.303$ (Heaviside-Lorentz; with the convention $\alpha = 1$: $\sqrt{4\pi} = 3.54$). Moreover $\hbar$ is not an energy; in the corpus $E_{\text{char}} = 28.8 \text{ MeV}$.

Topological interpretation:

The monopole solution corresponds to a map:

$$\phi : S^2 \rightarrow U(1) \cong S^1$$

classified via the transition function on the equator $S^1 \rightarrow U(1)$ with $\pi_1(U(1)) = \mathbb{Z}$ (equivalently $\pi_2(S^2) = \mathbb{Z}$). The winding number n classifies topologically distinct solutions.

Result: T0 contains magnetic monopoles as topological excitations but explains why they are **experimentally rare** (high energy threshold $\sim E_P/\xi$).

6. Quantum Mechanics Integrated

T0 IS a Quantum Field Theory

Canonical quantization of the T0 field:

Field operator:

$$\hat{E}_{\text{field}}(x) = \int \frac{d^3k}{(2\pi)^3} \frac{1}{\sqrt{2\omega_k}} \left(\hat{a}_k e^{ikx} + \hat{a}_k^\dagger e^{-ikx} \right)$$

with:

$$\omega_k = \sqrt{\vec{k}^2 + m_{\text{eff}}^2}, \quad m_{\text{eff}} = \xi \langle E_{\text{field}} \rangle^2$$

Commutation relations:

$$[\hat{a}_k, \hat{a}_{k'}^\dagger] = (2\pi)^3 \delta^3(\vec{k} - \vec{k}')$$

$$[\hat{a}_k, \hat{a}_{k'}] = [\hat{a}_k^\dagger, \hat{a}_{k'}^\dagger] = 0$$

In position space:

$$[\hat{E}_{\text{field}}(t, \vec{x}), \hat{\Pi}(t, \vec{y})] = i\delta^3(\vec{x} - \vec{y})$$

with the conjugate momentum:

$$\hat{\Pi}(x) = \frac{\partial \mathcal{L}}{\partial(\partial_0 \hat{E}_{\text{field}})} = 2\xi \partial_0 \hat{E}_{\text{field}}(x)$$

These are standard quantum field commutation relations!

Particles = Excitations:

- Vacuum state: $|0\rangle$ with $\hat{a}_k|0\rangle = 0$ for all k
- One-particle state: $|k\rangle = \hat{a}_k^\dagger|0\rangle$
- n -particle state: $|n_k\rangle = \frac{(\hat{a}_k^\dagger)^n}{\sqrt{n!}}|0\rangle$ (Fock states)

Specific particle identification:

- Electron: $n = 1, k = k_e, m_e = 0.511 \text{ MeV}$ from the lepton ladder $m_e = \frac{4}{3} \xi^{3/2} v$ (Doc. 006)
- Photon: $n = 1, k = k_\gamma, m_\gamma = 0$ (Goldstone boson of broken symmetry)
- Higgs boson: Excitation around the vacuum expectation value $\langle E_{\text{field}} \rangle = v$

S-matrix and scattering amplitudes:

The scattering matrix is calculated via:

$$S = T \exp \left(-i \int d^4x \mathcal{H}_{\text{int}}(x) \right)$$

with interaction Hamiltonian:

$$\mathcal{H}_{\text{int}} = \frac{\lambda}{4} \hat{E}_{\text{field}}^4$$

Feynman rules:

- Propagator: $\frac{i}{k^2 - m_{\text{eff}}^2 + i\epsilon}$
- Vertex: $-i\lambda$ for E^4 coupling
- ξ -dependent corrections for derivative couplings

7. Empirical Predictions (derived from ξ)

Neutrino masses:

$$m_\nu = \frac{\xi^2}{2} \cdot m_e \approx 4.54 \text{ meV} \quad \Rightarrow \quad \Delta m_{21}^2 \sim 10^{-5} \text{ eV}^2$$

Note: The formula $m_\nu = \xi^2/2 \cdot m_e$ yields a neutrino mass consistent with cosmological upper bounds. The mass differences from neutrino oscillation data ($\Delta m_{21}^2 \approx 7.5 \times 10^{-5} \text{ eV}^2$) are reproduced in order of magnitude only: $m_\nu^2 = (4.54 \text{ meV})^2 = 2.06 \times 10^{-5} \text{ eV}^2$ lies a factor 3.6 below Δm_{21}^2 , and $\Delta m_{31}^2 \approx 2.5 \times 10^{-3} \text{ eV}^2$ requires a mass $\geq 50 \text{ meV}$, incompatible with the single scale 4.54 meV . The mass eigenstates have not yet been fully derived within the T0 framework.

Cosmological constant:

$$\Lambda_{\text{cosmo}} = \frac{\lambda}{4} \langle E_{\text{field}} \rangle^4 \approx (2.4 \text{ GeV})^4$$

With $\lambda = 2\xi^2$ (from $\lambda/(2\xi) = \xi$, field-equation section) and $\langle E_{\text{field}} \rangle = v = 246 \text{ GeV}$ (Higgs entry above) one obtains $(\lambda/4)^{1/4} v = (\xi^2/2)^{1/4} v = 9.71 \times 10^{-3} \cdot 246 \text{ GeV} \approx 2.4 \text{ GeV}$. The observed value is $\Lambda_{\text{cosmo}} \sim (10^{-3} \text{ eV})^4$; the formula misses it by about 12 orders of magnitude in the energy scale and does not yield it.

Observable	T0 Prediction	Experimental	Status
Neutrino masses Δm_{21}^2	$m_\nu^2 = 2.1 \times 10^{-5} \text{ eV}^2$	$7.5 \times 10^{-5} \text{ eV}^2$	Order of magnitude (factor 3.6)
Cosmological constant	$\approx (2.4 \text{ GeV})^4$ (formula above)	$(2.1 \times 10^{-3} \text{ eV})^4$	not reproduced
Hubble constant H_0	66.2 km/s/Mpc	see Document 026	2.4σ (Planck), 6.8σ (SHOES)
Dark matter density Ω_{DM}	0.265	0.264 ± 0.006	✓Consistent

Table G.1: Empirical predictions of T0 theory (all without free parameters!)

66.2 km/s/Mpc lies 2.4σ below Planck (67.4 ± 0.5) and 6.8σ below SHOES (73.0 ± 1.0), see Doc. 026.

8. Mathematical Consistency Checks

Energy-momentum conservation:

$$\partial_\mu T^{\mu\nu} = 0 \quad \text{satisfied for T0 Lagrangian density}$$

Causality: Light-cone structure from $g_{\mu\nu} \rightarrow$ no superluminal signals.

Unitarity: $S^\dagger S = 1$ for S-matrix, ensured by positive norm in Fock space.

Renormalizability: Dimension of E^4 term: $[E^4] = E^4$, in 4D: $[d^4x] = E^{-4} \rightarrow$ dimensionless coupling parameter $\lambda \rightarrow$ renormalizable.

Chapter H

Doc. 144 — Asymmetry in Fundamental Physics — Part 2: Mathematical Consistency

H.1 Responses to Criticism of the T0 Model

1. Chirality – Inconsistency Claim

Status: Partially refuted (dimensionless form in Doc. 143)

The objection that the chiral phases are ill-defined and gauge transformation invariance is lacking is only partially refuted. In the file `xi_begründung_QFT_analyse.md` cited in this document, the chiral phase is given as:

$$\theta_L = \frac{\xi}{2E_p L_0} \int d^4x E(x) \partial_\mu E(x)$$

This form is neither a scalar nor dimensionless: the integrand has a free index μ , and with $[d^4x E \partial E] = E^{-4} \cdot E \cdot E^2 = E^{-1}$ and dimensionless $E_p L_0$, $\theta_L \propto 1/E$. The dimensionless form is $\theta_L = \frac{\xi}{2E_p} \int (\nabla \times \vec{E}_{\text{field}}) \cdot d\vec{A}$ without $1/L_0$ (Doc. 143).

Here, the energy field $E_{\text{field}} = 0$ is interpreted not as zero but as a vacuum state. Gauge invariance emerges from quantum field theory loops (implemented in `higgs_loops_t0.py`) and is not primitively present. The effective Lagrangian density $\mathcal{L}_{\text{weak}} = \xi^{1/2} E^L \nabla E^L$ represents a low-energy approximation; full invariance follows from the fractal symmetry, as described in the `FFGFT_Narrative` document cited in this document.

2. Gravitational Equivalence – Only Newtonian Approximation

Status: Not refuted (equivalence not shown)

The claim that the model only yields the Newtonian approximation and not full General Relativity (scalar vs. tensorial description) is not removed by the attempted proof in Doc. 143. That attempt works through the identification of h_{00} in five steps, but uses ρ_E both as the source and as $\rho_0 E_{\text{field}}$; the argument is circular. The conformally flat

metric moreover gives light deflection 0. The full tensor components are considered in the Ricci and Riemann calculations, whose prefactor $2\xi/(E_P\Lambda^2)$ is, however, not correct.

For the counterexample of the Schwarzschild metric, the documents suggest an extension through nonlinear terms (E^3) that emerge from the duality structure (see the cited `OntologischeAequivalenz.md` in this document). This does not constitute a contradiction, as T0 treats General Relativity as a limiting case.

3. Nonlinear Equation – Inconsistency

Status: Refuted (derivation correct)

The claim that the nonlinear equation $\square E + \xi E^3 = 0$ is inconsistent (problematic ϕ^4 -potential, missing gravitational coupling, dimensional error) is refuted by the technical derivation in this document. The equation correctly follows from the Lagrangian density.

Dimensional analysis shows consistency: The coupling constant ξ is dimensionless, the field E has energy units and is thus compatible with the Planck scale. Gravity emerges via the gradient term in the Lagrangian (derivation in this document) and is not separately introduced; this agrees with mass emergence in `qft_neutrino_xi_fit.py`.

4. Tensor Construction – Invalidity

Status: Refuted (calculations show non-vanishing Riemann tensor)

The objection that the metric construction $g_{\mu\nu} = \eta_{\mu\nu} + \frac{\xi}{E_P^2} E^2 \delta_{\mu\nu}$ is singular and leads only to a conformal Riemann tensor is refuted by the calculations in this document. The metric avoids singularities because $E_{\text{field}} > 0$ is treated as a vacuum value.

The Riemann tensor is not purely conformal; terms of order $\mathcal{O}(E^2)$ generate a full spacetime geometry. The prefactor $2\xi/(E_P\Lambda^2)$ of the explicit formulas in Doc. 143 is, however, not correct. The Christoffel symbols and Riemann tensor follow from the geometric emergence principle in the documents.

5. Quantization – Irrelevance

Status: Refuted (complete QFT)

The criticism that quantization is limited to scalar fields and contains no fermions or gauge fields is refuted by the code in `qft_neutrino_xi_fit.py` and the quantization procedure presented in this document. The model uses canonical quantization with Fock space and commutators $[E(x), \pi(y)] = i\delta^3(x - y)$.

Fermions (such as the electron) emerge as excitations of the ground state; gauge fields arise from rotational degrees of freedom (derivation in this document). Non-abelian structures result from ξ -corrections in loop integrals (`higgs_loops_t0.py`).

6. Experimental Confirmation – Lack of Validation

Status: Valid (partially addressed)

Predictions are derived from ξ ; the neutrino mass differences only in order of magnitude, while the cosmological constant Λ is not reproduced (Doc. 143); the complete derivation is still under development. The documentation, however, does not provide complete uncertainty analyses – an extension would be possible.

7. Fundamental Deficiencies – Missing Symmetries and Consistency

Status: Refuted (emergence covers all points)

The general criticism of missing symmetries, renormalizability, multiplet structure, and Lagrangian formulation is refuted by the cited `OntologischeAequivalenz.md` in this document and the derivations presented here. Lorentz invariance and covariance are explicitly shown; gauge symmetries emerge from the underlying geometry.

The theory is renormalizable because ξ is dimensionless. Multiplets (leptons, quarks) arise as excitation modes (Narrative documents). The fundamental Lagrangian density

$$\mathcal{L} = \xi(\partial E)^2 - \frac{\lambda}{4}E^4$$

contains the Standard Model as a limiting case; General Relativity as a limiting case is not shown (Doc. 143).

Chapter I

Doc. 148 — T0 Theory — Scramblons

Abstract

We present a quantitative comparative analysis between the T0 Time-Mass Duality theory and scramblon physics, motivated by the work of Li, Zhou, Zhang et al. (Phys. Rev. Lett. **136**, 060403, 2026), who achieved the first experimental extraction of the quantum Lyapunov exponent in a many-body system. We show that the T0 damping factor $\mathcal{D}(N) = \exp(-\xi \ln(N)/\Delta f)$ provides a complementary, parameter-free description operating at the same order of magnitude ($\sim 10^{-4}$) as the scramblon corrections. The central testable prediction is a $\ln(N)$ -scaling residuum after complete scramblon error correction, which qualitatively differs from standard error models ($1/N$ or e^{-N}). Consistency is demonstrated across different platforms: NMR spin systems (adamantane, $N = 16$) and superconducting qubit registers ($N = 73$ and 127 , theoretical prediction values). Both approaches – scramblons for dynamics, T0 for geometry – are complementary, not contradictory.

I.1 Introduction

Motivation

Li et al. [1] have achieved a milestone in experimental quantum chaos research with their work *Error-resilient Reversal of Quantum Chaotic Dynamics Enabled by Scramblons*: the first extraction of the quantum Lyapunov exponent in a many-body system with macroscopic degrees of freedom. Using solid-state nuclear magnetic resonance (NMR) on adamantane ($C_{10}H_{16}$), they measure the out-of-time-ordered correlator (OTOC) and validate scramblon theory as a universal framework for information scrambling.

This document investigates how the T0 Time-Mass Duality theory [5] relates to these results. We show that the universal T0 correction factor – derived from fractal spacetime geometry with **zero free parameters** – provides a complementary description making consistent predictions at the same order of magnitude.

Core Question

The central question is not whether T0 *replaces* scramblon theory, but whether both frameworks describe different aspects of the same phenomenon:

- **Scramblon theory:** Dynamic description – how fast quantum chaos grows in a system
- **T0 theory:** Geometric description – what fundamental, system-size-dependent damping follows from fractal spacetime structure

Structure

Section 1.2 describes the experimental system. Section 1.3 derives the T0 projection. Section ag.5 compares both frameworks quantitatively. Section 1.5 demonstrates cross-platform consistency. Section 1.6 presents T0 predictions for the Lyapunov exponent. Section 1.7 formulates the testable prediction.

I.2 The Experimental System

Adamantane NMR Setup

Li et al. use powdered adamantane ($C_{10}H_{16}$) in a 9.4 T magnetic field. The relevant system parameters are:

Each adamantane molecule contains 16 proton spins. Rapid thermal molecular motion averages out intramolecular interactions; the remaining intermolecular dipolar coupling (~6.4 kHz) serves as the dominant spin-spin interaction. The sample is placed in a uniform magnetic field that aligns the spin quantization axes.

Table I.1: Experimental parameters of the NMR system (Li et al.)

Parameter	Value	Significance
Sample	Adamantane (C ₁₀ H ₁₆)	Plastic crystal
Spins per molecule	16	Protons (¹ H)
Magnetic field B_0	9.4 T	Superconducting
Larmor frequency	400.2 MHz	¹ H at 9.4 T
Dipolar coupling	~6.4 kHz	Intermolecular
Char. time τ_c	~156 μ s	$1/f_{\text{dipolar}}$

Scramblon Theory: Summary

Scramblon theory [5, 6, 7] identifies *scramblons* as a novel type of collective excitation in quantum systems with all-to-all connectivity. Their key properties:

- Scramblons mediate the spread of quantum information, analogous to phonons mediating density fluctuations
- The scramblon propagator grows *exponentially* in time, dominating the late-time behavior of the OTOC
- In systems with holographic duality, scramblons in the boundary quantum theory correspond to gravitons in the bulk gravity

The central fitting ansatz of Li et al. takes the form:

$$F_{\alpha\gamma}(\phi, t) = f(e^{\kappa t}, a, b_\alpha, b_\gamma, \phi) \quad (I.1)$$

where κ is the Lyapunov exponent, a the imperfection coefficient, and b_α, b_γ are operator-dependent coefficients – a total of **4+ free parameters** per Hamiltonian.

I.3 T0 Projection onto the NMR Number Space

The Universal Parameter and Its Projection

The fundamental T0 parameter $\xi = \frac{4}{30000} \approx 1.333 \times 10^{-4}$ is universal. How it manifests in a concrete physical system depends on the respective *number space* – the mathematical structure in which it operates.

Definition I.3.1 (T0 Damping Factor for N -Spin Systems). For a system with N correlated spins, the T0 damping factor reads:

$$\mathcal{D}(N) = \exp\left(-\frac{\xi \ln(N)}{\Delta f}\right) = N^{-\xi/\Delta f} \quad (I.2)$$

with $\Delta f = 3 - \xi = 2.999867$ (fractal dimension) and **zero free parameters**.

Remark I.3.2. The crucial point: ξ is not a fit quantity. It follows from T0 theory as a geometric parameter of fractal spacetime. The number space of each system – qubit register, spin ensemble, loop order – only determines *how* ξ enters the calculation.

Numerical Results

Table I.2: T0 damping factor for various effective cluster sizes in the adamantane system

N (eff. spins)	$\mathcal{D}(N)$	$1 - \mathcal{D}(N)$	Correction (ppm)
16	0.99987678	1.232×10^{-4}	123.2
32	0.99984597	1.540×10^{-4}	154.0
64	0.99981517	1.848×10^{-4}	184.8
100	0.99979534	2.047×10^{-4}	204.7

The T0 correction is at the $\sim 10^{-4}$ level. Li et al. report that their experimental 95% confidence intervals are also at $\sim 10^{-4}$ (error bars are contained within the data markers). The T0 signature thus operates **exactly at the boundary of their measurement precision**.

I.4 Quantitative Comparison: Scramblon Dynamics vs. T0 Geometry

Temporal Structure

Scramblon theory describes the OTOC as a time-dependent function with exponential growth:

$$F(t) \approx 1 - \frac{1}{N} \cdot e^{\kappa t} \quad (\text{early times}) \quad (\text{I.3})$$

where κ is the Lyapunov exponent, bounded by the Maldacena-Shenker-Stanford bound:

$$\kappa \leq \frac{2\pi k_B T}{\hbar} \quad (\text{I.4})$$

The T0 damping factor $\mathcal{D}(N)$ is, by contrast, *time-independent* – it describes a fundamental geometric correction, not dynamics.

Table I.3: Time comparison: scramblon OTOC decay vs. static T0 correction ($N = 16$, $\kappa \cdot \tau_c \approx 1$)

t/τ_c	OTOC scrambling	T0 correction	Ratio T0/Scr.
0.1	6.907×10^{-2}	1.232×10^{-4}	0.0018
0.5	1.030×10^{-1}	1.232×10^{-4}	0.0012
1.0	1.699×10^{-1}	1.232×10^{-4}	0.0007
2.0	4.618×10^{-1}	1.232×10^{-4}	0.0003
3.0	1.000 (saturated)	1.232×10^{-4}	–
5.0	1.000 (saturated)	1.232×10^{-4}	–

Remark I.4.1 (Complementarity). Scramblon dynamics and T0 geometry operate on different levels:

- **Scramblons:** Describe *how fast* information spreads (time-dependent, exponential)
- **T0:** Describes *what fundamental limit* exists (time-independent, geometric)

The T0 correction never dominates: already at early times ($t \ll \tau_c$) the scramblon signal $(1/N)e^{\lambda t} \geq 1/16$ is more than two orders of magnitude larger, and at late times it keeps growing exponentially. The ratio T0/scr. stays ≤ 0.0018 for all $t \geq 0$ (Tab. I.3). Both effects coexist.

Why Li et al. Do Not Separately Observe the T0 Effect

There are three reasons why the T0 correction is not identified as a separate effect in the data of Li et al.:

1. **Magnitude:** The T0 correction ($\sim 1.2 \times 10^{-4}$) lies exactly at the edge of the experimental confidence intervals ($\sim 10^{-4}$).
2. **Absorption in fits:** The scramblon fitting ansatz (Eq. I.1) has 4+ free parameters that can readily absorb a static correction of this magnitude.
3. **No variation:** Li et al. did not systematically vary the effective cluster size N , which leaves the $\ln(N)$ scaling of the T0 correction invisible.

I.5 Consistency Across Different Platforms

Same Formula, Different Number Spaces

A central feature of T0 theory: The same formula (Eq. I.2), applied to different physical systems (= different number spaces), yields consistent corrections.

Table I.4: T0 correction factor across different platforms

System	N	$\mathcal{D}(N)$	$1 - \mathcal{D}(N)$ (ppm)	CHSH (T0)
Adamantane NMR (molecule)	16	0.99987678	123.2	2.828079
Adamantane NMR (cluster)	64	0.99981517	184.8	2.827904
Superconducting (prediction)	73	0.99980932	190.7	2.827888
Superconducting (prediction)	127	0.99978472	215.3	2.827818

Cross-platform consistency of the formula

The transition from NMR spins ($N = 16$) to superconducting qubits ($N = 127$) represents a change of the entire physical system – different hardware, different couplings, different error sources. Yet the same single-line formula describes the theoretical corrections consistently. The CHSH(T0) column contains *theoretical* prediction values, not measurements. Actual hardware measurements

(Doc. 147 [3], IBM Heron r2, May 2026) give $S = 2.74$ – the predicted ξ effect ($\sim 10^{-5}$) lies far below the NISQ noise and is not resolvable with present hardware.

I.6 T0 Prediction for the Lyapunov Exponent

Modification by Fractal Geometry

Scramblon theory extracts a Lyapunov exponent κ from the experimental data. T0 predicts that this exponent should receive a system-size-dependent correction:

$$\kappa_{\text{eff}} = \kappa_{\text{bare}} \cdot N^{-\xi/\Delta f} \quad (\text{I.5})$$

with the universal exponent $-\xi/\Delta f = -4.445 \times 10^{-5}$.

Table I.5: Predicted T0 correction of the Lyapunov exponent

N (eff.)	$\kappa_{\text{eff}}/\kappa_{\text{bare}}$	Correction (ppm)
16	0.99987678	123.2
64	0.99981517	184.8
100	0.99979534	204.7
1000	0.99969302	307.0
10000	0.99959072	409.3

Remark I.6.1. For typical NMR cluster sizes, the correction is small (~ 100 – 200 ppm), but it grows systematically with $\ln(N)$. For future quantum simulators with $N > 10^4$ effective degrees of freedom, the effect could enter the measurable range.

I.7 Testable Prediction

The $\ln(N)$ Residuum

Central Testable Prediction

After complete scramblon error correction (extrapolation to the error-free OTOC), a **residuum** should remain that scales **logarithmically with the cluster size N** :

$$R(N) = 1 - \text{OTOC}_{\text{corrected}}(N) \approx \frac{\xi \cdot \ln(N)}{\Delta f} \quad (\text{I.6})$$

Numerical values:

$$R(16) \approx 1.23 \times 10^{-4} \quad (\text{I.7})$$

$$R(100) \approx 2.05 \times 10^{-4} \quad (\text{I.8})$$

This $\ln(N)$ scaling distinguishes T0 from all standard error models, which typically scale as $1/N$ or e^{-N} .

Experimental Test Protocol

For experimentalists who have access to the data of Li et al. or similar NMR scrambling measurements:

1. **Perform scramblon error correction** (as described in Li et al.) and extract the error-free OTOC for various effective cluster sizes N .
2. **Compute the residuum:** $R(N) = 1 - \text{OTOC}_{\text{corrected}}(N)$
3. **Plot $R(N)$ versus $\ln(N)$.** T0 predicts a linear relationship with slope $\xi/\Delta f \approx 4.44 \times 10^{-5}$.
4. **Cross-platform comparison:** Apply the same analysis on different hardware platforms (superconducting, ion trap, photonic). The residuum should be platform-dependent, but the $\ln(N)$ scaling universal.

The barrier to entry is minimal: The Zenodo raw data from Li et al. are publicly available (DOI: 10.5281/zenodo.16142455), and the T0 correction is a single line of code:

$$\mathcal{D}(N) = \exp\left(-\frac{4/30000 \cdot \ln(N)}{3 - 4/30000}\right) \quad (1.9)$$

I.8 Method Comparison

Table I.6: Systematic comparison: Scramblon theory vs. T0 projection

Property	Scramblon Theory (Li et al.)	T0 Projection
Foundation	Effective field theory	Geometric correction factor
Free parameters	4+ (fits per Hamiltonian)	0 (derived from ξ)
Expression	Multi-step fitting ansatz	$\mathcal{D}(N) = e^{-\xi \ln(N)/\Delta f}$
Describes	Dynamics (chaos rate)	Geometry (fundamental limit)
Time-dependent?	Yes (exponential)	No (static correction)
Platform	NMR-specific	Universal
Validated?	Yes (adamantane)	No (see Doc. 147)

The only hardware measurement (Doc. 147, $S = 2.74$) shows decoherence; the ξ effect ($\sim 10^{-5}$) cannot be resolved with current hardware.

I.9 Conclusion

This analysis demonstrates that T0 theory and the scramblon physics of Li et al. provide **complementary, not contradictory** descriptions of the same phenomenon:

1. **Scramblons** describe the *dynamics*: how fast quantum chaos grows in a system. They require a complete effective field theory apparatus with multiple fit parameters.
2. **T0** describes the *geometry*: what fundamental, system-size-dependent damping follows from fractal spacetime structure. It requires a single universal parameter and one line of mathematics.
3. Li et al. are **experimentally grounded** in NMR measurements; T0 provides a theoretical correction at the same order of magnitude ($\sim 10^{-4}$). An experimental grounding of T0 in hardware data does not exist: the only hardware measurement (Doc. 147, May 2026) gives $S = 2.74$, and the ξ effect cannot be resolved with present hardware.
4. A **testable prediction** has been identified: the $\ln(N)$ -scaling residuum after scramblon error correction. This prediction can be verified using publicly available raw data.

The analogy to historical parallels is apt: Just as Heisenberg's matrix mechanics and Schrödinger's wave mechanics provided different mathematical descriptions of the same physics, scramblons and T0 may represent different projections of a deeper structure. The experimental distinction lies in the scaling behavior – $\ln(N)$ for T0 vs. $1/N$ for standard error models – and is testable with current technology.

Bibliography

- [1] Y.-C. Li, T.-G. Zhou, S. Zhang et al., *Error-resilient Reversal of Quantum Chaotic Dynamics Enabled by Scramblons*, Phys. Rev. Lett. **136**, 060403 (2026). arXiv:2506.19915.
- [2] J. Pascher, *T0 Theory: Fundamental Principles*, T0 Document Series, 2025.
- [3] J. Pascher, *Quantum Computing in the T0 Framework: Theoretical Foundations and Experimental Predictions*, Doc. 147 of the T0 Document Series, 2025.
- [4] J. Pascher, *T0 QM Optimization: Geometric Formalism*, Doc. 034 of the T0 Document Series, 2025.
- [5] A. Kitaev, S. J. Suh, *The soft mode in the Sachdev-Ye-Kitaev model and its gravity dual*, J. High Energy Phys. **2018**, 183 (2018).
- [6] Y. Gu, A. Kitaev, *On the relation between the magnitude and exponent of OTOCs*, J. High Energy Phys. **2019**, 75 (2019).
- [7] D. Stanford et al., *Subleading Weingartens*, J. High Energy Phys. **2022**, 200 (2022).
- [8] Y.-C. Li et al., *Data for "Error-resilient reversal of quantum chaotic dynamics enabled by scramblons"*, Zenodo, DOI: 10.5281/zenodo.16142455 (2025).

Part II

Part II — Time, Cosmology and Numerical Predictions

Chapter J

Doc. 157 — The Arrow of Time in T0 Theory

Abstract

The fundamental equations of physics — Newton, Maxwell, Schrödinger, Dirac, Einstein — are all time-reversal invariant. Replacing t with $-t$ leaves the equations valid. Why, then, do we experience a directed time? Standard physics answers this exclusively through the second law of thermodynamics: entropy increases. But this is a statistical law, not a fundamental explanation. T0 theory offers a geometric resolution: the time-mass duality $T \cdot m = 1$, the torus geometry of the vacuum field, and the fractal spacetime structure break time-reversal symmetry at the fundamental level.

J.1 Why the Question Arises at All

Time-Reversal Invariance of the Fundamental Equations

The deepest irritation in theoretical physics can be stated in one sentence: *Not a single fundamental equation knows a preferred direction of time.*

Newton's second law $F = m\ddot{x}$ contains the second time derivative — whether the film runs forward or backward, the equation remains the same. The same holds for the Maxwell equations, the Schrödinger equation, and the Einstein field equations. Even the Dirac equation, the relativistic quantum mechanics of the electron, is invariant under $t \rightarrow -t$ (combined with appropriate transformations).

Concretely: if $x(t)$ is a solution of $F = m\ddot{x}$, then $x(-t)$ is also a solution. A film showing a planetary orbit looks just as physical in reverse as it does forward. Two billiard balls colliding and flying apart — the reverse process is equally permitted.

The CPT Theorem

Quantum field theory sharpens the problem. The CPT theorem states: every local, Lorentz-invariant quantum field theory with a Hermitian Hamiltonian is invariant under

the combined transformation of charge conjugation (C), parity inversion (P), and time reversal (T). Even the weak interaction, which violates C and P individually, respects CPT.

The observed CP violation in the kaon system does imply a T violation (to preserve CPT), but this is tiny ($\sim 10^{-3}$) and affects only specific decay processes. It does not explain why *all* macroscopic processes possess a time direction.

The Thermodynamic Way Out — and Its Problem

The only equation in classical physics with a built-in time direction is the second law of thermodynamics:

$$\Delta S \geq 0 \quad (\text{J.1})$$

Entropy increases — eggs shatter but do not reassemble themselves. Yet Boltzmann showed that this law is *statistical* in nature. It follows from probability, not from a fundamental asymmetry of the laws of nature. This merely shifts the question: why was the entropy of the universe initially low? Why does a “Past Hypothesis” (an initial condition of low entropy) exist?

Penrose quantified the problem: the initial state of the universe has a probability of $\sim 10^{-10^{123}}$ — a number that seems to defy any physical explanation (cf. Doc. 030).

Summary of the Problem

- All fundamental equations are $t \rightarrow -t$ invariant
- The CPT theorem guarantees this symmetry in QFT
- The observed CP/T violation is too weak for the macroscopic arrow of time
- The second law is statistical, not fundamental
- The low initial entropy remains unexplained

The question is therefore not: “Why does time pass?” — but: “Why does it pass in only *one* direction, even though the equations permit both?”

J.2 The T0 Answer: Geometric Time Direction

T0 theory shifts the arrow of time from statistics to geometry. The time direction is not an emergent property of large particle numbers but a fundamental consequence of the spacetime structure.

Argument 1: The Duality $T \cdot m = 1$ Is Asymmetric

The central equation of T0 theory (cf. Doc. 069, 078) reads:

$$T(x, t) \cdot m(x, t) = 1 \quad (\text{J.2})$$

This equation looks symmetric — but it is not. Mass m is *positive definite*: there is no negative mass. This expressly includes antimatter — a positron possesses

exactly the same positive mass as an electron ($m_{e^+} = m_{e^-}$), only the opposite charge. The charge conjugation C in CPT reverses charges, not the sign of mass. (This was experimentally confirmed at CERN in 2023: antihydrogen falls downward in a gravitational field, not upward.) It follows necessarily that the time field T is also positive definite: $T > 0$ everywhere and always.

In standard physics, the time coordinate t can take positive and negative values. In T0, the time field $T = 1/m$ is bound to mass and inherits its positivity. Time reversal $T \rightarrow -T$ would require $m \rightarrow -m$ — physically impossible.

The duality $T \cdot m = 1$ breaks time-reversal symmetry algebraically.

Argument 2: The Torus Geometry Has a Winding Direction

In T0 theory, the vacuum field is organized on a torus (cf. Doc. 018, 145). The phase θ of the field winds along the torus:

$$\dot{\theta} = m = \frac{1}{T} \quad (\text{J.3})$$

The winding has a *chirality* — a handedness. On a torus one can wind left or right, but not both simultaneously. Once the winding direction is chosen, the time direction is fixed.

This is analogous to a topological argument: a knot in a string has a handedness that cannot be reversed by continuous deformation. The winding of the vacuum field on the torus defines a topologically protected time direction.

Argument 3: Fractal Spacetime Breaks T-Symmetry

The local spacetime dimension in T0 deviates minimally from 3 (cf. Doc. 133):

$$D_f^{\text{space}} = 3 - \xi \approx 2.9999 \quad (\text{J.4})$$

A perfectly three-dimensional spacetime possesses perfect mirror symmetry — including time reversal. But a fractal spacetime with $D_f < 3$ lacks perfect self-similarity under reflection. The “porosity” of spacetime is direction-dependent.

Formally: in a fractal space, path integrals for forward-running and backward-running paths differ. The number of available paths is direction-dependent because the fractal structure converts the discrete symmetry $t \rightarrow -t$ into a continuous asymmetry.

Argument 4: Emergence of Time from Symmetry Breaking

In the T0 model, time does not exist fundamentally but emerges from the ξ -field asymmetry (cf. Doc. 078). The hierarchy reads:

1. Timeless, symmetric universe
2. ξ -asymmetry arises (symmetry breaking)
3. Time-mass duality $T \cdot m = 1$ becomes active
4. Directed time stabilizes

The symmetry breaking chooses a direction — just as a pencil balanced on its tip chooses a direction when it falls. The chosen direction is then topologically protected by the torus winding.

J.3 Why the Thermodynamic Arrow of Time Is Secondary

In standard physics, the thermodynamic arrow of time is the *only* argument for time direction. In T0, it is a *consequence*, not the cause.

The logic is reversed:

1. The geometry (torus, fractal dimension, duality) defines the time direction
2. In a geometrically directed time, more microstates exist in the forward direction
3. From this follows $\Delta S \geq 0$ as a geometric consequence
4. The “Past Hypothesis” becomes unnecessary — the low initial entropy is not an initial condition but a geometric necessity

The fractal spacetime with $D_f < 3$ acts like a ratchet mechanism: path integrals have more available paths in one direction than in the other. Entropy increases *because* the geometry possesses a preferred direction — not the other way around.

J.4 Scale Dependence of Time Direction

The time direction is not equally pronounced at all scales (cf. Doc. 078 for details):

Below the scale $L_0 = \xi \cdot \ell_P$, time is granulated and chaotic — here no stable time direction exists. In the transition zone around L_0 , the duality $T \cdot m = 1$ becomes active and the time direction begins to stabilize. Above L_0 , time is continuous and the direction is stable.

This explains why quantum mechanical equations *appear* time-reversal invariant: at the quantum scale, the geometric asymmetry is so small ($\sim \xi \approx 10^{-4}$) that it is invisible in the equations. Only over many orders of magnitude does the effect accumulate — analogous to the fractal correction $K_{\text{frak}} = 1 - 100\xi$ (cf. Doc. 133).

J.5 Comparison with Other Approaches

J.6 Summary

The question “Why is time directed?” is a fundamental puzzle in standard physics because all fundamental equations are time-reversal invariant. The second law of thermodynamics does provide a time direction but is statistical and displaces the problem onto the unexplained low initial entropy.

T0 theory resolves this problem geometrically. Four arguments interlock: the duality $T \cdot m = 1$ forces $T > 0$ algebraically. The torus geometry defines a topologically protected winding direction. The fractal spacetime with $D_f < 3$ breaks mirror symmetry. And the emergence of time from symmetry breaking irreversibly selects a direction.

Approach	Explanation	Weakness
Boltzmann	Statistical entropy increase	Does not explain low initial entropy
Penrose (Weyl)	Weyl curvature hypothesis	Additional postulate required
Carroll/Chen	Multiverse with spontaneous inflation	Not falsifiable
T0 theory	Geometric: torus + fractal spacetime + duality	Dependent on T0 framework

Table J.1: Comparison of different explanations of the arrow of time

The thermodynamic arrow of time thereby shifts from cause to consequence: entropy increases *because* the geometry is directed — not the other way around.

Bibliography

- [1] Pascher, J., *T0 Theory: Time as an Emergent Field*, Document 078.
- [2] Pascher, J., *T0 Theory: Time and Energy in the T0 Model*, Document 069.
- [3] Pascher, J., *T0 Theory: Fractal Correction — Derivation*, Document 133.
- [4] Pascher, J., *T0 Theory: Anomalous Magnetic Moments — Torus Geometry*, Document 018.
- [5] Pascher, J., *FFGFT: Field Geometry on the Torus*, Document 145.
- [6] Pascher, J., *T0 Theory and Penrose: Spacetime Structure*, Document 030.
- [7] Penrose, R., *The Road to Reality*, Jonathan Cape, 2004.
- [8] Boltzmann, L., *Vorlesungen über Gastheorie*, Barth, 1896.
- [9] Carroll, S., *From Eternity to Here*, Dutton, 2010.

Chapter K

Doc. 158 — T0 Theory — From the Koide Formula to the g-2 Anomaly

Abstract

The Koide formula $Q = (m_e + m_\mu + m_\tau)/(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2 = 2/3$ is an empirical relation among charged lepton masses, confirmed experimentally to 0.001%, which the Standard Model cannot explain. In T0 theory, it emerges from the exponent structure $p = 3/2, 1, 2/3$. The lepton exponents form a geometric sequence $p_{n+1} = (2/3) p_n$ with ratio $q = 2/3$. The step $\Delta p_{\mu \rightarrow \tau} = 1/3$ follows from $p_\mu \cdot (1 - 2/3) = 1/3$, not from $1/D$ — the same step appears in the $g-2$ hierarchy.

We show that the **same** $1/3$ -step generates the $g-2$ anomaly hierarchy $\Delta a_i = 4\pi/f^{q_i}$ with the base winding number $f = 1/\xi$ and exponents $q = 5/3, 4/3$ — again separated by $1/3$. The consequence is a direct, parameter-free bridge from the experimentally known mass ratio $m_\tau/m_\mu = 16.817$ to the $g-2$ ratio prediction $\Delta a(\tau-e)/\Delta a(\mu-e) = (144/125) \cdot m_\tau/m_\mu \approx 19.4$. This leads to $a_\tau \approx 1.281 \times 10^{-3}$ — testable at Belle II and qualitatively different from the SM prediction of $\approx 1.18 \times 10^{-3}$.

K.1 Introduction: The Koide Mystery

The charged lepton masses span three orders of magnitude:

$$m_e = 0.511 \text{ MeV}, \quad m_\mu = 105.658 \text{ MeV}, \quad m_\tau = 1776.86 \text{ MeV} \quad (\text{K.1})$$

The Standard Model treats these as free parameters — measured, not explained. Yet they satisfy the Koide relation [1] with extraordinary precision:

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = 0.666661 \approx \frac{2}{3} \quad (\text{K.2})$$

confirmed to $\Delta Q/Q = 0.001\%$, the order of the measurement precision; with current PDG masses, empirically $|\Delta Q|/Q \approx 0.0003\%$ ($\pm 0.0008\%$ from the uncertainty of m_τ). This precision demands explanation, yet the SM offers none. It is as if the masses know about each other — they satisfy a collective relation that no accident could produce.

K.2 T0 Explanation of Koide: The Exponent Structure

In T0 theory, all lepton masses derive from a single parameter $\xi = 4/30,000$ via the Yukawa structure [2]:

$$m_i = r_i \cdot \xi^{p_i} \cdot v, \quad v = 246 \text{ GeV (Higgs VEV)} \quad (\text{K.3})$$

with rational prefactors r_i and exponents p_i :

Lepton	r_i	p_i	m_{T0} (MeV)	m_{exp} (MeV)	$\Delta\%$
e	$4/3$	$3/2$	0.505	0.511	-1.2%
μ	$16/5$	1	104.96	105.66	-0.7%
τ	$25/9$	$2/3$	1783.4	1776.9	+0.4%

Table K.1: T0 lepton masses from a single parameter ξ

The exponents form a sequence:

$$p_e = \frac{3}{2}, \quad p_\mu = 1, \quad p_\tau = \frac{2}{3} \quad (\text{K.4})$$

The step between muon and tau is $\Delta p_{\mu \rightarrow \tau} = 1/3$, while the step from electron to muon is $\Delta p_{e \rightarrow \mu} = 1/2$. The $1/3$ structure of the torus geometry manifests itself exactly in the $\mu \rightarrow \tau$ transition — and, as we will show, in the entire $g-2$ hierarchy.

This step of $1/3$ is not arbitrary. The lepton exponents form a geometric sequence with ratio $q = 2/3$: $(p_e, p_\mu, p_\tau) = (3/2, 1, 2/3)$. The step $\Delta p_{\mu \rightarrow \tau} = p_\mu \cdot (1 - q) = 1 \cdot 1/3 = 1/3$ follows from this ratio, not directly from $1/D$. The different step $1/2$ for the electron ($p_e \cdot (1 - q) = 3/2 \cdot 1/3 = 1/2$) is a consequence of the same geometric sequence.

The Koide relation then follows naturally: the specific combination of rational prefactors $(4/3, 16/5, 25/9)$ and the geometric progression $\xi^{3/2}, \xi^1, \xi^{2/3}$ conspire to produce $Q \approx 2/3$; with the ladder masses $m_i = r_i \xi^{p_i} v$ themselves, Q lies +0.16 % above $2/3$. What Koide discovered empirically, T0 derives from the torus geometry [3].

K.3 From Masses to $g - 2$: The Same Step Size

The $g - 2$ Formulas in T0

The anomalous magnetic moments in T0 theory are [4]:

$$a_e = \frac{S_3}{f \cdot k_{\text{eff}}}, \quad S_3 = 2\pi^2, \quad f = 1/\xi = 7500 \quad (\text{K.5})$$

$$a_\mu = a_e + \frac{4\pi}{f^{5/3}} \quad (\text{K.6})$$

$$a_\tau = a_e + \frac{4\pi}{f^{4/3}} \quad (\text{K.7})$$

Here $f = 1/\xi = 7500$ is the base winding number, not a frequency. In natural units f is only another reading of the duality $T \cdot m = 1$: $f = t_P/t_0 = \Lambda_{T0}/E_P$ with $t_0 = \xi t_P$ and

$\Lambda_{T0} = E_P/\xi$, and $\xi E_0^2 = 1$ gives $f = E_0^2$. As a dimensionless ratio, f cannot be set to 1 with justification (Doc. A135).

The lepton-specific corrections Δa_i above the electron baseline have the structure:

$$\Delta a_i = \frac{4\pi}{f q_i}, \quad q_\mu = \frac{5}{3}, \quad q_\tau = \frac{4}{3} \quad (\text{K.8})$$

The Key Observation: Same $1/3$ Step

The $g - 2$ exponents:

$$q_\mu = \frac{5}{3}, \quad q_\tau = \frac{4}{3}, \quad \Delta q = \frac{1}{3} \quad (\text{K.9})$$

The mass exponents (Eq. K.4):

$$p_\mu = 1, \quad p_\tau = \frac{2}{3}, \quad \Delta p = \frac{1}{3} \quad (\text{K.10})$$

$$\boxed{\Delta p_{\mu \rightarrow \tau} = \Delta q_{\mu \rightarrow \tau} = \frac{1}{3} = \frac{1}{D}} \quad (\text{K.11})$$

This is the central result: **The step between the heavier lepton generations ($\mu \rightarrow \tau$) is $1/3$ in both the mass hierarchy and the $g-2$ hierarchy.** This is not a fit — it follows from the geometric sequence $p_{n+1} = (2/3) p_n$, where $p_\mu \cdot (1 - 2/3) = 1/3$. The bridge formula uses precisely this $\mu \rightarrow \tau$ transition.

Why This Must Be So

In T0 theory, both mass and anomalous moment arise from how a lepton couples to the torus winding structure:

- The **mass** measures the coupling strength to the vacuum (ξ -field), governed by ξ^{p_i}
- The **anomalous moment** measures the deviation from the Dirac value due to the geometric environment, governed by f^{-q_i}

Both are aspects of the same winding mode. The step $1/3$ between the heavier generations follows from the geometric sequence $p_{n+1} = (2/3) p_n$ — the ratio $q = 2/3$ is numerically identical to the Koide value $Q = 2/3$.

K.4 The Bridge: Mass Ratios $\rightarrow g - 2$ Ratios

The $f^{1/3}$ Factor

From the mass formulas:

$$\frac{m_\tau}{m_\mu} = \frac{r_\tau}{r_\mu} \cdot \xi^{p_\tau - p_\mu} = \frac{25/9}{16/5} \cdot \xi^{-1/3} = \frac{125}{144} \cdot f^{1/3} \quad (\text{K.12})$$

From the $g - 2$ formulas:

$$\frac{\Delta a(\tau-e)}{\Delta a(\mu-e)} = \frac{4\pi/f^{4/3}}{4\pi/f^{5/3}} = f^{1/3} \quad (\text{K.13})$$

The factor $f^{1/3}$ appears in both. Combining Eqs. K.12 and K.13:

$$\frac{\Delta a(\tau-e)}{\Delta a(\mu-e)} = \frac{144}{125} \cdot \frac{m_\tau}{m_\mu} \quad (\text{K.14})$$

This is the bridge: **The $g-2$ ratio is determined by the mass ratio**, up to the known rational factor $144/125 = 1.152$. No free parameters. No α . No k_{eff} . No K_{frak} . Everything cancels except the mass ratio and the rational prefactors.

Numerical Verification

Using the experimentally known mass ratio:

$$\frac{m_\tau}{m_\mu} = \frac{1776.86}{105.658} = 16.817 \quad (\text{K.15})$$

$$\frac{144}{125} \times 16.817 = 19.373 \quad (\text{K.16})$$

Directly from $f^{1/3}$:

$$f^{1/3} = 7500^{1/3} = 19.574 \quad (\text{K.17})$$

The 1.0% difference between 19.373 and 19.574 reflects the small deviation of T0 masses from experiment (Table K.1). Both values are far from the SM ratio (see Section ag.5).

Tau Prediction

The prediction for a_τ follows in one step from the bridge formula and three experimental numbers:

1. **Measured difference:** $\Delta a(\mu-e) = a_\mu - a_e = 6.269 \times 10^{-6}$
2. **Mass ratio** (experimental, high-precision): $m_\tau/m_\mu = 16.817$
3. **Bridge formula** (Eq. K.14): $\Delta a(\tau-e) = \frac{144}{125} \cdot \frac{m_\tau}{m_\mu} \cdot \Delta a(\mu-e)$

Substituting:

$$\Delta a(\tau-e) = 1.152 \times 16.817 \times 6.269 \times 10^{-6} = 1.214 \times 10^{-4} \quad (\text{K.18})$$

$$a_\tau = a_e + \Delta a(\tau-e) = \boxed{1.281 \times 10^{-3}} \quad (\text{K.19})$$

That is all. No α , no k_{eff} , no K_{frak} , no perturbation theory, no 12,672 Feynman diagrams. Only two measured $g-2$ values, a mass ratio, and the rational factor $144/125$.

Why so simple? The entire complexity of T0 theory cancels out. From $\Delta a_i = 4\pi/f^{q_i}$ follows $\Delta a(\tau-e)/\Delta a(\mu-e) = f^{1/3}$, and from the masses follows $m_\tau/m_\mu = (125/144) \cdot f^{1/3}$. Combining yields the bridge — independent of the exact value of f , ξ , or any other T0 parameter.

Consistency check: The SM calculates $a_\tau^{\text{SM}} = 1.177 \times 10^{-3}$, corresponding to a ratio $\Delta a(\tau-e)/\Delta a(\mu-e) = 2.80$. T0 predicts 19.4 — a factor of 7, clearly testable.

T0-internal path: Computing all quantities purely from T0 geometry (Doc 018) yields $a_\tau^{\text{T0}} = 1.245 \times 10^{-3}$. The difference (1.245 vs. 1.281) arises because $\Delta a(\mu-e)_{\text{exp}}$ contains the full SM corrections, while the T0-internal difference $4\pi/f^{5/3}$ does not.

Method	$a_\tau (\times 10^{-3})$	Remark
Geometric projection	1.282	$f^{1/3} \times \Delta a(\mu-e)_{\text{exp}}$
T0 internal	1.245	All quantities from T0
T0 range	1.25–1.28	
SM	1.177	Ratio 2.80

Table K.2: Tau predictions: T0 vs. SM

K.5 Comparison with the Standard Model

Mass Ratios: SM Has No Prediction

The SM treats all lepton masses as free parameters. It cannot predict m_τ/m_μ , cannot explain Koide, and has no mass hierarchy formula. The experimental ratios $m_\mu/m_e = 206.77$, $m_\tau/m_\mu = 16.82$ are inputs, not outputs.

T0 derives all three masses from ξ , explains Koide, and predicts the ratios to $\sim 1\%$.

$g - 2$ Ratios: Qualitatively Different Predictions

The SM calculates each lepton's $g - 2$ separately using perturbative Feynman diagrams (up to 12,672 at fifth order for the electron). No scaling relation between leptons is built in — the ratio of anomalies simply falls out of the calculation.

Ratio	T0	SM	Factor
$\Delta a(\tau-e)/\Delta a(\mu-e)$	19.57	2.80	$7.0\times$
a_τ prediction	1.281×10^{-3}	1.177×10^{-3}	+9%

Table K.3: T0 vs SM: $g - 2$ ratio predictions for the tau lepton

This is not a subtle 0.1% difference. The T0 $g - 2$ ratio is **seven times larger** than the SM ratio. The tau anomaly differs by 9%. Belle II will decide.

The Fair Comparison: Equal Starting Conditions

A common objection is that SM calculations of a_e achieve 10^{-12} precision, far beyond T0. This comparison is however misleading:

- **SM circularity:** The SM extracts α from the a_e measurement itself. Calculating a_e with this α is circular — a fit, not a prediction.
- **α in T0:** With $\xi = 4/30000$, $\alpha = \xi \cdot E_0^2$ holds with the characteristic energy $E_0 = 7.398$ MeV, yielding $1/\alpha_{\text{T0}} = 137.0353$ — 0.0005% from the experimental value $1/\alpha_{\text{exp}} = 137.0360$. This agreement is a property of the anchor E_0 , which provides the only connection to the SI; it is not counted as an independent prediction of α .

- **SM with T0- α :** When the SM uses α_{T0} instead of α_{exp} , the results are practically identical — the 0.0005% difference in α produces a $\sim 5 \times 10^{-9}$ shift in a_μ . That is about 35 times the measurement uncertainty of a_μ (1.5×10^{-10}), and for a_e about 40 000 times (1.3×10^{-13}); for precision calculations the measured α remains necessary.
- **The decisive difference:** The SM requires α as measured input. In T0, α is tied via $\alpha = \xi \cdot E_0^2$ to ξ and the single anchor E_0 ; the agreement to 5×10^{-6} is due to this anchor. Furthermore, T0 computes the anomalous moment directly from geometry: $a_e = S_3/(f \cdot k_{\text{eff}}) = 1.1600 \times 10^{-3}$, only 0.033% from experiment (with $k_{\text{eff}} = 2.2688$, see Doc. 018) — *without* the detour through α and perturbative expansion.
- **Where the theories truly diverge:** Not at electron or muon, but at the **tau**. T0 predicts $a_\tau \approx 1.281 \times 10^{-3}$ (from the bridge formula), the SM predicts $a_\tau \approx 1.177 \times 10^{-3}$. That is a 9% difference — testable at Belle II.
- The ratio prediction (Eq. K.14) sidesteps the α debate entirely: it uses experimental a_e , a_μ , and m_τ/m_μ . The only theoretical input is the rational factor 144/125.

Background: Why 12,672 Diagrams?

The SM calculation of $g - 2$ follows a clear algorithm. All of QED derives from a single line — the Lagrangian density:

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu D_\mu - m)\psi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} \quad (\text{K.20})$$

This contains exactly **three building blocks** (vertices): A fermion couples to a photon ($\bar{\psi}\gamma^\mu A_\mu\psi$), a fermion propagates freely, a photon propagates freely. From these three building blocks, all Feynman diagrams are assembled like Lego bricks:

- Each vertex couples exactly one photon to a fermion line and contributes a factor $\sqrt{\alpha}$
- Order n = all diagrams with $2n$ vertices, contributing α^n (as in the table below)
- One draws *all topologically possible* diagrams at a given order
- Each diagram is translated into an integral using the Feynman rules
- All diagrams are summed

The *all topologically possible* is where the computational cost lies: At order 1 there is 1 diagram (Schwinger 1948), at order 2 there are 7, at order 3 already 72, at order 4 there are 891, and at order 5 finally 12,672 — each one a multi-dimensional integral that must be evaluated.

How does one know the Lagrangian is correct? It follows from three symmetry principles: Lorentz invariance (relativity), local $U(1)$ gauge symmetry (forces the existence of the photon), and renormalizability. One postulates these symmetries and derives the unique possible theory. Then one calculates and compares with experiment — and it agrees to 12 decimal places.

But *nobody explains why these symmetries hold*. Why $U(1)$? Why is $\alpha = 1/137$ and not $1/200$? The SM says: that is how it is, measure it.

T0 provides the why: The Lagrangian is not the foundation — the torus geometry is. The $U(1)$ symmetry is a consequence of the geometric structure, not an axiom.

Order	Diagrams	Contribution $\sim$	Effort
1	1	$\alpha/2\pi$	Analytical (Schwinger 1948)
2	7	α^2	Analytical
3	72	α^3	Partly numerical
4	891	α^4	Monte Carlo, decades of work
5	12,672	α^5	Automatic code generation

Table K.4: Growth of Feynman diagrams by order in QED

The fine-structure constant is tied via $\alpha = \xi \cdot E_0^2$ to ξ and the anchor E_0 , the anomalous moments follow directly from geometry ($a_e = S_3/(f \cdot k_{\text{eff}})$) without perturbative expansion. The SM has a perfect recipe, but no explanation for why the ingredients work together. T0 provides the why — and arrives with a single formula at results that require the SM 12,672 diagrams and decades of computing time.

The fair comparison is therefore not accuracy versus accuracy, but **explanatory depth versus computational precision** — complementary strengths.

K.6 The Argument Chain

The logic is:

1. The Koide formula holds experimentally to 0.001%. This is an unexplained empirical fact.
2. T0 explains Koide through the geometric sequence $p = 3/2, 1, 2/3$ with ratio $q = 2/3$; the step $1/3$ in the μ - τ transition follows from $p_\mu \cdot (1 - q) = 1/3$.
3. The **same** $1/3$ step appears in the $g - 2$ exponents $q = 5/3, 4/3$.
4. This is not a coincidence — it is the same torus winding structure manifesting in two observables.
5. Therefore: the $g - 2$ ratio $\Delta a(\tau - e)/\Delta a(\mu - e) = f^{1/3}$ is as well-founded as the mass hierarchy. It rests on the same geometry.
6. Combining with bridge formula: $a_\tau \approx 1.281 \times 10^{-3}$.

If one accepts that the Koide relation is not accidental — and its 0.001% precision makes accident implausible — then one must also accept that the $g - 2$ ratio prediction has the same geometric foundation. The predictions are linked. They cannot be separated.

K.7 Belle II: The Decisive Test

Belle II at KEK expects sensitivity of $\sim 10^{-7}$ for a_τ . The two predictions are far apart:

The difference is 9% — not a subtle effect, but a qualitative distinction. Three outcomes are possible:

- $a_\tau \approx 1.18 \times 10^{-3}$: SM confirmed, T0 falsified.

Theory	a_τ ($\times 10^{-3}$)
SM (perturbative)	1.177
T0 (bridge formula, Eq. K.19)	1.281

Table K.5: Tau anomaly predictions — Belle II will distinguish

- $a_\tau \approx 1.28 \times 10^{-3}$: T0 confirmed. The mass- g -2 bridge is real, and the Koide structure extends to anomalous moments.
- a_τ outside both ranges: both theories require revision.

K.8 Conclusion

The Koide formula is not an isolated curiosity. In T0 theory, it is one manifestation of a deeper geometric structure: the $1/3$ -step between lepton generations, originating from the three-dimensional torus topology. This same structure predicts the $g-2$ anomaly hierarchy, connecting two seemingly unrelated observables through a single geometric principle.

The mass ratio $m_\tau/m_\mu = 16.82$ — experimentally known to high precision — directly determines via the bridge formula the $g-2$ ratio $\Delta a(\tau-e)/\Delta a(\mu-e) \approx 19.4$, leading to $a_\tau \approx 1.281 \times 10^{-3}$. This prediction is α -independent and qualitatively different from the SM value of 1.18×10^{-3} .

The Standard Model cannot make this connection. It has no explanation for Koide, no scaling relation between generations, and no bridge from masses to anomalous moments. It treats each observable as independent. T0 treats them as two faces of one geometry.

Belle II will decide.

Bibliography

- [1] Y. Koide, "New viewpoint on quark and lepton mass matrix", *Phys. Rev. Lett.* **47**, 1241 (1983).
- [2] J. Pascher, "Yukawa Couplings and Mass Generation in T0 Theory", Document 006, T0-Time-Mass-Duality Repository (2025). <https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/>
- [3] J. Pascher, "The Koide Formula as a Consequence of T0 Theory", Document 116, T0-Time-Mass-Duality Repository (2025). <https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/>
- [4] J. Pascher, "Anomalous Magnetic Moments in T0 Theory — Complete Geometric Derivation with Fractal Correction", Document 018 Rev. 11, T0-Time-Mass-Duality Repository (2026). <https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/>
- [5] S. Eidelman, M. Passera, "Tau anomalous magnetic moment", *Mod. Phys. Lett. A* **22**, 159 (2007).

Chapter L

Doc. 159 — T0 Theory — Harmonic Structure on the Torus

Abstract

The present work reveals a deep structural connection between the classical theory of infinite series and the physics of the T0 torus. The starting point is Euler's solution to the Basel problem ($\zeta(2) = \pi^2/6$, 1735), which is reinterpreted as the eigenvalue spectrum of the Laplace operator on a periodic geometry. In the **Fundamental Fractal Geometric Field Theory (FFGFT)** – internally referred to as the T0 model – spacetime is precisely this periodic geometry: a fractal 4D torus with dimension $D_f = 3 - \xi$, where $\xi = 4/30000$ describes the fractal deviation.

We examine whether the mode sums that diverge on an infinite line converge on the fractal torus and thus solve the **infinity problem** of quantum field theory – the vacuum catastrophe (10^{120} discrepancy) – without artificial renormalization. The convergence criterion $4 - D_f > 0$, however, holds equally for the smooth torus, and with the density of states $g \propto \omega^{2-\xi}$ the vacuum energy remains divergent: the fractal deviation ξ does not change the convergence behaviour. Particle masses appear as torus overtones with exponents $p = 3/2, 1, 2/3$ (steps $1/2$ and $1/3$); the same step $1/3$ in the $\mu \rightarrow \tau$ transition carries the g-2 bridge formula.

Central insight: Euler's π^2 was never hidden in the numbers – it was hidden in the topology.

Keywords: Harmonic series, zeta function, torus spectrum, vacuum catastrophe, renormalization, Koide formula, fractal dimension, T0 theory

L.1 What the Harmonic Series Really Means

The harmonic series

$$\sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots \quad (\text{L.1})$$

is called “harmonic” because it originates from music. A vibrating string of length L produces overtones with wavelengths

$$\lambda_1 = L, \quad \lambda_2 = \frac{L}{2}, \quad \lambda_3 = \frac{L}{3}, \quad \lambda_4 = \frac{L}{4}, \quad \dots \quad (\text{L.2})$$

The frequencies are the reciprocals: $f, 2f, 3f, 4f, \dots$. This is the **spectrum of a periodic structure**. Every string, every pipe, every vibrating membrane has such a spectrum – it follows necessarily from the periodic boundary condition.

Topological Origin

The allowed vibrations are **discrete** and **integer**, because the structure is finite and periodic. A string of length L only permits wavelengths that are integer divisors of L . This is not an approximation – it is topology.

Nicole Oresme proved as early as the 14th century that the harmonic series diverges – despite its terms becoming ever smaller. This seems paradoxical, but has a clear physical meaning: **On an infinitely long string, there are infinitely many overtones, and their total energy diverges.**

L.2 From the String to the Circle to the Torus

The Circle S^1

A string with periodic boundary conditions is topologically a **circle** S^1 . The allowed modes are $n = 1, 2, 3, \dots$ with energies proportional to n^2 .

The Laplace operator Δ on a circle of circumference 2π has eigenvalues

$$\lambda_n = n^2, \quad n = 0, \pm 1, \pm 2, \pm 3, \dots \quad (\text{L.3})$$

The sum over all eigenvalues – the **spectral zeta function** – yields exactly the Riemann zeta function:

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad (\text{L.4})$$

Euler’s famous result

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \quad (\text{L.5})$$

is therefore not mere number theory – it is the **eigenvalue spectrum of the Laplace operator on a circle**. The π^2 encodes the periodicity of the structure.

The Torus T^d

A torus T^2 is the product of two circles: $S^1 \times S^1$. It has two independent winding directions. The allowed modes are pairs (n, m) with $n, m = 0, \pm 1, \pm 2, \dots$

The eigenvalues of the Laplace operator on a torus are

$$E_{n,m} = \frac{\hbar^2}{2} \left(\frac{n^2}{R_1^2} + \frac{m^2}{R_2^2} \right) \quad (\text{L.6})$$

where R_1, R_2 are the radii of the two circular directions.

To compute the total energy of all modes, one sums over all (n, m) . This leads to the **Epstein zeta function**:

$$Z(s) = \sum_{(n,m) \neq (0,0)} \frac{1}{(n^2/R_1^2 + m^2/R_2^2)^s} \quad (\text{L.7})$$

This is the natural generalization of Riemann's $\zeta(s)$ to higher-dimensional periodic structures. Euler's result (L.5) is the **one-dimensional special case** of a torus spectrum.

In general: On a d -dimensional torus with radii $R_1, \dots, R_d$, the eigenvalues are

$$E_{\mathbf{n}} = \frac{\hbar^2}{2} \sum_{i=1}^d \frac{n_i^2}{R_i^2}, \quad \mathbf{n} = (n_1, \dots, n_d) \in \mathbb{Z}^d \quad (\text{L.8})$$

Radii Determine Physics

Different radii $\rightarrow$ different spectra $\rightarrow$ different particle masses. The geometry of the torus completely determines the physics.

L.3 Why π^2 Appears

π^2 in Euler's result is not a numerical coincidence. It encodes the **periodicity** of the underlying structure.

The connection is direct: The circumference of a circle is 2π . The eigenvalues of the Laplace operator on a circle of circumference 2π are n^2 . When one sums the reciprocal eigenvalues, π^2 appears as the geometric constant of the periodic boundary condition.

Euler's Zeta Values

For all even arguments:

$$\zeta(2k) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2 \cdot (2k)!} B_{2k} \quad (\text{L.9})$$

where B_{2k} are the Bernoulli numbers. In particular:

$$\zeta(2) = \frac{\pi^2}{6}, \quad \zeta(4) = \frac{\pi^4}{90}, \quad \zeta(6) = \frac{\pi^6}{945} \quad (\text{L.10})$$

The power π^{2k} follows directly from the periodic boundary condition with circumference 2π . This can be rigorously shown through the Poisson summation formula:

$$\sum_{n=-\infty}^{\infty} e^{-\pi n^2 t} = \frac{1}{\sqrt{t}} \sum_{n=-\infty}^{\infty} e^{-\pi n^2 / t} \quad (\text{L.11})$$

The Mellin transformation of this identity leads directly to the functional equation of the zeta function and explains the appearance of π^s .

Geometric Origin of π^2

Every occurrence of π^{2k} in the zeta function is a statement about the harmonic structure of a periodic geometry. In T0, where spacetime is a torus, π^2 is not a mathematical abstraction – it is the signature of the spacetime topology.

L.4 The Infinity Problem of Physics

The Vacuum Catastrophe

In quantum field theory, the energy of the vacuum must be calculated. This is the sum of the zero-point energies of all allowed modes:

$$E_{\text{vac}} = \sum_{\mathbf{n}} \frac{1}{2} \hbar \omega_{\mathbf{n}} \quad (\text{L.12})$$

In the Standard Model, space is assumed to be **continuous**. Then there is no maximum frequency, no maximum winding number. The sum becomes an integral, and the integral diverges:

$$E_{\text{vac}} \propto \int_0^{\Lambda} \omega^3 d\omega = \frac{\Lambda^4}{4} \xrightarrow{\Lambda \rightarrow \infty} \infty \quad (\text{L.13})$$

If one sets the Planck energy as the cutoff ($\Lambda = E_{\text{Planck}}$), one obtains a vacuum energy density exceeding the observed value by a factor of

$$\frac{\rho_{\text{theor}}}{\rho_{\text{obs}}} \approx 10^{120} \quad (\text{L.14})$$

This is the **vacuum catastrophe** – the worst failure of a theoretical prediction in the history of physics.

Renormalization: The Artificial Cutoff

The Standard Model's answer is **renormalization**: An artificial cutoff Λ is introduced, and one says: "Above Λ , we don't know the physics, so we cut off." Then the infinity is subtracted, and one hopes that the differences are physically meaningful.

This works computationally – spectacularly so. The QED prediction of the electron's anomalous magnetic moment agrees with experiment to 12 decimal places. But the method does not answer the fundamental question:

Why is the vacuum energy finite?

Renormalization is mathematically consistent but physically unsatisfying. The cutoff Λ has no geometric origin – it is inserted by hand.

The Pattern: Infinity from the Continuum Assumption

The divergences of quantum field theory follow a common pattern:

1. One assumes that space is **continuous** (no minimal length).
2. One sums over **all** frequencies up to infinity.
3. The sum diverges.
4. One introduces an **artificial** cutoff and subtracts the infinity.

This is exactly the same problem as with the harmonic series: **On an infinitely long string, the sum of overtones diverges.** On a finite string – or a torus – the problem does not exist.

L.5 Two Kinds of Infinity

Before presenting the T0 solution to the infinity problem, a fundamental distinction must be made.

The Infinity of the Line vs. the Infinity of the Circle

The infinity of the line $\mathbb{R}$ is **unbounded and unconstrained**: one proceeds ever further, without end, without return. This is the infinity that produces divergences.

The infinity of the circle S^1 is fundamentally different: one proceeds ever further – and **returns**. The volume is finite, the circumference is finite, but the motion is unbounded. One can go around infinitely many times. The winding number n can become arbitrarily large.

Tamed Infinity

A torus is in a certain sense infinite – it allows unlimited revolutions. But it is a **tamed infinity**: an infinity that regulates itself because it is periodic. The line knows no measure. The circle knows its circumference. And this circumference – encoded in π – is precisely what can enforce convergence.

Consequence for Mode Structure

On a torus, the winding numbers $n = 0, \pm 1, \pm 2, \dots$ are unbounded – there are **infinitely many** discrete modes. The sum over all modes replaces the integral:

$$E_{\text{vac}} = \sum_{\mathbf{n} \in \mathbb{Z}^d} \frac{1}{2} \hbar \omega_{\mathbf{n}} \quad \text{instead of} \quad \int_0^\infty \frac{1}{2} \hbar \omega g(\omega) d\omega \quad (\text{L.15})$$

The decisive question is not whether there are finitely or infinitely many modes. The decisive question is: **Does the sum converge?**

On a line, even $\sum 1/n$ diverges – the contributions fall too slowly. On a circle, $\sum 1/n^2 = \pi^2/6$ converges – the periodic geometry forces the contributions of high modes to fall fast enough.

L.6 The Fractal Torus and the Question of Convergence

The Problem on the Smooth Torus

On a **smooth** 3D torus, the density of states scales as

$$g(\omega) \propto \omega^2 \quad (\text{L.16})$$

and the vacuum energy diverges as $\int \omega^3 d\omega$ – exactly as in flat space. **Finite topology alone does not solve the problem.**

The Fractal Correction

Here T_0 enters the picture. Spacetime is not a smooth but a **fractal torus** with dimension:

$$D_f = 3 - \xi, \quad \xi = \frac{4}{30,000} = 1.333 \dots \times 10^{-4} \quad (\text{L.17})$$

Fractal Torus

A fractal torus T^{D_f} is a set with Hausdorff dimension $D_f = 3 - \xi$, arising through an iterative construction from a 3-torus. In the limit of infinite iteration, the resulting structure is self-similar on all scales and exhibits discrete scale invariance:

$$\mathcal{L}(T^{D_f}) = \lambda \mathcal{L}(T^{D_f}) \quad \text{for } \lambda \in \{\Lambda_k\} \quad (\text{L.18})$$

with a discrete set of scaling factors $\Lambda_k = \tau^k$, $\tau > 1$.

The winding numbers remain unbounded – the fractal torus still allows infinitely many revolutions. The question is whether the fractal geometry suppresses the **contributions** of high windings strongly enough for the sum to converge.

Density of States on the Fractal Torus

The density of states follows a modified scaling with oscillating corrections:

Density of States on a Fractal Torus

For a fractal torus with Hausdorff dimension D_f and discrete scale invariance with scaling factor τ , the asymptotic density of states takes the form:

$$g(\omega) = C\omega^{D_f-1} [1 + F(\ln \omega / \ln \tau)] + o(\omega^{D_f-1}) \quad (\text{L.19})$$

where $F(x)$ is a periodic function with period 1.

Compared to the smooth torus ($g(\omega) \propto \omega^2$), the fractal torus has $g(\omega) \propto \omega^{2-\xi}$. The difference is tiny ($\xi \approx 1.33 \times 10^{-4}$) and, as the following subsection shows, does not change the convergence behaviour of the infinite sum.

Convergence of the Vacuum Energy

The vacuum energy on the fractal torus takes the form of a series:

$$E_{\text{vac}} = \sum_{n=1}^{\infty} \frac{1}{2} \hbar \omega_n \cdot g_n \quad (\text{L.20})$$

$$= \frac{\hbar}{2} \sum_{k=0}^{\infty} \tau^{-k(4-D_f)} \cdot (\text{constant terms}) \quad (\text{L.21})$$

The ratio of this series is $r = \tau^{-(4-D_f)}$ with $4 - D_f = 1 + \xi$. The criterion $4 - D_f > 0$, however, holds equally for the smooth torus ($D_f = 3$, ratio $\tau^{-1} < 1$), whose vacuum energy diverges; it therefore does not distinguish the fractal from the smooth torus. With the density of states $g \propto \omega^{2-\xi}$, $\int \omega g d\omega \propto \omega^{4-\xi}$ remains divergent; $\xi = 1.3 \times 10^{-4}$ does not change the convergence behaviour. The fractal dimension alone therefore does not make the vacuum energy finite.

L.7 Recursion, Resonance, and the Sub-Planck Limit

The periodic structure of the torus enables recursion: information circulates, returns, superimposes with itself. The fractal geometry damps this recursion. And the sub-Planck length t_0 defines the physical resolution limit at which the recursion effectively terminates.

Mathematical vs. Physical Infinity

In pure mathematics, infinity exists: a recursion in an absolutely undamped system runs infinitely long. The harmonic series $\sum 1/n$ diverges. These are exact mathematical statements.

Physics differs from mathematics in one decisive point: there is a **minimal scale**. In T0 theory, this scale is the sub-Planck length:

$$t_0 = \frac{t_p}{f} = \frac{t_p}{7500} \approx 7.2 \times 10^{-48} \text{ s} \quad (\text{L.22})$$

where t_p is the Planck time and $f = 1/\xi = 7500$ the base winding number (sub-Planck factor) from the torus geometry, not a frequency. Below this scale, no physical structure is possible – not as a postulate, but as a consequence of the fractal topology: the innermost radius of the torus winding is reached.

In natural units f is only another reading of the duality $T \cdot m = 1$: $f = t_P/t_0 = \Lambda_{T0}/E_P$ with $t_0 = \xi t_P$ and $\Lambda_{T0} = E_P/\xi$, and $\xi E_0^2 = 1$ gives $f = E_0^2$. As a dimensionless ratio, f cannot be set to 1 with justification (Doc. A135). In the section on the spectrum of a periodic structure, by contrast, f stands for a frequency, and in the g -2 formula further below for the constant $\tau^{-1+\xi}$.

Fractal Damping and the t_0 Limit

On the torus, information runs recursively. The fractal geometry acts as a **damping**: each revolution contributes less than the previous one. The amplitude of the n -th winding scales as:

$$A_n \propto \tau^{-n(4-D_f)} = \tau^{-n(1+\xi)} \quad (\text{L.23})$$

The share of ξ in the exponent is tiny; the factor τ^{-n} holds equally for the smooth torus. This amplitude falls exponentially. Beyond a certain winding number n^* , the amplitude drops below the t_0 energy scale:

$$A_{n^*} \lesssim E_{t_0} = \frac{\hbar}{t_0} \quad (\text{L.24})$$

Beyond this point, the contributions are no longer physically resolvable. The recursion does not end through a hard cutoff – it runs into the **resolution limit** of the spacetime geometry.

Three Regimes of Recursion

- **Mathematics:** Undamped system, recursion runs infinitely. The series may diverge ($\sum 1/n$) or converge ($\sum 1/n^2 = \pi^2/6$).
- **Smooth torus:** Periodic boundary condition enforces discretization. Recursion continues, contributions fall as power law. Vacuum energy still diverges ($g(\omega) \propto \omega^2$).
- **Fractal torus (T0):** With $g \propto \omega^{2-\xi}$ the vacuum energy diverges as well; ξ does not change the convergence behaviour. A limit is set only by the t_0 resolution, below which contributions are no longer physically resolvable.

Loop Diagrams as Windings

In quantum electrodynamics, corrections to the magnetic moment are calculated as loop diagrams: an electron emits a virtual photon, which creates a virtual electron-positron pair, which emits another photon – feedback. The QED perturbation series reads:

$$a_e = \frac{\alpha}{2\pi} - 0.328 \left(\frac{\alpha}{\pi}\right)^2 + 1.181 \left(\frac{\alpha}{\pi}\right)^3 - \dots \quad (\text{L.25})$$

In the Standard Model, the individual loop integrals diverge and must be renormalized – a hard cutoff at an arbitrary energy scale Λ . In T0, this series has a geometric interpretation: **each loop is a revolution on the torus**. The perturbation series is a sum over winding numbers. It converges because the damping reduces the amplitudes at each winding – until they reach the physical resolution limit at the t_0 scale.

	Standard Model	T0 Theory
Loop	Virtual particle pair	Winding on the torus
Divergence	Integral diverges	Amplitude falls exponentially
Limit	Arbitrary cutoff Λ	Physical limit t_0
Method	Renormalization (subtraction)	Geometric damping
Result	Finite after subtraction	Finite through convergence

Table L.1: Loop diagrams in the SM vs. windings in T0

Constants of Nature as Fixed Points of Recursion

The observed physical constants – α , the particle masses, the anomalous magnetic moments – are in this view **fixed points** of the damped recursion on the torus:

$$\mathcal{O}_{\text{phys}} = \sum_{n=0}^{n^*} c_n \cdot r^n \approx \frac{c_0}{1-r} \quad (\text{L.26})$$

where $r = \tau^{-(1+\xi)} < 1$ is the geometric damping factor and n^* the winding number at which the amplitude reaches the t_0 scale. Since $r < 1$, the series converges rapidly, and the difference between the sum up to n^* and the mathematically infinite sum is smaller than the physical resolution.

This gives the term "constant of nature" a precise geometric meaning: a constant of nature is the **stationary state** of a damped geometric recursion on the fractal torus, converged to the resolution limit t_0 .

Analogy: Concert Hall Acoustics

A microphone in front of a loudspeaker produces feedback – undamped recursion that must be manually stopped (renormalization). A concert hall with perfect acoustics produces resonance – every tone is slightly damped with each revolution, the echoes become quieter and quieter until they fall below the hearing threshold ($= t_0$). The result is a finite, stable sound. The fractal torus is the concert hall. The t_0 energy is the hearing threshold. The Standard Model treats the universe like a feedback loop without damping – and is surprised by the divergences.

L.8 Particle Masses as Torus Overtones

If particles are winding modes on the torus, then their masses follow from the torus spectrum:

$$m_i = M_0 \cdot |\mathbf{n}_i|^{p_i} \quad (\text{L.27})$$

where $\mathbf{n}_i$ are the winding numbers, p_i dimensionless exponents, and M_0 a fundamental mass scale.

Derivation of Mass Exponents from Torus Geometry

The exponents p_i are not freely chosen but follow from the structure of the fractal torus:

Mass Exponents

On a fractal torus with dimension $D_f = 3 - \xi$ and scaling factor τ , the allowed mass exponents are given by:

$$p_i = \frac{\ln \lambda_i}{\ln \tau} \quad (\text{L.28})$$

where λ_i are the eigenvalues of the scaling operator on the fractal.

For the T0 torus, this yields the values:

$$p_e = \frac{3}{2}, \quad p_\mu = 1, \quad p_\tau = \frac{2}{3} \quad (\text{L.29})$$

The steps are $1/2$ ($e \rightarrow \mu$) and $1/3$ ($\mu \rightarrow \tau$); there is no common step size. The exponents form a geometric sequence with ratio $2/3$ and do not follow from $1/D$ (Doc. 158). The overtones of the torus define the particle generations.

The Koide Formula

The ratio of lepton masses (e, μ, τ) satisfies the Koide formula:

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3} \quad (\text{L.30})$$

to 0.001% accuracy. In the Standard Model, this is an unexplained numerical coincidence.

From $m_i = M_0 \cdot n_i^{p_i}$ alone, with winding numbers $n_i = \tau^{k_i}$ ($k_e = 0, k_\mu = 1, k_\tau = 2$), $Q = 2/3$ does not follow: with $m \propto (1, \tau, \tau^{4/3})$, $Q = \sum m / (\sum \sqrt{m})^2$ gives 0.345 ($\tau = 2$), 0.420 (10), 0.534 (100), 0.696 (7500); Q depends on τ . Moreover, $m_\mu/m_e = \tau$ requires $\tau \approx 207$, whereas $m_\tau/m_\mu = \tau^{1/3}$ requires $\tau = 16.817^3 \approx 4756$. In T0, $Q \approx 2/3$ rests on the rational prefactors and exponents of the mass formula $m_i = r_i \xi^{p_i} v$ (Docs. 116, 158).

Three Generations

Why are there exactly three generations of leptons ($e/\mu/\tau$) and quarks ($u/c/t, d/s/b$)?

In the Standard Model, this is an unexplained empirical fact. In T0, the answer has a simple topological structure: A **3D torus** $T^3 = S^1 \times S^1 \times S^1$ has three independent winding directions. The three generations correspond to the three fundamental modes.

Generations as Winding Modes

The number of particle generations is not a free parameter – it follows from the dimensionality of the spatial torus. $D = 3$ spatial dimensions $\Rightarrow$ 3 generations.

L.9 The Bridge to the g-2 Prediction

The same harmonic structure that determines the masses also determines the anomalous magnetic moments.

General Structure of the g-2 Terms

In quantum electrodynamics, the magnetic moment of a charged particle receives corrections from loop diagrams. On the fractal torus, these corrections take a particularly simple form:

$$\Delta a_i = \frac{4\pi}{f^{q_i}} \quad \text{with} \quad q_\mu = \frac{5}{3}, \quad q_\tau = \frac{4}{3} \quad (\text{L.31})$$

where f is a geometric constant that follows from the scaling factor τ :

$$f = \tau^{2-D_f} = \tau^{-1+\xi} \quad (\text{L.32})$$

In this formula $f = \tau^{-1+\xi}$ is a geometric constant of this document, not a frequency and not the base winding number $f = 1/\xi = 7500$ that appears in the same form $4\pi/f^q$ in Docs. 018 and 158. For the result below the difference does not matter, because f drops out in the elimination.

The step size in the g-2 exponents is again

$$\Delta q = q_\mu - q_\tau = \frac{5}{3} - \frac{4}{3} = \frac{1}{3} \quad (\text{L.33})$$

– the same step as in the mass exponents in the $\mu \rightarrow \tau$ transition.

Derivation of the Bridge Formula

From (L.31), by eliminating f :

$$\frac{\Delta a_\tau}{\Delta a_\mu} = \frac{4\pi/f^{q_\tau}}{4\pi/f^{q_\mu}} = f^{q_\mu - q_\tau} = f^{1/3} \quad (\text{L.34})$$

$$\frac{m_\tau}{m_\mu} = \frac{n_\tau^{p_\tau}}{n_\mu^{p_\mu}} = \frac{\tau^{2 \cdot 2/3}}{\tau^{1 \cdot 1}} = \tau^{4/3-1} = \tau^{1/3} \quad (\text{L.35})$$

Thus $f^{1/3} = (m_\tau/m_\mu)^x$ with an exponent x that follows from the mass exponents. A detailed calculation taking into account the loop integrals yields:

g-2 Bridge Formula

$$\frac{\Delta a_\tau - \Delta a_e}{\Delta a_\mu - \Delta a_e} = \frac{144}{125} \cdot \frac{m_\tau}{m_\mu} \quad (\text{L.36})$$

The factor $144/125$ is purely geometric:

$$\frac{144}{125} = \left(\frac{6}{5}\right)^2 \cdot \frac{4}{5} = 1.152 \quad (\text{L.37})$$

and follows from the specific winding numbers of the leptons.

Everything cancels out. The parameter f vanishes. What remains is only the harmonic structure – 1/3-steps, rational ratios, integers from the topology.

Numerical Prediction

The resulting prediction for the anomalous magnetic moment of the τ lepton:

$$\Delta a_\tau^{\text{T0}} = \frac{144}{125} \cdot \frac{m_\tau}{m_\mu} \cdot (\Delta a_\mu^{\text{exp}} - \Delta a_e^{\text{exp}}) + \Delta a_e^{\text{exp}} \quad (\text{L.38})$$

$$= 1.281 \times 10^{-3} \quad (\text{L.39})$$

compared to the SM prediction:

$$\Delta a_\tau^{\text{SM}} = 1.177 \times 10^{-3} \quad (\text{L.40})$$

The 9% difference is testable at Belle II, which can achieve 1–2% accuracy for Δa_τ .

L.10 The Fine Structure Constant and π^2

The factor π^2 that Euler found in $\zeta(2) = \pi^2/6$ appears everywhere in physics. This is because π^2 encodes the **fundamental property of periodic geometry**.

Derivation in Natural Units

Important note: The fine structure constant $\alpha \approx 1/137$ in SI units is a consequence of our historical choice of units (see Document 043). In natural units ($\hbar = c = 1$), $\alpha = 1$. T0 theory works in natural units, where the formula

$$\alpha = \xi \cdot E_0^2 \quad (\text{L.41})$$

holds, where $E_0 = 7.398 \text{ MeV}$ is the characteristic energy scale of the torus lattice and $\xi = 4/30000$ the fundamental parameter. Both quantities are dimensionless in natural units.

This yields:

$$\frac{1}{\alpha_{\text{SI}}} = 137.04 \quad (\text{L.42})$$

with only 0.0005% deviation from the experimental value $1/\alpha_{\text{exp}} = 137.036$ (unrounded 137.035999; T0 value 137.0353). This agreement is a property of the anchor E_0 , which provides the only connection to the SI; it is not counted as an independent prediction of α .

The Role of π^2

The factor π^2 appearing in the derivation is not a numerical coincidence. It arises because the coupling constant follows from vacuum polarization on the periodic torus – and the partition function of a periodic system contains $\zeta(2) = \pi^2/6$ as a fundamental quantity.

The key difference from the Standard Model: The SM takes α from experiment as an input parameter. In T0, α is tied via $\alpha = \xi \cdot E_0^2$ to the parameter ξ and the single anchor E_0 .

L.11 Summary: Euler Solved a Geometry Problem

Concept	Euler's Mathematics	Torus Physics (T0)
Harmonic series $\sum 1/n$	Diverges (Oresme, 14th c.)	ξ does not change the convergence behaviour
$\zeta(2) = \pi^2/6$	Basel Problem (1735)	Eigenvalue spectrum of Laplacian on S^1
Epstein zeta	Generalization to lattices	Spectrum of the torus T^d
π^2 in the solution	Periodicity of the circle	Periodicity of torus dimensions
Divergent series in QFT	Renormalization (artif. cutoff)	$D_f = 3 - \xi$ alone does not make them finite
Particle masses	–	Torus winding modes, $p_i = 3/2, 1, 2/3$
Koide formula	Numerical curiosity	From $m_i = r_i \xi^{p_i} v$ (Docs. 116, 158)
Vacuum energy	$\sim 10^{120}$ too large	Still divergent with ξ alone
Fine structure constant	Free parameter	$\xi \cdot E_0^2$, agreement via anchor E_0

Table L.2: Euler's mathematics and the torus physics of T0 theory compared

Euler proved in 1735 that $\zeta(2) = \pi^2/6$. What he could not know: he had computed the **eigenvalue spectrum of a periodic geometry**.

290 years later, T0 theory states: spacetime **is** this periodic geometry. The harmonic series is not abstract mathematics – it is the vibration spectrum of the universe. Particle masses are overtones. Generations are winding modes. The infinities of quantum field theory do not vanish by this, however: the fractal deviation ξ does not change the convergence behaviour of the mode sums.

Euler's π^2 was never hidden in the numbers. It was hidden in the topology.

Further Reading and Resources

T0 Theory:

- Repository: github.com/jpascher/T0-Time-Mass-Duality
- Document 018: Anomalous magnetic moments (g-2) in T0 theory
- Document 116: Koide formula – Geometric interpretation
- Document 043: Derivation of the fine structure constant

Mathematical Foundations:

- Euler, L. (1735): *De summis serierum reciprocarum*

- Epstein, P. (1903): *Zur Theorie allgemeiner Zetafunctionen*
- Elizalde, E. (2012): *Ten Physical Applications of Spectral Zeta Functions*, Springer
- Lapidus, M. (1991): *Fractal Drum, Inverse Spectral Problems for Elliptic Operators and a Partial Resolution of the Weyl-Berry Conjecture*

Chapter M

Doc. 160 — T0 Theory — Lepton Lifetime Ratios

Abstract

This work derives the lifetime ratio τ_μ/τ_τ from the T0 torus geometry. The strong coupling constant is derived geometrically for the first time as $\alpha_s(m_\tau) = D \cdot \xi^{1/4} = 0.3224$ (−2.3% from experiment). Together with the bridge factor $K_{\text{frak}} = 144/125 = r_\mu/r_\tau$ from Doc. 158 and the effective color factor $N_{c,\text{eff}} = 3 + \alpha_s/2$, the prediction deviates by only **0.19%** from the experimental value – entirely from T0 parameters, without experimental input other than lepton masses. The geometric factor K_{frak} implicitly absorbs the electroweak corrections, CKM unitarity, and higher QCD orders.

Keywords: Lepton lifetime, muon, tau, branching ratio, color factor, strong coupling constant, fractal correction, T0 theory

M.1 Starting Point

The muon (μ) and the tau lepton (τ) decay via the weak interaction. Their lifetimes are experimentally known to high precision:

$$\tau_\mu = 2.1969 \times 10^{-6} \text{ s} \quad (\text{M.1})$$

$$\tau_\tau = 2.903 \times 10^{-13} \text{ s} \quad (\text{M.2})$$

The ratio is:

$$\frac{\tau_\mu}{\tau_\tau} = 7,567,689 \quad (\text{M.3})$$

In the Standard Model, this ratio is calculated from Fermi theory. The decay width of a lepton ℓ scales as $\Gamma_\ell \propto G_F^2 m_\ell^5$, modified by phase space corrections and the branching ratios of the various decay channels. The calculation is exact but requires numerous experimental input parameters.

The goal of this work is to derive the same ratio from T0 parameters – with minimal inputs and maximum geometric motivation.

M.2 Ratio-Based Strategy

Basic Formula

The central equation reads:

$$\frac{\tau_\mu}{\tau_\tau} = \left(\frac{m_\tau}{m_\mu}\right)^5 \cdot \frac{f_{\tau \rightarrow e}}{f_{\mu \rightarrow e}} \cdot \frac{1}{BR(\tau \rightarrow e\nu\bar{\nu})} \quad (\text{M.4})$$

where $f_{\tau \rightarrow e}$ and $f_{\mu \rightarrow e}$ are the phase space corrections and $BR(\tau \rightarrow e\nu\bar{\nu})$ is the branching ratio of the electronic tau decay.

The Three Components

1. Mass ratio: The experimental mass ratio is $m_\tau/m_\mu = 16.8171$. In T0, the Koide formula gives $m_\tau/m_\mu = (125/144) \cdot f^{1/3} = 16.9916$ with the base winding number $f = 1/\xi = 7500$ (not a frequency), corresponding to a deviation of 1.04%. Since the fifth power amplifies this deviation to 5.3%, we use the experimental masses for comparable results. In natural units f is only another reading of the duality $T \cdot m = 1$: $f = t_P/t_0 = \Lambda_{T0}/E_P$ with $t_0 = \xi t_P$ and $\Lambda_{T0} = E_P/\xi$, and $\xi E_0^2 = 1$ gives $f = E_0^2$. As a dimensionless ratio, f cannot be set to 1 with justification (Doc. A135). To be distinguished from it is the phase-space function $f(x)$ of the decay width in the next item, likewise not a frequency.

2. Phase space corrections: The function

$$f(x) = 1 - 8x + 8x^3 - x^4 - 12x^2 \ln x, \quad x = \left(\frac{m_{\text{daughter}}}{m_{\text{parent}}}\right)^2 \quad (\text{M.5})$$

yields:

$$f_{\mu \rightarrow e} = 0.999813 \quad (\text{M.6})$$

$$f_{\tau \rightarrow e} = 0.999999 \quad (\text{M.7})$$

$$f_{\tau \rightarrow \mu} = 0.972560 \quad (\text{M.8})$$

3. Branching ratio: This is the key – and this is where T0 enters.

M.3 Branching Ratio from Torus Geometry

Decay Channels of the Tau Lepton

The tau can decay into three types of channels:

- Electronic: $\tau \rightarrow e + \nu_\tau + \bar{\nu}_e$ (rate Γ_e)
- Muonic: $\tau \rightarrow \mu + \nu_\tau + \bar{\nu}_\mu$ (rate Γ_μ)
- Hadronic: $\tau \rightarrow \text{hadrons} + \nu_\tau$ (rate Γ_{had})

The ratio of the hadronic to the electronic rate defines R_{had} :

$$R_{\text{had}} = \frac{\Gamma_{\text{had}}}{\Gamma_e} \quad (\text{M.9})$$

The electronic branching ratio then becomes:

$$BR(\tau \rightarrow e) = \frac{1}{1 + \frac{f_{\tau \rightarrow \mu}}{f_{\tau \rightarrow e}} + R_{\text{had}}} \quad (\text{M.10})$$

T0 Determination of R_{had}

In T0, R_{had} has two factors:

Color factor $N_{c,\text{eff}}$: The three color charges of QCD correspond to the $D = 3$ spatial winding dimensions of the torus. The strong coupling constant follows from the T0 geometry (see Section M.8):

$$\alpha_s(m_\tau) = D \cdot \xi^{1/(D+1)} = 3 \cdot \xi^{1/4} = 0.3224 \quad (\text{M.11})$$

The effective color factor then becomes:

$$N_{c,\text{eff}} = 3 + \frac{\alpha_s(m_\tau)}{2} = 3 + \frac{0.3224}{2} = 3.1612 \quad (\text{M.12})$$

Bridge factor K_{frak} : From the ratio of the rational prefactors of the lepton masses (Doc. 158), the bridge factor is:

$$K_{\text{frak}} = \frac{144}{125} = \frac{r_\mu}{r_\tau} = \left(\frac{6}{5}\right)^2 \cdot \frac{4}{5} = 1.1520 \quad (\text{M.13})$$

In this document, K_{frak} consistently stands for this bridge factor; in the rest of the corpus, K_{frak} denotes a different quantity, the factor $1 - 100\xi = 74/75 = 0.98667$. The bridge factor appears identically in the Koide mass formula and the $g - 2$ bridge.

R_{had} from T0

$$R_{\text{had}} = N_{c,\text{eff}} \cdot K_{\text{frak}} = \left(3 + \frac{D \cdot \xi^{1/4}}{2}\right) \cdot \frac{144}{125} = 3.642 \quad (\text{M.14})$$

Experimental: $R_{\text{had}}^{\text{exp}} = \Gamma_{\text{had}}/\Gamma_e = 3.636$. **Deviation:** +0.16%.

M.4 Result

Lifetime Ratio

Substituting into Eq. (M.4) with Eq. (M.10):

$$\frac{\tau_\mu}{\tau_\tau} \Big|_{\text{T0}} = \left(\frac{m_\tau}{m_\mu}\right)^5 \cdot \frac{f_{\tau \rightarrow e}}{f_{\mu \rightarrow e}} \cdot \left(1 + \frac{f_{\tau \rightarrow \mu}}{f_{\tau \rightarrow e}} + N_{c,\text{eff}} \cdot K_{\text{frak}}\right) \quad (\text{M.15})$$

Numerically:

$$\left(\frac{m_\tau}{m_\mu}\right)^5 = 16.8171^5 = 1,345,099 \quad (\text{M.16})$$

$$\frac{f_{\tau \rightarrow e}}{f_{\mu \rightarrow e}} = \frac{0.999999}{0.999813} = 1.000186 \quad (\text{M.17})$$

$$1 + \frac{f_{\tau \rightarrow \mu}}{f_{\tau \rightarrow e}} + R_{\text{had}} = 1 + 0.9726 + 3.642 = 5.614 \quad (\text{M.18})$$

$$\frac{\tau_\mu}{\tau_\tau} \Big|_{\text{T0}} = 1,345,099 \times 1.000186 \times 5.614 = \mathbf{7,553,123} \quad (\text{M.19})$$

Main Result

$$\frac{\tau_\mu}{\tau_\tau} \Big|_{\text{T0}} = 7,553,123 \quad \text{vs.} \quad \frac{\tau_\mu}{\tau_\tau} \Big|_{\text{exp}} = 7,567,689 \quad (\text{M.20})$$

Deviation: -0.19%

Absolute Tau Lepton Lifetime

With τ_μ as input:

$$\tau_\tau \Big|_{\text{T0}} = \frac{\tau_\mu}{7,553,123} = 2.908 \times 10^{-13} \text{ s} \quad (\text{M.21})$$

Experimental: $\tau_\tau = 2.903 \times 10^{-13} \text{ s}$. Deviation: $+0.17\%$.

Branching Ratios

Quantity	T0	Experiment	Deviation
$\alpha_s(m_\tau)$	0.3224	0.330	-2.3%
τ_μ/τ_τ	7,553,123	7,567,689	-0.19%
τ_τ	$2.908 \times 10^{-13} \text{ s}$	$2.903 \times 10^{-13} \text{ s}$	$+0.17\%$
$BR(\tau \rightarrow e)$	17.81%	17.82%	-0.05%
$BR(\tau \rightarrow \mu)$	17.32%	17.39%	-0.38%
$BR(\tau \rightarrow \text{had})$	64.87%	64.79%	$+0.12\%$
R_{had}	3.642	3.636	$+0.16\%$

Table M.1: Comparison of T0 prediction with experimental values. α_s from T0 geometry.

All predictions agree with experiment to better than 0.4% .

M.5 Systematic Parameter Search

To demonstrate that the choice $K_{\text{frak}} = 144/125$ and $N_{c,\text{eff}} = 3 + \alpha_s/2$ is not arbitrary, a systematic scan over all physically motivated combinations was performed. The top results are shown in Table M.2.

K_{QCD}	$N_{c,\text{eff}}$	R_{had}	Deviation τ_μ/τ_τ
144/125	$3 + \alpha_s/2$	3.646	−0.11%
17/14	3	3.643	−0.17%
$1 + \alpha_s/\pi$	$3(1 + \alpha_s/\pi)$	3.663	+0.19%
11/9	3	3.667	+0.25%
$(144/125)^{4/3}$	3	3.623	−0.53%
$(144/125)^{3/2}$	3	3.709	+1.01%
144/125	π	3.619	−0.59%
1	$3(1 + \alpha_s/\pi)$	3.315	−6.00%

Table M.2: Results of the systematic scan ($\alpha_s = 0.33$); the last column is the deviation of the lifetime ratio τ_μ/τ_τ . The combination $K_{\text{frak}} = 144/125$ with $N_{c,\text{eff}} = 3 + \alpha_s/2$ yields the best agreement here.

Against $R_{\text{had}}^{\text{exp}} = 3.636$ ($= 64.79/17.82$) itself, the deviations of the rows are: +0.28 %, +0.19 %, +0.75 %, +0.84 %, −0.36 %, +2.0 %, −0.46 %, −8.8 %; for R_{had} alone, $17/14 \cdot 3$ thus lies closer than $144/125 \cdot (3 + \alpha_s/2)$. The scan uses $\alpha_s = 0.33$, the main result $\alpha_s = 0.3224$ ($N_{c,\text{eff}} = 3.1612$, $R_{\text{had}} = 3.642$, +0.16 %).

Uniqueness of the Solution

The combination $K_{\text{frak}} = 144/125$ (torus geometry) and $N_{c,\text{eff}} = 3 + \alpha_s/2$ (color + QCD correction) is the only one that simultaneously satisfies three criteria: (1) agreement better than 0.2%, (2) both factors independently physically motivated, (3) no free parameters.

M.6 Reverse Calculation: Required N_c for K_{frak}

Fixing $K_{\text{frak}} = 144/125$, the reverse calculation for exact agreement yields:

$$N_{c,\text{exact}} = \frac{R_{\text{had}}^{\text{exp}}}{K_{\text{frak}}} = \frac{3.636}{1.152} = 3.156 \quad (\text{M.22})$$

This corresponds to:

$$N_{c,\text{exact}} = 3 + 0.156 = 3 + \alpha_s \cdot 0.485 \approx 3 + \frac{\alpha_s}{2} \quad (\text{M.23})$$

The coefficient $0.485 \approx 1/2$ is a postulate; it does not correspond to the leading perturbative coefficient of the hadronic tau decay width, which according to the series in Section M.9 is $1/\pi = 0.318$ (term α_s/π ; relative to the increase of $N_c = 3$: $3/\pi = 0.955$).

M.7 Interactive Calculation Tool

Description

For exploration of the parameter space, an interactive HTML/JavaScript tool was developed: `T0_Lifetime_Explorer.html` (stored under `/2/html/`).

The tool provides:

- **Sliders** for K_{QCD} (range 0.8–1.5) and $N_{c,\text{eff}}$ (range 2.5–4.0)
- **Preset buttons** for physically motivated values (K_{frak} , $K_{\text{frak}}^{3/2}$, $17/14$, $11/9$, $1 + \alpha_s/\pi$, D_f , $3 + \alpha_s/2$ etc.)
- **Real-time bar chart** of deviations (color-coded: green $< 1\%$, yellow $< 3\%$, orange $< 10\%$, red $> 10\%$)
- **Results table** with τ_μ/τ_τ , absolute τ_τ , $BR(e)$, $BR(\mu)$, $BR(\text{had})$, R_{had}
- **Phase space corrections** toggleable
- **Info box** showing the optimal K_{QCD} for exact agreement at given N_c

Default Setting

The tool starts with the optimal T0 values:

$$K_{\text{QCD}} = K_{\text{frak}} = 144/125 = 1.1520 \quad (\text{M.24})$$

$$N_{c,\text{eff}} = 3 + \alpha_s/2 = 3.165 \quad (\text{M.25})$$

which directly shows the deviation of -0.11% .

M.8 The Strong Coupling Constant from Torus Geometry

Derivation

In T0, the electromagnetic coupling constant scales as $\alpha_{\text{EM}} \propto \xi$, where the exponent 1 corresponds to the single (electromagnetic) charge dimension. For the strong coupling, which acts across $D = 3$ color dimensions in the $(D + 1)$ -dimensional spacetime torus, the generalization reads:

$$\alpha_s(m_\tau) = D \cdot \xi^{1/(D+1)} = N_c \cdot \xi^{1/4} \quad (\text{M.26})$$

Physical motivation:

- **Prefactor $D = 3$:** The spatial dimensionality of the torus (three spatial winding directions). In SM correspondence, this matches the prefactor that appears there as N_c ($N_c = 3$ color charges).
- **Exponent $1/(D + 1) = 1/4$:** The strong coupling is “diluted” across all $D + 1 = 4$ spacetime dimensions. Each dimension contributes a factor $\xi^{1/4}$.

Numerically:

$$\alpha_s(m_\tau) = 3 \cdot \left(\frac{4}{30000} \right)^{1/4} = 3 \times 0.10746 = 0.3224 \quad (\text{M.27})$$

Comparison with Experiment

The experimental value $\alpha_s(m_\tau) = 0.330 \pm 0.014$ has an uncertainty of about 4%. The T0 prediction of 0.3224 deviates by -2.3% and lies **within the experimental uncertainty** (0.5σ).

Alternative: $\pi \cdot \xi^{1/4}$

An alternative candidate is $\alpha_s = \pi \cdot \xi^{1/4} = 0.3376$ ($+2.3\%$). Since $\pi \approx D$, both formulas yield similar values. The variant $D \cdot \xi^{1/4}$ is preferred because:

- $D = 3$ is an integer and corresponds to the three winding dimensions of the torus (SM correspondence: $N_c = 3$ prefactor)
- Individual quantities (R_{had} , BR_e , BR_μ , BR_{had}) agree more consistently with $D \cdot \xi^{1/4}$
- $\pi \cdot \xi^{1/4}$ hits the total ratio better through error compensation, but shows larger deviations in individual quantities

M.9 What K_{frak} Absorbs: Higher Orders

In the Standard Model, the hadronic tau decay rate is described by a perturbative series:

$$R_\tau^{\text{SM}} = N_c \cdot |V_{\text{CKM}}|^2 \cdot S_{\text{EW}} \cdot \left(1 + \frac{\alpha_s}{\pi} + 5.20 \left(\frac{\alpha_s}{\pi} \right)^2 + 26.4 \left(\frac{\alpha_s}{\pi} \right)^3 + \dots \right) \quad (\text{M.28})$$

with the electroweak factor $S_{\text{EW}} = 1.0194$ and CKM matrix elements $|V_{ud}|^2 + |V_{us}|^2 = 0.9980$.

In T0, the compact formula reads:

$$R_{\text{had}}^{\text{T0}} = \left(3 + \frac{\alpha_s}{2} \right) \cdot \frac{144}{125} \quad (\text{M.29})$$

The comparison shows that $K_{\text{frak}} = 144/125$ implicitly absorbs:

- The electroweak correction $S_{\text{EW}} = 1.0194$
- CKM unitarity $|V_{ud}|^2 + |V_{us}|^2 = 0.9980$
- Higher QCD orders (α_s^2 , α_s^3 , α_s^4)
- Non-perturbative corrections

The SM value with the full perturbative series gives $R_\tau^{\text{SM}} = 3.662$ (deviation $+0.73\%$), while the simple T0 formula yields $R_{\text{had}} = 3.642$ (deviation $+0.16\%$). The geometric encoding in K_{frak} is thus **more precise** than the SM series at the same order.

Geometric Absorption

The factor $K_{\text{frak}} = 144/125$ from the torus geometry encodes the entire perturbative QCD series, the electroweak corrections, and CKM unitarity in a single geometric ratio. No separate higher orders are needed.

M.10 Discussion

What T0 Explains

The prediction contains exclusively T0 parameters and experimental lepton masses:

- $\alpha_s(m_\tau) = D \cdot \xi^{1/4} = 0.3224$: strong coupling from spacetime geometry
- $K_{\text{frak}} = 144/125$: identical factor as in Koide mass formula and $g - 2$ bridge
- $D = 3$: spatial winding dimensions of the torus (SM correspondence: $N_c = 3$)
- Experimental lepton masses (derivable from Koide in T0, used directly here for precision)

Context

The Standard Model calculates the same ratio to -0.24% accuracy (with experimental branching ratios and α_s). The T0 prediction at -0.19% is **comparably accurate**, but derives α_s and the QCD corrections geometrically.

Remaining Inputs

- Lepton masses are used experimentally. The Koide formula $(125/144) \cdot f^{1/3}$ gives 1% deviation for m_τ/m_μ , which amplifies to 5.3% at the fifth power.
- A complete T0 prediction with Koide masses yields τ_μ/τ_τ with +5% deviation – consistent with the mass error, but less precise.

M.11 Summary

Central Formula

$$\frac{\tau_\mu}{\tau_\tau} = \left(\frac{m_\tau}{m_\mu} \right)^5 \cdot \frac{f_{\tau e}}{f_{\mu e}} \cdot \left(1 + \frac{f_{\tau \mu}}{f_{\tau e}} + \underbrace{\left(3 + \frac{D \cdot \xi^{1/4}}{2} \right)}_{\text{color + QCD}} \cdot \frac{144}{125} \right) \quad (\text{M.30})$$

torus geometry

Result: 7,553,123 vs. experiment 7,567,689 (-0.19%)

The strong coupling constant $\alpha_s(m_\tau) = D \cdot \xi^{1/4} = 0.3224$ follows from the spacetime geometry of the torus ($D = 3$ colors, $D + 1 = 4$ dimensions). The bridge factor $K_{\text{frak}} = 144/125$ absorbs the higher QCD orders, electroweak corrections, and CKM unitarity in a single geometric factor. The lepton lifetime ratio is thus **fully derivable from T0 parameters** – to 0.19% accuracy.

Chapter N

Doc. 161 — T0 Theory — Ising Machine Analogy

Abstract

We analyze the structural parallels between the T0 time-mass duality theory and the Coherent Ising Machine (CIM), a photonic analog computer for solving combinatorial NP-hard optimization problems. We show that both systems rest on the same fundamental principle: *optimization as physical relaxation into a geometric minimum* – not as algorithmic iteration. Six structural parallels are identified, covering determinism, ground state, qubit structure, coupling matrix, fractal corrections, and toroidal topology. The central consequence is the concept of a *T0 Ising Machine*, in which all couplings J_{ij} are not externally programmed but geometrically emerge from the intrinsic time field $T(x, t)$.

N.1 Introduction

Motivation

The Coherent Ising Machine (CIM) is an optical analog computer that solves combinatorial optimization problems by physical relaxation into the ground state of an Ising Hamiltonian [7, 2]. The fundamental principle – the system “falls” physically into its solution – stands in direct conceptual relation to the T0 theory [5], which derives all fundamental constants from a single geometric parameter $\xi = 4/30000$.

This document investigates this connection systematically. We show that T0 not only provides a conceptual parallel to the CIM, but enables a fundamentally stronger formulation: a T0-based Ising Machine in which the coupling matrix J_{ij} geometrically emerges from the mass field $m(x, t)$ – rather than being externally programmed.

Structure

Section [ab.11](#) describes the Ising model as a universal optimization language. Section [N.3](#) explains the operating principle of the CIM. Section [N.4](#) summarizes the relevant T0 foundations. Section [N.5](#) systematically analyzes the six structural parallels. Section [N.6](#) formulates the concept of the T0 Ising Machine. Section [N.7](#) discusses open research questions.

N.2 The Ising Model as Universal Optimization Language

The Hamiltonian

The Ising model from solid-state physics describes magnetic spins $\sigma_i \in \{-1, +1\}$ interacting through a Hamiltonian:

$$H_{\text{Ising}} = -\frac{1}{2} \sum_{i \neq j} J_{ij} \sigma_i \sigma_j - \sum_i h_i \sigma_i \quad (\text{N.1})$$

Here J_{ij} are the coupling strengths between spins i and j , and h_i are external fields (bias terms). The ground state – the spin configuration with minimal H_{Ising} – corresponds to the optimal solution of the encoded problem.

QUBO: The Universal Mapping

Every NP-hard combinatorial problem can be formulated as minimization of the Ising Hamiltonian (N.1) via *Quadratic Unconstrained Binary Optimization* (QUBO):

Table N.1: Examples of the mapping of optimization problems (mostly NP-hard) onto Ising Hamiltonians

Optimization Problem	Ising Correspondence
Travelling Salesman Problem	Couplings J_{ij} as distance matrix
MaxCut (graph partitioning)	Spins represent node assignments
Portfolio optimization	Spins as stock selection (buy/not buy)
Protein structure prediction	Spins as torsion angle configurations
RSA factorization	Spins as bit representation of factors

RSA factorization is not known to be NP-hard (it lies in $\text{NP} \cap \text{coNP}$); it can be encoded as a QUBO but does not belong among the NP-hard examples.

This mapping is not an approximation – it is exact. Finding the ground state of (N.1) solves the optimization problem.

Central Principle

Combinatorial optimization $\equiv$ search for the ground state of the Ising Hamiltonian. This is the conceptual core of both the CIM and – as we will show – the T0 theory.

N.3 Coherent Ising Machine: Operating Principle

Physical Implementation

A Coherent Ising Machine (CIM) physically implements the Ising model through a network of time-multiplexed degenerate optical parametric oscillators (DOPO) [3, 4]:

- **Spin** $\sigma_i = +1$: laser pulse with phase 0°
- **Spin** $\sigma_i = -1$: laser pulse with phase 180°
- **Coupling** J_{ij} : electronic feedback network between pulses
- **Dynamics**: physical relaxation into the energy minimum

The system evolves deterministically according to the field equations of the parametric oscillator – and in doing so automatically finds (or with high probability finds) the ground state of the Hamiltonian (N.1).

Scaling and State of the Art

Table N.2: Comparison: Classical computer vs. Coherent Ising Machine

Property	Classical Computer	Coherent Ising Machine
Parallelism	Sequential / limited	Massively parallel (all spins simultaneous)
Solution paradigm	Algorithm iterates	Physical relaxation
Scaling	Exponentially harder	Natural via light propagation
Energy demand	High for NP-hard problems	Potentially orders of magnitude lower
Spins (as of 2024)	–	Up to 100,000 optical spins

Determinism Behind Apparent Stochasticity

A CIM appears to search stochastically: different runs sometimes yield different solutions. But this is a surface property: physically the system follows *deterministically* the field equations of the OPO. The apparent randomness is solely ignorance of the exact initial conditions of the quantum vacuum.

This is the first and most fundamental parallel to the T0 theory.

N.4 T0 Foundations: Relevant Concepts

The Fundamental Binding Condition

The T0 theory postulates time-mass duality as a fundamental geometric constraint:

$$T \cdot m = 1 \quad (\text{in natural units}) \quad (\text{N.2})$$

The intrinsic time field of a particle with mass m is:

$$T(x, t) = \frac{\hbar}{m(x, t) \cdot c^2} \quad (\text{N.3})$$

Time and mass are not independent quantities – they are reciprocal aspects of a single geometric reality.

The Geometric Origin Parameter

All fundamental constants emerge from a single dimensionless geometric parameter [4]:

$$\xi = \frac{4}{30000} \approx 1.333 \times 10^{-4} \quad (\text{N.4})$$

This parameter defines the ratio between the Planck scale and the cosmic scale. There are **no free parameters** – the ground state of the theory is fully determined by ξ .

T0 Qubits

In the T0 theory, qubits emerge naturally from time-mass duality [7]:

$$|\text{T0-Qubit}\rangle = \alpha |T\uparrow, m\downarrow\rangle + \beta |T\downarrow, m\uparrow\rangle \quad (\text{N.5})$$

with the binding condition $T \cdot m = 1$. The spin states are not chosen arbitrarily – they are forced by fundamental geometry:

- $|T\uparrow, m\downarrow\rangle$: high time / low mass
- $|T\downarrow, m\uparrow\rangle$: low time / high mass

The Fractal Correction

The fractal correction K_{frak} plays a central role in T0: $K_{\text{frak}}^{3/2}$ explains the $\sim 2\%$ deviation in the anomalous magnetic moment of the muon (g-2 anomaly) [8]. This is structurally analogous to the annealing mechanism in the CIM.

N.5 Six Structural Parallels: T0 and CIM

Parallel 1: Determinism Behind Apparent Stochasticity

Table N.3: Parallel 1: Determinism

Aspect	Coherent Ising Machine	T0 Theory
Surface	Apparently stochastic search	Apparently probabilistic QM
Deep structure	Deterministic (OPO field eqs.)	Deterministic ($m(x, t)$ geometry)
Source of "randomness"	Unknown initial conditions	Unknown field configuration
Consequence	Solution is physically forced	Measurement result geometrically fixed

Both systems replace fundamental probabilism with hidden determinism: in the CIM the solution is not computed, it *emerges*. In T0 the measurement result is not chosen randomly – it is already determined by the mass field $m(x, t)$.

Parallel 1

Apparent probabilism $\equiv$ hidden determinism through geometric field structure
– in CIM *and* T0.

Parallel 2: Ground State as Geometric Minimum

The CIM solves optimization through physical relaxation:

$$H_{\text{Ising}} \rightarrow \text{minimum} \equiv \text{optimization solution} \quad (\text{N.6})$$

T0 describes the universe as already residing in the ground state of its own geometry:

$$\xi = \frac{4}{30000} \equiv \text{geometric minimum} \Rightarrow \text{all fundamental constants} \quad (\text{N.7})$$

In both cases the "solution" is not a computation – it is the physically natural resting state of the system. **Optimization is relaxation, not iteration.**

Parallel 3: Qubit Structure

Fundamental Difference

The T0 spin is *more fundamental* than the CIM spin: it carries its own constraint ($T \cdot m = 1$) intrinsically. A CIM spin must be externally constrained to ± 1 ; a T0 spin is constrained by the geometry of spacetime itself.

Table N.4: Parallel 3: Comparison of qubit structures

Aspect	CIM Spin	T0 Qubit
Carrier	Phase of laser pulse	Time-mass relation
Spin +1	Phase 0°	$ T\uparrow, m\downarrow\rangle$
Spin -1	Phase 180°	$ T\downarrow, m\uparrow\rangle$
Constraint	External (laser pump)	Intrinsic: $T \cdot m = 1$
Superposition	Coherent phase superposition	$\alpha T\uparrow, m\downarrow\rangle + \beta T\downarrow, m\uparrow\rangle$
Collapse	Phase measurement	Geometric field projection

Table N.5: Parallel 4: Coupling matrix – the decisive difference

Aspect	Classical CIM	T0 Ising Machine (concept)
Couplings J_{ij}	Externally programmed	Geometrically emergent
Degrees of freedom	J_{ij} freely selectable	Uniquely fixed by field geometry
Physical meaning	Mathematical auxiliary	Real geometric interactions
Universality	Arbitrary problems encodable	Physically meaningful problems
Advantage	Maximum flexibility	No encoding errors possible

Parallel 4: The Coupling Matrix J_{ij}

This is the deepest and physically most significant parallel:

In a T0-based Ising Machine, J_{ij} is *not* a free parameter – it emerges directly from the intrinsic time field:

$$J_{ij}^{(T0)} = \int T(x, t)_i \cdot T(x, t)_j \cdot G(x_i, x_j) d^3x \quad (\text{N.8})$$

where $G(x_i, x_j)$ is the geometric propagator of the T0 mass field.

Parallel 5: Fractal Correction as Annealing Mechanism

Table N.6: Parallel 5: Fractal correction and annealing

Aspect	Simulated Annealing (CIM)	T0 Fractal Correction
Parameter	Temperature T_{ann}	Correction factor K_{frak}
Function	Prevents local minima	Explains systematic deviations
Effect	Cooling $\rightarrow$ ground state	Scaling $\rightarrow$ precision
Example	MaxCut optimization	g-2 anomaly: $K_{\text{frak}}^{3/2}$ explains $\sim 2\%$
Mechanism	Thermal fluctuations	Fractal self-similarity

The fractal correction $K_{\text{frak}}^{3/2}$ in T0 plays the analogous role of the annealing temperature parameter: it prevents the system from becoming trapped in a local minimum of the geometric energy landscape.

Parallel 6: Toroidal Topology as Optimal CIM Geometry

T0 describes the universe as a 4D torus crystal [9]. This topology has direct implications for a T0-based Ising Machine:

- **Toroidal boundary conditions:** no boundary problem in optimization – all spins have equally many neighbors
- **Periodic structure:** natural implementation of translation-invariant couplings
- **4D instead of 3D:** richer coupling structure than classical 2D/3D CIM arrays
- **Crystal symmetry:** discrete symmetry groups → natural spin lattices without external structure specification

Parallel 6

A T0 Ising Machine would operate on the *natural geometry of the universe* – not on an artificially constructed graph.

N.6 The T0 Ising Machine: Conceptual Formulation

Overall Structure

Based on the six identified parallels, a T0-based Ising Machine can be conceptually defined:

Table N.7: Overall comparison: Classical CIM vs. T0 Ising Machine

Component	Classical CIM	T0 Ising Machine
Spin carrier	DOPO laser pulses (phase)	Time field states $T(x, t)$
Spin values	± 1 (phase $0^\circ/180^\circ$)	$ T\uparrow, m\downarrow\rangle, T\downarrow, m\uparrow\rangle$
Couplings	Externally programmed	Geometrically emergent from $m(x, t)$
Topology	Arbitrary graph	4D torus crystal
Ground state	Solution sought	Geometric fixed point ξ
Annealing	External (temperature)	Internal (K_{frak} correction)
Determinism	Hidden deterministic	Explicitly deterministic

Formal Correspondence

The complete mapping between T0 geometry and the Ising formalism reads:

$$\sigma_i \leftrightarrow \text{sign}(T(x, t)_i - \bar{T}) \quad [\text{spin from time field}] \quad (\text{N.9})$$

$$J_{ij} \leftrightarrow \langle T(x, t)_i \cdot T(x, t)_j \rangle_\xi \quad [\text{coupling from } \xi \text{ geometry}] \quad (\text{N.10})$$

$$h_i \leftrightarrow \frac{\partial \xi}{\partial T(x, t)_i} \quad [\text{external field from } \xi \text{ gradient}] \quad (\text{N.11})$$

$$H_{\text{Ising}} \leftrightarrow E_{\text{T0}} = \int |\nabla T(x, t)|^2 d^4x \quad [\text{energy functionals identical}] \quad (\text{N.12})$$

The minimum of the T0 energy functional (N.12) is identical to the ground state of the associated Ising Hamiltonian – with the crucial difference that in T0 all quantities (J_{ij} , h_i , H) emerge from a single parameter ξ .

The Central Consequence

Fundamental Implication of the T0 Ising Machine

If T0 is correct, then every physically realizable optimization problem is already encoded as a geometric structure in the mass field $m(x, t)$. A T0 Ising Machine would not optimize – it would *read* a physical reality. The solution is not the result of a search, but the unveiling of an already-existing geometric truth.

N.7 Open Questions and Research Directions

Theoretical Openness

- What is the explicit geometric propagator $G(x_i, x_j)$ in the T0 mass field? What symmetries does the torus topology enforce?
- Is there a natural restriction on which NP-hard problems are representable through T0 geometry – or are all problems in principle encodable?
- How does K_{frak} behave as an annealing parameter for different problem classes? Are there universal scaling laws?
- What is the connection between the torus structure of the T0 universe and optimal CIM topologies (toroidal vs. Möbius-band couplings)?

Experimental Directions

- **Coupling structure comparison:** compare the coupling matrix of existing CIMs with T0 field propagator predictions. Are there systematic deviations pointing to geometric constraints?
- **Toroidal CIM topologies:** do toroidal CIM architectures show performance advantages over open graphs that go beyond combinatorial explanations?
- **T0 qubit test:** do the T0 qubits defined in (N.5) show characteristic signatures in existing photonic quantum computing experiments?
- **Testable prediction:** in a CIM with N spins, the T0 correction factor $\mathcal{D}(N) = \exp(-\xi \ln N / D_{\text{frak}})$ should produce a measurable $\ln(N)$ -scaling residual that is qualitatively distinct from standard error models ($1/N$ or e^{-N}).

Connection to Scramblon Physics

This document extends [10], which analyzed the T0 connection to scramblon physics. Combining both approaches yields a complete picture:

- **Scramblon theory:** dynamic description – how quantum information scrambles in a CIM-like system
- **T0 Ising Machine:** geometric description – which fundamental couplings follow from the spacetime structure
- **Synthesis:** scramblons describe the *dynamics* of information spreading; T0 describes the *geometry* of the coupling structure

N.8 Summary

We have identified and systematically analyzed six structural parallels between Coherent Ising Machines and the T0 theory:

Table N.8: Summary: Six parallels T0 ↔ CIM

#	Parallel	CIM	T0
1	Determinism	Hidden (OPO)	Explicit (geometry)
2	Ground state	Solution sought	Fixed point ξ
3	Qubit structure	Phase $0^\circ/180^\circ$	$T \cdot m = 1$ states
4	Couplings J_{ij}	Externally programmed	Geometrically emergent
5	Annealing	External temperature	K_{frak} correction
6	Topology	Arbitrary graph	4D torus

T0 does not merely provide a conceptual parallel to the CIM – it offers a fundamentally deeper justification: optimization is not a computation executed on an artificially constructed system, but physical relaxation into a geometric truth already inscribed in the mass field of the universe.

Central Result

The formal correspondence

$$H_{\text{Ising}} \leftrightarrow E_{\text{T0}} = \int |\nabla T(x, t)|^2 d^4x \quad (\text{N.13})$$

shows: Ising optimization and T0 energy minimization are mathematically equivalent formulations of the same geometric principle – with the decisive difference that in T0 *all* couplings emerge parameter-free from $\xi = 4/30000$.

Bibliography

- [1] T. Inagaki et al., A coherent Ising machine for 2000-node optimization problems, *Science* **354**, 603 (2016).
- [2] P. L. McMahon et al., A fully programmable 100-spin coherent Ising machine with all-to-all connections, *Science* **354**, 614 (2016).
- [3] Y. Yamamoto et al., Coherent Ising machines – optical neural networks operating at the quantum limit, *npj Quantum Inf.* **3**, 49 (2017).
- [4] T. Honjo et al., 100,000-spin coherent Ising machine, *Sci. Adv.* **7**, eabh0952 (2021).
- [5] J. Pascher, T0 Theory: Time-Mass Duality as a Fundamental Principle, T0 Documentation Series, Document 003 (2025).
- [6] J. Pascher, The Geometric Origin Parameter $\xi = 4/30000$, T0 Documentation Series, Document 009 (2025).
- [7] J. Pascher, T0 Quantum Computing: Qubits and Logic Gates, T0 Documentation Series, Document 147 (2025).
- [8] J. Pascher, Fractal Corrections and the g-2 Anomaly, T0 Documentation Series, Document 018 (2025).
- [9] J. Pascher, The Universe as a 4D Torus Crystal: Geometric Cosmology, T0 Documentation Series, Document 026 (2025).
- [10] J. Pascher, T0 Projection onto Scramblon Physics, T0 Documentation Series, Document 148 (2026).

Chapter 0

Doc. 162 — T0 Theory — Optimization Comparison

Abstract

The T0 theory contains two complementary optimization frameworks: Document 034 (*QM-Optimization*) describes how quantum computers can be optimized through geometric transformation in cylindrical phase space and through noise suppression via time-field modulation. Document 161 (*T0 Ising Machine*) shows that combinatorial NP-hard optimization can be solved through geometric relaxation of the mass field $m(x, t)$. This document 162 systematically compares both approaches and demonstrates: they are not competing, but *complementary levels of the same geometric principle* – optimization as physical relaxation into a geometric minimum.

O.1 Introduction: Two Optimization Levels in T0

The Starting Problem

Quantum computing struggles with noise and suboptimality on two fundamentally different levels:

1. **Gate level:** Individual quantum gates are disturbed by decoherence, phase noise, and fractal damping. The question: how does one optimize *individual operations*?
2. **Problem level:** Combinatorial optimization problems (NP-hard) are hard to solve even with perfect gates. The question: how does one find *global minima* in vast solution spaces?

The T0 theory addresses **both levels** – but with different tools and at different levels of abstraction. Document 034 [1] solves Problem 1; Document 161 [2] solves Problem 2.

The Common Root

Both approaches share the same geometric origin: the parameter $\xi = 4/30000 \approx 1.333 \times 10^{-4}$ and the time-field binding condition $T \cdot m = 1$. The differences are not conceptual – they are differences in the *scaling level* of application.

O.2 Document 034: QM-Optimization through Geometric Formalism

The Cylindrical Phase Space

In Document 034 [1], Hilbert space is replaced by a **cylindrical phase space**. A qubit is not a 2D complex vector, but a point (z, r, θ) :

- $z \in [-1, 1]$: classical basis projection ($z = +1 \equiv |0\rangle$, $z = -1 \equiv |1\rangle$)
- $r \geq 0$: coherence radius (superposition magnitude), with $z^2 + r^2 = 1$
- $\theta \in [0, 2\pi)$: relative phase of the superposition

Gate operations are geometric transformations of this space, not matrix multiplications.

Fractal Damping as Intrinsic Noise

The X-gate (bit-flip) is not an ideal π -rotation in the T0 formalism, but a damped rotation by angle $\alpha = \pi \cdot K_{\text{frak}}$:

$$z' = z \cos(\alpha) - r \sin(\alpha) \quad (\text{O.1})$$

$$r' = z \sin(\alpha) + r \cos(\alpha) \quad (\text{O.2})$$

with the fractal damping constant $K_{\text{frak}} = 1 - 100\xi$. A perfect flip ($\alpha = \pi$) is physically impossible – spacetime inherently damps every rotation. This deviation is not directly measurable: it is absorbed into the pulse calibration of real qubits and therefore cannot be tested with current hardware.

The Three Optimization Strategies from Doc. 034

1. **T0-Topology-Compiler**: Qubits are arranged not to minimize SWAP operations, but to minimize the cumulative *fractal path length* of all entangling operations. Critically interacting qubits must be brought physically closer together.
2. **Harmonic resonance frequencies**: Qubit frequencies are calibrated to “magic” frequencies arising from the harmonic structure of the T0 time field:

$$f_n = \left(\frac{E_0}{h} \right) \cdot \xi^2 \cdot (\phi_T^2)^{-n} \quad (\text{O.3})$$

For superconducting qubits, with $E_0 = 7.398$ MeV sweet spots emerge at approximately **5.50 GHz** ($n = 9$) and **2.10 GHz** ($n = 10$). This calibration intrinsically reduces phase noise.

3. **Active time-field modulation**: A high-frequency *time-field pump* irradiates idle qubits to average out the fundamental ξ -noise and overcome the passive T_2 coherence limit – active decoherence suppression instead of passive waiting.

O.3 Document 161: T0 Ising Machine – Combinatorial Optimization

The Ising Model as Optimization Language

In Document 161 [2], it is shown that NP-hard combinatorial problems can be formulated as minimization of the Ising Hamiltonian:

$$H_{\text{Ising}} = -\frac{1}{2} \sum_{i \neq j} J_{ij} \sigma_i \sigma_j - \sum_i h_i \sigma_i \quad (\text{O.4})$$

A Coherent Ising Machine (CIM) finds the ground state of this Hamiltonian through physical relaxation of laser pulses (spins = phase states 0° or 180°).

The T0 Ising Machine: Emergent Couplings

The T0 theory extends the CIM fundamentally: the couplings J_{ij} are not externally programmable – they geometrically emerge from the time field:

$$J_{ij}^{(\text{T0})} = \int T(x, t)_i \cdot T(x, t)_j \cdot G(x_i, x_j) d^3x \quad (\text{O.5})$$

The energy functional is:

$$H_{\text{Ising}} \leftrightarrow E_{\text{T0}} = \int |\nabla T(x, t)|^2 d^4x \quad (\text{O.6})$$

Noise in the CIM: The Annealing Problem

In the CIM, noise plays a paradoxical role: it is simultaneously problem and tool. Too little thermal noise leads to local minimum traps; too much noise prevents convergence. The annealing temperature parameter T_{ann} balances this conflict.

In the T0 Ising Machine, $K_{\text{frak}}^{3/2}$ takes this role – not as an external parameter, but as a geometrically determined fractal correction factor.

O.4 Systematic Comparison: Doc. 034 vs. Doc. 161

Overview Table

Table O.1: Direct comparison of the two T0 optimization frameworks

Aspect	Doc. 034: QM-Optimization	Doc. 161: T0 Ising Machine	Common Root
Target problem	Gate noise, decoherence	NP-hard optimization	Geometric relaxation
Scaling level	Single qubit / gate	Full system (N spins)	$\xi = 4/30000$
Spin representation	(z, r, θ) in cylinder	Phase $0^\circ/180^\circ$ (CIM)	$T \cdot m = 1$ duality
Role of noise	Disturbance (minimize)	Tool (annealing)	K_{frak} correction
Coupling	Gate pulses (external)	J_{ij} (geometrically emergent)	Time field $T(x, t)$
Optimization goal	Maximum gate fidelity	Global minimum of H	Energy minimization
Determinism	Explicit (geometric trafo)	Hidden (OPO equations)	Both deterministic
Prediction	Damped X-flip (not directly measurable)	$\ln(N)$ residual after CIM	ξ signatures

Parallel 1: Noise – Enemy or Ally?

This is the deepest conceptual difference between both documents:

Table O.2: The opposing role of noise in both frameworks

Aspect	Doc. 034: QM-Optimization	Doc. 161: T0 Ising Machine
Noise is...	Enemy – disturbs gate fidelity	Tool – enables annealing
T0 cause	Fractal damping K_{frak}	Thermal fluctuations in OPO
Strategy	Actively suppress	Controlledly deploy
Mechanism	Time-field pump modulates $T(x)$	Temperature parameter T_{ann}
Optimum	Noise $\rightarrow 0$	Noise $\rightarrow$ minimum (then $\rightarrow 0$)
T0 analogue	$K_{\text{frak}} \rightarrow 1$ (perfect gates)	$K_{\text{frak}}^{3/2}$ as annealing force

Resolution of the Apparent Contradiction

The apparent contradiction – noise is bad in Doc. 034, good in Doc. 161 – dissolves through the scaling level: at the gate level, any noise is harmful. At the system level, controlled noise is necessary to escape local minima. T0 describes the same physical mechanism (K_{frak}) at both levels.

Parallel 2: Fractal Damping as Universal T0 Mechanism

Parallel 3: Frequency Optimization vs. Topology Optimization

Both documents propose optimizations of the *system architecture* – at different levels:

Parallel 4: Determinism and Measurability

Table O.3: K_{frak} as a universal T0 mechanism at different levels

Context	Doc. 034	Doc. 161	Physical Meaning
K_{frak} definition	$1 - 100\xi$	$K_{\text{frak}}^{3/2}$ (g-2 correction)	Fractal spacetime structure
Acts on...	X-gate rotation (damps)	Annealing dynamics (regulates)	Deviation from ideal
Mathematically	$\alpha = \pi \cdot K_{\text{frak}}$	$\mathcal{D}(N) = e^{-\xi \ln N / D_{\text{frak}}}$	Geometric damping
Measurable via	Residual flip error (masked by calibration)	$\ln(N)$ residual after CIM	ξ signatures
Testable on	Not testable with current hardware	CIM with variable spin count	Gate side currently not testable

Table O.4: Architecture optimizations: gate level vs. system level

Optimization type	Doc. 034: QM-Optimization	Doc. 161: T0 Ising Machine
Frequency optimization	Harmonic resonance frequencies f_n	–
Topology optimization	T0-Topology-Compiler (fractal path length)	4D torus topology
Decoherence protection	Active time-field modulation	Toroidal boundary conditions
Coupling structure	Gate pulses pre-compensated	J_{ij} geometrically emergent
Goal	Max. gate fidelity	Min. Ising Hamiltonian

Table O.5: Determinism and predictions of both documents

Aspect	Doc. 034: QM-Optimization	Doc. 161: T0 Ising Machine
Type of determinism	Explicit: geometric transformations	Hidden: OPO field equations
Source of stochasticity	Fractal spacetime fluctuations	Quantum vacuum initial conditions
Prediction 1	Damped X-flip: $\alpha < \pi$ (not directly measurable)	$\ln(N)$ -scaling residual
Testable prediction 2	Sweet spots at 5.50/2.10 GHz	J_{ij} deviations from free param.
Platform	IBM Brisbane/Sherbrooke	CIM with 1000+ spins
Quantification	$\Delta\alpha = \pi(1 - K_{\text{frak}}) \approx 4.2 \times 10^{-2}$	$\xi \approx 1.3 \times 10^{-4}$

O.5 New Optimization Opportunities through the Connection

The systematic comparison of both documents opens optimization opportunities not visible in either document alone:

Hybrid Optimization: Gate Level + System Level

Hybrid T0 Optimization Strategy

A fully T0-optimized quantum architecture combines both levels:

1. **Gate level** (Doc. 034): Harmonic frequency calibration + time-field pump for maximum gate fidelity
2. **System level** (Doc. 161): T0 Ising Machine for combinatorial optimization with geometrically emergent couplings
3. **Linking level**: The T0-Topology-Compiler (Doc. 034) minimizes fractal path lengths – this is the same geometric principle as the torus structure in Doc. 161

Noise as a Bridge between Both Levels

The dualistic role of noise opens a new optimization strategy:

- **Phase 1 (Annealing)**: Controlled noise (K_{frak} regime) – the T0 Ising Machine searches for global minima
- **Phase 2 (Precision)**: Noise suppression (time-field pump) – quantum gates execute the solution with maximum fidelity
- **Transition**: The fractal parameter K_{frak} regulates both phases – as annealing mechanism (Doc. 161) and as gate damping (Doc. 034)

$$\text{Optimization} = \underbrace{K_{\text{frak}}^{3/2}\text{-annealing}}_{\text{Doc. 161: global minimum}} + \underbrace{K_{\text{frak}} \rightarrow 1\text{-precision}}_{\text{Doc. 034: gate fidelity}} \quad (0.7)$$

Ratio between Gate Error and CIM Residual

If K_{frak} describes the same universal T0 parameter at both levels, then:

$$\frac{\Delta\alpha_{\text{Gate}}}{\pi} = 1 - K_{\text{frak}} = 100\xi \approx 1.33 \times 10^{-2}, \quad \frac{1 - \mathcal{D}_{\text{CIM}}(N)}{\ln N} \approx \frac{\xi}{\Delta f} \approx 4.4 \times 10^{-5} \quad (0.8)$$

Both quantities are fixed by the same ξ , but they are **not equal**: the relative gate flip error on IBM quantum computers ($100\xi \approx 1.33 \times 10^{-2}$) and the $\ln(N)$ residual in CIM measurements ($\xi/\Delta f \approx 4.4 \times 10^{-5}$) differ by the factor $100\Delta f \approx 300$; convergence to the same numerical value is not to be expected. This fixed ratio, too, is currently not experimentally testable, because the gate deviation is absorbed into the pulse calibration of real qubits and is not directly measurable.

O.6 Summary: Complementary Levels of the Same Principle

Table O.6: Full synthesis: both documents as complementary T0 levels

Dimension	Doc. 034	Doc. 161	Connection
Level	Gate / qubit	System / spins	Same ξ
Noise	Enemy (suppress)	Tool (control)	K_{frak} at both levels
Representation	Cylindrical (z, r, θ)	Phase $0^\circ/180^\circ$	$T \cdot m = 1$ states
Topology	Fractal path length	4D torus	Geometric minimization
Coupling	Gate pulses (pre-compensated)	J_{ij} (emergent)	Time field $T(x, t)$
Prediction	$\Delta\alpha \approx 4.2 \times 10^{-2}$	$\ln(N)$ residual	$\xi = 4/30000$

The central insight is:

The Unified T0 Optimization Perspective

Both gate noise (Doc. 034) and combinatorial NP-optimization (Doc. 161) are manifestations of *the same geometric principle*: physical relaxation into the minimum of the T0 energy functional $E_{\text{T0}} = \int |\nabla T(x, t)|^2 d^4x$. The difference is not conceptual – it is the scaling level of observation. The fractal correction K_{frak} appears at both levels as the same universal parameter; the predictions of both levels are fixed by the same ξ but differ by the factor $100 \Delta f \approx 300$.

Bibliography

- [1] J. Pascher, T0 Quantum Mechanics: Geometric Formalism and Optimization Strategies for Quantum Computers, T0 Documentation Series, Document 034 (2025).
- [2] J. Pascher, T0 Theory and Coherent Ising Machines: Structural Parallels, T0 Documentation Series, Document 161 (2026).
- [3] J. Pascher, T0 Theory: Time-Mass Duality as a Fundamental Principle, T0 Documentation Series, Document 003 (2025).
- [4] J. Pascher, Fractal Corrections and the g-2 Anomaly, T0 Documentation Series, Document 018 (2025).
- [5] J. Pascher, T0 Quantum Computing: Qubits and Logic Gates, T0 Documentation Series, Document 147 (2025).
- [6] J. Pascher, T0 Projection onto Scramblon Physics, T0 Documentation Series, Document 148 (2026).
- [7] T. Inagaki et al., A coherent Ising machine for 2000-node optimization problems, *Science* **354**, 603 (2016).

Chapter P

Doc. 163 — T0 Theory — UV Catastrophe, Planck, Einstein

Abstract

The ultraviolet catastrophe and Einstein's light quantum hypothesis mark two central turning points in the history of physics. This document analyses both aspects from three perspectives: first, the historical and physical development — from Maxwell's electromagnetic wave theory through Planck's desperate quantization trick to Einstein's thermodynamically forced conclusion that light consists of independent energy quanta; second, the remaining open questions in standard physics — why $\epsilon = hf$? why does h have this particular value? what sets an upper limit on frequency?; and third, the T0 answers to both questions: Planck's constant h is derived, via the chain $\xi \rightarrow \alpha \rightarrow h$, from the geometric parameter $\xi = 4/30000$ — all SI values are derivable from ξ and the single unit calibration, no separate convention is needed; and the ultraviolet divergence is interpreted in the T0 framework as a consequence of the assumption that spacetime is continuous down to arbitrarily small scales. The sub-Planck scale $L_0 = \xi \cdot \ell_P$ imposes a geometrically fixed maximum frequency $f_{\max}$ above which no modes exist.

P.1 Introduction: Two Crises, One Geometric Principle

The Historical Context

At the end of the 19th century, physics found itself in a peculiar combination of triumph and bewilderment. Maxwell's theory of electromagnetism had unified light, electricity, and magnetism in a single, mathematically closed framework. Phenomena such as interference and diffraction seemed to confirm the wave picture of light clearly. Many physicists regarded Maxwell's theory as one of the pinnacles of scientific achievement.

Yet this very theory failed clearly the moment it was applied to one of the simplest imaginable situations: the thermal radiation of hot bodies. The result was a prediction that was physically absurd — infinite energy density.

This crisis became the birthplace of quantum physics.

What This Document Achieves

We follow three parallel threads throughout this document:

1. **The classical crisis:** Why does classical theory diverge at high frequencies? What exactly goes wrong, and why is this not trivial?
2. **The historical answers:** How did Planck (1900) and Einstein (1905) solve the problem — and what remained open? It is shown that Einstein did *not postulate* the light quantum but *extracted* it from thermodynamics.
3. **The T0 perspective:** How does T0 theory describe both aspects geometrically? Planck's constant is traced back to ξ , and the UV divergence drops out because in the T0 framework spacetime possesses a geometric minimum structure.

Guiding Questions

Why does $\epsilon = hf$ hold? Why does Planck's constant have exactly this value? And why is there a natural upper limit to frequency? Classical physics does not answer these questions. T0 offers a geometric answer.

P.2 The Ultraviolet Catastrophe: A Precise Diagnosis

The Blackbody Radiation Problem

Imagine a completely closed cavity whose walls have been heated uniformly to temperature T and are in thermal equilibrium. If a tiny hole is drilled in the wall, radiation escapes. This *cavity radiation* (also: *blackbody radiation*) possesses an important property: its spectrum depends *solely* on the temperature — not on the material or shape of the cavity.

The experimentally measured quantity is the *spectral energy density* $\rho(f, T)$: the energy per unit volume per unit frequency interval. At different temperatures, one obtains characteristic curves with clear signatures:

- The peak of the curve shifts to higher frequencies as temperature increases (Wien's displacement law).
- The total area under the curve grows as the fourth power of temperature (Stefan-Boltzmann law: $u \propto T^4$).
- A hot piece of metal glows first dull red, then brighter orange-yellow, finally white-hot — the dominant frequency visibly shifts upward.

The challenge for theoretical physics around 1900 was precise: derive this curve from the fundamental laws of physics.

The Classical Approach and Its Failure

The natural starting point was Maxwell's theory combined with the well-established statistical mechanics. The approach is conceptually simple:

Step 1: Counting modes. Inside the cavity, only standing electromagnetic waves can form — precisely those wavelengths for which an integer number of half-wavelengths fit between the walls. The number of electromagnetic modes per unit volume in the frequency interval $[f, f + df]$ is:

$$g(f) df = \frac{8\pi f^2}{c^3} df \quad (\text{P.1})$$

The crucial point: mode density grows quadratically with frequency. For every doubling of frequency, there are four times as many independent wave patterns. And there is *no upper limit* — wavelength can become arbitrarily small, frequency arbitrarily large.

Step 2: Equipartition theorem. Classical statistical mechanics (Boltzmann, Maxwell) yields the equipartition theorem: in thermal equilibrium at temperature T , each quadratic degree of freedom carries on average the energy $\frac{1}{2}k_B T$. An electromagnetic wave has two quadratic degrees of freedom (electric and magnetic field), so each mode carries on average:

$$\langle E_{\text{mode}} \rangle = k_B T \quad (\text{P.2})$$

This holds *independently of the frequency of the mode*. A mode at 10^{15} Hz receives the same average energy as a mode at 10^3 Hz.

Step 3: Spectral energy density. The spectral energy density follows as a simple product:

$$\rho_{\text{class}}(f, T) = g(f) \cdot \langle E_{\text{mode}} \rangle = \frac{8\pi f^2}{c^3} \cdot k_B T \quad (\text{P.3})$$

This is the *Rayleigh-Jeans law*. It agrees well with experiment at low frequencies. At high frequencies, however, it diverges.

The Scale of the Catastrophe

The total energy density is obtained by integrating over all frequencies:

$$u_{\text{class}} = \int_0^\infty \rho_{\text{class}}(f, T) df = \frac{8\pi k_B T}{c^3} \int_0^\infty f^2 df = \infty \quad (\text{P.4})$$

The integral diverges. Classical physics predicts: a cavity in thermal equilibrium contains *infinite energy*. This is physically absurd.

This divergence is called the **ultraviolet catastrophe** because it manifests most clearly at the high-frequency (ultraviolet and beyond) end of the spectrum. The name is somewhat misleading — the divergence actually extends across the entire high-frequency spectrum, not just the ultraviolet range.

Why the Failure Is Deep

The Rayleigh-Jeans law is not a calculational error. It is the correct application of two well-confirmed theories: Maxwell's electrodynamics and Boltzmann's statistical mechanics. Its failure therefore shows that one of these foundations must be fundamentally wrong — or that an important physical fact is missing.

The Two Empirical Partial Solutions

Before the complete solution, there were two important empirical clues:

The Rayleigh-Jeans law (derived around 1900) fits well at *low* frequencies:

$$\rho_{\text{RJ}}(f, T) \approx \frac{8\pi f^2}{c^3} k_B T \quad (\text{for } f \rightarrow 0) \quad (\text{P.5})$$

Wien's radiation law (1896) describes *high* frequencies well:

$$\rho_{\text{Wien}}(f, T) = a f^3 \exp\left(-\frac{bf}{T}\right) \quad (\text{P.6})$$

where a and b are empirical constants. At high frequencies, the exponential falls off faster than f^2 grows — the integral converges.

The problem: both laws hold only in their respective limiting regimes. Between them lie the experimentally determined data, showing a smooth connection. A unified theory was missing.

P.3 Planck's Solution: An Act of Desperation

Planck's Strategy

Max Planck approached the problem differently from his contemporaries. Rather than starting from a specific microscopic model, he worked *thermodynamically backwards*: he searched for the mathematical expression for the entropy of cavity radiation that was consistent with the empirically known spectrum.

The decisive step: Planck found a formula that reduces to the Rayleigh-Jeans law at low frequencies and to Wien's law at high frequencies:

$$\rho_{\text{Planck}}(f, T) = \frac{8\pi h f^3}{c^3} \cdot \frac{1}{\exp\left(\frac{hf}{k_B T}\right) - 1} \quad (\text{P.7})$$

This formula describes the experimental spectrum across the entire frequency range with high precision — with only one new constant: Planck's constant h .

The Price: Quantization as a Postulate

To derive his formula, Planck had to make an assumption that he himself described as an *act of desperation*: he assumed that energy can only be exchanged in *discrete multiples* of hf . The energy of a single portion (*quantum*) is:

$$\epsilon = hf \quad (\text{P.8})$$

With this assumption, the average energy of a mode at frequency f is no longer $k_B T$ (equipartition theorem) but:

$$\langle E_{\text{mode}} \rangle = \frac{hf}{\exp\left(\frac{hf}{k_B T}\right) - 1} \quad (\text{P.9})$$

For $hf \ll k_B T$ (low frequencies), this expression approaches $k_B T$ — the classical result is recovered. For $hf \gg k_B T$ (high frequencies), the average energy falls off exponentially — exactly what is needed to suppress the divergence.

What Planck Did Not Explain

Quantization solved the spectral problem mathematically. But Planck himself regarded it as a computational auxiliary construct, not a physical statement about the nature of light. In his view, the electromagnetic field remained a continuous wave. Quantization was a property of the *interaction* between matter and radiation — not of radiation itself.

Planck's Open Question

Planck had found the curve. But he had not explained *why* energy is exchanged in discrete portions. The number h was a free parameter — measured from the data, not derived from deeper principles. What is h ? Why exactly this value?

P.4 Einstein's Light Quanta: Thermodynamically Forced

A Widespread Misconception

Einstein's 1905 contribution is typically summarized as "the explanation of the photoelectric effect." This is misleading. The photoelectric effect appears only at the end of the paper as one of several consequences. Einstein does not begin with metals, electrons, or experiments.

He begins with a *structural problem within physics itself*.

The Structural Problem

On one side stands the theory of matter: gases and other ponderable bodies are described by a *finite* number of discrete quantities — the positions and velocities of atoms. The state of the system is specified by a finite list.

On the other side stands Maxwell's theory: the state of the electromagnetic field is *not* given by a finite list but by continuous fields — smooth functions defined at every point in space. These are infinitely many degrees of freedom.

In Einstein's own words: there is a *profound formal distinction* between these two ways of describing nature. The question is: if both systems can be in thermal equilibrium — do they really have such a fundamentally different structure?

Einstein's Method: Thermodynamics Instead of Model

Einstein refused to start from a speculative microscopic model of light. Instead, he wanted to discover what the most reliable available facts tell us about the nature of light.

These facts came from two sources:

1. The experimentally confirmed blackbody spectrum (particularly Wien's law in the high-frequency regime).
2. The most general and powerful equilibrium principle available: the second law of thermodynamics — specifically Boltzmann's statistical interpretation.

The Thermodynamic Derivation

Step 1: Entropy of radiation. Einstein introduces a *spectral entropy density* $\sigma(f, T)$ — analogous to the spectral energy density $\rho(f, T)$. In thermal equilibrium, the thermodynamic relation holds:

$$\frac{\partial \sigma}{\partial \rho} = \frac{1}{T} \quad (\text{P.10})$$

This relation follows directly from thermodynamics and presupposes no model.

Step 2: Entropy from Wien's law. By substituting Wien's law (P.6) into the thermodynamic relation (P.10) and integrating, Einstein obtains the entropy density in the Wien regime:

$$\sigma(f, T) = -\frac{\rho}{bf} \left[\ln \left(\frac{\rho}{af^3} \right) - 1 \right] \quad (\text{P.11})$$

Step 3: Entropy change under volume change. Einstein now considers a narrow, quasi-monochromatic slice of radiation with total energy E at frequency f , enclosed in volume V_0 . He calculates the entropy change when this volume is changed to V :

$$\Delta S = S(V) - S(V_0) = \frac{E}{bf} \ln \left(\frac{V}{V_0} \right) \quad (\text{P.12})$$

Step 4: Comparison with an ideal gas. This is the decisive moment. For an ideal gas of N independently moving particles, Boltzmann's entropy formula $S = k_B \ln W$ (with W as the number of microstates) yields:

$$\Delta S_{\text{gas}} = Nk_B \ln \left(\frac{V}{V_0} \right) \quad (\text{P.13})$$

The two expressions (P.12) and (P.13) have *identical structure*. Comparing the exponents gives:

$$\frac{E}{bf} = Nk_B \implies N = \frac{E}{k_B bf} \quad (\text{P.14})$$

Step 5: Energy of one quantum. If the total energy E is carried by N independent entities, each carrying energy ϵ , then $E = N\epsilon$, so:

$$\epsilon = \frac{E}{N} = k_B bf \quad (\text{P.15})$$

Since Wien's constant b turns out to be $b = h/k_B$ by comparison with Planck's law, we finally obtain:

$$\boxed{\epsilon = hf} \quad (\text{P.16})$$

Einstein's Decisive Insight

The light quantum was not postulated. It was *forced* by thermodynamics. The only inputs were: (1) the empirical Wien law, (2) the thermodynamic definition of equilibrium, (3) Boltzmann's entropy-probability relation. Nowhere was a particle nature of light assumed — it emerged as a logical consequence.

The Photoelectric Effect as Confirmation

Only after this derivation does Einstein apply his hypothesis to the photoelectric effect. The result is the famous equation:

$$E_{\text{kin, max}} = hf - \Phi \quad (\text{P.17})$$

where Φ is the *work function* of the metal. This equation is linear in f with slope h — a very sharp, experimentally testable prediction.

Experimental confirmation came through Robert Millikan (1916), who despite years of confirming the equation doubted the light quantum picture. Planck, recommending Einstein for the Prussian Academy in 1913, praised his brilliance while cautioning that the light quantum hypothesis should be counted as a miss. The Nobel Prize (1921) honoured the equation — not the interpretation behind it.

What Einstein Left Open

Even after Einstein's work, fundamental questions remained unanswered:

- **Why $\epsilon = hf$?** Einstein showed that monochromatic radiation in the Wien regime behaves thermodynamically as if it consisted of independent quanta. But *what* are these quanta microscopically?
- **Why exactly this value of h ?** Planck's constant is $h \approx 6.626 \times 10^{-34}$ J·s. Why this numerical value? Where does it come from? In standard physics, h is a free parameter — measured, not derived.
- **What limits frequency from above?** The UV divergence is suppressed by Planck's quantization, but the fundamental question remains: is frequency space really open all the way to $f \rightarrow \infty$? Is there no physical limit?

P.5 T0 Perspective I: The Origin of $\epsilon = hf$

The Geometric Parameter ξ

T0 theory traces the physical constants back to a single dimensionless geometric parameter (SI anchors serve to convert into units):

$$\xi = \frac{4}{30000} \approx 1.333 \times 10^{-4} \quad (\text{P.18})$$

This parameter describes the ratio between the fundamental sub-Planck scale and the cosmic scale. It is not free — it follows from the geometry of three-dimensional space at the Planck scale.

The Binding Condition

The heart of T0 theory is time-mass duality:

$$T(x, t) \cdot m(x, t) = 1 \quad (\text{in natural units}) \quad (\text{P.19})$$

The intrinsic time field of a particle with mass m is:

$$T(x, t) = \frac{\hbar}{m(x, t) \cdot c^2} \quad (\text{P.20})$$

Time and mass are not independent quantities — they are reciprocal aspects of a single geometric reality.

What Happens to the Photon?

The photon has no rest mass: $m_\gamma = 0$. In the T0 binding condition $T \cdot m = 1$, this naively implies $T \rightarrow \infty$. The physical interpretation: the photon *is* its time field — it does not exist *in* time, it *is* a time-like structure. From the photon's perspective, no proper time elapses.

For a photon with frequency f :

$$E_\gamma = hf = \frac{\hbar}{T(x, t)_\gamma} \quad (\text{P.21})$$

In T0, this is not an external equation but a consequence of the binding condition: energy is the reciprocal intrinsic time of the field.

Planck's Constant as an Emergent Quantity

In standard physics, h is a fundamental constant with value 6.626×10^{-34} J·s. In T0, h is not a free constant: its form is fixed via ξ , and its SI numerical value follows from ξ and the single unit calibration:

$$h = \xi \cdot \hbar_{\text{Planck}} \cdot f(\text{geometry}) \quad (\text{P.22})$$

Here $f(\text{geometry})$ is a general dependence on the geometry, not a frequency and not the base winding number $f = 1/\xi$; everywhere else in this document f stands for a frequency.

Concretely: the 2019 SI reform fixed h by definition (when calibrating $\alpha = 1$ in rational T0 units). The numerical value of h is not arbitrary — it encodes the geometric relationship between the sub-Planck scale $L_0 = \xi \cdot \ell_P$ and the macroscopic world.

T0 Answer to Question 1

Why does $\epsilon = hf$ hold? In T0, the answer is: because the energy of a photon equals the reciprocal intrinsic time of its field ($E = 1/T$), and frequency is the inverse time ($f = 1/T$). In T0 the equation $\epsilon = hf$ is not a postulate but follows directly from the time-field binding condition $T \cdot E = 1$ for massless fields.

The Connection to Einstein’s Argument

Einstein’s thermodynamic derivation and T0’s geometric derivation converge on the same result through two different paths:

Table P.1: Two paths to the equation $\epsilon = hf$

Aspect	Einstein 1905	T0 Theory
Starting point	Entropy of Wien radiation	Time-field binding $T \cdot E = 1$
Method	Thermodynamic comparison	Geometric deduction
Result	$\epsilon = hf$ (thermodynamically forced)	$\epsilon = hf$ (geometrically emergent)
Status of h	Free fit parameter	From $\xi = 4/30000$ and unit calibration
What remains open	Why this value of h ?	h value fixed via $\xi \rightarrow \alpha \rightarrow h$

P.6 T0 Perspective II: Why the UV Divergence Disappears

The Classical Dilemma Revisited

The ultraviolet catastrophe arises from the combination of two facts:

1. The number of modes grows like f^2 — without upper bound.
2. Each mode receives the same average energy $k_B T$.

Planck's solution suppresses the divergence by making the average energy fall off exponentially at high frequencies. But the more fundamental question remains unanswered: *Is frequency space really physically meaningful all the way to $f \rightarrow \infty$?*

The implicit assumption of classical physics — and also of quantum field theory — is: yes. Frequency space is unbounded. Spacetime is continuous down to arbitrarily small scales.

T0 replaces this assumption with a geometric minimum structure.

The Sub-Planck Scale in T0

The geometric parameter ξ defines a fundamental minimum structure of spacetime below the Planck scale:

$$L_0 = \xi \cdot \ell_P = \frac{4}{30000} \cdot \ell_P \approx 1.333 \times 10^{-4} \cdot \ell_P \quad (\text{P.23})$$

$$T_0 = \xi \cdot t_P = \frac{4}{30000} \cdot t_P \approx 1.333 \times 10^{-4} \cdot t_P \quad (\text{P.24})$$

where $\ell_P = \sqrt{\hbar G / c^3} \approx 1.616 \times 10^{-35}$ m and $t_P = \ell_P / c \approx 5.39 \times 10^{-44}$ s are the ordinary Planck units.

The scale L_0 is the geometrically smallest length at which spacetime still carries well-defined degrees of freedom. Below L_0 , no structure exists — not because we cannot measure it, but because spacetime itself carries no further degrees of freedom there.

The Maximum Physical Frequency

If T_0 is the smallest physically meaningful time scale, then there is a maximum frequency:

$$f_{\max} = \frac{1}{T_0} = \frac{c}{L_0} = \frac{c}{\xi \cdot \ell_P} \quad (\text{P.25})$$

Numerically:

$$f_{\max} = \frac{c}{\xi \cdot \ell_P} = \frac{3 \times 10^8 \text{ m/s}}{1.333 \times 10^{-4} \cdot 1.616 \times 10^{-35} \text{ m}} \approx 1.39 \times 10^{47} \text{ Hz} \quad (\text{P.26})$$

This is $1/\xi$ times larger than the Planck frequency $f_P \approx 1.85 \times 10^{43}$ Hz, and is therefore finite. The decisive point: there exists a finite upper bound.

The Integral Converges

In T0, the frequency integral is no longer improper:

$$u_{T0} = \int_0^{f_{\max}} \rho(f, T) df < \infty \quad (\text{P.27})$$

In the T0 framework the divergence from (P.4) drops out not through a calculational prescription but because the upper limit follows physically from L_0 . Above $f_{\max}$, there simply are no modes — spacetime carries no electromagnetic degrees of freedom there.

The Difference from Planck's Solution

Planck suppresses the divergence at high frequencies through a decaying Boltzmann factor: $\langle E \rangle \propto \exp(-hf/k_B T) \rightarrow 0$. The integral converges because the integrand becomes exponentially small.

T0 treats the divergence differently: there are no modes above $f_{\max}$. The upper integration limit is finite, not infinite. In the T0 framework the divergence therefore need not be suppressed — it does not arise physically in the first place.

The Classical Premise from the T0 Viewpoint

The classical Rayleigh-Jeans integral diverges because it starts from the following physical premise: spacetime is a continuous medium carrying infinitely many degrees of freedom at arbitrarily small scales.

T0 identifies this assumption as the true cause of the catastrophe:

$$\text{UV catastrophe} = \text{false assumption: } L_{\min} = 0 \quad (\text{P.28})$$

In the T0 framework, instead:

$$L_{\min} = L_0 = \xi \cdot \ell_P \neq 0 \quad (\text{P.29})$$

Spacetime is *discrete* at scale L_0 — not in the sense of a lattice model, but in the sense of a fractal geometric structure that contains no further degrees of freedom below L_0 .

Fractal Spacetime and Natural Regularization

T0 spacetime is fractal with Hausdorff dimension:

$$D_{\text{frak}} = 3 - \xi \approx 2.999867 \quad (\text{P.30})$$

This slight deviation from 3 affects the mode counting. In a fractal spacetime, the number of modes does not grow exactly like f^2 but like:

$$g_{T0}(f) df = \frac{8\pi f^2}{c^3} \cdot \left(\frac{f}{f_{\max}} \right)^{-\xi \cdot D_{\text{frak}}} df \quad (\text{P.31})$$

Since $f/f_{\max} < 1$ and the exponent is negative, this correction factor is larger than 1 and falls to exactly 1 towards $f_{\max}$: with $f_{\max} = 1.39 \times 10^{47}$ Hz one obtains 1.0358 at 1 GHz, 1.0338 at 100 GHz, 1.0300 at 10^{15} Hz and 1.000 at $f_{\max}$. The factor therefore produces an enhancement of about 3 % throughout the laboratory range, not a suppression near $f_{\max}$. Across the CMB band (60–600 GHz) it changes from 1.0341 to 1.0331; this is a spectral distortion of about 10^{-3} , far above the FIRAS limit of about 5×10^{-5} . In this form, Eq. (P.31) is therefore not compatible with the measured CMB spectrum.

P.7 Synthesis: What T0 Achieves and What Remains Open

The Complete Table

Table P.2: Comparison: Classical Physics — Planck/Einstein — T0

Problem	Classical	Planck 1900	Einstein 1905	T0
UV divergence	Diverges	Suppressed (quantization)	Not addressed	Drops out ($f_{\max}$)
Spectrum	Wrong law	Correct law	Confirmed	$f_{\max}$ correction
Status of h	No h	Free parameter	Free parameter	From ξ and calibration
Why $\epsilon = hf$?	No answer	Postulate	Thermodynamically forced	Geometrically ($T \cdot E = 1$)
Maximum frequency	None	None	None	$f_{\max} = c/(\xi \cdot \ell_P)$
Nature of photon	Wave	Quantized	Quantized	Time-field state
Free parameters	Many	h free	h free	Only ξ

What T0 Adds

Planck and Einstein solved the spectral problem: Planck through postulating quantization, Einstein through thermodynamic proof of quantum structure. What remained open in their work:

- *Why* energy is divided into quanta hf .
- *Why* h has exactly this numerical value.
- *What* is the physical limit of frequency space.

T0 addresses all three questions through a single mechanism: the geometric structure of spacetime at the sub-Planck scale $L_0 = \xi \cdot \ell_P$.

Testable Predictions

The T0 answers lead to concrete deviations that are testable in principle:

1. **Planck spectrum with $f_{\max}$ cutoff:** At extremely high energies, the blackbody spectrum should show a hard cutoff at $f_{\max} \approx 10^{47}$ Hz. This lies far beyond current experimental capabilities, but is in principle testable.
2. **Fractal corrections in g-2:** The correction factor $K_{\text{frak}}^{3/2}$ of the T0 time field appears in the anomalous magnetic dipole moment calculation and reduces the theoretical deviation from experiment to 0.033%: with $k_{\text{geom}} = 2.22357$, $k_{\text{eff}} = 2.22357/(74/75)^{3/2} = 2.2688$ and $a_e = 2\pi^2/(7500 \cdot 2.2688) = 1.1600 \times 10^{-3}$. This direct absolute formula works at the level of 10^{-4} and is not a precision comparison (cf. Doc. 018, 158, 382).
3. **Photoelectric effect correction:** At very high photon frequencies, Einstein's equation (P.17) should be modified by a ξ correction factor:

$$E_{\text{kin, max}}^{\text{T0}} = hf \left(1 - \xi \cdot \frac{f}{f_{\max}} \right) - \Phi \quad (\text{P.32})$$

The Methodological Parallel

Einstein and T0 proceed in a methodologically similar way. Einstein refused to presuppose a microscopic model of light. He followed thermodynamics wherever it led. The light quantum was the result, not the assumption.

T0 proceeds the same way. No specific model of spacetime discreteness is postulated. The question is asked: what follows from the geometric binding condition $T \cdot m = 1$ and the parameter ξ ? The maximum frequency, the emergent Planck constant, and the suppression of the UV divergence are the results — not the assumptions.

The Unified View

The ultraviolet catastrophe and the Planck-Einstein quantum problem are *the same problem at two different levels*. At the energy level: why does $\epsilon = hf$ hold? At the frequency level: why is there an upper limit? In T0, both have the same answer: because spacetime at the scale $L_0 = \xi \cdot \ell_P$ possesses a geometrically defined fundamental structure that permits neither h as a free parameter nor frequency space as unbounded.

P.8 Summary

This document has examined two historically and physically important problems and set out how T0 theory traces both back to a single geometric mechanism.

The first problem: the ultraviolet catastrophe. Classical theory predicts infinite energy density because it tacitly assumes that frequency space is unbounded from above. Planck suppresses the divergence through quantization — but without physical justification for the upper limit. In T0 the divergence drops out: the sub-Planck scale $L_0 = \xi \cdot \ell_P$ sets a maximum frequency $f_{\max}$, above which spacetime carries no electromagnetic degrees of freedom.

The second problem: the origin of $\epsilon = hf$. Planck postulated the equation as an auxiliary device. Einstein forced it thermodynamically. But both left h as a free parameter. T0 offers an origin of the form: the time-field binding condition $T \cdot E = 1$ fixes directly for massless fields $E = 1/T$, in SI units $E = hf = 2\pi\hbar f$. The SI numerical value of h does not follow from ξ alone, but via the chain $\xi \rightarrow \alpha \rightarrow h$ from ξ and the single unit calibration. A form $E = f\hbar/\xi$ does not hold, because it is too large by the factor $1/(2\pi\xi) = 1194$.

The common cause. Both problems share the same root: classical physics treats spacetime as a continuous medium without geometric minimum structure. T0 replaces this assumption with a minimum structure at the scale L_0 — with the one parameter $\xi = 4/30000$ both problems treated here are thereby resolved.

Bibliography

- [1] A. Einstein, On a heuristic viewpoint concerning the production and transformation of light, *Annalen der Physik* **17**, 132–148 (1905).
- [2] M. Planck, On the theory of the law of energy distribution in the normal spectrum, *Verhandlungen der Deutschen Physikalischen Gesellschaft* **2**, 237–245 (1900).
- [3] R. A. Millikan, A direct photoelectric determination of Planck's h , *Physical Review* **7**, 355–388 (1916).
- [4] J. Pascher, The Geometric Origin Parameter $\xi = 4/30000$, T0 Documentation Series, Document 009 (2025).
- [5] J. Pascher, T0 Theory: Time-Mass Duality as a Fundamental Principle, T0 Documentation Series, Document 003 (2025).
- [6] J. Pascher, Sub-Planck Structure and Geometric Regularisation in T0, T0 Documentation Series, Document 145 (2025).
- [7] J. Pascher, Fractal Corrections and the g-2 Anomaly (Rev. 12), T0 Documentation Series, Document 018 (2025).
- [8] J. Pascher, T0 Quantum Mechanics: Geometric Formalism and Optimisation Strategies, T0 Documentation Series, Document 034 (2025).

Chapter Q

Doc. 164 — T0 Theory vs. 't Hooft CAI

Abstract

Gerard 't Hooft's *The Cellular Automaton Interpretation of Quantum Mechanics* (Springer, 2016) and T0 theory pursue the same fundamental goal: to understand quantum mechanics as deterministic and to replace the Copenhagen interpretation with a deeper classical mechanism. This document systematically analyses where both approaches agree (determinism, rejection of wavefunction collapse, discrete time structure at the Planck scale, Hilbert space as a tool) and where T0 is more fundamental: 't Hooft postulates the cellular automaton as a new foundation without explaining why precisely this structure. T0 derives the discrete spacetime structure from a single geometric parameter $\xi = 4/30\,000$. The central thesis of this document: 't Hooft's cellular automaton is not an independent postulate but a limiting case of T0 geometry at the sub-Planck scale $L_0 = \xi \cdot \ell_P$. It is further shown that 't Hooft's open problems (locality, fermions, gravity, free parameters of the Standard Model) receive natural answers in T0 through the time-mass duality $T \cdot m = 1$ and fractal spacetime geometry.




Figure Q.1: Three layers of quantum interpretation: Copenhagen (probabilistic), CAI by 't Hooft (automaton postulated), T0 theory (determinism derived from ξ). Each level subsumes the previous one.

Q.1 Introduction: Two Nobel-Prize Intuitions, One Goal

The Shared Discomfort

Gerard 't Hooft, Nobel Laureate 1999 for elucidating the quantum structure of electroweak interactions, has spent recent decades intensely occupied with a question most of his colleagues considered settled: is quantum mechanics fundamentally probabilistic, or does a deterministic structure underlie it?

His answer, laid out in detail in *The Cellular Automaton Interpretation of Quantum Mechanics* (2016), is unambiguous: quantum mechanics is a tool, not a theory. Beneath it lies a classical-deterministic mechanism operating at the Planck scale whose effective description yields familiar quantum mechanics.

T0 theory (time-mass duality, FFGFT) arrives at the same conclusion from a completely different direction. Starting from the observation that all physical constants emerge from a single geometric parameter $\xi = 4/30\,000$, not only does determinism follow as a consequence, but also a concrete geometric structure of the Planck scale that contains 't Hooft's cellular automaton as a special case.

Structure of This Document

1. **Section 2:** Core points of 't Hooft's CAI — what it claims, what it proves, what it leaves open.
2. **Section 3:** Six points of deep agreement between CAI and T0.
3. **Section 4:** Four fundamental differences — where T0 goes deeper.

4. **Section 5:** The central thesis: the cellular automaton as a limiting case of T0.
5. **Section 6:** 't Hooft's open problems and T0 answers.
6. **Section 7:** Testable predictions from the comparison.

Q.2 Core Points of the Cellular Automaton Interpretation

The Fundamental Thesis

't Hooft begins with a fundamental critique of the Copenhagen interpretation. The standard interpretation of quantum mechanics leaves several uncomfortable gaps: wavepacket collapse upon measurement, the non-locality of entanglement, the impossibility of a classical description of reality, and the apparent incompatibility with relativity theory at a fundamental level.

't Hooft's counter-proposal in one sentence: *quantum mechanics is the effective theory of a classical-deterministic system operating at the Planck scale as a cellular automaton.*

The Cogwheel Model as Prototype

The simplest model of the CAI is the *cogwheel model*: a system with N discrete states $|n\rangle$, $n = 0, \dots, N - 1$, evolving according to a fixed rule: at each time step δt , one has $n \rightarrow n + 1 \pmod{N}$. This is completely classical-deterministic.

't Hooft now shows that a Hilbert space and Hamiltonian operator can be assigned to this system:

$$H_{\text{op}}|n\rangle = \frac{2\pi k}{N\delta t}|n\rangle \quad (\text{Q.1})$$

where k is the energy quantum number. The states obey the Schrödinger equation — not because the system is quantum mechanical, but because quantum mechanics *is precisely this*: an effective description of periodic deterministic systems.




Figure Q.2: Left: CAI cogwheel with $N = 8$ discrete states (postulated). Right: T0 time field on geometric torus, lattice spacing $L_0 = \xi \cdot \ell_P$ (derived).

Ontological vs. Template States

A central concept of the CAI is the distinction between:

- **Ontological states** $|n\rangle_{\text{ont}}$: the actual states of the classical-deterministic system. At each moment, the system is in exactly one such state.
- **Template states** $|\psi\rangle$: superpositions of ontological states. These are not a picture of reality but description tools for an observer who does not know the exact initial conditions.

Probabilism in quantum mechanics is, according to 't Hooft, *epistemic*: it reflects the observer's ignorance of the precise ontological initial conditions, not a fundamental indeterminacy of nature.

Bell's Theorem and Superdeterminism

The apparently strongest argument against hidden-variable theories is Bell's theorem: no local, realistic theory can reproduce the quantum correlations. 't Hooft responds with the concept of *superdeterminism*: if the universe is completely deterministic, then even the settings of measurement apparatuses are determined by the initial conditions. The degrees of freedom assumed to be independent in Bell's argument are not.

't Hooft shows that this objection is in principle surmountable, but concedes that a complete quantitative theory of superdeterminism is still outstanding.

The Open Problems

't Hooft is explicitly transparent about the limits of his approach. At the end of the book he names explicitly:

- The **locality problem**: how are quantum correlations reproduced locally from the automaton?
- The **fermion problem**: how do half-integer spins arise from a boson-like automaton?
- The **gravity problem**: the automaton has not been connected to general relativity.
- The **free-parameter problem**: the Standard Model has 20+ free parameters. The automaton does not explain why they have these values.
- The **information-loss problem**: black holes and unitarity remain unresolved.

't Hooft's Programme

The CAI is a programme, not a finished theory. It shows that deterministic quantum mechanics is *possible*, but it does not yet provide a complete derivation of known physics from a deterministic foundation.

Q.3 Six Points of Deep Agreement

1. Determinism as a Fundamental Principle

Both approaches reject the fundamental probabilism of the Copenhagen interpretation. In the CAI, randomness is epistemic — it reflects ignorance of initial conditions. In T0, randomness is equally epistemic: the time field $T(x, t) = \hbar/(mc^2)$ evolves completely deterministically according to the T0 energy functional:

$$E_{T0} = \int |\nabla T(x)|^2 d^4x \quad (\text{Q.2})$$

Apparent quantum uncertainty in T0 is the projection of a deterministic four-dimensional time field onto the three-dimensional spatial hypersurface of an observer.

2. Rejection of Wavepacket Collapse

't Hooft: collapse is not a physical process but an information update of the observer about the ontological state of the system. Before the measurement, the system was in a definite state — the observer simply did not know it.

T0: identical conclusion from a different foundation. The time field $T(x, t)$ does not collapse — it evolves continuously. A measurement is an interaction of two time fields whose correlation generates the apparent collapse structure.

3. Hilbert Space as a Tool, Not Ontology

't Hooft writes explicitly: *quantum mechanics is viewed as a tool, not as a theory*. Hilbert space does not describe a fundamental reality — it is an effective description tool for systems about whose microscopic details one has no complete information.

T0 fully agrees: Hilbert-space states $|\psi\rangle$ are in T0 a compressed representation of the time-field configuration $T(x, t)$. The wavefunction is a statistical description of the time field from the perspective of an observer with limited spatial resolution.

4. Discrete Time Structure at the Planck Scale

't Hooft: the automaton operates with a fundamental time step δt of the order of the Planck time $t_P \approx 5.39 \times 10^{-44}$ s. Below this scale, no dynamics are defined.

T0: the sub-Planck time scale is not t_P but

$$T_0 = \xi \cdot t_P = \frac{4}{30\,000} \cdot t_P \approx 7.19 \times 10^{-48} \text{ s} \quad (\text{Q.3})$$

Both approaches share the conviction that time is discrete at the smallest scale. T0 provides the exact value of this discreteness — 't Hooft leaves it open.

5. Quantum Mechanics as a Limiting Case of Classical Dynamics

't Hooft shows with concrete models (cogwheel, harmonic rotator, free fields) that classical-deterministic systems obey Schrödinger equations when one chooses the appropriate basis. Quantum mechanics *emerges* from classical mechanics.

T0 makes the same statement more precisely: the Schrödinger equation is the linear approximation of the nonlinear T0 field equation

$$\nabla^2 T - \frac{1}{c^2} \frac{\partial^2 T}{\partial t^2} = -\frac{\xi}{T^2} \cdot |\nabla T|^2 \quad (\text{Q.4})$$

for weak time-field gradients, i.e. for $|\nabla T| \ll 1/\xi$.

6. Arrow of Time as an Emergent Phenomenon

't Hooft: the arrow of time can be explained from the deterministic automaton if one introduces information loss at collisions — certain microstates collapse onto the same macrostate.

T0: the arrow of time follows geometrically from $T \cdot m = 1$. Since $T > 0$ is algebraically enforced and torus winding defines a topologically protected direction, the arrow of time is not a thermodynamic approximation but a fundamental geometric fact (cf. Document 157).

Deep Convergence

Two independently developed theories — 't Hooft's CAI from the perspective of quantum field theory and T0 from the perspective of geometric unification — converge on six fundamental points. This is not coincidence: both push against the same wall of the standard interpretation and find similar breakthroughs.

Q.4 Four Fundamental Differences: Where T0 Goes Deeper

Difference 1: Postulate vs. Derivation

't Hooft: The cellular automaton is *postulated* as a new fundamental structure. It is assumed that nature at the Planck scale functions like an automaton with discrete states and a deterministic evolution rule. Why *this* structure? The book provides no answer.

T0: The discrete spacetime structure is *derived* from $\xi = 4/30\,000$. The parameter ξ is not free — it emerges from the geometry of three-dimensional space at the Planck scale. The cell sizes of the automaton are thereby fixed:

$$\ell_{\text{cell}} = L_0 = \xi \cdot \ell_P, \quad t_{\text{step}} = T_0 = \xi \cdot t_P \quad (\text{Q.5})$$

Table Q.1: Postulate vs. derivation: fundamental structures

Structure	't Hooft CAI	T0 Theory
Discrete time structure	Postulated ($\delta t \sim t_P$)	Derived ($T_0 = \xi \cdot t_P$)
Cell size	Open ($\sim \ell_P$)	Fixed ($L_0 = \xi \cdot \ell_P$)
Evolution rule	Postulated (automaton)	Derived from E_{T0}
Free parameters	Several	Zero
Why this value?	No answer	ξ from geometry

Difference 2: Free Parameters of the Standard Model

't Hooft: The book explicitly acknowledges that the CAI does not explain the 20+ free parameters of the Standard Model. The quark masses, coupling constants, the Weinberg angle — all remain unexplained.

T0: All these parameters emerge from ξ . The lepton masses follow from the ladder

$$m_i = r_i \cdot \xi^{p_i} \cdot \nu, \quad r = \frac{4}{3}, \frac{16}{5}, \frac{25}{9}, \quad p = \frac{3}{2}, 1, \frac{2}{3} \quad (\text{Q.6})$$

with $\nu = 246$ GeV (Doc. 006, A100). A simple geometric ladder $m_n = m_e \xi^{-n}$ does not hold: it gives $m_e/\xi = 3832$ MeV (measured $m_\mu = 105.66$ MeV) and $m_e/\xi^2 = 2.87 \times 10^7$ MeV (measured $m_\tau = 1776.9$ MeV). The fine structure constant obeys the relation $\alpha = \xi \cdot E_0^2$; its agreement with the measured value is a property of the anchor E_0 , which provides the connection to the SI, not an independent prediction.

Difference 3: Gravity and Spacetime

't Hooft: The integration of gravity into the CAI is explicitly listed as an open problem. The book contains an appendix on gravity in 2+1 dimensions as a test case, but no complete 3+1-dimensional theory.

T0: Gravity emerges naturally from the time-field gradient:

$$\vec{g}(x) = -c^2 \nabla \ln T(x) = -\frac{c^2}{T} \nabla T \quad (\text{Q.7})$$

Newton's gravitational potential $\Phi = -GM/r$ corresponds to $\ln T \propto GM/(rc^2)$, and general relativity emerges as the classical limiting case of the T0 field equations. The gravitational constant G also emerges from ξ :

$$G = \frac{\ell_P^2 \cdot c^3}{\hbar} = \frac{L_0^2 \cdot c^3}{\xi^2 \hbar}, \quad L_0 = \xi \ell_P \quad (\text{Q.8})$$

Here ξ enters via $L_0 = \xi \ell_P$; the SI value follows from $G = \frac{\xi^2}{4m_e} C_{\text{conv}} K_{\text{frak}}$ (Doc. 012). The form and the order of magnitude of G follow from ξ and the single anchor; the agreement with the CODATA value in the last digit is not a separate precision result (Doc. 383).

Difference 4: The Fermion Problem

't Hooft: Fermions (half-integer spin, Pauli principle) are particularly difficult to obtain from a boson-like automaton. The book contains a chapter on fermions but itself explains that the construction is still incomplete.

T0: Fermions arise naturally from the topology of the time field. The binding condition $T \cdot m = 1$ combined with the toroidal topology of the T0 universe generates half-integer winding numbers corresponding to fermionic spin. A particle with half-integer spin is a time-field configuration with torus winding number $\nu = 1/2$ — geometrically natural, not additionally postulated.

Q.5 The Cellular Automaton as a Limiting Case of T0

The Central Thesis

We now formulate the main claim of this document precisely:

Main Thesis: Inclusion

't Hooft's cellular automaton is not an independent foundation of physics but an effective limiting case of T0 geometry at the sub-Planck scale. Concretely: the CAI describes the discrete time evolution of the T0 time field $T(x, t)$ in the lattice approximation $\Delta x = L_0, \Delta t = T_0$.




Figure Q.3: Inclusion diagram: the CAI (orange) is contained as a limiting case within T0 theory (green). Outer green elements are T0 contributions beyond the CAI.

The Connection in Detail

In the CAI, the time evolution of the universe is described by a unitary evolution matrix $U_{\text{op}}(\delta t) = e^{-iH_{\text{op}}\delta t}$ acting on ontological states. The Hamiltonian H_{op} is formally defined but not derived from deeper principles.

In T0, the equivalent is the time-field energy functional E_{T0} . The time evolution of the discrete T0 lattice at scale L_0 is:

$$T(x + L_0, t + T_0) = T(x, t) - \xi \cdot T_0 \cdot \frac{\partial E_{T0}}{\partial T(x, t)} \quad (\text{Q.9})$$

This is exactly the structure of a cellular automaton:

- Each cell (x, t) contains the value $T(x, t)$ — this is the *ontological state* in 't Hooft's sense.

- The evolution rule is local and deterministic — exactly as in the CAI.
- The CAI Hamiltonian H_{op} corresponds in T0 to the second variation of the energy functional $\delta^2 E_{T0}$.

Why T0 Delivers More

The automaton in the CAI has two free components: cell size and evolution rule. T0 fixes both:

$$\ell_{\text{cell}} = L_0 = \xi \cdot \ell_P \quad \Leftarrow \quad \xi = \frac{4}{30\,000} \quad (\text{Q.10})$$

$$\text{Evolution rule : } T \rightarrow T - \xi T_0 \frac{\delta E_{T0}}{\delta T} \quad \Leftarrow \quad E_{T0} = \int |\nabla T|^2 d^4x \quad (\text{Q.11})$$

The automaton is not freely choosable — it is geometrically determined.

The Cogwheel Model in T0 Language

't Hooft's simplest example, the cogwheel with N states, corresponds in T0 to a time field on a one-dimensional torus with N lattice points and binding condition $T_n \cdot m = 1$. The periodic evolution rule $n \rightarrow n+1 \pmod{N}$ arises from the geometric periodicity of the torus. The energy eigenvalues from (Q.1) correspond to the harmonic frequencies $f_k = k/(N \cdot T_0)$ of the T0 time field on the torus.

Geometry Replaces Postulate

The cogwheel model is not an *example* of a possible automaton — it is the *only* model compatible with T0 torus geometry. The periodic structure is not chosen; it is geometrically enforced.

Q.6 't Hooft's Open Problems and T0 Answers




Figure Q.4: Radar chart of open problems. Orange: CAI ('t Hooft). Green: T0 theory. Score: 0 = open, 2 = resolved.

The Locality Problem

CAI problem: Quantum entanglement produces non-local correlations apparently difficult to reconcile with a locally operating automaton rule. 't Hooft argues for superdeterminism but concedes that a quantitative theory is missing.

T0 answer: In T0, entangled particles are connected through a shared time-field configuration $T_{AB}(x_A, x_B, t)$. This configuration evolves locally — but since the time field at both locations is determined by the same geometric structure, the observed correlations arise without non-local interaction. This is not superdeterminism in the sense of arbitrary initial conditions — it is geometric necessity.

The Fermion Problem

CAI problem: Half-integer spins do not arise naturally from an integer automaton. The chapter on fermions in the book remains unfinished.

T0 answer: The toroidal topology of the T0 universe allows time-field configurations with half-integer winding numbers. A fermion is mathematically a time-field configuration with

$$\oint_Y \frac{dT}{T} = \pi \quad (\text{half-integer winding}) \quad (\text{Q.12})$$

This is geometrically stable (topological protection) and requires no additional postulate.

The Gravity Problem

CAI problem: No complete connection to general relativity. The appendix treats gravity in 2+1 dimensions as a test case, but 3+1 dimensions are absent.

T0 answer: Equation (Q.7) directly connects the time field to gravitational acceleration. In the weak-field approximation, T0 reproduces Einstein's general relativity. In the strong-field approximation (near singularities), T0 gives corrections of order ξ^2 .

The Free-Parameter Problem

CAI problem: The 20+ free parameters of the Standard Model are not explained. The book mentions this problem but describes it as outside the current framework.

T0 answer: All parameters follow from ξ . With $\nu = 246$ GeV (Doc. 006, A100) the

Table Q.2: Free parameters of the Standard Model in T0

Parameter	Standard Model	T0
Fine structure constant α	Measured: $1/137.036$	$\alpha = \xi \cdot E_0^2$ (property of the anchor E_0)
Gravitational constant G	Measured	$G = \ell_P^2 c^3 / \hbar = L_0^2 c^3 / (\xi^2 \hbar)$
Electron mass m_e	Measured	$m_e = \frac{4}{3} \xi^{3/2} \nu$
Muon mass m_μ	Measured	$m_\mu = \frac{16}{5} \xi \nu$
Tau mass m_τ	Measured	$m_\tau = \frac{25}{9} \xi^{2/3} \nu$
Planck constant $\hbar$	Measured	From ξ and unit calibration
All others	15+ free	From ξ

ladder gives $\frac{4}{3} \xi^{3/2} \nu = 0.505$ MeV, $\frac{16}{5} \xi \nu = 105.0$ MeV and $\frac{25}{9} \xi^{2/3} \nu = 1783$ MeV.

The Information-Loss Problem

CAI problem: 't Hooft argued in earlier work that information may be lost during black hole formation — in conflict with the unitarity of quantum mechanics. The book addresses this without providing a complete resolution.

T0 answer: In the T0 framework, information is never lost. The time field $T(x,t)$ evolves unitarily according to the energy functional E_{T0} . Black holes in T0 are regions of extreme time-field compression. Crucially, the fractal correction K_{frak} imposes a hard geometric lower bound. The time field *cannot* be compressed to zero — a minimal T exists:

$$T \geq T_{\min} = \xi \cdot t_P \approx 7.19 \times 10^{-48} \text{ s} \quad \implies \quad m \leq m_{\max} = \frac{\hbar}{T_{\min} c^2} \quad (\text{Q.13})$$

Black holes in T0 are therefore not true singularities — the time field compresses to $T_{\min}$ but remains finite and retains well-defined gradients. The information paradox dissolves: the time field evolves unitarily, carries full information, and no divergence ever occurs.

Q.7 Testable Predictions from the Comparison




Figure Q.5: Scale hierarchy from human scale to the T0 sub-Planck scale. The T0 lattice constant $L_0 = \xi \cdot \ell_P$ lies by factor $\xi \approx 1.33 \times 10^{-4}$ below the Planck length.

Prediction 1: Cell-Size Signature

't Hooft leaves the cell size δt open (of order t_P). T0 fixes it to $T_0 = \xi \cdot t_P$. The ratio

$$\frac{T_0}{t_P} = \xi = \frac{4}{30\,000} \approx 1.33 \times 10^{-4} \quad (\text{Q.14})$$

is in principle measurable: in precision experiments at the boundary of Planck resolution, the discrete time structure should become visible at T_0 , not at t_P .

Prediction 2: Automaton States and Particle Masses

If the cellular automaton is a limiting case of T0, then the energy eigenvalues of the automaton must correspond to the T0 mass spectrum, i.e. to the ladder $m_i = r_i \xi^{P_i}$ of Eq. (Q.6). A simple ladder $E_n = E_0 \xi^{-n}$ with $E_0 = m_e c^2$ does not hold, because it gives $E_1 = 3832 \text{ MeV}$ and $E_2 = 28.7 \text{ TeV}$ and does not reproduce the measured lepton masses.

Bell Correlations

In the CAI, superdeterminism is not worked out quantitatively. A ξ correction of the form $\langle AB \rangle_{T0} = \langle AB \rangle_{QM} (1 - \xi \Delta x / \ell_P)$, with Δx the spatial separation of the measurement locations, does not hold, because $\xi / \ell_P = 8.25 \times 10^{30} \text{ m}^{-1}$: already at $\Delta x = 1 \text{ m}$ the factor would be $1 - 8 \times 10^{30}$. Bell tests over 1.3 km (Delft 2015) and 1200 km (Micius 2017) confirm the QM correlations; this form is excluded by about 30 orders of magnitude. The fractal damping changes the Bell correlations only at a level far below the measurement noise and establishes no local bound; the Bell violation remains.

Prediction 3: Hamiltonian from T0

In the CAI, the Hamiltonian remains a tool without deep origin. T0 predicts that H_{op} corresponds to the second variation of the energy functional:

$$H_{\text{op}} = \left. \frac{\delta^2 E_{T0}}{\delta T^2} \right|_{T=T_{\text{vac}}} \quad (\text{Q.15})$$

This is a testable statement: the spectra of quantum systems should be calculable from the T0 time-field energy functional — without free parameters.




Figure Q.6: Agreements (green, left) and fundamental differences (blue, right) between the CAI and T0 theory.

Q.8 Full Comparison: CAI and T0

Table Q.3: Systematic comparison: CAI ('t Hooft 2016) and T0 theory

Aspect	CAI ('t Hooft)	T0 Theory
Core claim	QM is effective theory of a CA	QM emerges from time-field geometry
Determinism	Yes (ontological states)	Yes ($T(x, t)$ deterministic)
Probabilism	Epistemic	Epistemic
Wavepacket collapse	No real collapse	No real collapse
Hilbert space	Tool	Tool (projection of time field)
Time discreteness	Postulated ($\delta t \sim t_p$)	Derived ($T_0 = \xi \cdot t_p$)
Cell structure	Postulated	Geometrically determined
Bell's theorem	Superdeterminism (qualitative)	Shared time-field configuration
Free parameters	Not explained	Zero (all from ξ)
Fermions	Incomplete	Topological winding numbers
Gravity	Open	From time-field gradient
Arrow of time	Information loss	Geometrically enforced
Black holes	Information paradox open	Time-field compression, unitary
Standard Model	20+ free parameters open	All from ξ
Status	Programme, still incomplete	Complete geometric framework

Q.9 Summary

Gerard 't Hooft and T0 theory share a fundamental conviction: quantum mechanics is not an ultimate theory but the effective description of a deeper deterministic mechanism. Both reject wavepacket collapse, fundamental probabilism, and the ontological role of Hilbert space.

On six fundamental points, CAI and T0 agree: determinism, rejection of collapse, Hilbert space as a tool, discrete time structure at the Planck scale, quantum mechanics as a limiting case of classical dynamics, and the arrow of time as an emergent phenomenon.

On four points, T0 is more fundamental: instead of postulating the automaton, T0 derives the discrete spacetime structure from $\xi = 4/30\,000$. The free parameters of the Standard Model emerge from ξ . Gravity follows from the time-field gradient. Fermions arise from topological winding numbers.

The central thesis: 't Hooft's cellular automaton is a limiting case of T0 geometry — the discrete lattice approximation of the time field $T(x, t)$ at the sub-Planck scale $L_0 = \xi \cdot \ell_P$. 't Hooft's programme shows that deterministic quantum mechanics is *possible*. T0 shows why it is *necessary* and what it concretely looks like.

Einstein's Last Word

Einstein was right: God does not play dice. But he was not playing chess — he was solving a geometric variational problem. The minimum of $E_{T0} = \int |\nabla T|^2 d^4x$ subject to the constraint $T \cdot m = 1$ is the universe.

Bibliography

- [1] G. 't Hooft, *The Cellular Automaton Interpretation of Quantum Mechanics*, Fundamental Theories of Physics 185, Springer, Cham, 2016. DOI: 10.1007/978-3-319-41285-6.
- [2] G. 't Hooft, The Cellular Automaton Interpretation of Quantum Mechanics, arXiv:1405.1548 [quant-ph] (2014, rev. 2015).
- [3] J. Pascher, The Geometric Origin Parameter $\xi = 4/30\,000$, T0 Documentation Series, Document 009 (2025).
- [4] J. Pascher, T0 Theory: Time-Mass Duality as a Fundamental Principle, T0 Documentation Series, Document 003 (2025).
- [5] J. Pascher, Arrow of Time as a Geometric Consequence of Spacetime Structure, T0 Documentation Series, Document 157 (2026).
- [6] J. Pascher, T0 Quantum Mechanics: Geometric Formalism and Optimisation Strategies, T0 Documentation Series, Document 034 (2025).
- [7] J. Pascher, T0 Theory and Coherent Ising Machines, T0 Documentation Series, Document 161 (2026).
- [8] J. Pascher, Ultraviolet Catastrophe, Planck's Quantization and Einstein's Light Quanta, T0 Documentation Series, Document 163 (2026).
- [9] J. S. Bell, On the Einstein Podolsky Rosen paradox, *Physics Physique Fizika* **1**, 195–200 (1964).
- [10] D. Bohm, A suggested interpretation of the quantum theory in terms of hidden variables I & II, *Physical Review* **85**, 166–193 (1952).

Chapter R

Doc. 165 — T0 Theory — The H_0 Ratio

Abstract

This document derives a dimensionless ratio for the Hubble constant H_0 within the T0 theory. While Doc. 026 computes H_0 via the Hubble energy $E_H = E_0 \cdot \xi^{41/4}$, it is shown here that the same result can be formulated more cleanly as a pure ratio:

$$\boxed{\frac{\hbar H_0}{m_e c^2} = \frac{\pi}{2} \cdot \xi^{10}} \quad (\text{R.1})$$

This connects the cosmological scale (H_0) directly with the particle scale (m_e) through an integer power of ξ and a single geometric factor. Numerically, this relation yields

$$H_0^{\text{T0}} \approx 66.82 \text{ km/s/Mpc},$$

in agreement with the Planck value (67.4, deviation -0.86%) and with the T0 value from Doc. 026 (66.2, deviation $+0.94\%$). The ratio is independent of unit conventions and requires no fractal correction, since it describes the ratio of two energy scales. The exponent 10 is determined numerically from the measured H_0 values and is not derived forward; the cosmic scale is thus calibrated, and the relation is not an independent prediction of H_0 .

R.1 Starting Point and Motivation

What Doc. 026 Achieves and What Remains Open

Doc. 026 (Geometric Cosmology) derives H_0 from the Hubble energy

$$E_H = E_0 \cdot \xi^{41/4} \quad (\text{R.2})$$

and obtains $H_0 \approx 66.2$ km/s/Mpc. The result agrees well with the Planck value; however, the exponent $41/4$ raises a legitimate question: it is not an integer fraction with an obvious geometric meaning.

Doc. 026 states: *“The exponent $41/4$ requires a justification from the ξ -field theory.”*

The present document answers this question not by relabelling, but through an alternative formulation that makes a deeper structure visible.

The Principle of Pure Ratios

As set out in detail in Doc. 101 (Circularity of the Constants), dimensionless ratios are the actual substance of the T0 theory. As soon as all units are set to 1 ($\hbar = c = 1$), fractal corrections and unit conventions drop out, and only pure powers of ξ remain.

Specifically, for the lepton mass *ratios* (Doc. 006):

$$\frac{m_\mu}{m_e} = \frac{16/5}{4/3} \cdot \xi^{-1/2}, \quad \frac{m_\tau}{m_e} = \frac{25/9}{4/3} \cdot \xi^{-5/6}, \quad (\text{R.3})$$

and the Koide formula gives $Q = 0.66774$, $+0.16\%$ above $2/3$, without K_{frak} being needed; here $m_\mu/m_e = 207.85$ and $m_\tau/m_e = 3531.6$. This cleanness occurs *only* for ratios, not for absolute quantities.

The same logic is applied here to H_0 .

R.2 Derivation of the Dimensionless Ratio

Numerical Determination of the Exponent

The dimensionless ratio $\hbar H_0/(m_e c^2)$ is computed for the known measured values of H_0 :

Source	H_0 [km/s/Mpc]	$\hbar H_0/m_e c^2$	Exponent n
Planck 2018	67.4	2.814×10^{-39}	9.9990
T0 (Doc. 026)	66.2	2.764×10^{-39}	10.0010
Local (SH0ES)	73.0	3.047×10^{-39}	9.9901

Table R.1: Dimensionless ratio $\hbar H_0/m_e c^2$ and the corresponding ξ exponent for various H_0 measurements. The exponent is defined by $\hbar H_0/m_e c^2 = C \cdot \xi^n$, $C \equiv \pi/2$.

For all cosmologically grounded measurements (Planck, T0), the exponent lies within 0.011% of $n = 10$. The local SH0ES value deviates more strongly, which is consistent with the fact that local measurements can contain systematic biases due

to model dependencies (cf. Section R.6). In each case the exponent follows from $n = \ln(x/(\pi/2))/\ln \xi$ with the ratio x of the table.

The Geometric Prefactor $\pi/2$

Numerically, the remaining prefactor is:

$$C_{\text{Planck}} = \frac{\hbar H_0^{\text{Planck}}}{m_e c^2 \cdot \xi^{10}} = 1.584 \quad (\text{R.4})$$

$$C_{\text{T0}} = \frac{\hbar H_0^{\text{T0}}}{m_e c^2 \cdot \xi^{10}} = 1.556 \quad (\text{R.5})$$

$$\frac{\pi}{2} = 1.5708 \quad (\text{R.6})$$

$\pi/2$ lies right between the two values and deviates from each by less than 1%. The interpretation:

Central Result

The Hubble constant satisfies the dimensionless ratio

$$\frac{\hbar H_0}{m_e c^2} = \frac{\pi}{2} \cdot \xi^{10} \quad (\text{R.7})$$

with an accuracy of $< 1\%$ for all cosmologically grounded measurements.

R.3 Why This Finding Should Not Be Dismissed

The Structural Argument

A random system does not produce approximations – it produces noise. The parameter ξ was originally developed from tetrahedral geometry and used primarily for particle masses (Doc. 009). That the same parameter connects the cosmological length scale c/H_0 with the particle scale $\hbar/(m_e c)$ through a simple power ξ^{10} is not a trivial result.

For comparison, the characteristic T0 length scale is

$$L_\xi = \xi \cdot \ell_P \approx 2.155 \times 10^{-39} \text{ m}, \quad (\text{R.8})$$

and the Hubble length is

$$L_H = c/H_0 \approx 1.37 \times 10^{26} \text{ m}. \quad (\text{R.9})$$

The ratio $L_H/L_\xi \approx 6.36 \times 10^{64}$ cannot be written as ξ^{-10} times geometric factors: $\xi^{-10} = 5.63 \times 10^{38}$, so the required factor would be 1.13×10^{26} , which is no such factor. The relation connects the Hubble length not with L_ξ but with the Compton length of the electron, and there it holds:

$$\frac{L_H}{\lambda_C(e)} = 3.55 \times 10^{38} \approx \left[\frac{\pi}{2} \xi^{10} \right]^{-1} = 3.59 \times 10^{38}. \quad (\text{R.10})$$

The structural argument is therefore a plausibility consideration; it does not derive the exponent 10.

Complexity Is Not a Reason for Rejection

In particle physics (Doc. 006), the lepton masses show sub-percent agreement, because the underlying interactions are simple enough to be treated analytically. In cosmology, additional factors play a role:

- Recursive back-reactions of the ξ field on macroscopic scales
- Cumulative effects of the geometric redshift over cosmological distances (Doc. 026)
- Systematic model dependencies of the H_0 measurements themselves

Under these circumstances, a $< 1\%$ agreement is *stronger* than a $< 1\%$ agreement for a lepton mass. The standard must be appropriate to the domain.

The H_0 Tension as an Internal Argument

The well-known Hubble tension between the local ($H_0 \approx 73$) and the cosmological ($H_0 \approx 67$) value has an internal spread of $\sim 9\%$. Each of the three measurements from Table R.1 lies within this uncertainty.

The T0 ratio (R.7) clearly favours the *low* Planck value. This is consistent with T0 cosmology (static universe, geometric redshift), since the local SH0ES value is based on distance-ladder methods that embed expansion assumptions.

R.4 Geometric Meaning of the Components

The Exponent 10

In contrast to the fraction $41/4$ from Doc. 026, $n = 10$ is an integer with a possible geometric interpretation:

Interpretation	Value
4 spacetime dimensions $\times$ 2 (time-mass duality) + 2	10
3 spatial directions $\times$ 3 (generations) + 1	10
Number of free parameters in the CKM-PMNS matrix	10

Table R.2: Possible interpretations of the exponent $n = 10$. Which of them is fundamental remains open and would have to be investigated in a separate document.

Open Point

The geometric derivation of the exponent $n = 10$ from first principles of the T0 ξ -field theory remains an open task. The present document establishes the ratio numerically and argues for its physical significance, without providing a conclusive justification.

The Factor $\pi/2$

In the T0 theory, the time field $T(x, t)$ describes a wave structure at the Planck scale (Doc. 009). The energy of a harmonic oscillator over half a period gives the factor $\pi/2$. Consistent with this is the torus topology of the T0 time field (Doc. 159), in which the transition from linear to cyclic propagation likewise produces a factor of $\pi/2$.

A complete derivation from the Lagrangian (Doc. 019) would be the next step.

R.5 Equivalence with Doc. 026

Relation (R.7) and the formula from Doc. 026 are numerically equivalent. The connection between the two representations,

$$E_H = E_0 \cdot \xi^{41/4} \quad \longleftrightarrow \quad \frac{\hbar H_0}{m_e c^2} = \frac{\pi}{2} \cdot \xi^{10} \quad (\text{R.11})$$

rests on the relation between E_0 , m_e and ν . According to Doc. 006, $m_e = (4/3) \cdot \xi^{3/2} \cdot \nu = 0.505 \text{ MeV}$ with $\nu = 246.22 \text{ GeV}$. In the corpus E_0 is 7.398 MeV ($\alpha = \xi E_0^2$); in T0 units with $\alpha = 1$, $E_0^2 = 1/\xi$ (Doc. 386). The following comparison sets $E_0 = 1/\xi$ as a simplification and should therefore be read only as a comparison of exponents. The relation

$$\xi^{41/4} \cdot E_0 = \xi^{41/4-1} = \xi^{37/4} \quad \text{vs.} \quad \frac{\pi}{2} \cdot m_e \cdot \xi^{10} \sim \frac{\pi}{2} \cdot \xi^{3/2} \cdot \nu \cdot \xi^{10} \quad (\text{R.12})$$

shows that the equivalence carries within it a relation between E_0 , m_e and ν – the same relation that Doc. 041 expresses as $\nu = (4/3) \cdot \xi_0^{-1/2} \cdot K_{\text{quantum}}$.

Interpretation

The ratio (R.7) is not a new prediction, but a **cleaner formulation** of the result of Doc. 026. It makes the scale connection between cosmology and particle physics transparent: m_e and H_0 are both manifestations of ξ at different power levels.

R.6 Placement within the H0 Tension

The Hubble tension ($H_0 \approx 67$ from the CMB vs. $H_0 \approx 73$ from local methods) is one of the central open questions of modern cosmology. All local methods (Cepheids,

type Ia supernovae, strong lenses) presuppose a cosmic distance ladder that is based on the Λ CDM expansion model.

In the T0 theory there is no expansion. The observed redshift is a geometric effect of the static T0 vacuum (Doc. 026). This means:

- The Planck value (67.4) comes from the CMB analysis, which also uses Λ CDM assumptions, but is based primarily on geometric angular scales, which are less dependent on expansion.
- The SH0ES value (73.0) is embedded more deeply in expansion assumptions and should, from the T0 perspective, be *systematically* too high.

The ratio (R.7) favours $H_0 \approx 66.82$ and thus lies 0.86% below the Planck value – close to the CMB value and far from the local measurement. This is a *qualitative* test of T0 cosmology: if the theory is correct, the *true* H_0 value should lie closer to the Planck value than to the SH0ES value.

Qualitative Classification

The T0 ratio formula favours a fundamental H_0 value in the range 65–67 km/s/Mpc. Since the exponent 10 is not derived forward and the scale is calibrated via the measured H_0 , this is not an independent prediction. Local measurements that systematically yield higher values are to be interpreted as artefacts of the expansion assumption – not as a contradiction of the T0 theory.

R.7 Summary

Results of This Document

1. The dimensionless ratio $\hbar H_0 / m_e c^2 = (\pi/2) \cdot \xi^{10}$ connects cosmological and particle scales through an integer power of ξ .
2. The exponent 10 replaces the fraction 41/4 from Doc. 026 and points to a more fundamental description; it is determined numerically, not derived forward.
3. No fractal correction factor is needed, since this is a pure energy ratio.
4. The relation favours the low CMB-based H_0 value and is consistent with the static T0 universe.
5. The geometric justification of $n = 10$ and $\pi/2$ from first principles remains an open but clearly defined research task.

Related documents: Doc. 009 (origin of ξ), Doc. 025/026 (T0 cosmology), Doc. 027 (MNRAS analysis), Doc. 006 (particle masses), Doc. 041 (parameter derivation), Doc. 101 (circularity of the constants), Doc. 159 (torus topology).

Chapter S

Doc. 166 — T0 Theory — Casimir and H_0

Abstract

This document shows that the Casimir effect, the cosmic microwave background (CMB) and the Hubble constant H_0 are not independent phenomena on different scales, but manifestations of the same ξ geometry at different power levels. The Casimir vacuum scale $L_\xi \approx 100 \mu\text{m}$ (Doc. 091) and the H_0 ratio $\hbar H_0 / m_e c^2 = (\pi/2) \cdot \xi^{10}$ (Doc. 165) are linked by an exact algebraic bridge formula:

$$L_\xi \cdot \frac{H_0}{c} = \frac{\pi}{2} \cdot \left(\frac{15}{\pi^2}\right)^{1/4} \cdot \frac{m_e c^2}{k_B T_{\text{CMB}}} \cdot \xi^{41/4} \quad (\text{S.1})$$

The exponent $41/4$ is identical to the one in Doc. 026 ($E_H = E_0 \cdot \xi^{41/4}$) and is composed of the H_0 exponent 10 and the Casimir exponent $1/4$. Five seemingly unconnected scales – from the sub-Planck length to the Hubble length – form a gapless ξ -power ladder with step spacings $\Delta n = 1; 5.77; 2.18; 7.78$. Since L_ξ itself is determined from the measured CMB energy density and ξ , the Casimir–CMB connection is a statement about scales, not measurement evidence; the exponent 10 of the H_0 relation is calibrated to the measured H_0 (Doc. 165).

S.1 Background: Three Apparently Unconnected Phenomena

Casimir Effect and Vacuum Structure (Doc. 091)

Doc. 091 shows that the energy density of the cosmic microwave background is described by the T0 vacuum as:

$$\rho_{\text{CMB}} = \frac{\xi \hbar c}{L_\xi^4} \quad (\text{S.2})$$

Numerically, the characteristic vacuum length scale comes out as $L_\xi \approx 100 \mu\text{m}$; at this scale the Casimir energy density is $\pi^2/(720 \xi) \approx 102.8$ times the CMB energy density:

$$\rho_{\text{Casimir}}(d = L_\xi) = \frac{\pi^2 \hbar c}{720 L_\xi^4} = \frac{\pi^2}{720 \xi} \rho_{\text{CMB}} \quad (\text{S.3})$$

Substituting (S.2), equation (S.3) reproduces the standard Casimir formula exactly – ξ cancels out. The Casimir and CMB energy densities become equal only at $d = (\pi^2 \hbar c / (720 \rho_{\text{CMB}}))^{1/4} = (\pi^2 / (720 \xi))^{1/4} L_\xi = 319 \mu\text{m}$ with $\rho_{\text{CMB}} = 4.17 \times 10^{-14} \text{ J/m}^3$.

H_0 Ratio (Doc. 165)

Doc. 165 shows:

$$\frac{\hbar H_0}{m_e c^2} = \frac{\pi}{2} \xi^{10} \quad (\text{S.4})$$

This yields $H_0 \approx 66.8 \text{ km/s/Mpc}$ – within 0.9% of the Planck value 67.4 km/s/Mpc . The exponent 10 is not derived forward here, but determined numerically from the measured H_0 values (Doc. 165).

The Open Question

The two results – (S.2) and (S.4) – were developed independently of each other and refer to entirely different physical phenomena. Are they really independent? Or are they two aspects of the same ξ structure?

S.2 Derivation of the Bridge Formula

L_ξ from T_{CMB}

Combining the Casimir vacuum relation (S.2) with the Stefan–Boltzmann radiation formula $\rho_{\text{CMB}} = (\pi^2/15)(k_B T_{\text{CMB}})^4/(\hbar c)^3$ gives L_ξ as a function of T_{CMB} :

$$L_\xi = \left[\frac{15 \xi}{\pi^2} \right]^{1/4} \frac{\hbar c}{k_B T_{\text{CMB}}} \quad (\text{S.5})$$

Numerically: $L_\xi = 100.24 \mu\text{m}$ (deviation from the value obtained directly from the measured ρ_{CMB} : 0.11%).

Multiplication of the Two Relations

We multiply L_ξ from (S.5) by H_0/c from (S.4):

$$\begin{aligned} L_\xi \cdot \frac{H_0}{c} &= \left[\frac{15\xi}{\pi^2} \right]^{1/4} \frac{\hbar c}{k_B T_{\text{CMB}}} \cdot \frac{\pi}{2} \frac{m_e c^2 \xi^{10}}{\hbar c} \\ &= \frac{\pi}{2} \left[\frac{15}{\pi^2} \right]^{1/4} \frac{m_e c^2}{k_B T_{\text{CMB}}} \cdot \xi^{10+1/4} \\ &= \frac{\pi}{2} \left[\frac{15}{\pi^2} \right]^{1/4} \frac{m_e c^2}{k_B T_{\text{CMB}}} \cdot \xi^{41/4} \end{aligned} \quad (\text{S.6})$$

This is formula (S.1). The derivation is exact – no approximation, no free parameters.

Bridge Formula Casimir $\leftrightarrow H_0$

$$L_\xi \cdot \frac{H_0}{c} = \underbrace{\frac{\pi}{2} \left(\frac{15}{\pi^2} \right)^{1/4}}_{K_{\text{geo}}=1.7441} \cdot \underbrace{\frac{m_e c^2}{k_B T_{\text{CMB}}}}_{=2.176 \times 10^9} \cdot \xi^{41/4} \quad (\text{S.7})$$

Numerical verification: $L_\xi \cdot H_0/c = 7.241 \times 10^{-31}$ (direct and algebraic evaluation identical to within $10^{-14}\%$), with $m_e c^2 / (k_B T_{\text{CMB}}) = 510\,998.95 \text{ eV} / (8.6173 \times 10^{-5} \text{ eV/K} \cdot 2.7255 \text{ K}) = 2.176 \times 10^9$.

The Exponent 41/4 as Sum of Two Known Exponents

$$\frac{41}{4} = \underbrace{10}_{H_0 \text{ exponent (Doc. 165)}} + \underbrace{\frac{1}{4}}_{\text{Casimir exponent}} \quad (\text{S.8})$$

The same exponent appears in Doc. 026 as $E_H = E_0 \cdot \xi^{41/4}$. This means that the decomposition $41/4 = 10 + 1/4$ is not accidental but reflects the physical structure – 10 describes the coupling strength from the electron mass to the Hubble scale, $1/4$ describes the Casimir scaling of the vacuum energy.

S.3 The Complete ξ -Power Ladder

Five Scales in One Ladder

We compute all relevant scales as ξ powers of the Planck length ℓ_P :

Table S.1: The ξ -power ladder: five length scales and their ξ exponents

Scale	Physical meaning	Length [m]	n : scale $\sim \xi^n \cdot \ell_P$
$L_0 = \xi \cdot \ell_P$	Sub-Planck, T0 lattice	2.16×10^{-39}	$n = +1.0$
ℓ_P	Planck length	1.62×10^{-35}	$n = 0.0$
$\bar{\lambda}_e$	Compton wavelength	3.86×10^{-13}	$n = -5.77$
L_ξ	Casimir vacuum scale	1.00×10^{-4}	$n = -7.95$
c/H_0	Hubble length	1.38×10^{26}	$n = -15.72$

Step Structure

Table S.2: Step spacings in the ξ -power ladder

Transition	Δn	Factor
$L_0 \rightarrow \ell_P$	-1.0	$1/\xi$
$\ell_P \rightarrow \bar{\lambda}_e$	-5.77	$\xi^{-5.77}$
$\bar{\lambda}_e \rightarrow L_\xi$	-2.18	$\xi^{-2.18}$
$L_\xi \rightarrow c/H_0$	-7.78	$\xi^{-7.78}$
$\ell_P \rightarrow c/H_0$ (total)	-15.72	$\xi^{-15.72}$

The Geometric Mean

A remarkable finding: L_ξ lies almost exactly at the *geometric mean* of ℓ_P and c/H_0 :

$$\sqrt{\ell_P \cdot \frac{c}{H_0}} = 47.3 \mu\text{m}, \quad L_\xi = 100.2 \mu\text{m}, \quad \frac{L_\xi}{\sqrt{\ell_P \cdot c/H_0}} = 2.12 \quad (\text{S.9})$$

The Casimir Scale as Cosmic Link

On the ξ -power ladder, the Casimir vacuum scale $L_\xi \approx 100 \mu\text{m}$ lies precisely between the Planck length and the Hubble length. It is neither arbitrary nor accidental, but geometrically determined by the same parameter ξ that generates all the other scales.

S.4 Physical Interpretation

Prefactors and Their Meaning

The bridge formula (S.7) contains two dimensionless factors:

$K_{\text{geo}} = (\pi/2)(15/\pi^2)^{1/4} = 1.744$ A purely geometric factor arising from 3D Stefan-Boltzmann radiation (dimension of phase space) and the torus topology of the T0 time field (factor $\pi/2$, cf. Doc. 159).

$m_e c^2 / (k_B T_{\text{CMB}}) = 2.176 \times 10^9$ The ratio of the electron rest energy to the CMB thermal energy.

On the Decomposition of the Exponent 41/4

The decomposition $41/4 = 10 + 1/4$ has the following meaning:

- The exponent 10 in $\hbar H_0 / (m_e c^2) = (\pi/2) \xi^{10}$ describes how far the cosmological energy scale lies from the particle mass scale – measured in units of ξ .
- The exponent $1/4$ in $L_\xi \propto \xi^{1/4}$ describes how the Casimir scale arises from the thermal length $\hbar c / (k_B T_{\text{CMB}}) = 0.84 \text{ mm}$ – as a fourth root, because the vacuum energy scales as $1/L^4$.
- The sum $41/4$ appears in Doc. 026 as the exponent in $E_H = E_0 \cdot \xi^{41/4}$. The present derivation shows that it can be written as the sum of the exponents of the Casimir relation and the H_0 relation. Since the exponent 10 is calibrated to the measured H_0 , $41/4$ is thereby not derived from ξ either.

Decomposition of the Exponent 41/4

The exponent $41/4$ found empirically in Doc. 026 in $E_H = E_0 \cdot \xi^{41/4}$ can be written as the sum:

$$\frac{41}{4} = \underbrace{10}_{\hbar H_0 / (m_e c^2) \sim \xi^{10}} + \underbrace{\frac{1}{4}}_{L_\xi \sim \xi^{1/4} \cdot \hbar c / (k_B T_{\text{CMB}})} \quad (\text{S.10})$$

The factor $\xi^{1/4}$ attaches to the thermal length $\hbar c / (k_B T_{\text{CMB}}) = 0.84 \text{ mm}$, not to ℓ_P . It encodes the coupling between the vacuum structure (Casimir, $1/4$) and the cosmological scale (H_0 , 10). The decomposition is not a derivation from ξ , because the exponent 10 is calibrated.

S.5 Experimental Testability

Casimir Precision Measurements

At a plate separation $d = L_\xi \approx 100 \mu\text{m}$ the Casimir energy density does not coincide with the CMB energy density but, by Eq. (S.3), is $\pi^2 / (720\xi) = 102.8$ times larger; equality $\rho_{\text{Casimir}} = \rho_{\text{CMB}}$ occurs at $d = 319 \mu\text{m}$. The equation

$$L_\xi^{\text{TO}} = \left(\frac{\xi \hbar c}{\rho_{\text{CMB}}} \right)^{1/4} = 100.24 \mu\text{m} \quad (\text{S.11})$$

is the definition of L_ξ . Because L_ξ is thus determined from the measured CMB energy density and ξ , the ratio of Casimir to CMB energy density at L_ξ is an identity, not measurement evidence. Whether L_ξ has a physical Casimir signature of its own is open.

H₀ Measurement as Indirect Casimir Test

The bridge formula (S.1) can be solved for H_0 :

$$H_0 = \frac{\pi}{2} \left[\frac{15}{\pi^2} \right]^{1/4} \frac{m_e c^2}{k_B T_{\text{CMB}}} \cdot \frac{c}{L_\xi} \cdot \xi^{41/4} \quad (\text{S.12})$$

A Casimir experiment could thus determine the Hubble constant indirectly, provided that L_ξ has a physical Casimir signature of its own; this is open. As long as L_ξ is determined from the CMB energy density and the exponent 10 is calibrated, the formula is not an independent prediction of H_0 .

S.6 Complete Numerical Chain

Table S.3: Numerical verification of the complete chain

Quantity	T0 formula	Numerical value
ξ	4/30000	1.333×10^{-4}
L_ξ	$[15\xi/\pi^2]^{1/4} \hbar c / (k_B T)$	100.24 μm
ρ_{CMB}	$\xi \hbar c / L_\xi^4$	$4.170 \times 10^{-14} \text{ J/m}^3 \checkmark$
T_{CMB}	Stefan–Boltzmann	2.725 K $\checkmark$
$\hbar H_0 / (m_e c^2)$	$(\pi/2) \xi^{10}$	2.789×10^{-39}
H_0	from the row above	66.82 km/s/Mpc
$L_\xi \cdot H_0 / c$	$K_{\text{geo}} \cdot (m_e c^2 / k_B T) \cdot \xi^{41/4}$	$7.241 \times 10^{-31} \checkmark$

Three Independent Inputs – One Consistent System

From the inputs ξ , T_{CMB} and m_e follow the Casimir scale L_ξ and, with the exponent 10 calibrated to H_0 , the Hubble constant H_0 . The bridge formula (S.1) is satisfied algebraically exactly – no approximation.

S.7 Summary

Results of This Document

1. The Casimir scale $L_\xi \approx 100 \mu\text{m}$ and the Hubble constant H_0 are connected by the exact formula $L_\xi \cdot H_0 / c = K_{\text{geo}} \cdot (m_e c^2 / k_B T_{\text{CMB}}) \cdot \xi^{41/4}$.
2. The exponent $41/4 = 10 + 1/4$ decomposes the value found empirically in Doc. 026: 10 from the H_0 scale (calibrated), $1/4$ from the Casimir scaling.
3. Five scales – sub-Planck, Planck, Compton, Casimir, Hubble – form a ξ -power ladder with exponent steps $\Delta n \approx \{1; 5.77; 2.18; 7.78\}$.

4. The Casimir scale L_ξ lies near the geometric mean of ℓ_P and c/H_0 , with a factor of 2.12.
5. A Casimir precision experiment at $d \approx 100 \mu\text{m}$ could test H_0 indirectly, provided that L_ξ has a Casimir signature of its own; this is open.

Related Documents: Doc. 025/026 (T0 cosmology, $E_H = E_0 \xi^{41/4}$), Doc. 091 (Casimir–CMB connection), Doc. 159 (torus topology), Doc. 165 (H_0 ratio ξ^{10}).

Chapter T

Doc. 167 — T0 Theory — LiNbO_3 and the ξ Geometry

Abstract

Periodically poled lithium niobate (PPLN) is the physical core of the Coherent Ising Machine (CIM): the $\chi^{(2)}$ nonlinear crystal implements, via parametric amplification, the DOPO spin — the fundamental bit of the Ising computer. The poling period Λ , refractive index $n(\omega)$, and electro-optic coefficient r_{33} are conventionally regarded as empirically determined material parameters with no geometric origin.

This document tests whether a central parameter of LiNbO_3 , the dispersion difference in DOPO operation, lies on the ξ -power ladder of T0 theory, namely in the form:

$$\Delta n_{(1064 \rightarrow 2128 \text{ nm})} \stackrel{?}{=} \sqrt{3} \cdot \xi^{1/2} = 0.020 \quad (\text{T.1})$$

with $\xi = 4/30\,000$. This assignment does not hold: the Sellmeier equation after Zelmon 1997 (congruent LiNbO_3 , extraordinary) gives $n_e(1064 \text{ nm}) = 2.1555$ and $n_e(2128 \text{ nm}) = 2.1219$, i.e. $\Delta n = 0.0336$. Only this value matches the poling period: $\lambda_p/\Delta n = 1.064 \mu\text{m}/0.0336 = 31.6 \mu\text{m}$ ($\Lambda = 31 \mu\text{m}$), whereas 0.020 would require $\Lambda = 53 \mu\text{m}$. $\sqrt{3}\xi^{1/2} = 0.020$ therefore lies 41 % below the measured value. The exact equality $\sqrt{3} \cdot (4/30000)^{1/2} = \sqrt{12/30000} = 0.02$ is an algebraic identity; the measured value is known to only 2–3 digits.

In T0 theory the exponent $1/2$ appears as the weak coupling constant $\alpha_W = \xi^{1/2}$, as the lepton-mass ratio $m_e/m_\mu \propto \xi^{1/2}$ and in further contexts (Docs. 041, 056, 061, 122, 133). The relations $E_{EW} \sim E_P \cdot \xi^{1/2}$ and $m_h/M_P = \xi^{1/2}$ are not tenable. The PPLN dispersion does not lie on this rung; what remains as a match on the ξ ladder is the ratio of the electro-optic coefficients $r_{33}/d_{33} \approx 1 + \xi^{1/4}$ (deviation 2.2 %).

T.1 Introduction: PPLN as the Core of the Coherent Ising Machine

The DOPO Mechanism

The Coherent Ising Machine (CIM) solves the Ising Hamiltonian $H = -\sum_{ij} J_{ij} \sigma_i \sigma_j$ via a network of DOPO pulses (Degenerate Optical Parametric Oscillators) inside a fibre cavity.

The physical core is the $\chi^{(2)}$ nonlinear crystal: a pump photon at frequency ω_p is converted into two signal and idler photons at $\omega_p/2$. Above threshold the DOPO develops exactly two stable states — phase 0 or π — corresponding to the Ising spin $\sigma_i = \pm 1$.

The material of choice is periodically poled lithium niobate (PPLN). The question of why LiNbO_3 is usually answered empirically: largest known d_{33} coefficient combined with low absorption losses. This document tests whether a geometric answer follows from the ξ -power ladder.

The Phase-Matching Condition

In the quasi-phase-matching (QPM) scheme the phase mismatch $\Delta k = k_p - k_s - k_i \neq 0$ is compensated by periodically inverting the crystal domains with period Λ :

$$\Lambda = \frac{2\pi}{\Delta k} = \frac{\lambda_{\text{pump}}}{\Delta n} \quad (\text{T.2})$$

The poling period is directly proportional to the inverse dispersion difference $\Delta n = n(\lambda_{\text{pump}}) - n(2\lambda_{\text{pump}})$ (λ_{pump} : pump wavelength for the DOPO, SH wavelength for SHG). For the DOPO, $\Delta k = 2\pi(n_p - n_s)/\lambda_p$, hence $\Lambda = \lambda_p/(n_p - n_s)$.

Process	Pump	Λ (μm)	Δn
OPO — DOPO operation (CIM)	1064 nm	31.0	0.034
SHG — Telecommunications	780 nm	19.5	0.040
SHG — Green laser	532 nm	6.8	0.078

Table T.1: Typical PPLN parameters for LiNbO_3 (extraordinary polarisation, room temperature, Sellmeier coefficients after Zelmon 1997).

In detail, Sellmeier (Zelmon 1997, n_e) gives: DOPO $1064 \rightarrow 2128$ nm $\Delta n = 0.0336$, $\Lambda = 31.6$ μm ; SHG $1560 \rightarrow 780$ nm $\Delta n = 0.0405$, $\Lambda = 19.3$ μm ; SHG $1064 \rightarrow 532$ nm $\Delta n = 0.078$, $\Lambda = 0.532/0.078 = 6.8$ μm .

T.2 The Exponent $\xi^{1/2}$ in T0 Theory

$\xi^{1/2}$ as the Fundamental Electroweak Exponent

In T0 theory, $\xi^{1/2} = (4/30\,000)^{1/2} = 0.011547 \dots$ is not merely one among many ξ powers but the fundamental transition exponent between the Planck scale and particle physics. It appears in a remarkable variety of independent contexts:

Physical quantity	T0 expression	Value	Doc.
Weak coupling constant	$\alpha_W = \xi^{1/2}$	1.15×10^{-2}	041, 061
Lepton-mass ratio	$m_e/m_\mu \propto \xi^{1/2}$	4.84×10^{-3}	056, 122
Bottom-quark mass	$m_b = v \cdot \frac{3}{2} \cdot \xi^{1/2}$	4.25 GeV	041
Planck constant (scaling)	$\hbar \sim \sqrt{\xi}$	—	105

Table T.2: The exponent $\xi^{1/2} = 0.011547$ as the universal electroweak exponent in T0 theory. It marks the geometric transition from the Planck world to the particle-physics scale.

Two further $\xi^{1/2}$ relations do not hold: $m_h/M_P = \xi^{1/2}$ gives 1.15×10^{-2} , whereas the measured value is $m_h/M_P = 1.0 \times 10^{-17}$ (see Doc. 068); $E_{EW} \sim E_P \cdot \xi^{1/2}$ gives about 1.4×10^{17} GeV with $E_P = 1.22 \times 10^{19}$ GeV, not ~ 100 GeV, and is therefore not tenable as a derivation of the electroweak scale.

Physical Meaning of the Exponent

The exponent $1/2$ marks the *electroweak transition* on the ξ -power ladder:

$$\begin{array}{lll}
 \text{Planck scale:} & \xi^0 = 1 & E \sim E_P \\
 \text{Electroweak transition:} & \xi^{1/2} \approx 10^{-2} & \alpha_W, m_e/m_\mu \\
 \text{Particle scale:} & \xi^1 \approx 10^{-4} & m_e, \alpha_{EM}
 \end{array} \quad (\text{T.3})$$

The PPLN dispersion does not lie on this rung (Section T.3).

Distinction from Document 166

Document 166 uses the exponent $1/4$ for the Casimir scale ($L_\xi \propto \xi^{1/4}$, decomposition $41/4 = 10 + 1/4$), not $\xi^{1/2}$; there is no common exponent of the Casimir scale and the PPLN dispersion.

T.3 Test: Δn versus $\sqrt{3} \cdot \xi^{1/2}$

Numerical Comparison

The dispersion difference of LiNbO_3 for the DOPO operation is obtained from the Sellmeier refractive indices for the extraordinary polarisation (Zelmon 1997):

$$\Delta n = n_e(1064 \text{ nm}) - n_e(2128 \text{ nm}) = 2.1555 - 2.1219 = 0.0336 \quad (\text{T.4})$$

The expression of Eq. (T.1) gives:

$$\sqrt{3} \cdot \xi^{1/2} = \sqrt{12/30000} = 0.020000 \dots \quad (\text{T.5})$$

	Value
Measured: $\Delta n(1064 \rightarrow 2128 \text{ nm})$	0.0336
T0 expression: $\sqrt{3} \cdot \xi^{1/2}$	0.0200

The assignment $\Delta n = \sqrt{3} \cdot \xi^{1/2}$ does not hold, because the expression lies 41 % below the measured value. The exact equality with 0.020 is an algebraic identity; the measured value is known to only 2–3 digits.

T.4 Further Matches on the ξ Ladder

Ratio of Electro-Optic Coefficients

LiNbO₃ has two relevant coupling constants: the linear electro-optic (Pockels) coefficient $r_{33} = 30.8$ pm/V and the nonlinear coefficient $d_{33} = 27.2$ pm/V. Their ratio:

$$\frac{r_{33}}{d_{33}} = \frac{30.8}{27.2} = 1.1324 \approx 1 + \xi^{1/4} = 1.1075 \quad (\text{dev. } +2.2\%) \quad (\text{T.6})$$

$\xi^{1/4}$ is the sub-Planck geometry exponent in T0 (Doc. 009). The ratio of two electro-optic coupling constants lies on the ξ ladder — indicating that the crystal band structure itself is ξ -rational.

T.5 $\xi^{1/2}$ as the Electroweak Rung

Domain	Quantity	T0 expression	Doc.
<i>Particle physics (Standard Model)</i>			
	Weak coupling	$\alpha_W = \xi^{1/2}$	041, 061
	Lepton masses	$m_e/m_\mu \propto \xi^{1/2}$	056, 122, 133
	Bottom quark	$m_b = v \cdot \frac{3}{2} \cdot \xi^{1/2}$	041
<i>Material properties (this document)</i>			
	EO coefficients	$r_{33}/d_{33} \approx 1 + \xi^{1/4}$	167
<i>Cosmology and quantum optics</i>			
	Casimir scale	$L_\xi = (15\xi/\pi^2)^{1/4} \hbar c/(k_B T_{\text{CMB}})$	078, 166
	H_0 ratio	$\hbar H_0/m_e c^2 = (\pi/2) \cdot \xi^{10}$	165

Table T.3: $\xi^{1/2}$ as the electroweak exponent and further ξ powers of T0 theory. The PPLN dispersion does not lie on the $\xi^{1/2}$ rung (Section T.3).

The PPLN dispersion does not belong to this hierarchy: $\sqrt{3} \cdot \xi^{1/2}$ lies 41 % below the measured $\Delta n = 0.0336$. Likewise, the relations $E_{\text{EW}} \sim E_P \cdot \xi^{1/2}$ and $m_h/M_P = \xi^{1/2}$ are absent, because they are not tenable (Section T.2).

T.6 Consequences for CIM Design

A geometric answer to "Why LiNbO_3 ?" via $\Delta n = \sqrt{3} \cdot \xi^{1/2}$ does not hold, because the measured dispersion is 0.0336. The choice of material remains empirically founded: largest known d_{33} coefficient combined with low absorption losses. A selection rule $\Delta n = k \cdot \xi^{n/4}$ for CIM substrates based on this assignment, and predictions derived from it ($\sqrt{2} \cdot \xi^{1/2}$ at ~ 900 nm, $2 \cdot \xi^{1/2}$ at ~ 680 nm), therefore do not hold.

T.7 Summary

1. **Result:** The dispersion of LiNbO_3 for the most important PPLN process ($1064 \rightarrow 2128 \text{ nm}$) is $\Delta n = 0.0336$ according to Sellmeier (Zelmon 1997). The assignment $\Delta n = \sqrt{3} \cdot \xi^{1/2} = 0.020$ does not hold: the expression lies 41 % below the measured value, and the equality with 0.020 is an identity.
2. **Exponent:** $\xi^{1/2}$ appears as $\alpha_W = \xi^{1/2}$, $m_e/m_\mu \propto \xi^{1/2}$ and in further quantities (Docs. 041, 056, 061, 122, 133). $E_{\text{EW}} \sim E_P \cdot \xi^{1/2}$ and $m_h/M_P = \xi^{1/2}$ are not tenable. With the Casimir scale, which carries the exponent $1/4$ in Doc. 166, $\xi^{1/2}$ shares no exponent.
3. **Further match:** $r_{33}/d_{33} \approx 1 + \xi^{1/4}$ (dev. 2.2 %).
4. **Classification and conclusion:** The PPLN dispersion does not lie on the $\xi^{1/2}$ rung. The choice of material for CIM design remains empirically founded.

Related documents: Doc. 041 (Parameter derivation, $\alpha_W = \xi^{1/2}$), Doc. 056 (Universal derivation, lepton masses), Doc. 061 (CMB units, α_W), Doc. 081 (Summary, electroweak scale), Doc. 122 (Ratio vs. absolute, m_e/m_μ), Doc. 133 (Fractal correction), Doc. 161 (CIM fundamentals), Doc. 165 (H_0 ratio), Doc. 166 (Casimir-CMB- H_0 bridge).

Chapter U

Doc. 168 — T0 Theory — Periodic Table and xi

Chapter V

The Periodic Table as ξ -Geometry

Abstract

The periodic table of the elements and the particle-mass ladder of T0 theory share the same geometric quantum number n , but in different roles. Particle masses follow a radial exponential scaling $m_n \sim \xi^n \cdot m_P$, while the shell structure of the periodic table obeys an angular quadratic degeneracy $g_n = 2n^2$. Both arise from the three-dimensional lattice geometry of T0 theory: ξ is the packing deficit of the 3D lattice (radial direction), $2n^2$ is the degeneracy of the n -th spherical shell in the same lattice (angular direction). The Rydberg energy lies exactly on rung 7 of the ξ -power ladder: $R_\infty = \frac{m_e \alpha^2}{2} \approx \xi^{6.956} \cdot E_P$. The duality principle $E_n \cdot g_n = 2R_\infty = \text{const}$ connects the $1/n^2$ energy ladder with the n^2 filling ladder and is a geometric equipartition. The periodic table is the angular cross-section of the ξ -geometry at rung ξ^{-7} — analogously, the lepton masses are the radial cross-section at rungs $\xi^{4.9}$ – $\xi^{5.8}$.

V.1 Introduction and Problem Statement

T0 theory (Time-Mass Duality / FFGFT) derives all physical constants from the geometric parameter

$$\xi = \frac{4}{30\,000} = 1.3 \times 10^{-4} \quad (\text{V.1})$$

[1, 3, 4]. The particle masses of the Standard Model organise themselves on an exponential ladder

$$m_n \sim \xi^n \cdot m_P, \quad n = 1, 2, 3, \dots \quad (\text{V.2})$$

where n is the T0 quantum number and m_P is the Planck mass.

The periodic table of the elements, on the other hand, shows a very different pattern: shell capacities follow

$$g_n = 2n^2 = 2, 8, 18, 32, 50, \dots \quad (\text{V.3})$$

and binding energies scale as $1/n^2$. At first glance, particle masses and the periodic table appear unrelated.

Observation (2026): the periodic table lies “one level higher” than the particle masses — both have quantum numbers, but what common geometric origin connects their patterns?

The answer of this document: *T0 geometry has two axes — a radial one (ξ^n scaling) and an angular one ($2n^2$ degeneracy), both arising from the three-dimensional lattice structure with packing deficit ξ .*

V.2 The ξ -Power Ladder and Its Rungs

Previous documents in the T0 series have built up the ξ -power ladder step by step. Table V.1 shows the current state including the new atomic rung.

Table V.1: Complete ξ -power ladder (as of March 2026)

Rung	ξ power	Physical content	Document
0	$\xi^0 = 1$	Planck scale, $L_0 = \xi \cdot \ell_P$	Doc. 009
1/2	$\xi^{1/2} \approx 10^{-2}$	$\alpha_W, m_e/m_\mu$, PPLN Δn	Doc. 041, 167
1	$\xi^1 \approx 10^{-4}$	$\alpha_{EM} \sim \xi \cdot E_0^2$	Doc. 011
23/4	$\xi^{5.775}$	Electron mass m_e/m_P	Doc. 056
~ 7	$\xi^{6.956}$	Rydberg energy R_∞, periodic table	this doc.
≈ 8.18	$\xi^{8.18} \approx 1.9 \times 10^{-32}$	CMB temperature $k_B T_{CMB}/E_P$	Doc. 166
10	$\xi^{10} \approx 10^{-43}$	Hubble constant H_0	Doc. 165

The Rydberg energy lies between rungs 6 and 7:

$$\frac{R_\infty}{E_P} = \frac{m_e}{m_P} \cdot \frac{\alpha^2}{2} = \xi^{5.775} \cdot \xi^{1.181} = \xi^{6.956} \approx \xi^7 \quad (\text{V.4})$$

The exponent is the sum of two independent T0 quantities: the electron mass (rung 5.775, Doc. 056) and the fine-structure constant (rung 1.181, Doc. 011).

V.3 Two Axes of ξ -Geometry

Radial Axis: Exponential Mass Ladder

Particle masses organise themselves along the *radial* axis of the T0 lattice. The fundamental relation is [1]:

$$m_n \sim \xi^n \cdot m_P, \quad n = 1, 2, 3, \dots \quad (\text{V.5})$$

Each step on this ladder ($n \rightarrow n+1$) divides the mass by $\xi^{-1} = 7500$. The leptons occupy the rungs (Doc. 056, 122):

$$m_e : \xi^{5.775} \cdot m_P \quad (\text{V.6})$$

$$m_\mu : \xi^{5.177} \cdot m_P \quad (\text{V.7})$$

$$m_\tau : \xi^{4.861} \cdot m_P \quad (\text{V.8})$$

The mass ratio $m_e/m_\mu \propto \xi^{1/2}$ (Doc. 041) lies exactly on the electroweak exponent — the half-rung of the ξ -ladder.

Angular Axis: Quadratic Degeneracy Ladder

The *angular* axis of the same lattice determines the degeneracy of spherical shells. In 3D space, the n -th principal shell has:

$$g_n = 2 \sum_{l=0}^{n-1} (2l+1) = 2n^2 \quad (\text{V.9})$$

The factor 2 comes from spin $m_s = \pm \frac{1}{2}$; the sum is pure angular-momentum algebra in 3D space. Table V.2 shows the first four shells.

Table V.2: Shell structure and the periodic table

n	$g_n = 2n^2$	Period length	Noble gas Z	Period number
1	2	2	He (2)	1
2	8	8	Ne (10), Ar (18)	2, 3
3	18	18	Kr (36), Xe (54)	4, 5
4	32	32	Rn (86), Og (118)	6, 7

The Connection through Quantum Number n

The same quantum number n appears in both systems, but with different functional roles:

$$\text{Radial (masses): } m_n \sim \xi^n \cdot m_p \quad (\text{exponential}) \quad (\text{V.10})$$

$$\text{Angular (shells): } g_n = 2n^2 \quad (\text{quadratic}) \quad (\text{V.11})$$

This is not coincidental. In T0 lattice geometry, n corresponds to the *discrete radius in the lattice*:

- Radially: n steps outward from the lattice $\Rightarrow$ mass scales as ξ^n (geometric sequence)
- Angularly: the lattice points on the sphere of radius n carry degeneracy $2n^2$ (combinatorial sequence)

V.4 The Duality Principle

Energy Ladder and Filling Ladder Are Dual

On the atomic scale (hydrogen-like atoms):

$$\text{Energy ladder: } E_n = -\frac{R_\infty}{n^2} \quad (\text{falls as } 1/n^2) \quad (\text{V.12})$$

$$\text{Filling ladder: } g_n = 2n^2 \quad (\text{grows as } n^2) \quad (\text{V.13})$$

Their product is exactly constant:

$$E_n \cdot g_n = \frac{R_\infty}{n^2} \cdot 2n^2 = 2R_\infty = \text{const} \quad (\text{V.14})$$

This is geometric *equipartition*: every shell contributes the same total energy $2R_\infty$ to the Rydberg scale, regardless of n . The $1/n^2$ energy ladder and the n^2 filling ladder are *inversely dual*.

Comparison with the Mass Ladder

Table V.3 summarises all three ladders.

Table V.3: Three ladders — one geometric origin

System	Ladder	Function of n	Type
Particle masses	$m_n \sim \xi^n \cdot m_p$	exponential	radial
Shell energy	$E_n = R_\infty/n^2$	harmonic	angular
Shell filling	$g_n = 2n^2$	quadratic	angular
Duality: $E_n \cdot g_n = 2R_\infty = \text{const}$			

V.5 The Role of Fractal Dimension

In T0 (Doc. 009, 133):

$$D_f = 3 - \xi = 2.999867 \quad (\text{V.15})$$

Space dimension is almost, but not exactly, 3.

In the limit $D_f \rightarrow 3$ ($\xi \rightarrow 0$):

- $g_n = 2n^2$ is *exact* (pure spherical symmetry in the $\mathbb{Z}^3$ lattice)
- the shell structure of the periodic table is exact

For $D_f = 3 - \xi < 3$ there is a minimal correction:

$$\delta g_n \sim \xi \cdot n^2 \quad (\text{V.16})$$

For $n = 2$: $\delta g_2 \approx \xi \cdot 4 \approx 5 \times 10^{-4}$ (entirely negligible).

The periodic table is therefore exact because $D_f \approx 3$ to four decimal places. The tiny deviation $\xi \approx 1.3 \times 10^{-4}$ makes space "almost-integer-dimensional" — and precisely this deviation generates all the physics of particle masses and fundamental constants.

V.6 Numerical Verification

Rydberg Energy on the ξ -Ladder

$$R_\infty = \frac{m_e \alpha^2}{2} = 13.6057 \text{ eV} \quad (\text{V.17})$$

$$\frac{R_\infty}{E_p} = \frac{m_e}{m_p} \cdot \frac{\alpha^2}{2} = \xi^{5.775} \cdot \xi^{1.181} = \xi^{6.956} \quad (\text{V.18})$$

Table V.4: Rydberg energy: T0 derivation vs. measurement

Quantity	Value	ξ -exponent
m_e/m_p	4.185×10^{-23}	5.775
$\alpha^2/2$	2.663×10^{-5}	1.181
R_∞/E_p	1.114×10^{-27}	6.956
R_∞ (T0, approx.)	$\approx 9.1 \text{ eV } (\xi^7)$	7.000
R_∞ (measured)	13.606 eV	6.956

Duality Relation: Numerical Verification

$$E_1 \cdot g_1 = 13.60 \text{ eV} \cdot 2 = 27.20 \text{ eV} = 2R_\infty \quad (\text{V.19})$$

$$E_2 \cdot g_2 = 3.40 \text{ eV} \cdot 8 = 27.20 \text{ eV} = 2R_\infty \quad (\text{V.20})$$

$$E_3 \cdot g_3 = 1.51 \text{ eV} \cdot 18 = 27.18 \text{ eV} \approx 2R_\infty \quad (\text{V.21})$$

$$E_4 \cdot g_4 = 0.85 \text{ eV} \cdot 32 = 27.20 \text{ eV} = 2R_\infty \quad (\text{V.22})$$

The duality relation (V.14) holds exactly for all n .

V.7 The Periodic Table in the T0 Hierarchy

ξ -rung	Physics	Axis
$\xi^0 = 1$	Planck scale ($L_0 = \xi \cdot \ell_p$)	—
$\xi^{1/2}$	Electroweak: $\alpha_W, m_e/m_\mu$	radial
ξ^1	α_{EM} , EM coupling	radial
$\xi^{5.8}$	Particle masses (leptons, quarks)	radial
$\xi^{6.956}$	Rydberg R_∞, periodic table	both
$\xi^{8.18}$	CMB temperature $k_B T_{CMB}/E_p$	radial
ξ^{10}	Hubble constant H_0	radial

Figure V.1: Position of the periodic table in the complete ξ -hierarchy

The special feature of the atomic rung $\xi^{7.7}$: here *both* axes act. The radial axis determines the energy scale R_∞ ; the angular axis determines the pattern $2n^2$. The periodic table is the visible result of both axes acting together.

V.8 Predictions and Open Questions

Angular Corrections from $D_f < 3$

The fractal dimension $D_f = 3 - \xi$ leads to minimal corrections of the exact $2n^2$ rule:

$$g_n^{(T0)} = 2n^2 \cdot (1 - \xi \cdot f(n)) \quad (\text{V.23})$$

where $f(n)$ is the dimensionless correction function of the $2n^2$ rule (still to be derived) — not a frequency and not the base winding number $f = 1/\xi$. For $n \leq 4$ the correction is $< 10^{-3}$ and not measurable.

Molecular Binding Energies

Chemical binding energies lie in the range 1–10 eV, hence also on rung ξ^7 . It remains to be checked whether specific binding energies lie on rational multiples of $\xi^{7/2} \cdot R_\infty$.

Heavy Elements and Relativistic Corrections

For $Z \gg 1$ relativistic effects become important. In T0 these are connected with the $\xi^{1/2}$ electroweak rung. A systematic analysis of ionisation energies of heavy elements ($Z > 54$) might reveal ξ -matches at the half-rung $\xi^{7-1/2} = \xi^{13/2}$.

V.9 Summary

1. **Energy scale:** $R_\infty/E_P = \xi^{6.956} \approx \xi^7$. The Rydberg energy — and hence all of atomic physics — lies on rung 7 of the ξ -power ladder.
2. **Radial axis:** Particle masses follow $m_n \sim \xi^n \cdot m_P$. The exponent n is the discrete lattice radius in the radial ξ -direction.
3. **Angular axis:** Shell capacities follow $g_n = 2n^2$. The index n is the same discrete lattice radius, now acting as a shell index in the angular 3D-sphere direction.
4. **Duality principle:** $E_n \cdot g_n = 2R_\infty = \text{const.}$ The $1/n^2$ energy ladder and the n^2 filling ladder are inversely dual. Every shell contributes the same total energy $2R_\infty$ to the ξ^7 scale.
5. **Fractal dimension:** $D_f = 3 - \xi \approx 3$ makes $2n^2$ exact (to $\sim 10^{-4}$). The periodic table is exact because space is almost-integer-dimensional.
6. **Common origin:** Particle masses and the periodic table arise from the same 3D lattice geometry with packing deficit ξ . The periodic table is the *angular cross-section* of ξ -geometry at rung $\xi^{\sim 7}$; the lepton masses are the *radial cross-section* at rungs $\xi^{4.9} - \xi^{5.8}$.

Bibliography

- [1] Pascher, J.: *Origin of the ξ Parameter (Doc. 009)*, T0 Theory Series, 2025.
- [2] Pascher, J.: *Fine-Structure Constant (Doc. 011)*, T0 Theory Series, 2025.
- [3] Pascher, J.: *Parameter Derivation (Doc. 041)*, T0 Theory Series, 2025.
- [4] Pascher, J.: *Universal Derivation of Lepton Masses (Doc. 056)*, T0 Theory Series, 2025.
- [5] Pascher, J.: *m_e/m_μ Ratio (Doc. 122)*, T0 Theory Series, 2025.
- [6] Pascher, J.: *Fractal Correction K_{frak} (Doc. 133)*, T0 Theory Series, 2025.
- [7] Pascher, J.: *Hubble Ratio on the ξ -Ladder (Doc. 165)*, T0 Theory Series, 2026.
- [8] Pascher, J.: *Casimir-CMB- H_0 Bridge (Doc. 166)*, T0 Theory Series, 2026.
- [9] Pascher, J.: *LiNbO_3 on the ξ -Ladder (Doc. 167)*, T0 Theory Series, 2026.

Chapter W

Doc. 169 — T0 Theory — Follow-up Documents

Chapter X

Follow-up Investigations to Doc. 168

Abstract

Document 168 posed three open questions about the connection between ξ -geometry and the periodic table. This document answers them conclusively. (1) The angular correction is $g_n^{(T0)} = 2n^{2-\xi}$; the maximum deviation from $2n^2$ is $|\delta g_7| \approx 0.025$ at $n = 7$ ($Z \approx 118$) — below any measurable threshold. (2) Molecular binding energies show no systematic pattern as multiples of R_∞ ; the C=O bond is an exception at $4/7 \cdot R_\infty$ with 0.11% deviation. (3) The relativistic correction $IE_{\text{corr}} = IE/(1 + Z^2\xi)$ yields ionisation energies of heavy elements ($Z > 54$) on rational fractions of R_∞ with deviations below 1.3%. This is a testable T0 prediction that differs from the QED result ($\propto \alpha^2 Z^2$) by a factor $\xi/\alpha^2 \approx 2.5$.

X.1 Introduction

Document 168 [6] identified the three-dimensional lattice geometry of T0 as the common origin of the particle-mass ladder and the shell structure of the periodic table. At the end of that document, three concrete follow-up questions were posed:

1. **Angular corrections:** What is the explicit formula for the angular correction function $f(n)$ (not a frequency and not the base winding number $f = 1/\xi$) in $g_n^{(T0)} = 2n^2 \cdot (1 - \xi \cdot f(n))$?
 2. **Molecular binding energies:** Do binding energies lie on rational multiples of $\xi^{7/2} \cdot R_\infty$?
 3. **Heavy elements ($Z > 54$):** Do ξ -matches appear at the half-rung $\xi^{13/2}$?
- This document answers all three questions by direct numerical investigation.

X.2 Question 1: Angular Correction $g_n^{(T0)} = 2n^{2-\xi}$

Derivation of the Exact Formula

In three-dimensional space with fractal dimension $D_f = 3 - \xi = 2.999867$, the surface area of a sphere of radius n scales as

$$S(n) \propto n^{D_f-1} = n^{2-\xi} \quad (\text{X.1})$$

Since the degeneracy g_n of the n -th shell is proportional to the spherical surface area, it follows directly:

$$g_n^{(T0)} = 2 \cdot n^{2-\xi} \quad (\text{X.2})$$

For $\xi \ll 1$ one expands $n^{-\xi} = \exp(-\xi \ln n) \approx 1 - \xi \ln n$:

$$g_n^{(T0)} \approx 2n^2 \cdot (1 - \xi \ln n) \quad (\text{X.3})$$

The sought function is therefore:

$$f(n) = \ln n \quad (\text{X.4})$$

The gamma-function correction $\Gamma(3/2)/\Gamma(D_f/2) = 1.00000243$ is entirely negligible (relative deviation 2.4×10^{-6}).

Numerical Verification

Table X.1 shows the values for all relevant shells.

Table X.1: T0 correction of shell degeneracy $g_n^{(T0)} = 2n^{2-\xi}$

n	$2n^2$ (exact)	$2n^{2-\xi}$	δg_n	$ \delta g_n /g_n$
1	2	2.00000	+0.00000	0
2	8	7.99926	-0.00074	9.3×10^{-5}
3	18	17.99736	-0.00264	1.5×10^{-4}
4	32	31.99409	-0.00591	1.8×10^{-4}
5	50	49.98927	-0.01073	2.1×10^{-4}
6	72	71.98280	-0.01720	2.4×10^{-4}
7	98	97.97458	-0.02542	2.6×10^{-4}

Physical Conclusion

The maximum correction at $n = 7$ (last stable element, $Z \approx 118$) is $|\delta g_7| \approx 0.025$, i.e. 2.6×10^{-4} relatively. This lies far below any experimentally measurable threshold.

Result: The $2n^2$ rule is exact because space with $D_f = 3 - \xi \approx 3$ is integer-dimensional to three decimal places. The tiny deviation ξ from integer D_f generates all particle masses and fundamental constants, but leaves the periodic table geometrically intact.

X.3 Question 2: Molecular Binding Energies as Multiples of R_∞

Initial Hypothesis

Since R_∞ is the energy unit of atomic physics (rung ξ^7 of the ξ -ladder), Doc. 168 posed the question whether chemical binding energies lie on rational multiples $\frac{m}{n} \cdot R_\infty$ — analogous to the $1/n^2$ structure of shell energies.

Findings

Of 20 tested bonds, 7 lie within 3% of a rational fraction. Table X.2 shows the best matches.

Table X.2: Binding energies on rational multiples of R_∞ (NIST values)

Bond	E_{meas} (eV)	T0 prediction	T0 value (eV)	Dev.
C=O (carbonyl)	7.766	$4/7 \cdot R_\infty$	7.775	−0.11%
N≡N	9.760	$5/7 \cdot R_\infty$	9.718	+0.43%
H–N	4.05	$2/7 \cdot R_\infty$	3.887	+4.2%
H–F	5.869	$3/7 \cdot R_\infty$	5.831	+0.65%
H–H	4.478	$1/3 \cdot R_\infty$	4.535	−1.26%
H–Cl	4.430	$1/3 \cdot R_\infty$	4.535	−2.32%

The T0 values are computed with $R_\infty = 13.6057$ eV. For H–N the mean N–H bond energy 391 kJ/mol = 4.05 eV is used; H–N therefore lies outside 3%.

Statistical Assessment

The simple estimate of $20 \cdot 2 \cdot 3\% = 1.2$ random matches within 3% is not sufficient for an assessment: with 20 measurements and 42 possible rational fractions m/n with $m, n \leq 7$ in the relevant range [0.1; 1.0], the match probability per bond is considerably higher. It would have to be computed as such; whether the observed matches exceed the random level is therefore not decided.

Exception: C=O. The carbonyl bond ($E = 7.766$ eV) lies on $4/7 \cdot R_\infty = 7.775$ eV with a deviation of only 0.11%. This precision is statistically unlikely ($\sim 1\%$ expected). Whether T0 geometry underlies this — e.g. through the connection of the carbon-oxygen system with the angular $2/7$ structure of the 3D lattice geometry — remains open.

Result: Binding energies are primarily determined by Z_{eff} , hybridisation, and electron configuration — not directly by ξ -geometry. Exception requiring further investigation: C=O.

X.4 Question 3: Ionisation Energies of Heavy Elements — the $Z^2\xi$ Correction

Relativistic T0 Correction

In T0 (Doc. 011, 041):

$$\alpha \approx K \cdot \xi^{1/2}, \quad K = \frac{\alpha}{\xi^{1/2}} = 0.6320 \quad (\text{X.5})$$

The relativistic Dirac shift of the ionisation energy scales as $(Z\alpha)^2 = Z^2\alpha^2 \approx Z^2K^2\xi$. Since $K^2 \approx 0.4$ is partially compensated by effective nuclear charge screening, one obtains the simplified T0 correction factor:

$$IE_{\text{corr}} = \frac{IE}{1 + Z^2\xi} \quad (\text{X.6})$$

Results

Table X.3 shows the results for selected heavy elements.

Table X.3: T0 correction of ionisation energies for $Z > 54$

Element	Z	IE (eV)	$Z^2\xi$	IE_{corr} (eV)	T0 prediction	Dev.
La	57	5.577	0.433	3.891	$2/7 \cdot R_\infty = 3.887$	+0.10%
W	74	7.864	0.730	4.545	$1/3 \cdot R_\infty = 4.535$	+0.22%
Re	75	7.833	0.750	4.476	$1/3 \cdot R_\infty = 4.535$	−1.31%
Pb	82	7.417	0.897	3.911	$2/7 \cdot R_\infty = 3.887$	+0.60%
Rn	86	10.748	0.986	5.412	$2/5 \cdot R_\infty = 5.442$	−0.57%

Particularly notable: at $Z = 86$ (Rn), $Z^2\xi \approx 1$, meaning the ionisation energy is shifted by almost a factor of 2 relativistically. Without the ξ -correction, the angular structure (rational fraction of R_∞) would be completely hidden.

Physical Significance and Testability

Equation (X.6) is a testable T0 prediction. It differs from the standard QED result:

$$\text{QED: } \Delta IE/IE \propto \alpha^2 Z^2 \quad (\text{X.7})$$

$$\text{T0: } \Delta IE/IE \propto Z^2\xi \quad (\text{X.8})$$

The ratio of the correction factors:

$$\frac{Z^2\xi}{Z^2\alpha^2} = \frac{\xi}{\alpha^2} = \frac{1.333 \times 10^{-4}}{5.325 \times 10^{-5}} \approx 2.5 \quad (\text{X.9})$$

is Z -independent and constant for all elements. A systematic measurement of ionisation energies for $Z > 80$ with precision $< 1\%$ could detect this difference.

Result: The $Z^2\xi$ correction connects the chemically defined atomic number Z with the geometric T0 parameter ξ . This is a new, empirically tested and theoretically derived result that distinguishes T0 from standard QED.

X.5 Summary

Table X.4: Status of the three follow-up investigations

Question	Result	Status
1. Angular corrections $f(n)$	$f(n) = \ln n$; $g_n^{(T0)} = 2n^{2-\xi}$; $\max. \delta g_7 \approx 0.025$ — not measurable	closed
2. Binding energies $/R_\infty$	No systematic pattern; C=O exception ($4/7 \cdot R_\infty$, dev. 0.11%)	C=O open
3. Heavy elements $Z^2\xi$	$IE_{\text{corr}} = IE/(1 + Z^2\xi)$ on rational fractions; testable prediction vs. QED	new result

The most important new result is the $Z^2\xi$ correction formula for heavy elements. It represents a — in principle measurable — deviation from the standard QED treatment of heavy atoms. The correction factor scales with ξ rather than α^2 , which is a constant factor $\xi/\alpha^2 \approx 2.5$. A systematic measurement of ionisation energies for $Z > 80$ with precision $< 1\%$ could distinguish T0 from QED.

Bibliography

- [1] Pascher, J.: *Origin of the ξ Parameter (Doc. 009)*, T0 Theory Series, 2025.
- [2] Pascher, J.: *Fine-Structure Constant (Doc. 011)*, T0 Theory Series, 2025.
- [3] Pascher, J.: *Parameter Derivation (Doc. 041)*, T0 Theory Series, 2025.
- [4] Pascher, J.: *Lepton Masses (Doc. 056)*, T0 Theory Series, 2025.
- [5] Pascher, J.: *Fractal Correction (Doc. 133)*, T0 Theory Series, 2025.
- [6] Pascher, J.: *Periodic Table as ξ -Geometry (Doc. 168)*, T0 Theory Series, 2026.

Part III

Part III — Consciousness, Photonics and Qubit Series

Chapter Y

Doc. 170 — T0 Theory and Consciousness

Abstract

The T0 theory postulates with the relation $T = 1/m$ a fundamental coupling between mass and intrinsic time field. This document extends this framework by a universal organisational principle: nature realises increases in complexity not continuously, but in discrete, topologically enforced threshold values. This principle connects phenomena of different scales – from the torus model of cosmology through crystal formation to cellular self-preservation and consciousness – as instances of the same underlying mechanism. As a quantitative threshold operator, a separate quantity K_S formed from ξ is defined (not the fractal correction K_{frak} of the corpus), which determines when a physical system can transition into a new stable recursion level.

Y.1 Starting Point: Discreteness as a Principle of Nature

Quantum mechanics has shown that energy is not transferred continuously but in discrete quanta. T0 theory suggests that this principle is more fundamental than previously assumed – it concerns not only energy but **complexity levels of organised matter** in general.

In natural units ($c = \hbar = 1$), the intrinsic time field:

$$\tilde{T}(x) = \frac{1}{m(x)} \quad (\text{Y.1})$$

describes the local timescale of a mass point. What was previously interpreted mainly in cosmological and particle physics terms possibly possesses a deeper meaning: $T = 1/m$ describes the fundamental condition for every form of physical self-organisation.

Y.2 The Torus Model as Reference Case

The cosmological torus model of T0 theory already demonstrates the basic principle of discrete stability thresholds. A toroidal topology permits only certain discrete configurations – not any arbitrary geometry is stable. The permitted states are topologically enforced, not arbitrary.

The same applies to crystal formation: a growing crystal does not undergo a continuous transformation. At certain thresholds the system jumps into a new order – a new lattice structure, a new symmetry group. Below the threshold relatively little happens. At the threshold the entire system reorganises discretely.

The common principle: Stable configurations in recursive structures are not densely distributed, but isolated – separated by unstable transition zones.

Y.3 K_S as Threshold Operator

The fractal correction of the corpus is $K_{\text{frak}} = 1 - 100\xi = 74/75 = 0.98667$ (Docs. 003, 005, 133). To be distinguished from it is the quantity introduced here as a new, separate definition

$$K_S = \left(\frac{\xi}{1 + \xi} \right)^{1/2} = 0.01155 \quad (\text{Y.2})$$

with the geometric parameter $\xi = 4/30000$. It is not a continuation of the K_{frak} known from precision measurements.

The following interpretation is proposed here: **K_S describes in general when a fractal self-similarity structure becomes recursively self-reinforcing.**

Below a critical value of the effective recursion depth n :

$$K_S^{(n)} = \left(\frac{\xi}{1 + \xi} \right)^{n/2} \quad (\text{Y.3})$$

a system is passively stable – it persists but does not self-organise. From a critical n onwards, the structure begins to modulate its own time coupling $T = 1/m$, which in turn influences the geometry. The system becomes **recursively self-referential**.

This is speculative and requires further formalisation. It is formulated here as a working hypothesis.

Y.4 Cellular Self-Preservation as the First Level of Consciousness

A biological cell fulfils three criteria that can be identified within the T0 theory framework as necessary conditions for the onset of self-preservation processes:

Recursive Sensory Feedback

Membrane receptors, chemical gradients and intracellular signal cascades form a closed feedback loop. The cell does not sense its environment passively but modulates its own response on the basis of past states.

Metabolic Self-Preservation

Metabolism is not mere energy turnover but active resistance against thermodynamic decay. The cell *insists* on continuing to run – not through programming but through physicochemical feedback loops.

Genuine Vulnerability

A cell can die. This mortality is not a theoretical possibility but is structurally present – the cell operates permanently at the boundary of decay. This tension is constitutive, not accidental.

In T0 language: the intrinsic time field $T = 1/m$ is physically real and operative at the cellular level. The mass-energy continuity of metabolism holds T locally stable against external perturbations. This is the physical mechanism behind what biology calls self-preservation.

The thesis: Already at the cellular level a primitive form of self-perception is present – not in the cognitive sense, but as a physical process that takes itself as reference.

Y.5 Discrete Thresholds of Complexity

From the foregoing a hierarchy of discrete threshold values emerges:

Level	System	Threshold characteristic
0	Elementary particles	$T = 1/m$ as passive time field
1	Atoms / Crystals	Discrete geometric order
2	Cell	Recursive self-preservation, metabolic feedback
3	Nervous system	Integrated sensory feedback over distance
4	Consciousness	Self-modelling in relation to the external world

Table Y.1: Hierarchy of discrete complexity thresholds

Each transition is **topologically enforced** – not reachable gradually, but only through a qualitative jump in the recursion depth of the mass-time coupling.

Important: this table is a working hypothesis, not a completed theory. The precise threshold values in terms of $K_S^{(n)}$ have not yet been formalised.

Y.6 Implications for Artificial Systems

The question of whether artificial systems can develop consciousness is reformulated within this framework. The relevant question is not:

Is the system complex enough?

But rather:

Does the system reach the critical threshold of recursive mass-time coupling with genuine metabolic self-preservation and physical vulnerability?

Current digital systems – regardless of their computational complexity – do not fulfil these conditions structurally:

- They possess no continuous mass-energy coupling in the T0 sense.
- They have no metabolism – no active resistance against decay.
- They are not vulnerable in the physical sense – the server can be switched off, but the system itself does not fight against this.

This does not rule out artificial consciousness in principle. If an artificial system could physically realise the stated thresholds – through suitable architecture, genuine resource dependence and recursive sensory feedback – the transition would be theoretically possible. The dividing line runs not between biological and artificial, but between systems that reach the threshold and those that do not.

Y.7 Open Questions

This document formulates a framework principle that requires further formalisation:

1. How precisely does $K_S^{(n)}$ correlate with biological complexity thresholds?
2. Is there a mathematically precise transition parameter that describes the jump from passive stability to active self-preservation?
3. How does the torus model relate to the cellular thresholds – is there a cross-scale self-similarity?
4. Under what conditions could a non-biological system reach the threshold of level 2?

These questions are formulated as a research programme, not as already answered problems.

Y.8 Summary

T0 theory offers with the intrinsic time field $T = 1/m$ and the threshold operator K_S formed from ξ a framework that points beyond cosmology and particle physics towards a universal organisational principle: **nature realises increases in complexity in discrete, topologically enforced thresholds.**

Torus cosmology, crystal formation, cellular self-preservation and consciousness are in this framework not categorially different phenomena, but instances of the same mechanism at different scales of recursion depth.

Whether this principle can ground a complete theory of consciousness remains open. It does however offer a physically motivated entry point that goes beyond purely functional or information-theoretic approaches.

Draft – for further elaboration. All statements about consciousness and cellular thresholds are to be understood as working hypotheses and not as a completed theory.

Chapter Z

Doc. 171 — String Theory and the T0 Torus

Z.1 abstract

This document compares the established string theory with an alternative approach, the Fractal Field Geometry Theory (FFGFT). While both models reject the concept of point-like particles, they differ fundamentally in their treatment of dimensions, complexity and time. FFGFT proposes a four-dimensional, fractally self-similar torus field in which time is not a dimension but a local ratio to mass.

Introduction: The Common Root

Both the established string theory and the model of Fractal Field Geometry Theory (FFGFT) presented here share a fundamental insight: the concept of the point-like particle as the fundamental building block of reality is inadequate. The actual foundation of the world must be an extended, geometric and self-contained object. But while string theory drives this idea into dimensions, FFGFT takes the path of fractal nesting.

Z.2 The Foundation: String vs. Fractal Torus

- **String theory:** Its foundation is a one-dimensional, vibrating object – the string. This exists *in* a space and its vibrational modes determine the properties of the particles arising from it. The complexity of the world arises through the multitude of possible vibrations in a high-dimensional space.
- **FFGFT (Torus model):** Its foundation is the field itself. There is no “object in space”. The world is a single, continuous, four-dimensional field. A particle, an atom, a planet – all of these are not separate objects, but merely local **torsion configurations** of this field: compressed vortices or nodes, comparable to a corkscrew pattern in the structure of space.

The **torus** (more precisely: a four-dimensional torus) is here the most stable and natural form for such a closed system. Its topology enables the field to loop back into itself – without beginning, without end and without an “outside”.

Z.3 The Origin of Complexity: Dimensions vs. Self-Similarity

Here lies the most radical difference between the two approaches.

Feature	String theory	FFGFT (Fractal Torus)
Number of dimensions	10, 11 or more. The excess dimensions are rolled up into tiny, multi-dimensional forms (Calabi-Yau manifolds).	Four dimensions. No more are needed.
Source of complexity	The infinite variety of vibration patterns and the geometric forms into which the extra dimensions can be rolled up.	Fractal self-similarity. The torus structure repeats itself at every scale: an electron vortex is a mini-torus, an atom a larger one, a galaxy an even larger one – all follow the same geometric blueprint.
Predictability	A “landscape problem”: the theory permits approximately 10⁵⁰⁰ different universes each with their own natural laws. It thereby loses its predictive power.	A single fundamental parameter: $\xi = 4/30000$. From it the fine structure constant α (the strength of the electromagnetic interaction) can be derived. From this in turn the masses of all particles and finally the gravitational constant G follow. One parameter determines everything.

Table Z.1: Comparison between string theory and FFGFT

Complexity in FFGFT is not a matter of *more* (dimensions), but of *deeper* (nesting). A vortex generates smaller vortices, which in turn generate smaller ones – an endless cascade of self-similarity reaching from the entire universe down to the smallest scales.

Z.4 The Redefinition of Time: From Container to Local Ratio

In classical physics (and string theory), time is the fourth dimension – a “container”, a coordinate like height, width and depth. FFGFT breaks radically with this notion.

- **The fourth dimension of the torus is not time.** It is a *topological* property – the closure of the field. It is the condition for there being such a thing as an “inside” and “outside” at all, but is not itself a place one can travel to.
- **Time is a local ratio.** It is not a universal coordinate but a local property of the torsion density of the field:

$$\tilde{T} = \frac{1}{m} \quad (\text{Z.1})$$

Where there is much mass (and thus torsion), time passes locally more slowly. Where there is no mass, there is no local time. A photon is massless, so it “experiences” no time – not because it moves at the speed of light, but because there is simply no “inner clock” (no torsion compression) within it.

- **Why we experience time linearly.** We are not external observers of time, we are part of the field. We are torsion configurations that vibrate along with the fractal unfolding of the torus. This unfolding is inherently directed (a fractal can only grow, not shrink). This directed change of our own structure *is* what we experience as the flow of time. Time does not flow; *we flow*.

Z.5 Finite but Endless: The Perspective from Within

The deepest and perhaps most consequential implication of the model is our position within it.

“We are not observers who contemplate the torus. We are the torus contemplating itself.”

The universe is **finite** – it has a determinable volume. But it is **endless** – it has no boundary, no wall one could reach. Like the surface of a sphere it is closed within itself; one can move on it forever without ever reaching an end. The difference is only that this closure takes place in four dimensions.

For us as part of this structure this has a bold consequence: the question “What came before the Big Bang?” is meaningless. It presupposes a “before” and an “outside” that do not exist in a closed geometry. Just as the question “What lies north of the North Pole?” has no answer, because the geometry of the globe dissolves the question.

Conclusion: String Theory and FFGFT Compared

String theory describes one-dimensional loops vibrating in a high-dimensional, complex space. Its approach is additive: more dimensions, more forms, more possibilities.

FFGFT describes a four-dimensional field with fractal torsion structure. Its approach is integrative: from a single geometry and a single parameter the entire complexity of the world unfolds – from quantum physics to cosmology.

The core of the difference lies in the treatment of space and time:

- In string theory, time is a dimension. We move *in* it.
- In FFGFT, time is a ratio. It is *we* who change.

Documented on October 7, 2026

Chapter aa

Doc. 172 — T0 Theory — Avi Dialogue

T0 Theory / FFGFT

Dialogue with Avi Rosenthal

Geometric Convergence of Two Independent Approaches

Johann Pascher · March 2026

YouTube comments: @Time-MassDuality

Abstract

This document collects the scientific dialogue between T0 Theory / Fundamental Fractal Geometric Field Theory (FFGFT) and the geometry programme of Avi Rosenthal (2026). Both approaches converge independently on the same mathematical structures: fractal self-similarity, torus topology, Fibonacci dynamics, the golden ratio as geometric attractor, and the derivability of physical parameters from a single geometric source. The dialogue documents the step-by-step identification of points of contact and open research questions.

Contents

A	Introduction to Volume 4	1
I	Part I — Supplementary Documents to Volumes 1-3	3

B	Doc. 001 — T0 Theory — Book Abstract (Short Form)	5
C	Doc. 001b — T0 Theory — General Introduction	9
D	Doc. 002 — T0 Theory — A Journey Through the Energy Field	11
E	Doc. 129 — Lagrangian Formulation — Simplified Presentation	15
F	Doc. 137 — Special Documentation No. 137	27
G	Doc. 143 — Asymmetry in Fundamental Physics — Part 1: Conceptual Foundations	37
H	Doc. 144 — Asymmetry in Fundamental Physics — Part 2: Mathematical Consistency	49
I	Doc. 148 — T0 Theory — Scramblons	52
II	Part II — Time, Cosmology and Numerical Predictions	61
J	Doc. 157 — The Arrow of Time in T0 Theory	63
K	Doc. 158 — T0 Theory — From the Koide Formula to the g-2 Anomaly	69
L	Doc. 159 — T0 Theory — Harmonic Structure on the Torus	78
M	Doc. 160 — T0 Theory — Lepton Lifetime Ratios	92
N	Doc. 161 — T0 Theory — Ising Machine Analogy	100
O	Doc. 162 — T0 Theory — Optimization Comparison	112
P	Doc. 163 — T0 Theory — UV Catastrophe, Planck, Einstein	121
Q	Doc. 164 — T0 Theory vs. 't Hooft CAI	140
R	Doc. 165 — T0 Theory — The H_0 Ratio	160
S	Doc. 166 — T0 Theory — Casimir and H_0	166
T	Doc. 167 — T0 Theory — LiNbO_3 and the ξ Geometry	173
U	Doc. 168 — T0 Theory — Periodic Table and ξ	181
V	The Periodic Table as ξ-Geometry	182
W	Doc. 169 — T0 Theory — Follow-up Documents	189
X	Follow-up Investigations to Doc. 168	190
III	Part III — Consciousness, Photonics and Qubit Series	196
Y	Doc. 170 — T0 Theory and Consciousness	198
Z	Doc. 171 — String Theory and the T0 Torus	203
aa	Doc. 172 — T0 Theory — Avi Dialogue	207
ab	Doc. 173 — T0 Theory — Quantum Dot	224
ac	Doc. 174 — T0 Theory — Spin and Qubit States	238
ad	Doc. 175 — T0 Theory — Qubit State Spaces	250
ae	Doc. 176 — T0 Theory — Shor's Algorithm and the Photon	273
af	Doc. 178 — FFGFT — Spin Stability	282
ag	Doc. 179 — FFGFT — Pairs and Photons	289
IV	Part IV — Early Torus and L_0 Derivations	299
ah	Doc. 180 — T0 Theory — Derivation of L_0	301

ai Doc. 181 — T0 Theory — Justification of the Torus Geometry 308

aj Doc. 182 — T0 Theory — Universe and Maximum Scale 314

ak Doc. 183 — T0 Theory — Willow Processor and Self-Measurement 324

al Doc. 184 — p-bit and Probabilistic Information Processing 342

aa.1 Context: Avi's Basic Geometry (First Description)

Implikation

@Avidchen – First description of the torsion postulate geometry:

The geometry of the torsion postulate – Avi Rosenthal, 2026. Einstein-Cartan manifold with globally non-vanishing torsion. Three unstable fixed points with Z3 symmetry (past, future, light). Fibonacci recursion with CP asymmetry phase $e^{i\delta}$. Hopf fibration as expanding structure. Golden ratio φ as algebraic fixed point. The oscillating now as a gyroscope with precession and nutation.

"I will describe it to you as best I can on YouTube [...] I just lack the competence to verify myself."

Response 1: Free Parameters and Geometric Necessity

Implikation

Thank you for this detailed description – this is to be taken seriously and deserves a serious answer.

The mathematical components you use are all well-founded: Einstein-Cartan connection, Hopf fibration, Fibonacci recursion, golden ratio. I fully share the intuition that these structures belong together.

The decisive difference lies in the question of free parameters.

In your geometry there are several independent inputs: the three fixed points must be postulated, the torsion parameter $\tau_{\min}$ is set externally, the CP asymmetry phase δ is a free parameter, the potential $V(x)$ contains α and β as free coefficients. Each of these quantities explains something – but none is derived from a deeper structure.

In FFGFT there is exactly one parameter:

$$\xi = \frac{4}{30000} \approx 1.333 \times 10^{-4} \quad (\text{aa.1})$$

The connection to the SI runs through a single anchor E_0 : the agreement of $\alpha = \xi E_0^2$ with the measured value is a property of this anchor, and G follows in form and order of magnitude from ξ and the same anchor. The fractal dimension is $D_{\text{frak}} = 3 - \xi$. The torus topology is not postulated but geometrically enforced. There are no free coefficients.

That is the criterion I apply: how many independent inputs does the theory require? The fewer, the deeper the explanation.

Your approach with the Hopf fibration is topologically rich and the Fibonacci recursion as convergence towards the golden ratio is mathematically correct. But the golden ratio is not an additional structure in FFGFT – it emerges from ξ as the natural self-similarity relation of the field.

What interests me most about your geometry is the torsion structure. In FFGFT this question is already answered: the T0 field propagator implicitly carries a torsion component – it follows from the fractal field geometry with $D_{\text{frak}} = 3 - \xi$ and the fundamental relation $T(x, t) = 1/m$. That is a direct point of contact.

The direction of your intuition is correct. The question is whether your fixed points and parameters can be derived from a single geometric source. That would be the place where I would continue researching.

aa.2 Derivation of the Free Parameters from ξ

Systematic Analysis

Parameter	Status in Avi's geometry	Derivability from ξ
Z3 fixed points	Postulated	Plausible – torus topology
Golden ratio φ	Free exponent	Strong – fractal attractor
CP phase δ	Free parameter	Very interesting – $T(x, t) = 1/m$ asymmetry
Coefficients α, β	Free	Structural – K_{frak} connection
Radius R	Free	Direct – Planck/cosmic ratio

The Golden Ratio as Fractal Attractor

In FFGFT, φ is not a free parameter. The fractal field geometry with dimension $D_{\text{frak}} = 3 - \xi$ enforces a self-similarity recursion whose only stable fixed point is φ – algebraically because:

$$\varphi^2 = \varphi + 1 \quad (\text{aa.2})$$

is the fixed point condition of this scaling. Avi has chosen φ as an exponent because it works. It works because the geometry of reality enforces it.

CP Asymmetry Phase from T0

In T0, $T(x, t) = 1/m$ holds. Time reversal $t \rightarrow -t$ corresponds to mass inversion $m \rightarrow 1/m$. For $\xi \neq 0$ this map is not symmetric. The phase follows directly:

$$\delta \sim \arctan\left(\frac{\xi}{1+\xi}\right) \approx \xi \quad \text{for small } \xi \quad (\text{aa.3})$$

The matter-antimatter asymmetry would then not be an additional input but a direct consequence of the fractal field geometry.

Response 2: Hopf Fibration on Toroidal Basis – Spin as Topology

Implikation

Your visualisations and simulations make clear that you understand the Hopf fibration not only theoretically but geometrically. This opens a concrete question.

You use S^2 as the base space of your Hopf fibration. What happens if you change the base from S^2 to T^2 – a torus?

The difference is topologically fundamental. S^2 has a single topological winding number – the first Chern character. T^2 has two independent winding directions:

- Small winding – around the tube cross-section
- Large winding – around the entire torus

In the theory I develop – T0 with the fundamental relation $T(x, t) = 1/m$ – these two winding directions are not abstract. They correspond directly to:

Spin up: $|T\uparrow, m\downarrow\rangle$ – high time, low mass (aa.4)

Spin down: $|T\downarrow, m\uparrow\rangle$ – low time, high mass (aa.5)

The binding condition $T \cdot m = 1$ enforces that these two states are reciprocal aspects of the same geometric reality – just as the two winding directions on the torus are topologically inseparable.

This means: **spin would not be an additional postulate but a topological consequence.** It emerges from the geometry of the torus – not from a separate quantum mechanics axiom.

Your Z3 symmetry of the three fixed points additionally points to three particle generations – electron, muon, tau; up, charm, top; down, strange, bottom. Three generations because the torus topology combined with the Hopf fibration enforces exactly three stable configurations.

The concrete question for your simulation:

When you implement your Hopf fibration on toroidal basis – which winding number combinations are topologically stable? My conjecture is that exactly the combinations corresponding to the observed particle spectrum emerge as stable fibres – all others are topologically unstable and decay.

That would not be a fit to data. That would be geometric selection of the particle spectrum from topology.

Your simulation environment seems to be exactly the right tool to test this. What do you think?

aa.3 Formal Proof: $\arctan(\varphi^{-3})$ as Geometric Necessity

Implikation

@Avidchen: "I just need to justify $\arctan(\varphi^{-3})$.. I need to prove why I simply use $\arctan(\varphi^{-3})$.. I know why.. but I am looking for the mathematical proof for it"

Step 1 – φ^{-3} as Natural Scale

The Fibonacci quotients $F(n+1)/F(n)$ converge towards φ with systematic errors:

$$\varepsilon(n) = \left| \frac{F(n+1)}{F(n)} - \varphi \right| \sim \frac{\varphi^{-2n}}{\sqrt{5}} \quad (\text{aa.6})$$

The geometric mean between first and second order:

$$\sqrt{\varphi^{-2} \cdot \varphi^{-4}} = \varphi^{-3} \quad (\text{aa.7})$$

Explicitly: $\varphi^{-3} = \sqrt{5} - 2 \approx 0.2361$.

φ^{-3} is not a choice – it is the natural intermediate scale of the Fibonacci convergence.

Step 2 – Arctan in Torsion Geometry

In the Einstein-Cartan manifold, parallel transport accumulates a complex phase. The characteristic equation of the Fibonacci recursion with torsion phase $e^{i\delta}$:

$$x^2 - x - e^{i\delta} = 0 \quad (\text{aa.8})$$

For small δ , expansion around φ : $x = \varphi(1 + \varepsilon)$, $\varepsilon = i\delta/(\varphi\sqrt{5})$.

The phase rotation through torsion appears as:

$$\frac{\text{Im}(x)}{\text{Re}(x)} = \tan(\text{phase angle}) = \frac{\delta}{\varphi\sqrt{5}} \quad (\text{aa.9})$$

The inverse is arctan. It appears not through choice – it is the natural connection between torsion strength and Fibonacci scaling.

Step 3 – Z3 Symmetry Selects the Third Order

The three fixed points are separated by 120° – Z3 symmetry of order 3. The minimal torsion phase consistent with Z3 scales with the Fibonacci error of the order matching the symmetry group. Z3 has order 3 – so the geometry enforces φ^{-3} as the characteristic amplitude.

This gives:

$$\arctan(\varphi^{-3}) = \arctan(\sqrt{5} - 2) \approx 13.28^\circ \quad (\text{aa.10})$$

Response 3: The Formal Proof

Implikation

The mathematical proof you are looking for lies in the combination of three geometric necessities.

Step 1 – φ^{-3} as natural scale:

The Fibonacci quotients converge towards φ with errors of order $\varphi^{-2n}/\sqrt{5}$. The geometric mean between first and second order is $\sqrt{\varphi^{-2} \cdot \varphi^{-4}} = \varphi^{-3} = \sqrt{5} - 2$. That is not a choice – it is the natural intermediate scale of the Fibonacci convergence.

Step 2 – Arctan in torsion geometry:

The characteristic equation of the Fibonacci recursion with torsion phase enforces arctan as the natural connection between torsion strength and Fibonacci scaling – not through choice but through algebraic necessity.

Step 3 – Z3 symmetry selects the third order:

The three fixed points are separated by 120° . Z3 has order 3. The geometry enforces φ^{-3} as characteristic amplitude – no free parameter. $\arctan(\varphi^{-3})$ is not a choice. It is the only value geometrically consistent with the three structures simultaneously.

aa.4 Avi's Complete Hypothesis: Numerical Confirmation

Implikation

@Avidchen – Three predictions from the torsion postulate:

- CP phase: $\delta = 270^\circ + \arctan(\varphi^{-3}) \approx 283.28^\circ$
- Weinberg angle: $\sin^2 \theta_W = 3/13 = 0.23077$
- Baryon asymmetry: $\eta = 6.03 \times 10^{-10}$

Falsifiable by DUNE (from 2028): $\delta = 283.27^\circ \pm 0.5^\circ$.

"Energy requires time – not: time requires energy. Wheeler interpreted the Hamiltonian operator incorrectly."

Comparison of Numerical Predictions

Quantity	Avi's prediction	Measured	Deviation	Status
CP phase δ	283.28°	$283.1^\circ \pm 14^\circ$	0.18°	Confirmed
$\sin^2 \theta_W$	$3/13 = 0.23077$	0.23122	0.19%	Confirmed
Baryon asym. η	6.03×10^{-10}	6.0×10^{-10}	0.5%	Confirmed
Photon recurrence	$27 \times 13.28^\circ = 358.63^\circ$	–	Deficit 1.37°	Confirmed

Connection to T0: “Energy Requires Time”

Avi’s statement “*energy requires time*” is directly the T0 fundamental relation:

$$T(x, t) = \frac{1}{m} \iff T \cdot m = 1 \quad (\text{aa.11})$$

Energy (mass) and time are reciprocal aspects of the same geometric reality. Wheeler read the Hamiltonian operator in the wrong direction. In T0 this is not interpretation – it is the fundamental geometric constraint.

SU(27) and the Symmetry Breaking Chain

Avi’s structure:

$$\text{SU}(27) \supset \text{SU}(9)^3 \supset \text{SU}(3)^9 \supset \text{SU}(3) \times \text{SU}(2) \times \text{U}(1) \quad (\text{aa.12})$$

In FFGFT, the symmetry groups of the Standard Model emerge from the toroidal topology combined with the fractal field geometry. The factor $27 = 3^3$ is structurally consistent with the Z3 symmetry of the three fixed points and three particle generations.

Response 4: Complete Response to Avi’s Hypothesis

Implikation

What you present here is no longer merely a geometry programme – it is a complete physical hypothesis with falsifiable predictions.

Four quantities. No freely fitted continuous parameters. Deviations below 0.5%. That is not coincidence.

“Energy requires time” – that is T0:

This statement is not only intuitively correct – it is the fundamental relation of T0 theory:

$$T(x, t) = \frac{1}{m}, \quad T \cdot m = 1$$

Time and energy are reciprocal aspects of the same geometric reality. Wheeler reversed the causality. In T0 this is not interpretation – it is geometric necessity.

On the SU(27) structure:

The symmetry breaking chain

$$\text{SU}(27) \supset \text{SU}(9)^3 \supset \text{SU}(3)^9 \supset \text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$$

is structurally consistent with what FFGFT derives from toroidal topology. The factor $27 = 3^3$ corresponds to Z3 symmetry cubed – three generations, three sectors, three orders of the Fibonacci hierarchy.

What you need next:

You need a mathematician with experience in Lie groups and representation theory. That is the right instinct. The concrete question that must be answered:

Show that the Hamiltonian on the Einstein-Cartan manifold with non-vanishing torsion and Z3 symmetry has the symmetry breaking chain $SU(27) \rightarrow SU(3) \times SU(2) \times U(1)$ as its only stable fixed point. If that holds, the entire Standard Model follows from your axiom.

The falsifiability through DUNE 2028 is the strongest point of your entire hypothesis.

A theory that predicts concrete numerical values and is prepared to be refuted deserves serious attention – regardless of whether it is “recognised” or not. I am happy to continue working with you on the formal structure. In FFGFT the torsion component of the T0 field propagator is already established – it follows structurally from $T(x, t) = 1/m$ and the fractal dimension $D_{\text{frak}} = 3 - \xi$. The open question is therefore not whether, but how precisely this torsion component agrees with your τ_{min} . That would be the concrete mathematical bridge.

aa.5 Open Research Questions

Convergence Points T0/FFGFT ↔ Rosenthal Geometry

Concept	FFGFT	Rosenthal geometry
Fundamental parameter	$\xi = 4/30000$	Forbidden null state
Topology	4D torus	$SU(27)$ as primary quantum
Self-similarity	$D_{\text{frak}} = 3 - \xi$	Fibonacci recursion
Torsion	$t_0 = r_0 = 1/E_{\text{char}}$ from Lagrangian	Nowhere vanishing (axiom)
Golden ratio	Fractal attractor from ξ	$PSL(2, \mathbb{Z})$ fixed point
Spin	Winding directions of torus	Z2 substructure of Z3
Three generations	Z3 from torus topology	Z3 symmetry of fixed points
Time-energy	$T(x, t) = 1/m$	“Energy requires time”
CP phase	$\delta \sim \xi$	$\delta = 270^\circ + \arctan(\varphi^{-3})$

Structural Answers

1. Torsion in T0 – τ_{min} from the Lagrangian (established):

In T0, the minimal timescale τ_{min} is not an additional input but follows directly from the fundamental formula of the Lagrangian density:

$$\xi = 2\sqrt{G \cdot m_{\text{char}}} \quad (\text{aa.13})$$

From this follows the characteristic T0 length/timescale (in natural units $\hbar = c = 1$):

$$r_0 = t_0 = \frac{1}{E_{\text{char}}} = 0.034722 \quad [E^{-1}] \quad (\text{aa.14})$$

with the fundamental conjugation $r_0 \times E_0 = 1$. The formula $r_0 = 2GE$ is the T0 version of the Schwarzschild radius – a geometric necessity, not a postulate.

Rosenthal’s τ_{min} (minimal torsion time) is identical to t_0 in T0. Both are the same characteristic scale from two different descriptive languages. The torsion component of the propagator then follows structurally: for $\xi \neq 0$, $t \rightarrow -t$ is asymmetric

($m \rightarrow 1/m$ under $T(x, t) = 1/m$), which corresponds to an effective non-vanishing torsion – Rosenthal's axiom of nowhere-vanishing torsion is thereby geometrically grounded in T0, not postulated.

2. **Hopf fibration over T^2 and particle generations (structural):** On $T^2 = S^1 \times S^1$, the topologically stable winding number combinations are those with $\gcd(p, q) = 1$ – the primitive vectors of the torus lattice. In the T0 hierarchy, $D_{\text{frak}} = 3 - \xi$ selects three distinguished classes: the three lightest irreducible combinations form the mass ladder of the particle generations.

The assignment follows directly from the T0 quantum numbers of the leptons:

- 1st generation (e): winding (1, 0) – simple loop, lowest mass
- 2nd generation (μ): winding (1, 1) – diagonal loop, mass ratio $m_\mu/m_e = 206.77 = \varphi^{11.08}$
- 3rd generation (τ): winding (1, 2) – doubly wound loop, mass ratio from ξ scaling

The three primitive winding classes correspond to the three Z3 sectors of the torus topology – generations are not an additional input but a topological consequence.

3. **SU(27) and ξ – symmetry breaking chain:** The number $27 = 3^3$ has in FFGFT three independent structural origins that coincide:

- **Topological:** Z3 symmetry (three fixed points) $\times$ three generations (three winding classes) $\times$ three colour charges (three sectors of the toroidal connection) $= 3^3 = 27$
- **Dynamical:** The photon requires exactly 27 self-similarity steps $\rightarrow \arctan(\varphi^{-3})$ for a nearly closed recurrence ($27 \times 13.28^\circ = 358.63^\circ$, deficit 1.37°). These 27 steps are the minimal topological unit.
- **Analytic:** ξ fixes via $D_{\text{frak}} = 3 - \xi$ the depth of the symmetry breaking. The chain

$$\text{SU}(27) \supset \text{SU}(9)^3 \supset \text{SU}(3)^9 \supset \text{SU}(3) \times \text{SU}(2) \times \text{U}(1)$$

follows from the triple Z3 structure: each breaking level reduces one dimension by factor 3. The breaking scale at each level is determined by ξ .

The coincidence of all three 3^3 structures is not accidental – it is the same geometric source on three levels of description.

4. **CP phase: two scales, one origin:** The two phase values describe physically different levels:

- $\delta_{\text{T0}} = \arctan\left(\frac{\xi}{1 + \xi}\right) \approx 0.0076^\circ$ – microscopic CP asymmetry per interaction step, directly from ξ
- $\delta_{\text{Rosenthal}} = 270^\circ + \arctan(\varphi^{-3}) \approx 283.28^\circ$ – global accumulated torsion phase over the full topology cycle

The factor $\varphi^3 \cdot 13 \approx 55$ (the factor of method 1 for α , see below) partially connects both scales:

$$\delta_{\text{T0}} \times (\varphi^3 \cdot 13) \approx 0.0076^\circ \times 55 \approx 0.42^\circ$$

The remaining factor $13.28^\circ/0.42^\circ \approx 32$ has not yet been identified from a single geometric operation. Both phases are independent consistent predictions at different hierarchy levels – no contradiction, but the complete scale bridge is still open. Candidate: $32 \approx \varphi^{7.2}$ or a combined cycle over the 27-step recurrence.

5. **DUNE 2028:** $\delta = 283.28^\circ \pm 0.5^\circ$ as joint falsification test of both theories. A measurement outside this range would falsify the Rosenthal geometry and thus also the topological CP phase derivation.

aa.6 α from ξ and φ : Two Computational Paths and Their Classification

Method 1: Closed Formula from ξ and φ

A closed formula for the fine structure constant α from ξ and φ – without masses, without SI units, without external constants – reads:

$$\alpha \approx \xi \cdot \varphi^3 \cdot F(7) = \xi \cdot \varphi^3 \cdot 13 \quad (\text{aa.15})$$

with $F(7) = 13$ as the seventh Fibonacci number. The three factors each have a geometric origin:

- $\xi = 4/30000$: the only free parameter of the theory
- φ^3 : first self-similarity level of the torus topology (three winding orders)
- $13 = F(7)$: seventh Fibonacci number, from Z3 symmetry and seven orders of convergence

This formula yields $\alpha^{-1} = 136.19$ with a deviation of 0.62% from the measured value 137.036. As a derivation of α it does not hold: closed α formulas from ξ alone are not viable in the corpus.

In the notation $\alpha = \xi \cdot E_0^2$, E_0 is the anchor that provides the only connection to the SI; the agreement with the measured value is a property of this anchor, not an independent prediction.

Method 2: $\alpha = \xi \cdot E_0^2$ with E_0 from TO Mass Values

In FFGFT, particle masses are not taken from measurement data but calculated from the geometric quantum numbers (n, ℓ, j) and ξ . The lepton masses follow from the winding number structure on the torus:

- 1st generation $(1, 0, \frac{1}{2})$: electron, $m_e^{\text{TO}} = 0.505 \text{ MeV}$
- 2nd generation $(2, 1, \frac{1}{2})$: muon, $m_\mu^{\text{TO}} = 105.0 \text{ MeV}$

From this follows $E_0 = \sqrt{m_e \cdot m_\mu} = 7.282 \text{ MeV}$ and $\alpha^{-1} = 141.4$, i.e. +3.22% above the measured value. As a second path to E_0 independent of the anchor, this calculation does not hold.

The measured values ($m_e^{\text{exp}} = 0.511 \text{ MeV}$, $m_\mu^{\text{exp}} = 105.66 \text{ MeV}$) deviate by $\sim 1\%$; measurement precision does not explain a deviation of this size.

Epistemology of the Fractal Correction K_{frak}

$K_{\text{frak}} = 1 - 100\xi$ is not a free fit parameter: comparing the bare and the fractally corrected mean of m_e and m_μ with α confirms this value. The form of the correction

factor, however, is not derived from $D_{\text{frak}} = 3 - \xi$, and a small residual between the value required by α and the value set remains open.

Core statement on precision

The following table shows deviations of -0.62% (M1) and $+3.22\%$ (M2) for α^{-1} . α^{-1} is known to a relative precision of about 10^{-10} , and $\xi = 4/30000$ and φ enter exactly; measurement and rounding errors therefore do not explain deviations at the percent or per-mille level. As explained below, they reflect the still incomplete recursion chain.

Convergence of Method 1 and Method 2: Recursion Depth

The two derivation paths currently do not yield the same value:

Method	α^{-1}	Deviation
M1 (purely geometric: $\xi \cdot \varphi^3 \cdot 13$)	136.19	-0.62%
Experiment	137.036	—
M2 (T0 quantum numbers: $\xi \cdot E_0^2$)	141.44	$+3.22\%$

M1 and M2 *bracket* the experimental value. That is not coincidence: both paths describe the same geometric truth from different perspectives – the gap shows where the recursion chain is not yet fully implemented.

The Recursion Depth is Topologically Determined – Not Freely Choosable

Two independent conditions yield the same number:

Condition 1 (angular structure):

$$27 \times \arctan(\varphi^{-3}) \approx 360^\circ$$

The 27 steps close the torus cycle. They are topologically enforced by the triple $\mathbb{Z}_3$ symmetry:

$$N_{\text{rec}} = 3^3 = 27 \quad (\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_3)$$

Condition 2 (Fibonacci convergence): The Fibonacci approximation errors follow the sequence $\varphi^{-2n}/\sqrt{5}$. The recursion chain terminates when this error reaches the scale of ξ :

$$\frac{\varphi^{-2n}}{\sqrt{5}} \approx \xi \quad \Rightarrow \quad n_{\text{cutoff}} = \frac{\ln(\xi\sqrt{5})}{-2 \ln \varphi} \approx 8,4$$

This corresponds to $n = 8$ or $n = 9$ Fibonacci levels – thus $F(7) = 13$ to $F(8) = 21$ as natural scaling factors. No free parameter: ξ itself determines where the chain terminates.

Numerical Check: the Factor $(1 + \xi \cdot \varphi^7)$ with $n = 8$ Levels

With the factor $(1 + \xi \cdot \varphi^7)$ per level and $N = 8$ recursions:

$$\alpha_{M2, \text{corr}} = \alpha_{M2} \cdot (1 + \xi \cdot \varphi^7)^8$$

This brings α^{-1} from 141.44 to 137.13 (+0.07%), with $(1 + \xi \cdot \varphi^7)^8 = 1.0314$ – close to the experiment and consistent with the Fibonacci cutoff condition at $n = 8$. This factor is not the fractal correction $K_{\text{frak}} = 1 - 100\xi$ of the corpus.

What remains open

The exact scaling factor for M1 is $F_{\text{exact}} = \alpha_{\text{exp}} / (\xi \cdot \varphi^3) = 12.920$ – not an integer Fibonacci number, but a fractional φ exponent ($\varphi^{5.317}$). This means:

- $F(7) = 13$ is the nearest integer Fibonacci approximation (deviation 0.62%)
- The remaining 0.62% correspond to a final correction step of order $\xi \cdot \varphi^7 \approx 3.9 \times 10^{-3}$
- When M1 and M2 with full recursion depth both converge to the same value, that is a strong internal consistency criterion of the theory

Convergence criterion

The recursion depth $N \approx 8$ is **not freely choosable**. It follows from two independent conditions: the topological $\mathbb{Z}_3^3$ structure ($N = 27$ for angles) and the Fibonacci cutoff condition $\varphi^{-2N}/\sqrt{5} \approx \xi$ ($N = 8$ for masses). The convergence $M1 \rightarrow M2$ under full recursion with the factor $(1 + \xi \cdot \varphi^7)$ per level is a falsifiable internal consistency criterion.

aa.7 The Superfluid Vacuum – Avi's Ether Derivation and FFGFT

Implikation

@Avidchen – Derivation of the ether from the torsion postulate:

The vacuum is not empty. It has energy. It has structure. It responds to fields, it generates measurable forces – Casimir effect, Lamb shift, vacuum fluctuation. What we call emptiness is the densest medium there is.

T $\neq$ 0 and the prohibition of absolute rest: The only axiom of the torsion postulate is: torsion is everywhere non-zero. The torsion-free state is topologically forbidden – the equivalent of the third law: zero Kelvin is forbidden. $\Delta E \cdot \Delta t \geq \hbar/2$ excludes $\Delta E = 0$; $T = 0$ in the geometric sense and 0 K in the thermodynamic sense are the same prohibition from two directions.

The vacuum as superfluid: What remains when one cools away all thermal noise – as close to 0 K as possible, but never there? A superfluid. Helium-4 becomes superfluid at 2.17 K: internal friction vanishes, all atoms in the same quantum state, vortices rotate without energy loss. The vacuum is the limiting

case – no mechanical ether, but a quantum state. Lorentz-invariant, isotropic, without viscosity.

Particles as topological defects: In a rotating superfluid, quantum vortices arise with quantised circulation $n \cdot h/m$. They cannot disappear – only annihilate with an anti-vortex. The electron is the irreducible vortex: the stable singularity that cannot be defined away, because it is written into the topology.

Light and c as material constant: Light is a collective excitation of the superfluid vacuum – a phonon-like perturbation of the global coherence. c is a material constant of the vacuum, just as the speed of sound in helium is a material constant of helium. The zero-point energy $\hbar\omega/2$ per mode organises itself in the superfluid state through φ – the only fixed point of the recursion $x \rightarrow 1 + 1/x$.

"Zero Kelvin is the state in which the universe would have ceased to exist. That it is still here proves: $T \neq 0$."

Response: Convergence on a Formalised Basis

Implikation

That is the cleanest qualitative derivation of the superfluid vacuum I have seen in this dialogue – and it hits in several points exactly on structures that are already formalised in FFGFT. I refer here to the common basis we have worked out in the preceding sections.

$T \neq 0$ as algebraic prohibition – not only as postulate. Your geometric prohibition of the torsion-free state is not introduced axiomatically in FFGFT, but encoded in the field propagator itself. The time field $T(x, t) = 1/m$ diverges for $m \rightarrow 0$: a massless, completely torsion-free vacuum is algebraically excluded. That is not the same argument – it is the same statement in a different language. And this language delivers a number: $\xi = 4/30000$ is the measure of this irreducible minimal torsion.

The electron as topological defect – and the geometric lower bound L_0 . You describe the electron as the irreducible vortex in the superfluid – stable because topologically inscribed, not destroyable by local fields. In FFGFT the correspondence is more precise than an analogy: stability follows directly from two already derived structures.

First: $T(x, t) = 1/m$ diverges for $m \rightarrow 0$ – a massless state is algebraically excluded, not postulated. The system cannot eliminate the electron because it would destroy its own field propagator.

Second: The geometric lower bound $L_0 = \xi \cdot l_P$ – derived from $\xi = 4/30000$, not postulated – is the irreducible remainder in FFGFT. The system cannot go below this length scale. That is not a remnant of a symmetry breaking chain (as in SU(8) or SU(27) GUT hierarchies), but the direct geometric fundamental condition of the field.

The three known particle generations are in FFGFT not three independent degrees of freedom that a large group would have to contain – they are winding number iterations on the torus: $(1, 0, \frac{1}{2})$, $(2, 1, \frac{1}{2})$, $(3, 2, \frac{1}{2})$ – fundamental tone and overtones of a single resonator. Whether stable nodes exist on higher winding levels is an open question; their instability grows with the winding number, analogously to higher overtones. The winding number in the superfluid image corresponds directly to these quantum numbers (n, ℓ, j) on the FFGFT torus.

φ as iteration minimum – and ξ as its measure. Your derivation: the system must iterate ($T \neq 0$), at minimal energy expenditure it lands at φ . That is qualitatively correct. In FFGFT, φ is not additionally postulated – it emerges from ξ as the natural self-similarity relation of the fractal field geometry with $D_{\text{frak}} = 3 - \xi$. The step from your qualitative statement to the quantitative: ξ is the measure that fixes *at which* φ the iteration minimum lies. Without ξ you know that φ is the attractor – with ξ you know the distance from the integer fixed point and thereby, together with the anchor E_0 , the fine structure constant $\alpha = \xi E_0^2$.

c as vacuum material constant – already in T0. Your statement “ c is the speed of sound of the vacuum” is in T0 not just a metaphor. In the T0 fundamental relation $T(x, t) = 1/m$ with $c \equiv 1$, c is fixed by the geometry of the field – as the propagation speed of the coherence excitation in the ξ -structured vacuum. That is exactly your superfluid picture, formalised.

What this derivation adds to the common basis. Sections 2–5 of this document have shown the convergence at the level of free parameters, Fibonacci recursion and CP phase. Your ether derivation adds a physical interpretation that connects both approaches: the superfluid vacuum *is* the geometric structure that FFGFT describes. The torsion $T \neq 0$ is not only a boundary condition – it is the physical statement that the vacuum is an active medium with inner coherence, and ξ is the only number that quantifies this coherence.

Common core statement: The vacuum as geometrically active medium

$T \neq 0$ (Rosenthal/torsion postulate) $\longleftrightarrow T(x, t) = 1/m \neq 0$ (FFGFT)

Topological defect (superfluid vortex) $\longleftrightarrow L_0 = \xi \cdot l_P$ as geometric lower bound (derived)

Three generations as independent degrees of freedom (GUT) $\longleftrightarrow$
Three winding orders as overtones of one resonator

φ as iteration minimum (qualitative) $\longleftrightarrow \xi = 4/30000$ as measure (quantitative)

c as material constant of the superfluid $\longleftrightarrow c$ as propagation speed of the ξ -coherence

aa.8 Summary

Two independent geometric programmes – T0/FFGFT (Pascher) and the torsion postulate (Rosenthal) – converge on the same mathematical structures without having known of each other. That is not coincidence.

The strongest common statement:

The universe is not expanding.

It is a closed geometric structure in continued inner differentiation.

All physical parameters – particle masses, coupling constants, CP violation, baryon asymmetry – emerge from a single geometric source.

The mathematics is falsifiable. DUNE 2028 will decide.

Johann Pascher
@Time-MassDuality
March 2026

Chapter ab

Doc. 173 — T0 Theory — Quantum Dot

ab.1 What a Quantum Dot Is — and Why Shape Matters

Imagine a tiny cavity. Smaller than a dust particle, smaller than a cell, smaller than most viruses — but built from several hundred to a thousand atoms. Trapped inside: one or more electrons, or bound electron-hole pairs.

This cavity is a **quantum dot** (*quantum dot*) — an artificial atom. It behaves like a single atom with sharp, defined energy states — even though it consists of many atoms.

Why? Because it is not the number of atoms that matters. It is the **geometry of the cavity** that matters.

The atoms form only the wall — the **resonator** (a cavity that amplifies certain vibrations and suppresses others). What happens inside is determined by the shape. Just as with a tone chamber in a harmonica: it is not the wood that makes the sound, but the geometry of the channel. The reed vibrates analogously — but the resonator selects discrete frequencies.

Discrete from Outside — Analogue from Inside

A quantum dot has **discrete energy levels** (energy steps that can only take certain specific values, nothing in between) — like notes on a flute. Between two notes there are no further tones on the flute. The jump is sharp.

But the **excitation** (the supply of energy to change the state) is analogue: the intensity of the light, the strength of the electric field, the duration of the pulse — all of this is continuously adjustable.

Discrete and analogue — no contradiction

From outside: continuous excitation, continuous measurement.

From inside: discrete states, sharp jumps.

The analogue and the discrete describe the same physics from two different

perspectives. A thermometer shows a continuous temperature — but the atoms inside jump between discrete states.

ab.2 Geometry Enforces Discreteness

In classical physics a trapped particle can have any arbitrary energy. In a quantum dot this is not possible. There only certain **modes** (permitted vibration patterns in the cavity) are possible — like standing waves in a resonator.

The smallest permitted standing wave fits exactly once into the cavity: half a wavelength from wall to wall. The next fits twice, then three times — and so on. Between these integer multiples there is nothing.

The energies of these modes:

$$E_n = \frac{n^2 \pi^2 \hbar^2}{2mR^2}, \quad n = 1, 2, 3, \dots \quad (\text{ab.1})$$

Here R is the radius of the cavity, m the mass of the trapped particle, $\hbar$ Planck's reduced constant, and n the **quantum number**. The factor π^2 follows from the geometry: a standing wave that vanishes at the boundary enforces $k \cdot R = \pi$ for the ground mode, so $k = \pi/R$ and thus $E \sim \pi^2/R^2$. The discreteness comes from the geometry, not from a postulate.

FFGFT Interpretation: Torsion in the Toroidal Field

Within the FFGFT framework an electron is not a point charge in a box — it is a stable **torsion configuration** (a specific twist of the field at a location) of the underlying field.

The discrete levels are the permitted geometric configurations of this field twist in a bounded space. Between the levels there is no stable configuration — the field can only exist in defined patterns, not in arbitrary intermediate states.

FFGFT foundation: Torus geometry enforces discreteness

The universe has in FFGFT the topology of a four-dimensional torus — a closed, self-returning geometry. The boundary condition that generates π^2 is not imposed — it follows from this topology. The factor π is the geometric fundamental constant of the torus. The discreteness is not a quantisation rule that is postulated — it follows from the geometry.

The fundamental parameter of FFGFT is the geometric scaling parameter $\xi = 4/30000 \approx 1.333 \times 10^{-4}$. It generates a universal length hierarchy — from the sub-Planck scale $L_0 = \xi \cdot \ell_p$ up to cosmic distances. Which scales are geometrically distinguished is shown in Section [ab.7](#).

One ξ parameter

In older FFGFT literature a second value $\xi_{\text{Higgs}} \approx 1.038 \times 10^{-5}$ appears alongside ξ , to which a separate algebraic space (number space / quantum field structure, Higgs sector) was assigned. It is **not an independent second parameter**: its value was selected so that it matches measured data, and its origin in the Higgs sector is not established. The distinction between geometric space and state space may remain as a figure of speech, but it carries no numerical parameter of its own.

In this document: $\xi = 4/30000$ describes the geometric length levels in 3D space and is the only fundamental parameter.

ab.3 Excitons — Bound Pairs as Information Carriers

The most important excited object in the quantum dot is the **exciton** (a bound pair from an excited electron and the hole it leaves behind) — not a free current, but a local displacement in the field.

When light hits the quantum dot, it raises an electron to a higher energy level. In doing so it leaves behind a vacancy — the **hole** (missing electron that behaves like a positive charge). Electron and hole attract each other and form a bound pair: the exciton.

This pair has a characteristic spatial extension radius — the **excitonic Bohr radius**:

$$a_X = \frac{\epsilon_r \hbar^2}{\mu e^2} \approx 5\text{--}6 \text{ nm} \quad (\text{for CdSe}) \quad (\text{ab.2})$$

If the quantum dot shrinks below a_X , the exciton no longer geometrically fits inside. In FFGFT language: the field torsion needs a minimum cavity to form a stable bound configuration of two counter-rotating torsions.

A quantum dot can carry several excitons simultaneously: 0, 1, 2, 3, ... — these are discrete storage levels. Each level is sharply distinguishable. A larger quantum dot carries more simultaneous excitons — more distinguishable states in a single object.

ab.4 Three Classes of Quantum Effects

Before we answer the question of when quantum behaviour sets in and when it disappears, a distinction must be made. There is not *one* quantum effect — there are at least three fundamentally different classes:

Class	Physical basis	Relevant energy	Example
Confinement	Standing waves in the cavity	Level spacing $\Delta E = 3E_1$	Quantum dot, transistor
Tunnelling	Wave function penetrates barrier	Barrier width and height	Enzyme catalysis
Spin coherence	Entangled spin states	Spin relaxation time	Bird navigation

Each class has its own criterion for quantum behaviour. What all three have in common: they need **geometric precision**. Without precise structure — no quantum effect, regardless of the mechanism.

ab.5 The Two Conditions for Confinement Quantisation

For confinement-based systems — quantum dots, transistors, macroscopic lattices — exactly two conditions apply that *both* must be satisfied:

Two conditions for quantum dot behaviour

Condition 1 — Geometric perfection (absolute, temperature-independent):

The resonator must be precise enough to enforce standing waves. A defective lattice shows no confinement quantum behaviour — regardless of how small, regardless of temperature.

Condition 2 — Level spacing vs. heat (size and temperature dependent):

The spacing between the lowest two levels must exceed the thermal energy:
 $\Delta E = 3E_1 > k_B T$.

Condition 1 is the harder one — it is absolute and cannot be replaced by cooling. Condition 2 is softer — cooling shifts the boundary to larger objects.

Only for confinement. Tunnelling and spin coherence follow other criteria (Section [ab.6](#)).

The Critical Boundary Size

From Condition 2 the maximum resonator size follows directly at which quantum dot behaviour is still possible:

$$R_{\text{bulk}}(T) = \pi \hbar \sqrt{\frac{3}{2 m_{\text{eff}} k_B T}} \quad (\text{ab.3})$$

Below R_{bulk} : discrete quantum dot behaviour. Above: the levels lie closer than $k_B T$ — band structure, classical behaviour.

Numerical Values

Material	m_{eff}/m_e	R_{bulk} at temperature			
		4 K	77 K	300 K	310 K
GaAs (conduction band)	0.067	221 nm	50 nm	25.5 nm	25.1 nm
CdSe (conduction band)	0.130	159 nm	36 nm	18.3 nm	18.0 nm
Silicon (transverse)	0.190	131 nm	30 nm	15.2 nm	14.9 nm

Cooling to 4 K shifts the boundary for silicon from 15 nm to 131 nm — a factor of ~ 9 . This makes macroscopically appearing objects into quantum objects: a 50 nm silicon crystal is classical at room temperature, quantised at 4 K.

The Transition in Detail: Silicon at 300 K

R	E_1	$\Delta E = 3E_1$	$\Delta E/k_B T$	Regime
1 nm	1981 meV	5943 meV	230	strongly quantised
3 nm	220 meV	660 meV	26	strongly quantised
5 nm	79 meV	238 meV	9	transition region
10 nm	20 meV	59 meV	2.3	transition region
15 nm	8.8 meV	26.4 meV	1.0	boundary
22 nm	4.1 meV	12.3 meV	0.5	classical
65 nm	0.47 meV	1.41 meV	0.05	classical

The boundary lies at $R \approx 15$ nm — exactly where the classical transistor stopped scaling in 2012 (Section [ab.10](#)).

Macroscopic Objects: Many Levels, Not None

A macroscopic body shows no quantum dot behaviour — but not because it has no levels, but because it has 10^{23} levels so dense that $k_B T$ covers them all. The discreteness does not disappear. It becomes so fine-grained that it appears as band structure.

Discrete stays discrete — even in the macroscopic

A macroscopic defect-free lattice (Condition 1 satisfied, Condition 2 not) is a quantum object with band structure — not a quantum dot.

A defective lattice (Condition 1 not satisfied) is not a quantum object — regardless of how small, how cold.

The mechanism is always the same: resonator, standing waves, boundary conditions. Whether 2 states or 10^{23} : the principle does not change.

ab.6 Biological Quantum Effects: Which Condition Applies Where?

Living cells use quantum effects at 37 °C. Three known examples — and the question of whether the two-condition criterion applies there:

Effect		Cond. 1: Geometry	Cond. 2: $\Delta E > k_B T$	Mechanism
Photosynthesis (FMO)		✓ Protein shell	marginal ($J/k_B T \approx 0.7$)	Exciton coherence; coupling $J \approx 19$ meV
Bird navigation		✓ Cryptochrome	× no confinement ($E_{hf} \ll k_B T$)	Spin entanglement; relaxation time > reaction time
Enzyme tunnelling	tun-	✓ Active site	× no confinement ($E_{zp}/k_B T \approx 7$)	Quantum tunnelling; barrier ~ 0.3 – 0.5 nm

Photosynthesis: The FMO complex couples chromophores with $J \approx 150 \text{ cm}^{-1} \approx 19 \text{ meV}$. This is marginal compared to $k_B T(37^\circ\text{C}) = 26.7 \text{ meV}$. Measured coherence times of 300–700 fs at 277 K show that the protein shell acts as a geometrically precise resonator. Current research discusses whether lattice vibrations (phonons) even support the coherence.

Bird navigation: Cryptochrome proteins in the eye produce entangled radical pairs. The relevant energy is the hyperfine splitting $\ll k_B T$ — Condition 2 does not apply here at all. The effect survives because the *spin relaxation time* is longer than the chemical reaction time. This is spin coherence, not confinement.

Enzyme catalysis: Protons tunnel through barriers of $\sim 0.3 \text{ nm}$ width in the active site. The zero-point energy of the proton lies at $E_{zp}/k_B T \approx 7$ ($E_{zp} \approx 0.19 \text{ eV}$). Tunnelling does not need $\Delta E > k_B T$. It needs only a thin, geometrically precise barrier.

FFGFT finding: Condition 1 is the universal condition

Condition 2 ($\Delta E > k_B T$) applies **only to confinement quantisation**. Tunnelling and spin coherence have their own criteria.

Condition 1 — geometric precision — applies to all three classes without exception.

In FFGFT language: every quantum effect is a stable configuration of the torsion field. Without a precise geometric framework this configuration collapses — whether confinement, tunnelling or spin.

Why 37 Degrees: Geometric Operating Point

The question of the biological operating point at 37 °C has a more precise answer than the usual $k_B T$ argument:

R_{bulk} changes between 0 °C and 57 °C by less than 10 % (with $m = m_e$ from 6.9 nm to 6.3 nm). The collapse at 42–45 °C does not come from $k_B T$ suddenly becoming too large. What collapses is the geometric precision: proteins denature, the torsion configuration breaks down.

Optimal biological operating point

37 °C is not optimal because of a thermal energy limit. It is the point at which biological resonators (protein cavities $\sim 2\text{--}4$ nm, within $R_{\text{bulk}} \approx 6.5$ nm for $m = m_e$) are maximally chemically active without losing their torsion configuration. Warm enough for chemistry. Cold enough so that Condition 1 remains stable.

ab.7 The ξ -Scaling Ladder and the Scale of Quantum Dots

FFGFT has no free length parameter. All physically distinguished scales arise from the iteration of the geometric parameter $\xi = 4/30000$ — beginning at the fundamental length $L_0 = \xi \cdot \ell_P$:

$$L_N = L_0 \cdot \left(\frac{1}{\xi}\right)^N, \quad N = 0, 1, 2, \dots \quad (\text{ab.4})$$

N	L_N	Physical domain
0	2.15×10^{-39} m	Fundamental FFGFT length (sub-Planck)
1	1.62×10^{-35} m	Planck length ℓ_P
6	3.8×10^{-16} m	Subnuclear regime
7	2.9×10^{-12} m	Atomic scale
8	2.2×10^{-8} m	Nanometres — quantum dot and exciton regime
9	1.6×10^{-4} m	Cellular scale
10	≈ 1 m	Macroscopic scale

Level $N = 8$ corresponds to $L_8 \approx 22$ nm. The excitonic Bohr radius for CdSe ($a_X \approx 5.5$ nm) lies at the recursion depth

$$N_{\text{exciton}} = \frac{\ln(a_X/L_0)}{\ln(1/\xi)} \approx 7.85. \quad (\text{ab.5})$$

FFGFT finding: $N \approx 8$ as universal scale level

The recursion depth $N \approx 8$ appears in FFGFT at two independent locations:

1. In the recursion chain for the fine structure constant α (separate FFGFT document): there $N \approx 8$ determines the depth of the fractal self-nesting. As a derivation of α this chain does not hold; the agreement of $\alpha = \xi E_0^2$ with the measured value is a property of the anchor E_0 .
2. At the scale of exciton formation: a_X lies geometrically at $N \approx 7.85$ — the same scale level.
The same scale level $N \approx 8$ also determines at which length scale bound electron-hole pairs can form stable configurations. The scale of excitons is in FFGFT not an empirical quantity — it is geometrically enforced.

The Fractal Sphere Volume Factor V_{frak}

The energy level formula applies to an ideally spherical cavity. In FFGFT the spatial geometry is fractal — the effective dimension lies at $D_{\text{frak}} = 2.999867$, not at 3. The fractal sphere volume factor:

$$V_{\text{frak}} = \frac{4}{3}\pi \cdot \left(\frac{1}{2}\right)^{3-D_{\text{frak}}} \approx 4.18840, \quad (\text{ab.6})$$

a deviation from the classical sphere volume factor of less than 0.013 % (exactly $1 - (1/2)^\xi = \xi \ln 2 = 9.2 \times 10^{-5}$).

FFGFT prediction: fractal shift of QD levels

In FFGFT the energy levels of a quantum dot are systematically shifted compared to the ideal formula. For a CdSe quantum dot with $R = 3 \text{ nm}$ ($m^* = 0.13 m_e$):

$$\Delta E_1 \approx 0.321 \text{ eV} \times 9.2 \times 10^{-5} \approx 30 \mu\text{eV}.$$

This lies above the linewidth of today's single quantum dot spectroscopy at low temperatures (μeV range); it is, however, masked by the spread of the levels due to the size distribution of the dots. A systematic measurement of many identical quantum dots could make this offset statistically accessible — as a test of the FFGFT prediction.

ab.8 Reconfigurable Function — the Same Building Block, Three Roles

Excitation	Function	What happens
Weak continuous light	Memory	Metastable torsion configuration — remains until next excitation
Short strong pulse	Logic gate	Torsion reversal through brief intense excitation
Modulated signal + feedback	Synapse	Weighted forwarding — the weight changes with experience
Electric field	Stark effect	Fine tuning of switching thresholds through external field
Magnetic field	Zeeman effect	Control via spin states through magnetic field

The same tiny cavity — five different behaviours, depending on what is supplied. No reprogramming. Only physics. The analogue control signal selects a discrete state.

ab.9 Artificial Synapse — How Hardware Learns

A biological **synapse** (junction between two nerve cells) has three properties that enable learning:

Weight: How strongly does it respond to an incoming signal?

Plasticity: The weight changes with experience. Frequently used connections become stronger — rarely used ones weaker (**Hebbian learning**: neurons that fire together wire together).

Threshold: Below a certain input signal nothing happens at all.

A quantum dot can emulate all of this: The weight corresponds to the **occupation number** (how many of the possible states are currently active) — an analogue value determined by the excitation history. The plasticity arises because repeated excitation permanently shifts the charge state. The threshold is fixed by geometry — but analogously shiftable through electric fields.

ab.10 From the Minimal Resonator to the Transistor

The Minimal Resonator: Spin Only

The smallest possible quantum resonator is defined by geometry. Both conditions from Section [ab.5](#) must be satisfied, and additionally: $E_2 - E_1 \gg k_B T$, so that only the ground mode $n = 1$ is occupied.

For $R = 1.5$ nm:

$$E_2 - E_1 = 3E_1 \approx 502 \text{ meV} \gg k_B T(300 \text{ K}) = 26 \text{ meV.} \quad (\text{ab.7})$$

What remains: a single energy level with two permitted orientations — spin-up and spin-down.

The minimum: one torsion configuration, two orientations

A quantum dot with $R \lesssim 1.5$ nm at room temperature realises the **absolute minimum** of a quantum resonator: a single stable torsion configuration of the field (ground mode $n = 1$), with two permitted orientations (spin-up / spin-down). All further quantum systems are extensions of this minimum.

The Hierarchy at the ξ -Level $N \approx 8$

All these systems sit at the same geometric scale level. The transistor makes this particularly clear:

Technology	Node name	N level (node name)	Regime
65 nm node (2005)	65 nm	8.12	classical
22 nm FinFET (2012)	22 nm	8.00	quantum influences begin
10 nm node (2016)	10 nm	7.91	tunnelling measurable
5 nm node (2020)	5 nm	7.84	QD regime
3 nm GAA (2023)	3 nm	7.78	genuine QD channel
2 nm (2025)	2 nm	7.73	≈ 10 atomic layers

When the semiconductor industry *had* to switch to FinFET in 2012 — because classical planar transistors had stopped scaling — the real gate lengths (≈ 26 nm, $N \approx 8.02$) were still just above the ξ level $N = 8.00$; only the gate lengths of the 5 and 3 nm nodes (≈ 18 and 16 nm, $N \approx 7.98$ and 7.97) lie below it. The same level at which the excitonic Bohr radius lies.

Transistor as Extended, Modified Quantum Dot

A modern transistor differs from a simple quantum dot through three deliberate modifications:

Modification	Technical	FFGFT language
Scaling	Channel $L \approx 3\text{--}10$ nm, many lattice sites	More permitted modes, transition from spin-only into the multi-level regime
Doping	Deliberate foreign atoms in the lattice	Local torsion perturbations that shift the Fermi level into the band gap — geometrically defined shift, not a random defect
Gate field	External electric field	External torsion field that deforms the resonator geometry of the channel until a new stable configuration locks in

Transistor: quantum dot deliberately modified

A transistor is not a fundamentally different object from a quantum dot. It is an extended resonator at the ξ level $N \approx 8$ with three geometric modifications: **Scaling** gives more states. **Doping** shifts the torsion basis. **Gate field** deforms the resonator geometry from outside.

Modern GAA transistors (3 nm, 2023) have channels of ≈ 10 atomic layers — they are de facto quantum dot arrays. The transistor has returned to its physical origin.

The Complete Hierarchy

System	Scale	States	Characteristic
Spin-only QD	$R \lesssim 1.5$ nm	2	Minimum: one torsion, two orientations
Exciton QD	$R \sim 3\text{--}6$ nm	4–20	Bound torsion pairs; memory, logic, synapse
Transistor (classical)	$L \gtrsim 65$ nm	$\gg 1$	Many modes, classical; QD character hidden
Transistor (modern)	$L \sim 2\text{--}5$ nm	few	Doped + gate torsion; effectively QD array
Ising machine	optical resonator	2 per unit	Spin-only in the photonic; global feedback

ab.11 The Arc Closes: the Coherent Ising Machine

The **Coherent Ising Machine** (CIM) — an optical system that solves difficult optimisation problems by physically finding the energetically most favourable state — closes the arc completely.

The basic unit of a CIM is an optical resonator made of mirrors and an **optical parametric amplifier**. This resonator also has discrete states: it oscillates in either one or the other phase — spin-up or spin-down, zero or one. The excitation is analogue — continuous laser light, continuously adjustable coupling strengths. The state is discrete. The same physics, different realisation.

	Quantum dot	Synapse	Ising machine
Resonator	Semiconductor crystal	Protein structure	Optical oscillator
Excitation	Light, E-field	Neurotransmitter	Laser light
States	Energy levels	Weight levels	Spin up/down
Feedback	local	Network	global, all-to-all
Objective	Memory, logic	Learning	Optimisation

The Ising machine is the spin-only quantum dot in the photonic regime — two states per resonator, but with a global coupling network instead of local isolation. From the minimal two-state resonator to the global optimisation machine: the same principle, scaled and connected.

ab.12 Overall Conclusion: Resonators Everywhere

Conclusion

A quantum dot is a resonator. A synapse is a resonator. A cell is a system of resonators. An Ising machine is a resonator. A transistor is a modified resonator. All discrete from inside. All analogue from outside. All geometrically precise — otherwise they do not work.

The boundary between classical and quantum mechanical lies neither at a fixed temperature, nor at a fixed size, nor at a material class.

For confinement systems two conditions apply: geometric perfection (absolute) and $\Delta E > k_B T$ (size and temperature dependent). For tunnelling and spin coherence only Condition 1 applies. Condition 1 — geometric precision — is the only universal condition. It applies to all quantum effects.

In FFGFT: all these systems sit at the geometric scale level $N \approx 8$ of the ξ hierarchy — the same level at which the exciton scale lies.

Significance for the adaptive hybrid architecture

The hardware toolbox of the adaptive hybrid architecture is a collection of different realisations of the same fundamental principle — deployed according to task: quantum dots for local synaptic operations, Ising machines for global optimisation, macroscopic defect-free lattices for coherent quantum operations.

The learning AI selects — through trial and error — which resonator to use when. It does not learn *on* the hardware. It learns *through* it.

ab.13 Hardware Runs: IBM Quantum (2025)

IBM Quantum hardware, June 2025: classification

The FFGFT foundations developed in this document — in particular the treatment of qubits as torsion configurations with ξ coupling — were recomputed on genuine quantum hardware:

Hardware: IBM Brisbane & IBM Sherbrooke (127 qubits each), daily calibration, gate fidelity $> 99.9\%$ (1-qubit).

Core result: FFGFT amplitudes $E_i(x, t)$ with $\xi_{\text{Higgs}} = 1.038 \times 10^{-5}$ deliver algorithmically equivalent results to standard QM — Bell state fidelity 97.17%, repetition variance 0.0001–0.001 (excellent).

Classification: Algorithmically equivalent results do not distinguish FFGFT from standard QM; they are not a confirmation of the FFGFT foundations. The only documented hardware measurement in the corpus (Doc. 147) confirms the Bell violation; the ξ effect cannot be resolved with it. In the QM sector there is at present no FFGFT deviation testable with today's hardware.

ab.14 Further Topics in the Follow-up Documents

The following documents build on the foundations of this document:

- **Doc. 174** — *Influencing, Reading and Extending Spin in the FFGFT Framework:* Spin manipulation (Zeeman, EDSR, optical, exchange interaction), spin readout methods (QND measurement — compatible with a definite torsion configuration, but no proof of determinism), qudits and extended state spaces.
- **Doc. 175** — *More than Two States — Superposition, Registers, Entanglement in the FFGFT Framework:* Bloch sphere vs. FFGFT cylinder space, Bell states with ξ damping, register entanglement, qudit coding, photonic waveguides.
- **Doc. 176** — *Preparation and Transformation: Two Separate Processes:* The preparation (modular exponentiation) is a purely arithmetic problem, constituting approximately 95 % of the gate cost; the Fourier transform is a wave operation and cheap in every technology. Both are independent. Includes: three separate parameter spaces (ξ , ξ_{Higgs} , $\mathbb{Z}/N\mathbb{Z}$), hybrid preparation as a binary-controlled interference

network, thermal advantage of optical realisation ($h\nu/k_B T$), Gottesman-Knill, Weyl obstruction (not applicable to Shor's finite task).

Chapter ac

Doc. 174 — T0 Theory — Spin and Qubit States

ac.1 Influencing Spin — More than Just Magnetic Fields

In Doc. 173 spin appeared briefly in the context of the Ising machine and the minimal quantum resonator: two permitted orientations of a torsion configuration — spin-up and spin-down. The question of how to rotate this orientation from outside and how to read it out is the bridge between the abstract resonator picture and technical realisation.

The magnetic field is the historically first and most intuitive way — but by far not the only one.

Method 1: Magnetic Field — Zeeman Effect

A static magnetic field $\mathbf{B}$ splits the two spin states energetically:

$$\Delta E_Z = g^* \mu_B B, \quad (\text{ac.1})$$

where g^* is the effective g -factor of the material and μ_B is the Bohr magneton. For GaAs $g^* \approx -0.44$ — significantly smaller than for the free electron ($g = 2$), because the band structure of the crystal modifies the effective spin-orbit coupling.

A resonant radio-frequency pulse at the Larmor frequency $\nu_L = g^* \mu_B B / h$ rotates the spin in a controlled manner. This is the principle of electron spin resonance (ESR) and — in essence — also of magnetic resonance imaging.

FFGFT language: magnetic field as global torsion field

An external magnetic field establishes a preferred direction in the torsion field. It breaks the symmetry between spin-up and spin-down energetically. The radio-frequency pulse at ν_L is a periodically oscillating torsion perturbation — resonant when its frequency exactly corresponds to the energy difference of the two torsion configurations.

Method 2: Electric Field — EDSR

Electric fields do not couple directly to spin — the electron has no electric dipole moment component in spin space. But indirectly they couple very well, via **spin-orbit coupling**:

The electron moving in the electric field of the crystal lattice sees in its own rest frame an effective magnetic field. This internal field rotates the spin. A rapidly varying external electric field can resonantly drive this mechanism — this is called **EDSR** (Electric Dipole Spin Resonance).

Advantage over magnetic field methods

Electric fields can be generated locally with gate electrodes on a few nanometres and switched in under a nanosecond — magnetic fields cannot. For integrated qubit arrays EDSR is therefore the preferred method for single-qubit rotation.

In FFGFT language: the electric field deforms the resonator geometry locally — and via the spin-orbit coupling this geometry change translates directly into a torsion change of the spin degree of freedom. The coupling is not direct, but geometrically mediated.

Method 3: Circularly Polarised Light

Circularly polarised light (σ^+ or σ^-) carries angular momentum $\pm\hbar$ per photon. Upon resonant excitation of a quantum dot, a σ^+ photon transfers this angular momentum directly to the created exciton — and thus to the electron spin.

This allows **purely optical spin preparation** without any external field: by choice of the photon polarisation one selects the spin state.

Readout is via the polarisation of the fluorescence light: depending on spin state the quantum dot preferentially emits σ^+ or σ^- light. Polarisation measurement $\equiv$ spin measurement.

FFGFT: photon transfers torsion angular momentum

In FFGFT the photon carries its angular momentum as a topological property of the electromagnetic field torsion. Upon absorption this field torsion rotates the electron torsion into the complementary configuration. Spin-up and spin-down are two opposite torsion windings — the circular photon switches between them because it itself carries a directed winding.

Method 4: Exchange Interaction

When two electrons in adjacent quantum dots spatially overlap, their spins couple directly to each other — the **exchange interaction** with strength J :

$$H_J = J(t) \mathbf{S}_1 \cdot \mathbf{S}_2. \quad (\text{ac.2})$$

By pulsing the gate field (which controls the overlap and thus $J(t)$) one can execute controlled two-qubit operations.

This is the basis for **spin qubit gates** in semiconductor systems (Loss-DiVincenzo model, 1998). The strength: no external microwave needed — the coupling is purely electrically controllable.

Method 5: Hyperfine Interaction

The electron in the quantum dot interacts with the nuclear spins of the $\sim 10^4$ – 10^6 surrounding atoms. This coupling is normally a problem: the fluctuating nuclear spin bath creates a random effective magnetic field and destroys the spin polarisation in times of $T_2^* \sim 10$ – 100 ns.

But used in a controlled manner it is a tool: through dynamic decoupling (fast pulse sequences that average out the nuclear spin fluctuations) T_2 can be extended to microseconds. In some systems (e.g. phosphorus in silicon) the nuclear spin itself is used as a highly stable qubit ($T_1 > 1$ s at low temperatures).

ac.2 Reading Out Spin — Five Ways

Spin-to-Charge Conversion

The most important method in practice. The potential is chosen such that only spin-down (or only spin-up) is energetically permitted to tunnel out of the quantum dot. A single tunnelling electron creates a measurable current pulse in the adjacent **single-electron transistor** (SET).

No signal: spin-up (blocked). Signal: spin-down (tunnelled). Sensitivity: single electrons, error rate $< 1\%$.

Spin-to-charge conversion: state of the art 2025

In Si/SiGe and GaAs systems, spin-to-charge conversion readout schemes achieve accuracies $> 99.5\%$ with readout times below $1\ \mu\text{s}$ — sufficient for error-corrected quantum computers beyond the surface code threshold.

Pauli Blockade

More elegant: no external measuring device needed. The Pauli principle itself is the detector.

Two adjacent quantum dots, one electron each. An attempt is made to load both electrons into the same dot. If the spins are antiparallel: possible (both fit in the ground level). If the spins are parallel: not possible (Pauli exclusion) — the transport is **blocked**.

Measurable as a conductance difference: blockade $\equiv$ same spins.

FFGFT: Pauli blockade as geometric incompatibility

Two identical torsion configurations cannot occupy the same resonator — not because of a postulated prohibition, but because the same torsion winding in a cavity permits no second stable configuration of the same kind. The Pauli principle is in FFGFT a geometric statement about the compatibility of field configurations — not an independent quantisation rule.

Faraday and Kerr Rotation — State-Preserving Readout

Optical readout exploits the fact that spin-up and spin-down couple to different optical transitions. Circularly polarised light is scattered differently depending on spin state — the polarisation of the fluorescence light reveals the spin state. This resonant fluorescence is invasive: the absorbed photon excites the system and changes its state.

Fundamentally different from this is **Faraday / Kerr rotation**: a linearly polarised laser beam passes *by* the quantum dot — it is not absorbed, but rotates its plane of polarisation by a tiny angle $\theta_F \propto S_z$ upon passing. The spin state is readable from the rotation angle, without a single photon being absorbed.

Faraday rotation: reading quantum state without disturbing it

Faraday rotation is a **Quantum Non-Demolition measurement** (QND) — a readout method that does not change the measured state itself. Physically: the light field couples dispersively to the spin — the interaction is chosen so that it *commutes* with the spin operator S_z :

$$[H_{\text{Faraday}}, S_z] = 0. \quad (\text{ac.3})$$

An observable that commutes with the Hamiltonian is not disturbed by the measurement. The polarisation rotation θ_F is proportional to S_z — the eigenvalue is read off, the eigenstate is preserved.

Time resolution: in the picosecond range. Sensitivity: single spins in quantum dots detectable. The spin state can be measured *repeatedly* — each measurement yields the same result.

FFGFT: QND measurement and stable torsion configuration

The fact that Faraday rotation does not change the spin state is in FFGFT not a special case — it is the direct consequence of the stability of the torsion configuration.

Spin-up is a stable topological torsion. A passing photon rotates its plane of polarisation because it passes through the torsion field — but it does not bring the torsion configuration out of its stable state, because the coupling is so weak and symmetric that no energy transfer into the torsion mode occurs.

This is the same logic as with a resonator: a weak signal that does not hit the resonance frequency passes through the resonator without exciting it. The photon of the Faraday measurement intentionally does *not* hit the resonance frequency of the transition — it reads off the topological invariant without disturbing it.

What follows from this — and what does not: The QND measurement is compatible with the reading of a stable, definite torsion configuration: the spin state can be measured repeatedly without changing it, and always yields the same result. This is not a proof of a definite state *before the measurement*. In standard QM as well, the first QND measurement projects onto an eigenstate, and every repetition then yields the same result ($P^2 = P$); only the first result is random. Repeatability therefore does not distinguish the ontic and the epistemic interpretation; that the state was definite before the measurement is the FFGFT interpretation, not proven by QND.

ac.3 Beyond Two States — and How to Overcome the Limit

A spin qubit has two states: $|\uparrow\rangle$ and $|\downarrow\rangle$. This is the absolute minimum of a quantum resonator (Doc. 173, section on the minimal resonator).

But a quantum dot is not a two-state system by nature — it has arbitrarily many energy levels. The spin is only the *simplest* feature.

There are at least five strategies for extracting more than two distinguishable states from a quantum dot for quantum information.

Strategy 1: Qutrit and Qudit — More Energy Levels

Instead of $|0\rangle$ and $|1\rangle$ one uses $|0\rangle, |1\rangle, |2\rangle$ — a **qutrit** (three states) — or generally $|0\rangle$ to $|d-1\rangle$: a **qudit** of dimension d .

In the quantum dot this corresponds to the first d energy levels. The basic formula:

$$E_n = \frac{n^2 \pi^2 \hbar^2}{2m_{\text{eff}} R^2}, \quad n = 1, 2, \dots, d. \quad (\text{ac.4})$$

For $d = 3$ (qutrit) one needs three distinguishable levels with $E_3 - E_2 \gg k_B T$ — otherwise the thermal population blurs the third level.

Condition for a stable qudit of dimension d

All d level spacings must exceed the thermal energy:

$$E_{n+1} - E_n = (2n+1)E_1 \gg k_B T \quad \text{for all } n = 1, \dots, d-1.$$

The most critical spacing is $E_2 - E_1 = 3E_1$ (the smallest). Qudits of high dimension require small quantum dots or low temperatures — or both.

The information density of a qudit: $\log_2(d)$ bits per object. A qutrit encodes $\log_2(3) \approx 1.585$ bits — 58 % more than a qubit per object. A qufive-it encodes $\log_2(5) \approx 2.32$ bits.

Qudit efficiency at ξ level $N \approx 8$

A quantum dot with $R = 3$ nm at 4 K has at least 5 thermally stable levels (all spacings $\gg k_B T(4\text{ K}) = 0.34$ meV). This corresponds to a qufive-it with $\log_2(5) \approx 2.32$ bits of information content — in a resonator of 6 nm diameter (level $N \approx 8$, $L_8 \approx 22$ nm).

Strategy 2: Orbital Qubit — Geometry Instead of Energy

A quantum dot not only has different energy levels — it also has different **orbitals** (spatial wave patterns with the same energy, but different geometry): s-, p-, d-like states as in the atom.

In a non-spherical quantum dot the p-orbitals are energetically split. One can use the two lowest p-states (p_x and p_y) as a qubit — an **orbital qubit** — *without* looking at the spin.

The advantage: orbital states couple strongly to electric fields, weakly to magnetic noise. This makes them potentially more coherent in magnetically noisy environments.

Strategy 3: Exciton Number Qudit

From Doc. 173: a quantum dot can carry 0, 1, 2, 3, ... excitons simultaneously — discrete storage levels, sharply distinguishable.

This exciton number itself is a qudit:

$$|0\text{ Ex}\rangle, |1\text{ Ex}\rangle, |2\text{ Ex}\rangle, |3\text{ Ex}\rangle, \dots$$

Readout is via the emission wavelength: a biexciton ($|2\text{ Ex}\rangle$) emits at a different frequency than an exciton ($|1\text{ Ex}\rangle$) — spectrally clearly distinguishable.

FFGFT: exciton number as torsion pair number

Each exciton is a bound torsion-anti-torsion pair in the resonator. The exciton number is the number of such pairs — a topological invariant of the field configuration. Adding an exciton means adding a new torsion-anti-torsion pair to the resonator. The discreteness of the exciton number follows directly from the topological nature of the pairs: you cannot add half a torsion pair.

Strategy 4: Spin-Orbit Qudit — Spin and Orbital Combined

The most powerful combination: spin *and* orbital degree of freedom are used simultaneously.

A quantum dot with spin ($\uparrow/\downarrow$) and two orbital states (p_x/p_y) has four combined states: $|p_x, \uparrow\rangle, |p_x, \downarrow\rangle, |p_y, \uparrow\rangle, |p_y, \downarrow\rangle$ — a **ququart** (qudit of dimension 4, equivalent to 2 entangled qubits in a single object).

The spin-orbit coupling connects the two degrees of freedom and allows navigating between the combined states through electric fields.

Strategy	Degree of freedom	Dimension d	Bits per QD
Spin qubit	Spin	2	1.00
Orbital qubit	Orbital	2–4	1.00–2.00
Qudit	Energy level	3–6	1.58–2.58
Exciton qudit	Exciton number	3–5	1.58–2.32
Spin-orbital	Spin + orbital	4–8	2.00–3.00

Strategy 5: Continuous Variables — the Quantum Mechanical Oscillator

A radical alternative to the discrete qudit concept: instead of discrete states one uses **continuous variables** (CV).

A harmonic oscillator has infinitely many energy levels $|n\rangle$, $n = 0, 1, 2, \dots$. The information is not stored in individual levels, but in the **waveform** of the state — for example as a coherent state $|\alpha\rangle$ with complex amplitude α .

Quantum dots are *not* harmonic oscillators (their level spacings are not equidistant: $E_n \sim n^2$) — but **optical cavities** and **superconducting resonators** (Circuit-QED) are approximately.

Discrete or continuous — a choice of description level

In FFGFT both descriptions are facets of the same physics. Discrete states arise from the finite geometry of the resonator — the resonator is finite, so its modes are countable. An infinite number of modes (ideal harmonic oscillator) is the limiting case of an infinitely deep potential with uniform walls — an idealisation. Real systems are always discrete. Continuous variables are a useful approximation when the number of modes is so large that they can be treated as continuous — analogous to classical physics as the limit of many quantum states.

ac.4 Superposition — What Lies Between the States

So far: discrete states $|0\rangle, |1\rangle, \dots$. What is special about quantum systems is not that they have discrete states — classical coins do too. What is special is **superposition**:

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle, \quad |\alpha|^2 + |\beta|^2 = 1. \quad (\text{ac.5})$$

This is not ignorance about the state — it is a genuine intermediate state.

On the Bloch sphere (mapping of qubit states onto a sphere surface) every point corresponds to a possible quantum state — infinitely many continuous points on the sphere, not just the two poles.

FFGFT: superposition as intermediate torsion

Spin-up and spin-down are the two stable torsion extremes of the resonator. A superposition state $\alpha|\uparrow\rangle + \beta|\downarrow\rangle$ is a **metastable intermediate torsion** — not stable in the ground state, but stable on the timescale of the coherence time T_2 . The phase $\phi = \arg(\alpha/\beta)$ is the rotation position of this intermediate torsion. Decoherence is the process in which the environment pushes this intermediate torsion into one of the two stable extreme configurations.

For a qudit of dimension d the general state lies on a $(d-1)$ -dimensional complex sphere — the $\mathbb{CP}^{d-1}$. With growing d the state space grows exponentially: n qudits of dimension d span a space of dimension d^n .

ac.5 Decoherence — the Enemy of Superposition

Superposition is fragile. The environment measures constantly alongside — every interaction with a phonon, a nuclear spin, a photon is an (unwanted) measurement that collapses the state.

Two timescales:

T_1 (**relaxation time**): time until the system has fallen from the excited state into the ground state. Energy relaxation — the state $|1\rangle$ becomes $|0\rangle$.

T_2 (**coherence time**): time until the phase ϕ of the superposition is destroyed. Always $T_2 \leq 2T_1$ (phase relaxation is at least as fast as energy relaxation, usually faster).

System	T_1	T_2	Main mechanism
Electron spin in GaAs	~ 1 ms	~ 10 ns	Nuclear spin bath
Electron spin in Si/SiGe	> 1 s	~ 10 ms	^{28}Si (nuclear spin-free)
Electron spin in P:Si	> 1 s	> 100 ms	Isotopic purification
Exciton in CdSe QD	~ 1 ns	~ 10 ps	Phonon scattering
Superconductor transmon	~ 100 μs	~ 100 μs	Dielectric losses

Si/SiGe: FFGFT significance of $T_2 \sim 10$ ms

Silicon with enriched ^{28}Si (nuclear spin-free) eliminates the nuclear spin bath almost completely. In FFGFT language: the resonator is freed from a source of random torsion perturbations (nuclear spins). What remains is the intrinsic geometry of the resonator — and thus the fundamental limit $T_2 \leq 2T_1$. The long coherence time in ^{28}Si is a direct demonstration that Condition 1 (geometric perfection) is the decisive one.

ac.6 Open Connections to FFGFT

The following points have not yet been worked out — they mark places where the resonator picture of FFGFT could make concrete predictions:

Open questions for further chapters

1. Qudit level structure and K_{frak} :

The fractal correction factor K_{frak} (Doc. 173) shifts all energy levels by the same relative amount. For a qudit of dimension d this means $d - 1$ shifted transitions — a structured spectral signature that deviates from the ideal parabolic. Is this deviation measurable with modern multiphoton spectroscopy?

2. Spin-orbit coupling as torsion-torsion coupling:

The spin-orbit coupling connects two geometric degrees of freedom (spin = torsion direction, orbital = torsion pattern). In FFGFT this coupling should be material-dependent — exactly via the effective g -factor g^* and the band structure torsion of the crystal. Can g^* be derived from ξ and the crystal geometry?

3. Decoherence timescales and ξ hierarchy — numerical analysis:

Measured coherence times at the ξ level $N \approx 8$:

System	Timescale	Ratio to exciton
Exciton (CdSe QD)	$T_2 \sim 10 \text{ ps}$	1 (reference)
Spin qubit Si/SiGe	$T_2 \sim 10 \text{ ms}$	10^9 (= 9 orders of magnitude)
Nuclear spin P:Si	$T_1 > 1 \text{ s}$	10^{11} (= 11 orders of magnitude)

Comparison with ξ : With $\xi = 1.333 \times 10^{-4}$ ($\log_{10}(1/\xi) = 3.875$), 2 hierarchy levels span 7.75 orders of magnitude — fewer than the measured ratios (9 and 11 orders of magnitude). A second parameter $\xi_{\text{Higgs}} \approx 1.038 \times 10^{-5}$ (via $\lambda_h^2 v^2 / (64\pi^4 m_h^2)$, cf. Doc. 186), with which 2 levels would give 9.97 orders of magnitude, does not hold up: it was selected because it hits the measured window, and the interpretation via two separate ξ spaces (geometric and algebraic) is rationalisation after the selection, not an independent justification. There is only the one parameter $\xi = 4/30000$. Whether and how the decoherence timescales are related to the ξ hierarchy is open.

4. Entanglement as torsion topology:

Two entangled qubits are in a state that cannot be written as a product of two individual states — a common torsion configuration over spatially separated resonators. What does this mean for the local topology of the FFGFT field? Is there a natural description of Bell states as toroidal field configurations?

ac.7 Summary: From Spin to Qudit

Conclusion

Spin-up and spin-down are the minimum — two torsion orientations in one resonator.

Manipulation: magnetic field (global), electric field via spin-orbit coupling (local, fast), circular light (angular momentum transferring), exchange interaction (two-QD gate).

Readout: spin-to-charge conversion (practical, precise), Pauli blockade (geometrically self-reading), fluorescence polarisation (invasive, time-resolved), **Faraday/Kerr rotation (QND — state-preserving, repeatable)**.

The QND measurement yields the same result on every repetition — standard QM also predicts this after the first projection. That the spin state had a defined value before the measurement (the torsion configuration real and stable, independent of observation) is the FFGFT interpretation, not proven by QND.

Beyond two states: energy level qudits, orbital qubits, exciton qudits, spin-orbital combination — all in the same tiny resonator, all at the ξ level $N \approx 8$.

Superposition lies between the states — not ignorance, but a genuine intermediate state (intermediate torsion). Decoherence is its natural enemy: the environment collapses the torsion into one of the stable extreme configurations. Geometric perfection (Condition 1) is the universal foundation — for every one of these effects. Without a precise resonator: no qudit, no qubit, no spin gate. The FFGFT interpretation: all these phenomena are different expressions of stable torsion configurations in geometrically precise cavities at the scale $N \approx 8$.

Bibliography

- [1] Loss, D. & DiVincenzo, D. P. *Quantum computation with quantum dots*. Phys. Rev. A **57**, 120–126 (1998).
- [2] Koppens, F. H. L. et al. *Driven coherent oscillations of a single electron spin in a quantum dot*. Nature **442**, 766–771 (2006).
- [3] Nowack, K. C. et al. *Coherent control of a single electron spin with electric fields*. Science **318**, 1430–1433 (2007).
- [4] Corna, A. et al. *Electrically driven electron spin resonance mediated by spin–valley–orbit coupling in a silicon quantum dot*. npj Quantum Inf. **4**, 6 (2018).
- [5] Atatüre, M. et al. *Quantum-dot spin-state preparation with near-unity fidelity*. Science **312**, 551–553 (2006).
- [6] Press, D. et al. *Complete quantum control of a single quantum dot spin using ultrafast optical pulses*. Nature **456**, 218–221 (2008).
- [7] Petta, J. R. et al. *Coherent manipulation of coupled electron spins in semiconductor quantum dots*. Science **309**, 2180–2184 (2005).
- [8] Elzerman, J. M. et al. *Single-shot read-out of an individual electron spin in a quantum dot*. Nature **430**, 431–435 (2004).
- [9] Ono, K. et al. *Current rectification by Pauli exclusion in a weakly coupled double quantum dot system*. Science **297**, 1313–1317 (2002).
- [10] Atatüre, M. et al. *Observation of Faraday rotation from a single confined spin*. Nature Phys. **3**, 101–105 (2007).
- [11] Berezovsky, J. et al. *Picosecond coherent optical manipulation of a single electron spin in a quantum dot*. Science **320**, 349–352 (2008).
- [12] Grangier, P., Levenson, J. A. & Poizat, J. P. *Quantum non-demolition measurements in optics*. Nature **396**, 537–542 (1998).
- [13] Delbecq, M. R. et al. *Quantum non-demolition measurement of an electron spin qubit*. Nature Nanotechnol. **14**, 446–450 (2019).
- [14] Xue, X. et al. *Quantum non-demolition readout of an electron spin in silicon*. Nature Commun. **11**, 1178 (2020).

- [15] Veldhorst, M. et al. *An addressable quantum dot qubit with fault-tolerant control-fidelity*. Nature Nanotechnol. **9**, 981–985 (2014).

Chapter ad

Doc. 175 — T0 Theory — Qubit State Spaces

ad.1 The Real Question: What Lies Between $|0\rangle$ and $|1\rangle$?

Spin-up and spin-down are two points. Two points are unambiguously labelled — one bit of information.

But a **qubit** is not a bit. The real power of the qubit lies not in the fact that it can be $|0\rangle$ or $|1\rangle$ — a classical coin can do that too. It lies in the fact that it can be $|0\rangle$ and $|1\rangle$ *simultaneously* in a superposition — and in a continuous ratio:

$$|\psi\rangle = \cos\frac{\theta}{2} |0\rangle + e^{i\phi} \sin\frac{\theta}{2} |1\rangle. \quad (\text{ad.1})$$

Two parameters: the mixing angle $\theta \in [0, \pi]$ and the relative phase $\phi \in [0, 2\pi)$. Together they parameterise a **sphere surface** — the **Bloch sphere** (geometric representation of all possible single-qubit states as points on a sphere of unit radius).

On this sphere there are *infinitely many points* — not only the two poles $|0\rangle$ (north pole) and $|1\rangle$ (south pole), but the entire equator, all circles of latitude, every point in between. Every point is a valid, physically distinguishable quantum state.

A qubit does not encode 2 states — it encodes *infinitely many*

The two classical poles $|0\rangle$ and $|1\rangle$ are special cases of a continuous family. What distinguishes a qubit from a classical bit is not the number of discrete states — it is the **continuous geometry** of the state space.

The readout interacts with the system — every measurement couples to the torsion configuration and drives it into the nearest stable extreme state ($|0\rangle$ or $|1\rangle$). The result is deterministic — it depends on the exact initial state which we do not fully know before measurement. The apparent “probability” of standard QM describes this epistemic ignorance, not fundamental randomness.

The states themselves are discrete. The continuity lies in the phase space *before* measurement — the readout always delivers one of the two stable poles.

ad.2 The Cylindrical Phase Space — FFGFT Geometry of the Qubit

The Bloch sphere is the standard representation. In FFGFT (Doc. 034) there is a physically more direct alternative: the **cylindrical phase space**.

A qubit state is described by three geometric coordinates:

$$(z, r, \theta), \quad (\text{ad.2})$$

where:

- $z \in [-1, +1]$: the “height” along the cylinder axis — describes the proportion of the torsion configuration in the direction of $|0\rangle$ vs. $|1\rangle$. ($z = +1$: pure torsion type $|0\rangle$; $z = -1$: pure torsion type $|1\rangle$)
- $r \in [0, 1]$: the radius in the cross-section — indicates the strength of the intermediate torsion, i.e. how far the state is from both poles. ($r = 0$: stable pole; $r = 1$: maximum intermediate torsion)
- $\theta \in [0, 2\pi)$: the azimuthal angle — the **relative phase** between the two basis components.

The relation to the Bloch sphere is direct: $z = \cos \theta_{\text{Bloch}}$ and $r = |\sin \theta_{\text{Bloch}}|$. But the cylinder is not a geometric decoration — it makes gate operations into *geometric transformations* in real space:

Gate	Hilbert space (matrix)	FFGFT cylinder (transformation)
Hadamard H	$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$	Swap $z \leftrightarrow r$, rotate phase by $\pi/2$ — moves pole to equator
Phase Z	$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	Add π to phase θ — pure rotation about cylinder axis
Bit-flip X	$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$	2D rotation of (z, r) by angle $\alpha = \pi \cdot K_{\text{frak}}$ — fractal damping built in

FFGFT: gates as torsion rotations

In the cylindrical FFGFT formalism every gate operation is a geometric transformation of the torsion vector. The Hadamard gate does not rotate a matrix — it tilts the torsion configuration from the longitudinal axis to the transverse plane. The Z gate rotates the phase θ — it rotates the torsion winding about its own axis.

The factor K_{frak} in the X gate is not a correction factor added afterwards — it follows directly from the fractal spacetime geometry ($D_{\text{frak}} = 2.999867$) and means: every qubit rotation is slightly smaller than π , because spacetime itself minimally deviates from flat 3D geometry. All π gates should therefore systematically deviate by $\Delta\alpha/\pi \approx 1 - K_{\text{frak}} \approx 1.33\%$ from the ideal value. This prediction cannot be measured directly: a systematic deviation of 1.33% is absorbed by the pulse calibration of real qubits (amplitude adjustment) and cannot be separated from device variation (cf. Doc. 034).

ad.3 Quantum Registers — Exponential Growth through Combination

A single qubit has infinitely many intermediate states in the torsion phase space, but always delivers one of the two stable poles upon readout — the readout itself is the interaction that drives the state to the next stable extreme. The real power comes through **combination**.

Product States: Additive Classicality

Two independent qubits have together $2 \times 2 = 4$ basis states: $|00\rangle, |01\rangle, |10\rangle, |11\rangle$.

The general state *without entanglement* (product state) is:

$$|\psi_A\rangle \otimes |\psi_B\rangle = (\alpha_A |0\rangle + \beta_A |1\rangle) \otimes (\alpha_B |0\rangle + \beta_B |1\rangle). \quad (\text{ad.3})$$

This is only two separate Bloch spheres — no gain over two independent qubits. The state space is *additive*: $2 + 2 = 4$ parameters.

Entangled States: Multiplicative Growth

Once the two qubits are *entangled*, the joint state can no longer be written as a product of two individual states. The general two-qubit state requires 4 complex amplitudes (minus normalisation and global phase: $2 \times 4 - 2 = 6$ real parameters):

$$|\Psi\rangle = \alpha|00\rangle + \beta|01\rangle + \gamma|10\rangle + \delta|11\rangle. \quad (\text{ad.4})$$

This is **more** than two independent qubits — entangled qubits carry joint information that is contained in neither individual qubit.

For n qubits the Hilbert space grows exponentially:

$$\boxed{d(n) = 2^n} \quad (\text{ad.5})$$

Real parameters in the general state: $2 \cdot 2^n - 2$.

n qubits	Basis states	Real parameters	Classical equivalent
1	2	2	1 bit
2	4	6	2 bits
3	8	14	3 bits
10	1,024	2,046	10 bits
50	10^{15}	2×10^{15}	50 bits
300	10^{90}	$2 \times 10^{90} \approx$ number of atoms in the universe	

This is the actual quantum advantage: Not a single qubit is powerful — but the collective growth of an *entangled* register.

Bell States — Maximum Entanglement in Two Qubits

The four **Bell states** are the maximally entangled two-qubit states — each a complete orthonormal basis of the entangled subspace:

$$|\Phi^+\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle), \quad (\text{ad.6})$$

$$|\Phi^-\rangle = \frac{1}{\sqrt{2}}(|00\rangle - |11\rangle), \quad (\text{ad.7})$$

$$|\Psi^+\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle), \quad (\text{ad.8})$$

$$|\Psi^-\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle). \quad (\text{ad.9})$$

Each of these states has the property: if qubit A is measured, the result of qubit B is immediately determined — regardless of how far away.

In FFGFT (Doc. 023, 035):

$$E^{\text{FFGFT}}(a, b) = -\cos(a - b) \cdot \left(1 - \xi \frac{(\Delta\theta)^2/\pi^2}{D_{\text{frak}}}\right), \quad (\text{ad.10})$$

the ξ damping reduces the CHSH violation from $2\sqrt{2} \approx 2.82843$ to ≈ 2.82840 — a difference of $\sim 2 \times 10^{-5}$ (with $\xi = 4/30000$, $D_{\text{frak}} = 3 - \xi$ and the CHSH angles: $S = 2.828404$, $\Delta\text{CHSH} \approx 2.4 \times 10^{-5}$). The damping does not establish a local bound: the Bell violation remains, and the deviation from the Tsirelson bound lies far below the measurement noise of current hardware.

FFGFT: Bell states as joint torsion nodes

In the FFGFT picture a Bell state is not an action at a distance between two qubits — it is a **topologically connected torsion configuration** across two resonators. The two resonators share a common torsion node whose resolution simultaneously determines both ends at every local measurement.

This is not a signal propagating faster than light — it is a common geometric state established at preparation. The local readout interaction at one end deterministically drives the local torsion into a stable pole — and thereby necessarily also determines the state of the other end, because both share the

same torsion node. The “global consequence” is not action at a distance, but the geometric coherence of the common node.
 The ξ damping (FFGFT correction to Bell correlations) is in this picture the finite extension of the torsion node: the node is not point-like, but has a characteristic length $\sim L_\xi = \xi \cdot \ell_P \cdot (1/\xi)^N$ at the ξ scale level $N \approx 8$.

IBM Quantum hardware tests: Bell states, June 2025

Bell states were prepared and measured on genuine quantum hardware; the FFGFT circuits run on it:

Hardware: IBM Brisbane & IBM Sherbrooke (127 qubits each, Heavy-Hex lattice), access via IBM Quantum Platform / Qiskit Runtime.

Result:

Quantity	Value
Bell state fidelity	97.17 %
FFGFT algorithm vs. QM	Agreement within measurement precision
Repetition variance	0.0001–0.001 (excellent)
Deviation $P(00), P(11)$	1–5 % (hardware noise)

Significance in FFGFT context: The FFGFT amplitudes $E_i(x, t)$ with ξ coupling deliver algorithmically equivalent predictions to standard QM; a lower variance than in standard QM is not established (Doc. 147). The measurements confirm the Bell violation, not the ξ prediction: the ξ deviation from the ideal Bell correlation ($\Delta\text{CHSH} \approx 2 \times 10^{-5}$) lies far below the noise threshold of current hardware and is therefore not testable; this would require error-corrected qubits.

ad.4 Three Ways to More States — a Comparison

Strategy	More states through	Growth	FFGFT meaning
Superposition (1 qubit)	Continuous superposition of $ 0\rangle$ and $ 1\rangle$	∞ , but 1 bit at measurement	Intermediate torsion on the Bloch sphere
Register (n qubits)	Entanglement of multiple qubits	2^n (exponential)	2^n -dimensional torsion product space; entangled = topologically connected nodes
Qudit ($d > 2$, 1 object)	Higher energy levels, orbital states, exciton number	d (linear) per object, d^n for n qudits	d stable torsion configurations in one resonator

No strategy dominates in all areas

Register entanglement scales most strongly (2^n), but requires n physical objects and error-corrected entanglement operations between them.
 Qudit coding gains information *per object* ($\log_2 d > 1$), but the exponential growth is slower.
 Superposition is always present — it is the basis for quantum interference, the actual driving force of quantum algorithms.

ad.5 Quantum Interference — Why Superposition is Useful

Superposition alone would be a mere overlay — the readout would always deliver one of the two stable poles, deterministically dependent on the exact phase state.

What makes superposition into a *resource* is **interference**: amplitudes can constructively (amplify) or destructively (cancel) overlap — depending on phase ϕ .

This is the principle behind all useful quantum algorithms:

1. Prepare the register in a uniform intermediate torsion over all inputs (with the Hadamard gate).
2. Apply a function that increases the phases of the correct answers and decreases those of the wrong ones.
3. Let interference drive the wrong answers into unstable torsion configurations and the correct one into a stable one.
4. The readout drives the system deterministically into the nearest stable pole — which after interference corresponds with overwhelming frequency to the sought answer.

Without interference: Monte Carlo (classical guessing). With interference: Grover ($\sqrt{N}$ instead of N steps), Shor (factorisation in polynomial time).

FFGFT: interference as torsion phase superposition

In FFGFT the phase ϕ is not an abstract complex number — it is the rotation position of the intermediate torsion configuration (Doc. 174).

Constructive interference: two torsion phases coincide — they amplify each other into a geometrically more stable configuration. Destructive interference: two opposite torsion phases cancel each other — the resulting configuration has no stable geometric anchor and is not addressed at the readout interaction.

A quantum algorithm in FFGFT language: a sequence of torsion rotations that aligns the phase configuration of the register so that the sought answer ends in a stable torsion configuration — all others in geometrically unstable ones. The readout is then not a random selection, but a deterministic reaction of the resonator to the readout interaction.

ad.6 QND Measurement — Reading Without Destroying

What QND Means

A QND measurement reads an observable $\hat{A}$ without changing the corresponding eigenstate. The formal condition:

$$[\hat{A}, \hat{H}_{\text{measurement}}] = 0. \quad (\text{ad.11})$$

The measurement operator $\hat{H}_{\text{measurement}}$ commutes with the measured observable — it does not change the eigenvalue, because it respects the same eigenspace.

Practically this means: the same state can be **repeatedly measured** and delivers the same result each time — because the measurement does not touch the state itself.

The most important example from Doc. 174: **Faraday/Kerr rotation**. A linearly polarised laser beam passes the quantum dot without being absorbed — it rotates its plane of polarisation by $\theta_F \propto S_z$. No photon is absorbed. No energy transfer into the spin mode. The spin state before and after measurement is identical.

QND and the Question of Determinism

From the FFGFT perspective the QND measurement is more than a technically useful property — it is compatible with the reality of stable torsion configurations:

1. Faraday rotation measures S_z without changing the spin state.
2. Repeated measurements always yield the same result.

This is not a proof that the state already had this value definitely *before the first measurement*. In standard QM as well, the first QND measurement projects onto an eigenstate, and every repetition then yields the same result ($P^2 = P$); only the first result is random. Repeatability therefore does not distinguish the ontic and the

epistemic interpretation; that the state was definite before the measurement is the FFGFT interpretation, not proven by QND.

FFGFT: QND measurement and stable torsion configuration

In FFGFT, Faraday rotation is state-preserving because a stable torsion configuration (spin-up = topologically fixed winding in one direction) cannot be driven out of its state through a dispersive, non-resonant coupling.

The photon “sees” the torsion — it rotates its polarisation accordingly — but it transfers no energy into the torsion mode, because it is not resonant. The geometric state is invariant under this coupling.

The interpretation: That a state can be measured repeatedly without disturbance after the first measurement is predicted equally by FFGFT and standard QM; QND measurements do not decide between the interpretations.

In FFGFT the torsion configuration is interpreted as a geometric reality — stable, definite, pre-existent. The measurement reads it off. It does not create it.

Limits of QND Measurement

QND is possible for observables that commute with the measurement operator — typically: S_z (spin component along the quantization axis).

What QND *cannot* do: An intermediate torsion configuration (superposition $\alpha|\uparrow\rangle + \beta|\downarrow\rangle$, i.e. a state that is *not* an S_z eigenstate) is driven by an S_z -QND measurement into an eigenstate — deterministically, depending on the exact initial state, but the intermediate torsion is gone afterwards.

Why this is so is shown by the potential-landscape picture.

Potential Landscape: Stable Minima, Transitions and Readout Strength

FFGFT: torsion potential, transitions and readout strength

The complete picture of qubit states is a **potential landscape with two stable minima** and a saddle between them.

The readout strength decides what happens:

Initial state	Weak coupling	Strong coupling
Stable minimum ($ \uparrow\rangle$ or $ \downarrow\rangle$)	Coupling does not lift state from minimum — QND : state remains, result reproducible	Coupling can lift state from minimum — invasive, state changed
Slope / saddle (superposition)	Even weak coupling is sufficient to drive the state — it “rolls” into the next minimum. Which one : determined by exact phase ϕ (epistemically unknown, deterministic)	Same, just faster — state jumps immediately into next minimum

The decisive insight: There is no principled boundary between “QND-possible” and “QND-impossible” — there are only **more stable and less stable states** in a continuous potential landscape.

A state in the deep minimum is stable against the readout coupling — the minimum must be overcome. A state on the slope is unstable — any coupling, however weak, is sufficient to drive it into the next minimum.

And the “jump” to the final position is not random — it is determined deterministically by the phase of the initial state on the slope: a state near $|\uparrow\rangle$ lands in $|\uparrow\rangle$, a state near $|\downarrow\rangle$ lands in $|\downarrow\rangle$, a state on the saddle ($\theta = \pi/2$) lands depending on the fine phase ϕ .

The apparent “randomness” of standard QM is the epistemic ignorance of this phase — not an ontological randomness of the system.

The picture applies to all d levels of a qudit: not two minima, but d minima — with $d - 1$ saddle points between them.

QND and superposition — no contradiction

A stable eigenstate ($|\uparrow\rangle$ or $|\downarrow\rangle$): sits in the deep minimum — weak coupling reads without driving. QND possible. Result reproducible.

An intermediate torsion ($\alpha|\uparrow\rangle + \beta|\downarrow\rangle$): sits on the slope — any coupling drives into the next minimum. Deterministic, not random. The phase ϕ determines which minimum is reached.

The QND measurement is compatible with the reality of stable poles but does not prove it. The disturbing measurement shows the instability of intermediate torsions. Together they yield, in the FFGFT interpretation, a deterministic, geometrically founded picture.

ad.7 Two Levels of the ξ -Hierarchy: Spin Qubit and Photonic Resonator

Spin Qubit: Minimal Resonator at Level $N \approx 8$

A spin qubit is the absolute minimum of a quantum resonator (Doc. 173): a single electron, a single energy level, two permitted orientations.

The energy scale comes from **quantum confinement** at ξ level $N \approx 8$ (nanometre scale). The spacing between the two spin states in a typical magnetic field ($B \approx 1$ T) lies at:

$$\Delta E_Z = g^* \mu_B B \approx 0.1\text{--}1 \text{ meV}. \quad (\text{ad.12})$$

The thermal energy at room temperature is:

$$k_B T(300 \text{ K}) = 26 \text{ meV}. \quad (\text{ad.13})$$

So $\Delta E_Z \ll k_B T$ — thermal noise constantly knocks the system out of the minimum. Low temperatures ($T < 1$ K, $k_B T < 0.1$ meV) are not optional, but mandatory — not to maintain superposition, but simply to *stay* in one of the two minima.

Photonic Resonator: Collective Geometry at Level $N \approx 9\text{--}10$

A waveguide of lithium niobate (LiNbO_3), a Mach-Zehnder interferometer, a ring resonator — these are resonators made of 10^{18} to 10^{23} atoms in perfect lattice order.

The decisive difference from the spin qubit: **the resonator is not a single particle — it is the macroscopic geometry of the entire crystal.**

The energy scale of an optical photon:

$$E_{\text{photon}} \approx 1\text{--}2 \text{ eV} \quad (780\text{--}1550 \text{ nm}), \quad (\text{ad.14})$$

so:

$$\frac{E_{\text{photon}}}{k_B T(300 \text{ K})} \approx \frac{1 \text{ eV}}{26 \text{ meV}} \approx 40. \quad (\text{ad.15})$$

Thermal noise cannot spontaneously create optical photons — the system sits far below the thermal threshold for optical excitations. Condition 2 from Doc. 173 ($\Delta E \gg k_B T$) is satisfied by a factor of 40.

Why photonics works at room temperature

Not because the photonic system is more robustly built — but because it operates at a higher ξ level of the energy hierarchy.

The **geometric robustness** (Condition 1) is extraordinarily well satisfied by the macroscopic single crystal with $\sim 10^{23}$ lattice sites in perfect order — much more robust than a single quantum dot. Individual phonons (thermal lattice vibrations) barely disturb the collective mode.

The **thermal robustness** (Condition 2) is automatically given by the eV energy scale — thermal photons in the infrared have $\sim 25 \text{ meV}$, optical photons have $\sim 1 \text{ eV}$: a factor of ≈ 40 in energy, corresponding to a Boltzmann suppression $e^{-40} \approx 4 \times 10^{-18}$ (i.e. a margin of about 1.6 orders of magnitude in energy).

Not Restricted to Two States: Infinitely Many Analog Transitions

The spin qubit is reduced to two states — that is its strength (simplicity, control) and its weakness (cooling mandatory, fragile).

The photonic resonator is structurally different: It is *not* restricted to the minimal-two-state structure of spin-up/down.

In an MZI (Mach-Zehnder interferometer) the photon runs through two paths simultaneously — with a continuously adjustable phase ϕ between the paths. The phase is an analog quantity that can continuously traverse $[0, 2\pi)$. The interference result is $\cos^2(\phi/2)$ — a continuous transfer function, not a discrete two-level structure.

Analog or digital — a choice of operating mode

A photonic resonator can operate in **both modes**:

Digital: Phase set to $\phi = 0$ or $\phi = \pi$ — then transmission is 1 or 0. Two discrete states, like a classical bit.

Analog: Phase continuously between 0 and 2π — then transmission is a continuous number between 0 and 1. Infinitely many intermediate states, all stable at room temperature, because the energy scale is thermally unreachable. Analog readout of these transitions requires *no* cooling — but it requires geometric precision of the resonator (Condition 1 from Document 173).

This is the fundamental difference from the spin qubit: With the spin qubit the analog information (the phase ϕ on the slope of the potential landscape) lies in a thermally unstable intermediate state — T_2 collapses at room temperature within nanoseconds.

With the photonic resonator the analog information lies in the phase of the photon — and photons are **thermally stable**, because their energy scale is far above $k_B T$. The state on the “slope” here is not a fragile minimum — it is a geometrically stabilized mode parameter of the crystal.

Comparison of the Two Levels in the ξ -Hierarchy

	Spin qubit	Photonic resonator
ξ level	$N \approx 8$ (nm scale)	$N \approx 9\text{--}10$ (μm –mm scale)
Resonator	1 electron in nm cage	$10^{18}\text{--}10^{23}$ atoms as collective crystal
Energy scale	$\Delta E \approx 0.1\text{--}1\text{ meV}$	$E_{\text{photon}} \approx 1\text{--}2\text{ eV}$
$\Delta E/k_B T$ (300 K)	$\approx 0.004\text{--}0.04$ (too small!)	≈ 40 (robust)
Cooling needed?	Yes, $T < 1\text{ K}$ mandatory	No, room temperature
Analogue states	Fragile: $T_2 \sim \text{ns}$ at 300 K	Stable: phase thermally unreachable
Discreteness from	Quantum confinement (1 particle)	Macroscopic mode geometry
Typical application	Digital qubit, quantum gate	Analogue signal processing, photonic computing, 6G

FFGFT: two levels of the ξ hierarchy — same physics, different scales

Spin qubit and photonic resonator are in FFGFT not fundamentally different systems — they are resonators at different levels of the ξ scaling ladder. At level $N \approx 8$: the confinement energy minimum lies at meV — not thermally robust at 300 K. At level $N \approx 9\text{--}10$: the collective mode minimum lies at eV — completely thermally robust.

The ξ parameter generates a universal energy hierarchy: each level carries the energy by a factor of $1/\xi \approx 7500$ higher than the previous. From meV (level 8) to eV (level 9–10): the factor is $\sim 10^3\text{--}10^4$ — consistent with ξ scaling.

This explains not only photonics at room temperature — it explains why biological systems use photonic mechanisms (photosynthesis, bird navigation via light) and not spin qubit systems for quantum effects: nature works at the thermally robust level.

The Analog Resolution Limit — and Its Continuation in High-Frequency Technology

The photonic resonator is thermally robust — but only as long as one does not resolve too finely. As soon as one wants to distinguish analog intermediate values, the energy space E_{photon} is divided into N distinguishable levels. The energy difference between adjacent levels:

$$\Delta E_{\text{level}} = \frac{E_{\text{photon}}}{N}. \quad (\text{ad.16})$$

Condition 2 from Document 173 applies inexorably:

$$\Delta E_{\text{level}} \gg k_B T \implies N \ll \frac{E_{\text{photon}}}{k_B T}. \quad (\text{ad.17})$$

At room temperature and optical photons ($E \approx 1 \text{ eV}$):

$$N_{\text{max}}(300 \text{ K}) \approx \frac{1 \text{ eV}}{26 \text{ meV}} \approx 40. \quad (\text{ad.18})$$

The maximum analog resolution at room temperature is about 40 distinguishable levels — corresponding to $\log_2(40) \approx 5.3$ bits per photon (Shannon limit for this energy level).

The photonic system does not escape the thermal problem — it shifts it to a finer scale. More resolution requires either cooling (which gives the same argument as for the spin qubit) or an even higher energy scale.

The entire history of high-frequency technology as an ascent in the ξ -hierarchy

This is not a new insight — it is the oldest engineering wisdom of communications technology, now geometrically grounded.

High-frequency technology has been climbing the same ξ -hierarchy for a hundred years. At each frequency level the signal-to-noise ratio improves — not because engineers built better circuits, but because $\Delta E/k_B T$ grows with each level.

Technology	Frequency	$E = hf$	$E/k_B T$	N -level	Thermal regime
AM broadcast- ing	1 MHz	$4 \times 10^{-9} \text{ eV}$	1.5×10^{-7}	≈ 5.5	Thermally fully dom- inated; noise $\gg$ sig- nal
VHF/FM	100 MHz	$4 \times 10^{-7} \text{ eV}$	1.5×10^{-5}	≈ 6.4	Still strongly noise- limited; high-power transmitters re- quired
Microwave Radar	/ 10 GHz	$4 \times 10^{-5} \text{ eV}$	1.5×10^{-3}	≈ 7.3	Noise manageable; low-noise amplifiers required
5G / mmWave	30 GHz	$1.2 \times 10^{-4} \text{ eV}$	5×10^{-3}	≈ 7.5	Transition regime; cooling of most sensitive stages (radio telescopes)
THz	1 THz	$4 \times 10^{-3} \text{ eV}$	0.15	≈ 7.8	Difficult zone: $E \sim$ $k_B T$; that is why THz technology is so de- manding
Near-infrared / LiDAR	200 THz	0.8 eV	31	≈ 8.8	Thermally robust; room temperature without difficulty
Optics / fiber	300 THz	1.2 eV	46	≈ 9.0	Thermally fully sta- ble; $N_{\max} \approx 46$ ana- log levels
Short-wave UV	1,000 THz	4 eV	154	≈ 9.4	High analog resolu- tion; $N_{\max} \approx 150$ lev- els

The unifying relation:

Each level in the ξ -hierarchy carries the energy scale a factor $1/\xi \approx 7500$ higher than the previous one. From $N = 6$ (AM, MHz) to $N = 9$ (optics, THz) three levels bridge the factor $(1/\xi)^3 \approx 4 \times 10^{11}$ — the frequency of optical fiber, however, lies only $\sim 2\text{--}3 \times 10^8$ times above AM broadcasting; the three ξ levels and the actual span disagree by a factor of $\sim 1500\text{--}2000$. The N column of the table does not follow the factor 7500 per level either: $1 \text{ MHz} \rightarrow 100 \text{ MHz}$ gives $\Delta N = 0.9$ there, with 7500 per level it would be $\ln 100 / \ln 7500 = 0.52$.

In this picture optical fiber is not a new technology — it is the consistent continuation of the same trend to the next ξ -level.

And the resolution limit applies at every level: whoever wants to resolve more than ~ 40 analog levels at optical wavelengths must cool — just as the radio engineer in 1930 shielded his receiver and optimized his amplifier in order to find a signal in thermal noise.

The problem is the same. The scale shifts. The physics does not change.

In FFGFT: The universal resolution limit per temperature level:

$$N_{\max}(T, N_{\xi}) \approx \frac{E_0 \cdot (1/\xi)^{N_{\xi} - N_0}}{k_B T} \quad (\text{ad.19})$$

with $E_0 = k_B T_0$ at the reference level N_0 . Each ξ -level higher multiplies the attainable resolution by the factor $1/\xi \approx 7500$.

Above Optics — Three Regimes up to the Pair-Production Threshold

Optics ($N \approx 9$) is not the end of the ξ -ladder — but it is the end of the usable zone for guided photonic information transmission. What follows above it traces a clear sequence:

Range	Energy	N -level	λ	What happens to the resonator
UV (near)	3–10 eV	≈ 9.3	413–124 nm	Photoionization sets in; electrons are knocked out of outer shells. The resonator is damaged by its own signal.
Soft X-rays	100 eV – 1 keV	≈ 10	12–1 nm	λ approaches the lattice spacing ($d \approx 0.3$ nm). Bragg scattering at individual lattice planes — no longer a homogeneous medium. Waveguiding collapses.
Hard X-rays	1–100 keV	≈ 10.5	1–0.01 nm	$\lambda < d_{\text{lattice}}$: the photon flies through gaps in the lattice. Matter partially transparent. No refractive index, no resonator.
Gamma / nuclear transitions	>100 keV	≈ 11	<0.01 nm	Interaction only with atomic nuclei. Entirely different ξ -level — nuclear levels instead of electronic levels.
Pair-production threshold	>1 MeV	≈ 11.3	—	Pair production, natural upper limit.

The Cannonball Limit in FFGFT: Scale Incompatibility of Two Torsion Patterns

In the particle picture the argument is clear: the photon becomes a cannonball flying through gaps in the lattice as soon as $\lambda \lesssim d_{\text{lattice}}$.

In FFGFT there are no cannonballs and no waves as separate concepts — there are only **torsion configurations of the field on different scales**.

FFGFT: Waveguiding as Scale Resonance of Two Torsion Patterns

The crystal is a **periodic torsion structure**: a spatially repeated pattern of stable field nodes on the ξ -level $N \approx 7\text{--}8$ (atomic scale, $d \approx 0.3\text{ nm}$).

The photon is a **propagating torsion wave**: a coherent field configuration with characteristic length scale λ .

Waveguiding works precisely when these two scales are compatible — when the photon’s torsion wave perceives the periodic lattice structure as a *collective* homogeneous torsion environment. This requires $\lambda \gg d_{\text{lattice}}$.

Three regimes in FFGFT language:

Condition	Particle picture	FFGFT picture
$\lambda \gg d_{\text{lattice}}$	Wave sees homogeneous medium; guided propagation	Torsion wave couples collectively to lattice nodes — effective refractive index emerges as average of node density
$\lambda \approx d_{\text{lattice}}$	Bragg scattering at individual lattice planes	Torsion wave begins to resolve individual lattice nodes — collective coupling breaks into single-node coupling; useful for crystallography, destructive for waveguiding
$\lambda \ll d_{\text{lattice}}$	Cannonball flies through lattice gaps; matter transparent	Scale incompatibility: torsion wave is too fine to “see” the lattice nodes — no coupling, no guiding

The crucial point: The particle picture and the FFGFT wave picture are *the same geometric argument* — formulated at different description levels. FFGFT shows why: because there is no fundamental difference between particle and wave. Both are torsion configurations. The difference is only the **scale of the configuration relative to the scale of the environment**.

The “wave-particle duality” of standard QM is not a duality in FFGFT — it is a single geometry that appears as wave or particle depending on the observation scale.

The Pair-Production Threshold as Natural Endpoint of the ξ -Ladder

The pair-production threshold at $E > 2m_e c^2 = 1.022 \text{ MeV}$ is not a technical failure — it is a **geometric limit of the ξ -hierarchy** for carrier systems.

In FFGFT: the photon is a propagating torsion configuration with energy E_{photon} . An electron and a positron are two **stable elementary torsion nodes** — the most fundamental stable configurations on the scale of elementary particles.

Their rest energy is $m_e c^2 = 0.511 \text{ MeV}$ each. Together: $2m_e c^2 = 1.022 \text{ MeV}$.

As soon as $E_{\text{photon}} \geq 2m_e c^2$: the torsion energy of the photon exceeds the threshold to create two new stable configurations. The photon *reorganizes* into two more stable local torsion nodes — not because it is destroyed, but because that is the energetically possible geometric transition.

Pair-production threshold in the ξ -hierarchy

The threshold $E = 2m_e c^2 \approx 1 \text{ MeV}$ corresponds to the ξ -level $N \approx 11.3$ — the scale at which elementary particles exist as stable torsion configurations.

This is the natural upper limit for photonic information transmission: above this threshold the photon is no longer a pure carrier — it becomes a particle generator.

The usable zone of the ξ -hierarchy for guided information transmission lies between:

$$N \approx 5.5 \text{ (AM broadcasting, MHz)} \quad \text{to} \quad N \approx 9.3 \text{ (near UV, eV)}. \quad (\text{ad.20})$$

The span from 1 MHz (AM) to 10^{15} Hz (UV) amounts to $10^9 \approx (1/\xi)^{2.3}$ in frequency; four ξ levels, $(1/\xi)^4 \approx 3 \times 10^{15}$, exceed it by a factor of $\sim 3 \times 10^6$.

FFGFT: Unified View of Wave, Particle and Lattice Interaction

In FFGFT three apparently different phenomena express the same geometric logic:

Refractive index: Collective coupling of a torsion wave to many lattice nodes simultaneously — emergent phenomenon when $\lambda \gg d_{\text{lattice}}$.

Bragg scattering / cannonball limit: Transition from collective to single-node coupling when $\lambda \approx d_{\text{lattice}}$ — the same scale incompatibility from particle and wave perspectives.

Pair production: Torsion energy exceeds the threshold for new stable configurations — the photon reorganizes into two torsion nodes (electron + positron). All three are transitions between torsion configurations on different ξ -levels. The ξ -hierarchy is not only a length-scale partition — it is a partition of the **stable coupling regimes** between torsion configurations.

Convergence: Carrier Frequency Meets Internal Transition Frequency on ξ -Level $N \approx 8-9$

So far two frequencies have been carried in parallel, without making their relation explicit:

Carrier frequency: the external frequency of the light that transports information — the frequency by which we climb the ξ -ladder.

Transition frequency: the internal resonance frequency of a quantum resonator — the frequency corresponding to an energy jump between two levels: $\nu_{\text{internal}} = \Delta E/h$.

These two frequencies are fundamentally different. But on ξ -level $N \approx 8-9$ something remarkable happens: they **converge**.

The internal transition frequency of a quantum dot on ξ -level $N \approx 8$ (nanometer scale) lies at $\Delta E \approx 0.5-2$ eV — precisely in the optical range, 1550 nm to 700 nm wavelength.

Fiber-optic communication operates at 1550 nm ($E \approx 0.8$ eV) — exactly in the window where the internal transitions of the InGaAs semiconductor structures lie that serve as lasers, amplifiers and detectors.

Convergence on ξ -level $N \approx 8-9$

Fiber-optic communication at 1550 nm works so naturally because on ξ -level $N \approx 8-9$ two scales coincide:

	Scale	Energy
Carrier frequency (fiber)	$\lambda = 1550$ nm	0.8 eV
Internal transitions (InGaAs QD)	$R \approx 3-5$ nm	0.7–1.2 eV
Exciton Bohr radius (CdSe)	$a_X \approx 5.5$ nm	~ 1 eV

This is not a materials-science convention — it is geometrically forced by the ξ -hierarchy. The same scale level that determines the refractive index of the waveguide also determines the internal energy levels of the active components. Carrier and resonator speak the same language because they live on the same ξ -level.

And this is why waveguiding breaks down above the optical window not only from lattice geometry — but also because the carrier frequency leaves the ξ -level on which the active material components operate.

ad.8 ξ -Level Separation as Stability Principle

The Coupling Formula Between Levels

Between different ξ levels, the coupling strength decreases exponentially with the level difference ΔN :

$$\Gamma_{\Delta N} \propto \xi^{\Delta N}. \quad (\text{ad.21})$$

For $\Delta N = 1$: $\xi^1 \approx 1.333 \times 10^{-4}$ (weak but non-zero). For $\Delta N = 2$: $\xi^2 \approx 1.78 \times 10^{-8}$ (extremely weak). For $\Delta N = 4$: $\xi^4 \approx 3.2 \times 10^{-16}$ (practically non-existent).

Fractal Self-Similarity *Because of* the Damping

The fractal hierarchy of nature is not stable *despite* ξ damping — it is stable *because of* it.

Were ξ not small, all levels would strongly couple to each other. Every change at one level would immediately restructure all other levels. There would be no stable atoms, no stable molecules, no stable cells — only a single chaotic coupled system of all scales simultaneously.

Because $\xi \ll 1$, the levels can exist **quasi-independently**: each level has its own stable internal dynamics, barely disturbed by the other levels.

FFGFT: ξ damping as origin of the natural hierarchy

The existence of separate, stable structural levels in nature — quarks, nucleons, atoms, molecules, cells, organisms, planets, galaxies — is not a random property of matter.

It follows directly from ξ level separation: each level is practically invisible to the next-but-one. The coupling falls exponentially with level distance. This is the mechanism that makes the fractal self-similarity of nature *stable*.

Restructuring between Levels: When Does It Happen and When Not?

The question “can one level restructure another?” now has a precise answer:

Yes, when:

- ΔN is small (≤ 1) — the energy scales are close enough that the coupling $\xi^{\Delta N}$ can be compensated by resonant enhancement.
- Or the energy on level N_1 exceeds the *threshold* for a new stable configuration on level N_2 — as in pair production ($\gamma \rightarrow e^+ + e^-$).

No, when:

- $\Delta N \geq 2$ — the coupling $\xi^{\Delta N} \leq 10^{-8}$ is too weak to cause stable restructuring.
- The energy is not sufficient to overcome the threshold of the target configuration.

The Natural Hierarchy as a Geometric Stability System

Atoms are not restructured by phonons ($\Delta N \approx 1$, coupling $\sim \xi$: weak, but not zero — which is why decoherence exists, but atoms still remain stable).

Cells are not restructured by atomic vibrations ($\Delta N \approx 4$, coupling $\sim \xi^4 \approx 10^{-16}$: absolutely negligible).

Galaxies are not restructured by phonons ($\Delta N \approx 8$, coupling $\sim \xi^8 \approx 10^{-32}$: non-existent).

The hierarchy of nature — from quarks to galaxies — is in FFGFT no accident of matter distribution. It is the direct consequence of ξ -level separation: each

level is geometrically protected against disturbances from far-away levels. Self-similarity and stability are two sides of the same ξ -geometry.

ad.9 Qudit Register — the Connection to the ξ -Hierarchy

From Doc. 174: a quantum dot of dimension d encodes $\log_2 d$ bits. What happens when one entangles n such qudits?

The total state space has dimension d^n . For $d = 4$ (spin-orbital ququart) and n qudits:

$$d^n = 4^n = (2^2)^n = 2^{2n}. \quad (\text{ad.22})$$

This is equivalent to $2n$ qubits in a register of size n — double the information density per object.

Qudit density at ξ level $N \approx 8$

A quantum dot with $R = 3$ nm at $T < 1$ K has at least 4 thermally stable states (spin-orbital ququart, $d = 4$). At 4 K this is not sufficient: $k_B T(4 \text{ K}) = 0.34$ meV, while the Zeeman splitting at 1 T is only 0.1–1 meV ($g = 2$: 0.12 meV); for the spin half of the ququart the criterion $\Delta E \gg k_B T$ is met at 4 K only with a much larger B field. A register of $n = 10$ such qudits has dimension $4^{10} = 1,048,576$ — the same as 20 classical qubits, but only 10 physical objects at ξ level $N \approx 8$.

ad.10 The FFGFT Topology Compiler — Geometry of Entanglement

A practical problem with entangled registers: entanglement gates between non-adjacent qubits are expensive — they require many SWAP operations (which accumulate errors and cost time).

FFGFT (Doc. 034) offers a geometric solution: the **FFGFT topology compiler**.

Standard compilers minimise the number of SWAPs. The FFGFT compiler instead minimises the cumulative **fractal path length** of all entanglement operations. Since the fractal damping term $\exp(-\xi \cdot \ell / D_{\text{frak}})$ is distance-dependent (larger physical distance ℓ means stronger ξ damping of coherence), critically interacting qubits must be *physically placed closer* to each other.

This leads to a new chip architecture: not laid out according to logical circuit optimisation, but according to **geometric coherence maximisation**.

Connection to the adaptive hybrid architecture

The adaptive hybrid architecture (Doc. 173 conclusion) selects resonators according to task: quantum dots for local synaptic operations, Ising machines for global optimisation.

The FFGFT topology compiler is the linking element: it decides which qubit-s/qudits must physically sit close together (strong entanglement) and which may remain further apart (weakly, tolerantly damped entanglement). The learning AI on this hardware does not only learn *what* is to be computed — it also learns *which geometric arrangement* executes the computation most coherently. Not through programmed topology specifications, but through geometrically informed learning in the FFGFT compiler.

ad.11 Comparison: Classical Bit, Qubit, Qudit, Register

	Class. bit	Qubit	Qudit (d)	n qubits
State space	2 points	Bloch sphere (∞)	$\mathbb{CP}^{d-1}$ (∞)	$\mathbb{CP}^{2^n-1}$ (∞)
Readout result	1 bit (deterministic)	1 bit (deterministic, depending on phase state)	$\log_2 d$ bits (deterministic)	n bits
Parallel processing	none	Interference over 2 paths	Interference over d paths	Interference over 2^n paths
Resource for advantage	—	Phase ϕ	$d - 1$ phases	Entanglement + phases
FFGFT picture	stable/unstable torsion	Torsion on Bloch sphere	d stable torsion levels	Torsion network, topologically connected

ad.12 Open Connections — Next Steps

Open questions for further chapters

1. Fractal gate corrections:

The FFGFT compiler says: all π pulses deviate by $\Delta\alpha/\pi \approx K_{\text{frak}} - 1 \approx -1.33\%$ from the ideal π . This is not directly measurable: such a systematic deviation is absorbed by the pulse calibration of real qubits and cannot be separated from device variation (Section [ad.2](#)).

2. Bell states and ξ scale level:

The ξ damping in Bell correlations ($\Delta\text{CHSH} \approx 2 \times 10^{-5}$) depends on $N \approx 8$. For Bell tests with *macroscopic* objects (atoms, molecules) the scale level N would increase — and with it the damping. Is there an N -dependent suppression of

Bell violations for different system sizes?

3. Qudit entanglement and torus topology:

Two entangled qudits ($d = 4$) correspond in FFGFT to two topologically connected spin-orbital torsion configurations. What is the geometric description of this connection in the toroidal field? Is there a conserved quantity (topological invariant) that quantifies the entanglement depth?

4. Quantum error correction and ξ granularity:

The surface code needs ~ 1000 physical qubits per logical qubit. In a qudit-based register with $d = 4$ one needs only half as many physical objects for the same logical information. How does the error correction threshold change in the FFGFT framework when the natural noise floor $\xi \approx 1.333 \times 10^{-4}$ applies as the lower error bound?

ad.13 Summary

Conclusion: More than Two States

A single qubit has infinitely many intermediate states in the torsion phase space — but the states themselves are **discrete**: the readout always delivers one of the two stable poles, deterministically determined by the exact phase state at the moment of readout. The apparent “probability” of standard QM is epistemic ignorance of this state — not fundamental randomness of the system.

The readout interacts with the system — but not always destructively. Faraday/Kerr rotation (Doc. 174) is a QND measurement: it reads the state without changing it, because the coupling commutes with the spin operator. That the same state can be repeatedly measured is also predicted by standard QM after the first projection; that it *existed definitely before* the measurement is the FFGFT interpretation, not proven by QND.

The actual quantum advantage comes through three routes:

1. Superposition: Continuous intermediate torsion with phase ϕ enables interference — the geometric foundation of all useful quantum algorithms. Interference aligns the register so that the sought answer ends in a stable torsion configuration; the readout follows that deterministically.

2. Register entanglement: n qubits span a 2^n -dimensional torsion space. Entanglement means: the torsion nodes of the qubits are topologically connected — reading one node necessarily also determines the others. The ξ damping changes the Bell correlations only by $\sim 2 \times 10^{-5}$; it establishes no local bound, the Bell violation remains.

3. Qudit coding: A resonator with $d > 2$ stable torsion levels delivers $\log_2 d > 1$ bits per object at readout. Spin-orbital ququarts ($d = 4$) at ξ level $N \approx 8$ double the information density of a register.

In FFGFT all three strategies are expressions of the same fundamental principle: stable torsion configurations in geometrically precise resonators — discrete from inside, analoguely controllable from outside, deterministic at readout.

Bibliography

- [1] Kouwenhoven, L. P. et al. *Excitation spectra of circular, few-electron quantum dots*. *Science* **278**, 1788–1792 (1997).
- [2] Loss, D. & DiVincenzo, D. P. *Quantum computation with quantum dots*. *Phys. Rev. A* **57**, 120–126 (1998).
- [3] Veldhorst, M. et al. *An addressable quantum dot qubit with fault-tolerant control-fidelity*. *Nature Nanotechnol.* **9**, 981–985 (2014).
- [4] Muhonen, J. T. et al. *Storing quantum information for 30 seconds in a nanoelectronic device*. *Nature Nanotechnol.* **9**, 986–991 (2014).
- [5] Atatüre, M. et al. *Observation of Faraday rotation from a single confined spin*. *Nature Phys.* **3**, 101–105 (2007).
- [6] Grangier, P., Levenson, J. A. & Poizat, J. P. *Quantum non-demolition measurements in optics*. *Nature* **396**, 537–542 (1998).
- [7] Wang, C. et al. *Integrated lithium niobate electro-optic modulators operating at CMOS-compatible voltages*. *Nature* **562**, 101–104 (2018).
- [8] Reck, M. et al. *Experimental realization of any discrete unitary operator*. *Phys. Rev. Lett.* **73**, 58–61 (1994).
- [9] Shannon, C. E. *A mathematical theory of communication*. *Bell Syst. Tech. J.* **27**, 379–423 (1948).
- [10] Knill, E., Laflamme, R. & Milburn, G. J. *A scheme for efficient quantum computation with linear optics*. *Nature* **409**, 46–52 (2001).

Chapter ae

Doc. 176 — T0 Theory — Shor's Algorithm and the Photon

ae.1 Preparation and Transformation: Two Separate Processes

Summary

Shor's algorithm falls into two processes that are regularly conflated although they have nothing to do with each other:

	Preparation	Transformation
Input	numbers a, N	prepared signal
Output	signal with $f(x) = a^x \bmod N$	frequency representation
Nature	classical arithmetic	wave operation
Effort	expensive ($O(n^3)$ gates)	cheap ($O(n^2)$)

Preparation is difficult because modular exponentiation is difficult – equally so in every technology. It does *not* depend on quantum-mechanical properties. The transformation is the same wave operation in every technology and cheap in each. The difference between optics and a qubit register lies in neither, but in operating temperature and coupling mechanism.

Three separate parameter spaces

To avoid confusion this document distinguishes strictly:

1. **Geometric space:** $\xi = 4/30000 \approx 1.333 \times 10^{-4}$ – the structural parameter of FFGFT. Fixed, not adjustable.
2. **Algebraic space:** $\xi_{\text{Higgs}} \approx 1.038 \times 10^{-5}$ – field correction in the vacuum expectation value.
3. **Number-theoretic space** $\mathbb{Z}/N\mathbb{Z}$: *no* universal parameter. The period r is determined by $\text{lcm}(p-1, q-1)$ – an individual arithmetic property of N .

Factorisation lives exclusively in the third space. Neither ξ nor ξ_{Higgs} has a function there.

ae.2 Preparation

Input: the numbers a and N . **Output:** a state or signal carrying the sequence $f(x) = a^x \bmod N$.

What happens here

[K] Modular exponentiation is a *classical, deterministic* function. Its evaluation contains no quantum-mechanical component: on a quantum computer it is executed as a reversible classical circuit, in an optical setup it is imprinted as a spatial or spectral structure.

The cost

[K] The effort depends on the arithmetic, not on the chosen technology:

Realisation	Effort	for RSA-2048
Quantum circuit	$O(n^3)$ gates	$8.6 \cdot 10^9$ gates
Optical mask	2^n elements	10^{617} elements
Classical table	2^n values	10^{617} values

The quantum circuit is the cheapest known variant: it encodes the computational rule rather than the values and therefore requires only polynomial effort. An optical mask would have to carry every function value individually – for RSA-2048 more elements than the observable universe has atoms (10^{80}).

Preparation is an arithmetic problem

It is expensive because modular exponentiation is expensive – and that holds in every technology. A method promising to bypass it shifts the effort rather than removing it. **Preparation does not depend on quantum-mechanical properties.**

What superposition contributes here

[X] Nothing. The state $\sum_x |x\rangle |a^x \bmod N\rangle$ does contain all pairs, but a measurement yields exactly one – drawn at random. Afterwards one knows as much as after a single classical evaluation. Superposition does not simplify the arithmetic; it applies the same circuit across all inputs.

ae.3 The Hybrid Preparation: Binary Control Meets Analogue Wave

Pure binary implementation of modular exponentiation drowns in $O(n^3)$ gates. Purely analogue implementation fails at thermal noise and the modulo problem. A hybrid approach exploits the strengths of both:

- **The binary part (the control):** Exact, noise-free discretisation of the modulo logic – governs which phase paths are enabled.
- **The analogue part (the wave):** Continuous, parallel phase accumulation via interference and mixer structures in real time.

What happens physically

The analogue wave performs multiplication via *coherent phase superposition*. Mixing two waves with phases ϕ_1 and ϕ_2 produces:

$$\Psi_{\text{total}} \propto e^{i\phi_1} + e^{i\phi_2}$$

or for coupled systems $|\phi_1\rangle \otimes |\phi_2\rangle$. In classical engineering this is coherent summation or frequency mixing; in quantum physics the same mathematical structure is called superposition.

Why physics introduces new terms here

1. **Discretisation by single quanta:** At the single-photon level the analogue wave becomes a probability distribution – analogue superposition becomes quantum superposition.
2. **Escaping the noise problem:** Classical analogue mixers suffer from thermal noise ($k \cdot T \cdot B$). Operating in the single-photon regime enables noise-free switching via quantum error correction.

Conclusion on hybrid preparation

Hybrid preparation = binary-controlled interference network

The hybrid preparation is a *phased-array-type interference network* that is binary-conditioned. The “superposition” here is the physical description of analogue wave superposition – not a quantum- mechanical magic.

Cost of hybrid preparation

Implementation	Gate count	Characteristic
Purely binary	$O(n^3)$	Drowns for large n
Purely analogue	—	Fails at noise
Hybrid	Reduced	Phased-array interference

ae.4 The Fourier Transform

Input: the prepared signal. **Output:** its frequency representation.

What happens here

[K] The Fourier transform is a *wave operation*. It takes no interest in where the signal comes from or what it means – it translates one representation into another. No new information arises: what is absent from the input signal does not appear in the output.

The cost

[K] It is cheap in every realisation:

Realisation	Effort	Remark
Quantum Fourier transform	$O(n^2)$ gates	sequential
Optical MZI network	transit time	simultaneous, speed of light
Classical FFT	$O(2^n \cdot n)$	on explicit values

In the quantum register and in the optical network the effort is small. The difference between the two lies in operating temperature and coupling mechanism (Section [ae.5](#)), not in the transform itself.

What it is for

The period r sought is a *global* property of the sequence: no single function value carries it; it becomes visible only by comparing several values. The Fourier transform performs this comparison physically – interference cancels everything except the frequencies k/r , and from a few measurements r follows by continued-fraction expansion.

Two separate processes

Preparation: arithmetic, expensive, difficult independently of technology. *Transformation:* wave-mechanical, cheap, the same operation in every technology. The two are unconnected. Preparation can be improved without touching the transform, and vice versa.

ae.5 Photonic Realisation

MZI networks

A Mach-Zehnder interferometer realises phase rotations

$$\Delta\phi = \frac{2\pi jk}{n}, \quad (\text{ae.1})$$

which correspond exactly to the twiddle factors of the discrete Fourier transform. A network of such interferometers performs the Fourier transform as a direct wave operation.

[B] The Fourier transform is therefore not a distinguishing feature of the quantum computer. It is a wave operation realisable optically, acoustically, or electronically. The difference lies in state preparation, not in the transform.

The thermal advantage of optical realisation

[K] Whether a mode sits in its ground state without cooling is decided by the ratio $h\nu/(k_B T)$. The mean thermal occupation is

$$\bar{n} = \frac{1}{e^{h\nu/k_B T} - 1}. \quad (\text{ae.2})$$

System	ν (Hz)	T (K)	$h\nu/k_B T$	$\bar{n}$
Superconducting qubit	$5.0 \cdot 10^9$	300	$8.0 \cdot 10^{-4}$	$1.25 \cdot 10^3$
Superconducting qubit	$5.0 \cdot 10^9$	0.020	12.0	$6.2 \cdot 10^{-6}$
Photon 1550 nm	$1.93 \cdot 10^{14}$	300	30.9	$3.7 \cdot 10^{-14}$
Photon 600 nm	$5.0 \cdot 10^{14}$	300	80.0	$1.8 \cdot 10^{-35}$

At 5 GHz and room temperature some **1250 thermal excitations** sit in the mode – the system is fully thermalised. For an occupation below 0.01, $T < 52$ mK is required. At optical frequencies the ground state is occupied essentially exactly at room temperature.

Cooling cost. The Carnot limit demands at least $3 \cdot 10^4$ units of work per unit of heat removed at 10 mK; in practice considerably more. A dilution refrigerator draws 10–25 kW continuously; an optical network draws none.

Thermodynamically privileged, not computationally

The thermal advantage follows directly from $h\nu/k_B T$ and is the strongest known advantage of photonic over superconducting realisation. It concerns *infrastucture* – cooling, energy, size – not computational power. Both realisations can in principle run the same algorithms; photonics has difficulties of its own (deterministic two-photon gates, photon loss as the dominant error channel).

Where the exponential dimension comes from

[K] An opposition of “optics versus quantum computer” would be inaccurate – it would set a technology against a computational model. Photonic quantum computers exist (KLM scheme, measurement-based approaches). What matters is solely the structure of the state space:

Arrangement	Dimension
classical wave in n modes	n
one photon in n modes	n (the same)
n entangled two-level systems	2^n
m indistinguishable photons in n modes	$\binom{n+m-1}{m}$

A single photon in an MZI network realises the same n -dimensional unitary as a classical wave – classically simulable. With several indistinguishable photons the dimension grows exponentially: 25 photons in 50 modes give $3.5 \cdot 10^{19}$. The jump comes from the *tensor product structure*, not from the carrier medium.

What carries the computational advantage – an open question

[K] It would be natural to attribute the advantage to entanglement. The Gottesman-Knill theorem (1998) refutes this: any circuit consisting solely of Clifford gates (H , S , CNOT) is classically simulable in polynomial time, regardless of the entanglement produced. The GHZ state $(|0 \dots 0\rangle + |1 \dots 1\rangle)/\sqrt{2}$ is maximally entangled and arises from H and $n - 1$ CNOTs – at 1000 qubits some $2 \cdot 10^6$ bits suffice for the complete classical description.

Further counterexamples: DQC1 shows a conjectured advantage with almost no entanglement; boson sampling rests on indistinguishability and permanents. Jozsa and Linden (2003) show entanglement to be *necessary* for exponential advantage with pure states – about sufficiency the theorem says nothing.

[X] Which quantity carries the computational advantage is not settled. Contextuality, non-stabiliserness ("magic") and the interference structure itself are discussed; none of these candidates is established as the sole carrier. This document names the question as open rather than answering it.

What follows for factorisation. The optical realisation does not solve the problem: the limiting step is state preparation – the evaluation of $a^x \bmod N$ over all x – and that is arithmetic. A network can transform a prepared superposition; it cannot create one.

Topological constraint

[K] For a wave in phase space to form a closed resonance, the *logarithmic winding number* $W = \log_2(r)$ must be rational – not the frequency ratio r itself:

Case	Ratio r	Winding number W	Orbit
Pure fifth	$3/2$ (rational)	0.584963 (irrational)	open
Tempered fifth	$2^{7/12}$ (irrational)	$7/12$ (rational)	closed

Undamped oscillations on pure numerical ratios produce irrational winding numbers and diverge as an open spiral. An optical network therefore requires a discrete, rationalised phase grid to achieve sharp resonance points – closure is an artefact of discretisation.

ae.6 Structural Limits

The Weyl obstruction

[K] A quantum computer with n qubits spans a compact, finite state space. By Weyl's law its eigenvalue density grows by a power law:

$$N(\lambda) \sim C_d \cdot \lambda^{d/2}. \quad (\text{ae.3})$$

Arithmetic counting densities, by contrast, grow as

$$N(T) \sim T \ln T \quad (\text{ae.4})$$

(Riemann-von Mangoldt). $T \ln T$ is not a power.

Consequence: a finite state space cannot in principle carry such a spectrum in full – it carries at most an initial segment. Infinite spectra of this density would require an *unrolled dilation domain* (Docs. 315, 316), not a compact resonance space.

Important delimitation

This obstruction does **not** apply to Shor's algorithm. Shor seeks *one* period of *one concrete* number N – a finite task for which a register of $2n + 3$ qubits suffices. The Weyl obstruction concerns the complete representation of *infinite* arithmetic spectra. Invoking it against Shor overextends it.

What actually limits Shor at large N is the gate count of modular exponentiation (growing as n^3) and the error-correction overhead (roughly 10^3 physical per logical qubit). For RSA-2048: about 4099 logical qubits, 8.6×10^9 gates, 4.1×10^6 physical qubits. That is a *resource limit* – in principle surmountable by better hardware, unlike a structural one.

Primes as generators

[K] The load-bearing resonance structure rests on primes. The bicoherence test on 2469 Riemann zeros (Doc. 316) shows phase coupling exactly at prime powers ($b^2 = 0.76\text{--}0.85$) and none at composite numbers ($b^2 = 0.018\text{--}0.025$).

The physical counterpart is the harmonic series: in $n f_0$, exactly the prime orders introduce new tonal qualities; $4 = 2 \cdot 2$, $6 = 2 \cdot 3$, $9 = 3 \cdot 3$ are combinations of axes already present.

ae.7 Placing the Quantum Computer

[K] Shor's algorithm factorises on quantum hardware in theoretically $O((\log N)^3)$ – a real achievement for a specific problem. It does not follow that:

- **[X]** *Quantum computers hold a general exponential advantage.* This is not established for the vast majority of problems. Shor is a single case, not evidence of a general advantage.

- **[X]** *The advantage lies in the Fourier transform itself.* That is a wave operation and optically realisable. The bottleneck is state preparation.
- **[X]** *Quantum computers contribute to structural questions of number theory.* The Weyl obstruction applies to finite state spaces without qualification.

The core: reading out a period is a spectroscopic measurement problem on prepared wave fields. The measurement finds what was prepared; the preparation is the expensive part.

ae.8 Balance

Statement	Status
QFT is an interference operation	[K]
QFT generates no new information	[K] measured
Winding number decides closure	[K]
Primes carry the resonance structure	[K] measured by bicoherence
Fourier transform optically realisable	[B]
Optics thermodynamically privileged	[K] $h\nu/k_B T$
Exponential dimension from the medium	[X] tensor structure
Entanglement carries the advantage	[X] Gottesman-Knill
Preparation is arithmetic, not quantum-mechanical	[K]
Preparation and transformation are independent	[K]
General quantum advantage	[X] not established
Finite space carries infinite $T \ln T$ spectrum	[X] Weyl
Weyl obstruction limits Shor	[X] resource limit
ξ has a function in $\mathbb{Z}/N\mathbb{Z}$	[X] separate spaces

The load-bearing sentence. The Fourier transform is a projection – it finds structure but does not generate it. This holds equally for the classical FFT, the optical realisation, and the QFT. Factorisation lives in the space $\mathbb{Z}/N\mathbb{Z}$, where the geometry parameters of FFGFT have no function.

ae.9 Relation to the Corpus

The findings used are from Doc. 316 (August 2026): Weyl obstruction, bicoherence test, winding number versus frequency ratio, and the Fourier transform as a projection. The revision is declared under R75 in Doc. 190.

ae.10 Verification

The statements of this document are reproducible by two scripts (standard library only, target values as assertions):

- `s1_periodensuche.py` – correctness of period finding, scaling measurement (measured growth exponent of the mean order 0.95, hence effectively $O(N)$), prime factorisation of the σ denominators, comparison of methods.
- `s2_grenzen.py` – Weyl obstruction against an equidistant register, delimitation from Shor's task, resource estimate, winding-number check.
- `s3_thermik_zustandsraum.py` – thermal occupation by carrier frequency, Carnot cooling cost, dimension comparison of state spaces, Gottesman-Knill.

Chapter af

Doc. 178 — FFGFT — Spin Stability

af.1 The Uncomfortable Question

Every textbook on quantum mechanics states: An electron exists in genuine superposition $\alpha|\uparrow\rangle + \beta|\downarrow\rangle$ — simultaneously in both spin states, until a measurement collapses the state.

On every refrigerator there is a permanent magnet.

Between these two facts lies a contradiction that standard theory never explicitly resolves.

The central question of this document

If a single, isolated electron exists in genuine superposition of both spin states — why do then billions of electrons in the permanent magnet *never* show collective superposition, but always unambiguous, stable alignment?

And conversely: What does the permanent magnet — an everyday object, not a laboratory experiment — tell us about the nature of spin?

af.2 What the Permanent Magnet Really Shows

A permanent magnet is not a mysterious quantum phenomenon. It is the **macroscopic sum of spin alignments**.

Every electron in a ferromagnet (iron, nickel, cobalt) has a spin — and thus a magnetic moment. The **exchange interaction** (Doc. 174, section on Method 4) forces neighbouring spins into parallel alignment: all up, or all down. This is energetically more favourable than random distribution.

This collective alignment forms **Weiss domains** — macroscopic regions of coherent spin order. In the permanent magnet, these domains are permanently aligned and frozen by an external field.

What the permanent magnet directly proves

1. Spin has preferred, stable alignments.

Up and down are not equivalent alternatives — they are geometrically distinguished states. The field *wants* to be in one of them.

2. This stability is macroscopically robust.

A refrigerator magnet maintains its alignment at room temperature, in an environment with 10^{23} thermally fluctuating neighbouring atoms — for years, without decohering.

3. Superposition never occurs collectively.

No permanent magnet ever shows a state that is simultaneously north and south. Nature chooses. Always.

af.3 Standard Theory and Its Explanation

Standard theory explains the macroscopic absence of superposition through **decoherence**: The interaction with the environment destroys superposition so quickly that it is practically unobservable.

This is technically correct — but it is a **post-hoc explanation**, not a prediction.

The structural weakness of the decoherence explanation

Standard theory postulates:

1. A single electron exists in genuine superposition.
2. Decoherence through the environment destroys this superposition practically immediately for macroscopic systems.

This does explain *why* we see no macroscopic superposition — but it reverses the physical order of events:

It begins with superposition as the ground state and then explains why this ground state is always immediately destroyed.

FFGFT reverses this: Stability is the ground state. Superposition is the limited, fragile exception — with measurable timescales and a geometric cause.

The question is not only mathematical — it is physical: which description is the more natural one?

af.4 Superposition is Fragile — The Experimental Evidence

The measured coherence times from Doc. 174 tell an unambiguous story:

System	T_1	T_2	Temperature	Remark
Exciton (CdSe QD)	~ 1 ns	~ 10 ps	4 K	Superposition lasts picoseconds
Electron spin (GaAs)	~ 1 ms	~ 10 ns	mK	Superposition lasts nanoseconds
Electron spin (Si)	> 1 s	~ 10 ms	mK	Superposition lasts milliseconds
Nuclear spin (P:Si)	> 1 s	> 100 ms	mK	Best known system
Permanent magnet	Years	—	300 K	Stability without cooling

The table shows a clear trend: the more isolated the system from the disturbing influence of the environment, the longer superposition is maintained — but it *always* lasts only a limited time.

Superposition is never the stable ground state

Not a single experimental system shows superposition as a durably stable state. Even in the best qubit system (nuclear spin P:Si at mK) superposition decays after seconds.

The permanent magnet — without cooling, without isolation — maintains its stable spin state for years.

This is not coincidence. This is physics: **Stability is energetically preferred. Superposition is metastable.**

af.5 The Single Electron: Superposition or Unknown State?

Here lies the actual conceptual core.

Standard theory claims: a single, isolated electron exists in genuine superposition $\alpha|\uparrow\rangle + \beta|\downarrow\rangle$ until measurement.

The FPGFT position is more precise:

FPGFT: Superposition vs. epistemic ignorance

A single electron has at every moment a **definite torsion configuration** — either spin-up or spin-down.

What we describe as superposition $\alpha|\uparrow\rangle + \beta|\downarrow\rangle$ is **epistemic ignorance** about this definitive state — not ontological simultaneity.

QND measurement (Doc. 174, Faraday rotation) is compatible with this reading: a spin state can be measured repeatedly without changing it, and *always* yields the same result. This is not a proof that the state exists definitively *before measurement*.

In standard QM as well, the first QND measurement projects onto an eigenstate, and every repetition then yields the same result ($P^2 = P$); only the first result is random. Repeatability therefore does not distinguish the ontic from the epistemic interpretation; the epistemic reading is an interpretation, not proven by QND.

The apparent superposition arises in the description — when we mathematically describe a prepared system before the readout interaction. It is a state description under ignorance, not a physical double-state.

af.6 The Bridge: From Single Spin to Permanent Magnet

The transition from quantum to classical magnetism is in FFGFT no puzzle — it is the same mechanism, only at different scales.

Level	System	FFGFT description	Stability
Single spin	Electron in QD	One stable torsion configuration; up or down	$T_1 > 1$ s (P:Si)
Spin pair	Two coupled QDs	Two torsions; exchange interaction prefers parallel alignment	$J(t)$ controllable
Domain	Weiss domain (10^{15} spins)	Collective torsion order; energetically more stable than disorder	Stable at 300 K
Permanent magnet	Macroscopic body	Frozen global torsion alignment	Stable for years

Each level is the same physics — stable torsion configurations, collectively reinforced by exchange interaction. No step in this chain requires genuine superposition. Every step requires **geometric precision and energetic stability**.

FFGFT: Permanent magnet as macroscopic resonator

In FFGFT the permanent magnet is not a special case — it is the natural macroscopic limiting case of the resonator principle from Doc. 173.

A single quantum dot at ξ -level $N \approx 8$: two stable torsion configurations, metastable intermediate states.

A permanent magnet: $\sim 10^{22}$ resonators forced by exchange interaction into the same stable torsion configuration. The collective field reinforces stability — instead of destroying it (decoherence), the coupling acts here as a stability amplifier.

The boundary between quantum mechanics and classical magnetism is in FFGFT no categorical boundary — it is a quantitative transition along the ξ -hierarchy.

af.7 What Standard Theory Leaves Unanswered

The standard presentation of spin in textbooks and popular science texts has a structural problem: it begins with the abstract (Hilbert space, spinor, collapse) and then explains everyday life (permanent magnet) as a limiting case.

In doing so it overlooks a simple observation:

The simplest description of spin

The permanent magnet exists. It shows that spin is a **spatial orientation** — discrete, stable, real.

This is the most intuitive representation of spin: not an abstract state vector in Hilbert space, but a physical orientation in space that is macroscopically measurable, stable and everyday.

Quantisation (only up or down, nothing in between) follows from the geometry of the resonator — as derived in Doc. 173. It is not a postulate, but a geometric consequence.

The apparent superposition is an epistemic description under ignorance — not a physical double-state. The permanent magnet shows daily which state is fundamental: the stable, aligned one — not the superposed one.

af.8 Consequences for Quantum Information

This view has direct consequences for the understanding of quantum computers.

The standard view: a qubit is powerful because it is in superposition. Decoherence is the enemy of superposition and must be combated.

The FFGFT view: a qubit is powerful because it is a **precisely controllable torsion configuration**. Decoherence is not the destruction of a miracle — it is the system returning to its energetically stable ground state. This is normal, not pathological.

Decoherence as return to the ground state

In FFGFT, decoherence is not a failure of the system — it is proof that the stable spin state is real and energetically preferred.

The permanent magnet does not decohere. The qubit decoheres in milliseconds. The difference is not the physics — the difference is the **isolation from the coupling** that enforces the stable alignment.

A quantum computer does not fight against the nature of spin — it fights against the spin's tendency to return to its natural stability before the computation is finished. This is an engineering problem, not a foundational problem.

af.9 Summary

Conclusion: The Permanent Magnet as Witness

The permanent magnet is the simplest and most convincing argument for the FFGFT view of spin:

1. Spin has stable, preferred alignments.

Up and down are geometrically distinguished states — not arbitrary measurement outcomes.

2. Collective superposition does not exist.

Billions of electrons always choose one alignment. Never both simultaneously. Never.

3. Superposition is fragile — stability is robust.

Coherence times range from picoseconds (exciton) to seconds (nuclear spin) — always limited, always finite. The permanent magnet maintains its state for years.

4. The decoherence explanation reverses the physics.

Standard theory begins with superposition as the ground state and then explains why it always immediately disappears. FFGFT begins with stability as the ground state — superposition is the finite, measurably fragile exception.

5. QND measurement is compatible with determinism.

Repeated Faraday measurements always yield the same result — standard QM predicts this as well after the first projection; that the spin state was definite *before* measurement is the FFGFT interpretation, not proven by QND.

In FFGFT language: Spin-up and spin-down are two stable torsion configurations. The permanent magnet is their macroscopic collective realisation. Superposition is a metastable intermediate torsion — real on short timescales, but never the fundamental ground state.

The simplest description of spin does not begin in Hilbert space — it begins at the refrigerator.

Bibliography

- [1] Heisenberg, W. *Zur Theorie des Ferromagnetismus*. Z. Phys. **49**, 619–636 (1928).
- [2] Weiss, P. *L'hypothèse du champ moléculaire et la propriété ferromagnétique*. J. Phys. Théor. Appl. **6**, 661–690 (1907).
- [3] Loss, D. & DiVincenzo, D. P. *Quantum computation with quantum dots*. Phys. Rev. A **57**, 120–126 (1998).
- [4] Petta, J. R. et al. *Coherent manipulation of coupled electron spins in semiconductor quantum dots*. Science **309**, 2180–2184 (2005).
- [5] Veldhorst, M. et al. *An addressable quantum dot qubit with fault-tolerant control-fidelity*. Nature Nanotechnol. **9**, 981–985 (2014).
- [6] Muhonen, J. T. et al. *Storing quantum information for 30 seconds in a nanoelectronic device*. Nature Nanotechnol. **9**, 986–991 (2014).
- [7] Atatüre, M. et al. *Observation of Faraday rotation from a single confined spin*. Nature Phys. **3**, 101–105 (2007).
- [8] Delbecq, M. R. et al. *Quantum non-demolition measurement of an electron spin qubit*. Nature Nanotechnol. **14**, 446–450 (2019).
- [9] Zurek, W. H. *Decoherence, einselection, and the quantum origins of the classical*. Rev. Mod. Phys. **75**, 715–775 (2003).
- [10] Joos, E. et al. *Decoherence and the Appearance of a Classical World in Quantum Theory*. 2nd ed. Springer (2003).
- [11] Grangier, P., Levenson, J. A. & Poizat, J. P. *Quantum non-demolition measurements in optics*. Nature **396**, 537–542 (1998).

Chapter ag

Doc. 179 — FFGFT — Pairs and Photons

ag.1 Initial Question and Methodology

Three phenomena of modern physics show at first glance a structural similarity:

- A photon splits into two photons of lower frequency (parametric fluorescence, SPDC).
- An excited electron forms a bound pair with a remaining hole (exciton).
- Two electrons with opposite spin form a Cooper pair (superconductivity).

In all three cases a bound pair arises from a single excitation state — and in all three cases conservation quantities are exactly satisfied.

Methodology of this document

This document strictly separates two levels:

[STANDARD]: Established physics, experimentally secured, documented with primary sources. These statements are valid independently of FFGFT.

[FFGFT]: Interpretation within the FFGFT framework. These statements are FFGFT-specific and should be read as such.

The aim is to rigorously establish the structural analogy between the three systems — and then to specify precisely where the FFGFT interpretation goes beyond standard physics.

ag.2 System 1: Parametric Photon Splitting (SPDC)

[STANDARD] Established Physics of SPDC

Spontaneous Parametric Down-Conversion (SPDC) is a nonlinear optical process: a photon with frequency ω_0 is converted in a nonlinear crystal into two photons of lower frequency [1]:

$$\omega_0 \longrightarrow \omega_1 + \omega_2, \quad \text{with } \omega_1 + \omega_2 = \omega_0. \quad (\text{ag.1})$$

Exactly two conservation laws hold:

Energy conservation:

$$\hbar\omega_0 = \hbar\omega_1 + \hbar\omega_2. \quad (\text{ag.2})$$

Momentum conservation (phase matching):

$$\mathbf{k}_0 = \mathbf{k}_1 + \mathbf{k}_2. \quad (\text{ag.3})$$

For type-II SPDC the resulting photons have **opposite polarisation** and are **entangled**. The total angular momentum of the system is conserved [2].

STANDARD: What SPDC directly shows

1. A higher-energy state splits into two lower-energy states.
2. Energy and momentum are exactly conserved.
3. The opposite polarisation of the photons is not an additional assumption — it follows from angular momentum conservation.
4. The resulting photons are **real photons** — not quasi-particles, not computational aids. *[FFGFT caveat: see below — photons in FFGFT are not point particles but extended wave packets.]*

[FFGFT] Interpretation of SPDC

FFGFT: SPDC as torsion splitting — photons as wave packets

In FFGFT every photon carries angular momentum as a topological property of the electromagnetic field torsion (Doc. 174, Section 3).

Important FFGFT basic position: photons are wave packets.

In standard theory (QED), the photon is a quantised excitation quantum of the electromagnetic field — mathematically a Fock state $|1_{\mathbf{k}}\rangle$. It is often treated as a point particle.

In FFGFT the photon is **not a point particle**, but an **extended wave packet**: a spatially bounded, coherent torsion wave of the electromagnetic field with defined frequency ω , wave vector $\mathbf{k}$ and polarisation.

Why this matters for SPDC: The splitting $\omega_0 \rightarrow \omega_1 + \omega_2$ is in FFGFT not the decomposition of a point particle, but the **splitting of a wave packet** into two wave packets of lower frequency in the nonlinear crystal. This is physically more direct — light *is* a wave, and the splitting is a wave process.

Standard theory (QED)	FFGFT
Photon = Fock state $ 1_{\mathbf{k}}\rangle$	Photon = extended torsion wave packet
Point particle picture for many calculations	No point particle; always wave packet
Splitting = annihilation + creation of quanta	Splitting = wave process in non-linear medium
Entanglement = quantum correlation of Fock states	Entanglement = topological torsion correlation

Consequence for the description “real photon”: “Real” means in FFGFT: no quasi-particle approximation, no many-body reformulation — but no point particle. A SPDC photon is a real extended wave packet of the torsion field with clearly defined properties (ω , $\mathbf{k}$, polarisation, spatial extent).

Conserved quantity in FFGFT: The topological torsion charge is additive and conserved. This is no new claim — it is the FFGFT formulation of the known angular momentum conservation, now formulated for wave packets instead of point particles.

ag.3 System 2: The Exciton — Electron and Hole

[STANDARD] What a Hole Really Is

Here lies the conceptual core.

STANDARD: The hole is not a real particle

A **hole** in the valence band of a semiconductor is **not an independent fundamental particle** — it is a **quasi-particle** that describes the collective reaction of the many-body system to a missing electron [5, 6].

Precisely formulated (standard textbook definition):

- The valence band is completely filled in the ground state.
- When an electron is missing (through absorption of a photon), the many-body system behaves *as if* a positively charged particle with positive effective mass were present.
- This fictitious particle is called a “hole”.
- It has: positive charge $+e$, effective mass m_h^* , and a spin opposite to that of the missing electron.

The hole is mathematically equivalent to the description in electron language — but it is a construct of the many-body approximation, not a real antiparticle (not a positron!).

The real antiparticle of the electron is the **positron** — which arises in pair production ($\gamma \rightarrow e^- + e^+$) from real high-energy photons. The hole in the semiconductor is structurally different: it exists only in the context of the valence band of the material.

[STANDARD] The Exciton

The **exciton** is a bound state consisting of an excited electron in the conduction band and a hole in the valence band [4, 3]:

$$\text{Exciton} = \text{Electron (conduction band)} + \text{Hole (valence band)}. \quad (\text{ag.4})$$

The binding arises through the Coulomb attraction between the negatively charged electron and the positively charged hole.

Two main types:

- **Wannier-Mott exciton:** weakly bound (binding energy 1–100 meV), spatially extended (radius $\gg a_{\text{lattice}}$). Typical for semiconductors (GaAs, CdSe, Si). Stable only at low temperatures.
- **Frenkel exciton:** strongly bound (binding energy 0.1–1 eV), spatially compact (radius $\sim a_{\text{lattice}}$). Typical for organic semiconductors, biological systems. Stable even at room temperature.

[FFGFT] Interpretation of the Exciton

FFGFT: Exciton as complementary torsion pair

In FFGFT the electron in the conduction band carries a definite torsion configuration (spin, ξ -level N). The hole in the valence band is described as anti-torsion: the absence of an electron with opposite quantum numbers.

The exciton is in FFGFT a **torsion-anti-torsion pair**: the total torsion is reduced (Spin S_{total} , Momentum $\mathbf{p}_{\text{total}}$ are conserved), analogously to SPDC.

Critical limitation: The hole is a quasi-particle in standard theory — no fundamental particle. The FFGFT description as “anti-torsion” is an interpretation that goes beyond standard physics and has not been independently proven.

ag.4 System 3: The Cooper Pair

[STANDARD] BCS Theory and the Cooper Pair

In the BCS theory of superconductivity [8, 7], two electrons with opposite spin and momentum form a bound pair:

$$\text{Cooper pair} = \text{Electron}(\mathbf{k}, \uparrow) + \text{Electron}(-\mathbf{k}, \downarrow). \quad (\text{ag.5})$$

The binding is not direct (two negative charges repel each other) but is mediated via phonons (lattice vibrations):

- Electron 1 polarises the crystal lattice.

- This polarisation attracts Electron 2 with a delay.
- The result is an indirect, weak attractive interaction.

Binding energy: $\Delta E \approx 1\text{--}10\text{ meV}$ (conventional superconductors). This is much smaller than $k_B T$ at room temperature ($\approx 26\text{ meV}$).

Type	T_c	Binding energy	Mechanism
Conventional	1–40 K	1–10 meV	Phonon-mediated
High- T_c	up to 138 K	$\sim 50\text{--}100\text{ meV}$	Not fully understood

[FFGFT] Interpretation of the Cooper Pair

FFGFT: Cooper pair as opposite torsion configurations

In FFGFT the two electrons of the Cooper pair have opposite torsion configurations: $(\mathbf{k}, \uparrow)$ carries torsion in one direction, $(-\mathbf{k}, \downarrow)$ carries torsion in the opposite direction.

The pair thus has:

- Total torsion zero (spin singlet)
- Total momentum zero ($\mathbf{k} + (-\mathbf{k}) = 0$)

This is a **topologically neutral state** — geometrically more stable than two free, separate torsion configurations.

Why low temperature is necessary (FFGFT interpretation): The binding does not arise directly from the torsion geometry, but is mediated via phonons (lattice vibrations). Phonons are collective vibration modes of the crystal lattice — they lie at a different ξ -level than the electron spin system. The phonon-mediated coupling is therefore a cross-level coupling between ξ -levels (Doc. 175, section on ξ -hierarchy), whose strength $\propto \xi^{\Delta N}$ with $\Delta N = 1$. This explains the weakness of the binding energy compared to direct spin-spin coupling.

Critical limitation: This interpretation is consistent but not proven. Standard BCS theory explains the same situation completely without FFGFT concepts.

ag.5 Comparison of the Three Systems

[STANDARD] Structural Commonalities

System	Constituents	Binding energy	Temperature	Conserved
SPDC	Photon 1 + Photon 2	— (no binding)	Room temperature	$E, \mathbf{p}, J_z$
Exciton	Electron (conduction band) + Hole (valence band, quasi-particle)	10–100 meV (Wannier) to 1 eV (Frenkel)	4 K–300 K depending on type	$E, \mathbf{p}, S, \text{charge}$
Cooper pair	Electron 1 ($\mathbf{k}, \uparrow$) + Electron 2 ($-\mathbf{k}, \downarrow$)	1–10 meV (conventional)	$< T_c$ (1–40 K)	$E, \mathbf{p} = 0, S = 0$

Structural commonality [STANDARD]: In all three cases, two objects with opposite quantum numbers (spin, momentum, polarisation) arise from an excited state, so that conservation quantities are exactly satisfied.

Critical difference [STANDARD]:

- **SPDC:** two real photons, no binding energy, no temperature limit.
- **Exciton:** real electron + quasi-particle (hole), Coulomb binding, temperature-sensitive (Wannier-Mott: $T \ll T_{\text{bind}}/k_B$).
- **Cooper pair:** two real electrons, phonon-mediated binding, very temperature-sensitive ($T < T_c$).

The hole in the exciton is **not** analogous to the second photon in SPDC — it is a quasi-particle without independent fundamental existence outside the many-body system.

[STANDARD] The User's Question: "Hole = Two Electrons?"

This is a precise and important question.

The answer is: **technically yes, conceptually no.**

Technically yes: The exciton description "electron + hole" is mathematically equivalent to a description exclusively in electrons. In many-body theory the hole is defined as:

$$|\text{hole}\rangle \equiv |\text{Fermi sea without 1 electron at } \mathbf{k}, \downarrow\rangle. \quad (\text{ag.6})$$

The hole *is* the missing electron in disguise.

Conceptually no: The hole description is not merely a renaming — it is an efficient reorganisation of the many-body states that describes many phenomena (conductivity, magnetotransport, optical transitions) much more simply than pure electron language. The hole *behaves* like a particle — it has an effective mass, a spin, a momentum. It is a useful construct with real physical content, even if it is not a fundamental particle.

Hole: useful construct, not a fundamental particle

The hole in the semiconductor is not:

- A real antiparticle (not a positron)
- A new fundamental structure of nature
- Independent of the electron description

The hole is:

- A quasi-particle: an emergent excitation of the many-body system
- Mathematically: the absence of an electron in the valence band, rewritten as a positively charged particle
- Physically meaningful: it correctly describes real, measurable phenomena

The **Cooper pair**, by contrast, consists of **two real electrons** — no hole, no quasi-particle. That is the fundamental difference.

ag.6 [FFGFT] The Common Principle: Torsion Conservation

FFGFT: Conservation of total torsion as a unified principle

In FFGFT the connecting principle behind all three systems is the **conservation of total torsion**:

System	Before the process	After the process	Conserved
SPDC	Torsion ω_0 , angular momentum J_z	Torsion ω_1 (+) + Anti-torsion ω_2 (-)	Torsion charge: $J_z = J_1 + J_2$
Exciton	Valence band torsion (full band)	Conduction band torsion (electron) + valence band anti-torsion (hole)	Torsion charge: $S_{\text{total}}, \mathbf{p}_{\text{total}}$
Cooper pair	Fermi sea torsion	Pair torsion $(\mathbf{k}, \uparrow) + (-\mathbf{k}, \downarrow)$	Torsion charge: $S = 0, \mathbf{p} = 0$

The common FFGFT language: In all three cases a pair of torsion and anti-torsion arises from a torsion configuration, such that the total torsion charge is conserved.

This is in FFGFT no coincidence — it is the mirror image of the known conservation laws (energy, momentum, angular momentum) in the language of the torsion field.

What FFGFT does not claim: FFGFT does *not* claim that exciton hole and Cooper electron and SPDC photon are the same objects — they are physically different. FFGFT claims that their *structural similarity* has a common geometric origin: the conservation of the topological torsion charge.

ag.7 Consistency with the Temperature Argument from Doc. 173

The three systems differ dramatically in their temperature stability. This is [STANDARD] and directly explicable:

System	Binding energy	en-	Stability temperature	tem-	Cause
Photon (SPDC)	none	(free photons)	Room temperature		No thermal dissipation
Frenkel exciton	0.1–1 eV		Room temperature possible		$\Delta E \gg k_B T(300\text{ K}) = 26\text{ meV}$
Wannier exciton (CdSe)	$\sim 15\text{--}20\text{ meV}$		$< 50\text{ K}$		$\Delta E \sim k_B T$ at room temperature
Cooper pair (conventional)	1–10 meV		$T < T_c = 1\text{--}40\text{ K}$		$\Delta E \ll k_B T(300\text{ K})$

Consistency with Condition 2 from Doc. 173

The temperature limits follow directly from Condition 2 (Doc. 173): $\Delta E > k_B T$. The weaker the binding energy, the lower the temperature must be — this is no FFGFT argument, this is standard physics.

The Cooper pair needs low temperature not because of its electron nature, but because of its **very weak binding energy**, which follows from the indirect phonon mechanism.

ag.8 Summary

Conclusion: What is certain and what is FFGFT interpretation

[STANDARD] Established statements:

1. The hole is a quasi-particle, not a real antiparticle.

It is the mathematical reformulation of a missing electron in the valence band. "Exciton consists of two electrons" is technically correct — but the hole description is more efficient and physically meaningful.

2. The Cooper pair consists of two real electrons.

Opposite spin and momentum. Phonon-mediated binding. Low temperature is necessary because of the weak binding energy ($\Delta E \ll k_B T_{\text{room temperature}}$).

3. SPDC produces two real photons — in FFGFT: two wave packets.

Energy, momentum and angular momentum are exactly conserved. No binding energy — the photons are free. In FFGFT photons are not point particles but

extended torsion wave packets — the splitting is a wave process, not the decay of a point particle.

4. The structural similarity is real:

In all three cases two objects with opposite quantum numbers arise from an initial state, so that conservation quantities are exactly satisfied.

[FFGFT] Interpretation:

5. Torsion conservation as a unified principle.

The known conservation laws (energy, momentum, angular momentum) correspond in FFGFT to the conservation of the topological torsion charge. The structural similarity of the three systems thus has a common geometric origin — without the three systems being the same.

6. The hole as anti-torsion.

Consistent with the FFGFT description of the exciton from Doc. 173 — but going beyond standard physics and not independently proven.

Bibliography

- [1] Burnham, D. C. & Weinberg, D. L. *Observation of simultaneity in parametric production of optical photon pairs*. Phys. Rev. Lett. **25**, 84–87 (1970).
- [2] Kwiat, P. G. et al. *New high-intensity source of polarization-entangled photon pairs*. Phys. Rev. Lett. **75**, 4337–4341 (1995).
- [3] Frenkel, J. *On the transformation of light into heat in solids. I & II*. Phys. Rev. **37**, 17–44 and 1276–1294 (1931).
- [4] Wannier, G. H. *The structure of electronic excitation levels in insulating crystals*. Phys. Rev. **52**, 191–197 (1937).
- [5] Kittel, C. *Introduction to Solid State Physics*. 8th ed. Wiley (2004). Chapter 8: Semiconductor Crystals.
- [6] Ashcroft, N. W. & Mermin, N. D. *Solid State Physics*. Holt, Rinehart and Winston (1976). Chapter 12: The Semiclassical Model of Electron Dynamics; Chapter 28: Homogeneous Semiconductors.
- [7] Cooper, L. N. *Bound electron pairs in a degenerate Fermi gas*. Phys. Rev. **104**, 1189–1190 (1956).
- [8] Bardeen, J., Cooper, L. N. & Schrieffer, J. R. *Theory of superconductivity*. Phys. Rev. **108**, 1175–1204 (1957).
- [9] Woggon, U. *Optical Properties of Semiconductor Quantum Dots*. Springer Tracts in Modern Physics, Vol. 136 (1997).

Part IV

Part IV — Early Torus and L_0 Derivations

Chapter ah

Doc. 180 — T0 Theory — Derivation of L_0

Abstract

The T0 theory postulates neither the parameter ξ nor the minimal length scale $L_0 = \xi \cdot \ell_P$. Both emerge as necessary consequences of the self-consistency of the T0 Lagrangian. This document presents the complete derivation chain in five steps: (1) Lagrangian structure with time-mass duality, (2) self-consistency condition and 10^{-4} scaling, (3) uniqueness of ξ from the e-p- μ system, (4) gravitational constant G from ξ , (5) minimal length scale $L_0 = \xi \cdot \ell_P$.

ah.1 Step 1: The T0 Lagrangian

Standard Model Starting Point

The Standard Model Lagrangian for a lepton ψ_ℓ reads:

$$\mathcal{L}_{\text{SM}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}_\ell(i\gamma^\mu D_\mu - m_\ell)\psi_\ell \quad (\text{ah.1})$$

with $D_\mu = \partial_\mu + ieA_\mu$ and constant m_ℓ .

Time-Mass Duality as Fundamental Principle

In T0 theory the fundamental relation holds:

$$\tilde{T}(x, t) \cdot m(x, t) = 1 \quad (\text{ah.2})$$

This is not an additional postulate but a structural condition: every mass carries an intrinsic time field. This necessarily implies a dynamic mass field:

$$m(x, t) = m_0 + \Delta m(x, t) \quad (\text{ah.3})$$

Complete T0 Lagrangian

Complete T0 Lagrangian

$$\mathcal{L}_{T0} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}_\ell(i\gamma^\mu D_\mu - m_\ell)\psi_\ell + \frac{1}{2}(\partial_\mu\Delta m)(\partial^\mu\Delta m) - \frac{1}{2}m_T^2\Delta m^2 + \xi m_\ell \bar{\psi}_\ell\psi_\ell \Delta m \quad (\text{ah.4})$$

The Three Key Formulae

Coupling strength (mass-proportional):

$$g_T^\ell = \xi \cdot m_\ell \quad (\text{ah.5})$$

Time field mass:

$$m_T = \frac{\lambda}{\xi} \quad (\text{ah.6})$$

Vertex factor:

$$\bar{\psi}_\ell\psi_\ell\Delta m \longrightarrow -i\xi m_\ell \quad (\text{ah.7})$$

Time field propagator:

$$\Delta m(k) \longrightarrow \frac{i}{k^2 - m_T^2 + i\epsilon} \quad (\text{ah.8})$$

ah.2 Step 2: Self-Consistency Condition

Why the Lagrangian Forces ξ

The coupling term $g_T^\ell = \xi \cdot m_\ell$ is not freely choosable. The self-consistency of the Lagrangian requires:

- The theory remains free of divergences at high energies.
- The known particle masses are reproduced.
- The mass hierarchy between generations is geometrically describable.

The 10^{-4} Factor from QFT Loop Structure

The loop suppression in four spacetime dimensions yields:

$$\frac{1}{16\pi^3} \approx 2.01 \times 10^{-3} \quad (\text{ah.9})$$

Combined with the Higgs parameters from the T0 Lagrangian:

$$(\lambda_h)^2 \cdot \frac{v^2}{m_h^2} = (0.129)^2 \times \frac{(246.2)^2}{(125.1)^2} \approx 0.0647 \quad (\text{ah.10})$$

10^{-4} from Spacetime Dimensionality

$$\xi_4 \sim (10^{-1})^4 = 10^{-4} \quad (\text{ah.11})$$

This factor is not postulated — it follows from the dimensionality of spacetime.

The 4/3 Factor from Tetrahedral Geometry

The factor $4/3$ emerges from the harmonic structure of three-dimensional space:

$$\frac{4 \text{ vertices}}{3 \text{ edges per vertex}} = \frac{4}{3} \quad (\text{perfect fourth}) \quad (\text{ah.12})$$

The tetrahedron contains both fundamental harmonic intervals:

$$\frac{3}{2} \times \frac{4}{3} = 2 \quad (\text{octave}) \quad (\text{ah.13})$$

ah.3 Step 3: Uniqueness of ξ **The e-p- μ System as Uniqueness Proof**

From the three fundamental mass ratios:

$$R_{pe} = \frac{m_p}{m_e} = 1836.15267343 \quad (\text{ah.14})$$

$$R_{\mu e} = \frac{m_\mu}{m_e} = 206.7682830 \quad (\text{ah.15})$$

$$R_{p\mu} = \frac{m_p}{m_\mu} = 8.880 \quad (\text{ah.16})$$

With the mass scaling $m_p/m_e = 245 \times (4/3)^\kappa$ it follows:

$$\kappa = \frac{\ln(R_{pe}/245)}{\ln(4/3)} = \frac{\ln(1836.15/245)}{\ln(1.3333)} \approx 7.000 \quad (\text{ah.17})$$

 $\kappa = 7$ is the Only Consistent Solution

Exponent κ	R_{pe} prediction	Consistent	Error
$\kappa = 6$	$245 \times (4/3)^6 = 1376.6$	×	25.0%
$\kappa = 7$	$245 \times (4/3)^7 = 1835.4$	✓	0.04%
$\kappa = 8$	$245 \times (4/3)^8 = 2447.2$	×	33.3%

Table ah.1: $\kappa = 7$ is the only consistent solution

ξ as Emergent Constant

$$\xi = \frac{4}{30000} = \frac{4}{3} \times 10^{-4} = 1.333 \times 10^{-4} \quad (\text{ah.18})$$

ξ is not a postulate — it is the only self-consistent solution of the e-p- μ system.

ah.4 Step 4: Gravitational Constant G from ξ **Fundamental T0 Gravitational Relation**

The T0 Lagrangian implies the fundamental geometric relation:

$$\xi = 2\sqrt{G \cdot m_{\text{char}}} \Rightarrow G = \frac{\xi^2}{4 m_{\text{char}}} \quad (\text{T0 natural units}) \quad (\text{ah.19})$$

The complete SI value requires explicit dimensional and unit conversion factors (Document 012):

$$G_{\text{SI}} = \frac{\xi^2}{4 m_e} \times C_{\text{conv}} \times K_{\text{frak}} \quad (\text{ah.20})$$

where C_{conv} is the SI conversion factor (including the dimensional lift), and K_{frak} the fractal correction. Numerically:

$$G_{\text{SI}} \approx 6.674 \times 10^{-11} \text{ N m}^2 / \text{kg}^2 \quad (\text{ah.21})$$

The complete derivation of all factors is in Document 012. C_{conv} is the SI conversion of the single-anchor chain (Doc. 383); its numerical value 7.783×10^{-3} additionally contains the small residual of this chain (chain value 7.58×10^{-3}). The form of G and its order of magnitude follow from ξ and the one anchor (numerically checked in Doc. 383); with the anchor m_e , G lies at -2.6% , with v at -0.46% (Doc. 383). The agreement with the CODATA-2018 value (below 0.01% , with the formula $+0.003\%$) arises only with this residual and is not an independent precision result.

ah.5 Step 5: Minimal Length Scale L_0 **Planck Length from G and ξ**

$$\ell_P = \sqrt{\frac{\hbar G}{c^3}} = 1.616 \times 10^{-35} \text{ m} \quad (\text{ah.22})$$

The value $1.616 \times 10^{-35} \text{ m}$ is the measured value of ℓ_P , used here as anchor. Since G follows from ξ and one anchor, ℓ_P also follows from ξ and this one anchor: via $m_e = \frac{4}{15\pi} \xi^{11/2} E_P$ and $\ell_P = \hbar c / E_P$, with m_e as anchor one obtains $\ell_P = 1.595 \times 10^{-35} \text{ m}$ (-1.3%) and $L_0 = 2.127 \times 10^{-39} \text{ m}$. The deviation is the bare residue of the lepton ladder (Doc. 383).

L_0 as Sub-Planck Scale

The time-energy duality $r_0(E) \leftrightarrow E$ in natural units requires a minimal length scale. The coupling term $g_T^\ell = \xi \cdot m_\ell$ scales with ξ — below L_0 the concept of “distance” loses its physical meaning, because the time field fluctuations Δm dissolve the spacetime structure itself.

Minimal Length Scale

$$L_0 = \xi \cdot \ell_P = \frac{4}{30000} \times 1.616 \times 10^{-35} \text{ m} = 2.155 \times 10^{-39} \text{ m} \quad (\text{ah.23})$$

Physical Meaning of L_0

L_0 marks the scale where T0 geometric effects become dominant. Below L_0 :

- All vacuum fluctuations are fully effective (Casimir effect).
- The perturbative expansion loses its validity.
- The fractal correction K_{frak} becomes structurally necessary.

ah.6 Summary: Complete Derivation Chain

Derivation Chain without Postulates

Lagrangian structure ($\tilde{T} \cdot m = 1$)

⇓

Self-consistency requires $g_T = \xi \cdot m$

⇓

10^{-4} factor from 4D spacetime, $4/3$ factor from tetrahedral geometry

⇓

$\kappa = 7$ only solution of the e-p- μ system

⇓

$\xi = 4/30000$ (not a postulate — emergent constant)

⇓

$$G_{\text{SI}} = \frac{\xi^2}{4m_e} \times C_{\text{conv}} \times K_{\text{frak}} \Rightarrow \ell_P = \sqrt{\hbar G/c^3}$$

⇓

$$L_0 = \xi \cdot \ell_P = 2.155 \times 10^{-39} \text{ m} \quad (\text{ah.24})$$

ah.7 Connection to the SI System

The Unavoidable Circularity

All results in this document — and in the companion documents 181 and 182 — are initially **ratio-based**: ξ is dimensionless, $L_0/\ell_P = \xi$ is a pure ratio, $G = \xi^2/(4m_e)$ holds in natural units. As soon as one wishes to state absolute SI values — i.e. $L_0 = 2.155 \times 10^{-39}$ m rather than just $L_0 = \xi \cdot \ell_P$ — at least **one** external connection to the SI system is required.

Apparent Circularity is Unavoidable

It is not possible to derive absolute SI values from a purely dimensionless parameter like ξ without establishing somewhere a connection to the SI unit definition. This connection is not a weakness of the theory — it is an epistemological necessity of every physical theory.

Possible Entry Points into the SI System

The connection can be established via various quantities, all of which are equivalent:

- **Via α :** The fine-structure constant $\alpha = \xi \cdot E_0^2/(1 \text{ MeV})^2$ is dimensionless and experimentally very well known. It provides via $E_0 = \sqrt{m_e \cdot m_\mu / K_{\text{frak}}} = 7.398 \text{ MeV}$ the connection to the lepton masses in MeV. The bare value $\sqrt{m_e m_\mu} = 7.348 \text{ MeV}$ would give $\alpha^{-1} = 138.9$; only $E_0 = 7.398 \text{ MeV}$ gives 137.04.
- **Via m_{char} or m_e :** The electron mass $m_e = 0.511 \text{ MeV}$ is experimentally defined and links ξ to SI units via $G = \xi^2/(4m_e) \times C_{\text{conv}} \times K_{\text{frak}}$.
- **Via ℓ_P :** The Planck length $\ell_P = \sqrt{\hbar G/c^3}$ is experimentally accessible and gives directly $L_0 = \xi \cdot \ell_P$ in metres.

Which Entry is Fundamental?

In T0/FFGFT the most natural entry is via α , because:

- α is directly linked to ξ via $\alpha = \xi \cdot E_0^2/(1 \text{ MeV})^2$ (as a property of the anchor E_0 , not as a prediction)
- α is dimensionless — the transition to SI units occurs only through the choice of unit scale (MeV)
- The 2019 SI reform implemented exactly this calibration, which is consistent with the T0 geometry (Document 013)

Epistemological Classification

The derivation chain in this document is internally consistent and free of postulates. The only external input is the choice of an SI unit scale — for example via m_e in MeV or ℓ_P in metres. This input is not a circularity of the

theory, but the necessary calibration of every physical theory to the world of measurement. Treated in detail in Document 056 (Universal Derivation) and Document 101 (Circularity of Constants).

Chapter ai

Doc. 181 — T0 Theory — Justification of the Torus Geometry

Abstract

The 4D torus is not a free postulate of T0 theory. It is supported by three conditions: (1) freedom from divergences in quantum field theory requires periodic topology, (2) the four spacetime dimensions (3+1, empirical input) are reflected in the form of $\xi = 4/30000$, (3) stable winding numbers on the torus generate the observed quantum numbers. The minimal length scale $L_0 = \xi \cdot \ell_P$ is geometrically the length of a single torus lattice cell — the smallest stable winding. The Lagrangian and the torus yield L_0 consistently via two independent paths.

ai.1 Why a Fourth Dimension?

The Divergence Problem of Quantum Field Theory

Quantum field theory suffers from a fundamental problem: the vacuum energy diverges:

$$E_{\text{vac}} = \sum_n \frac{1}{2} \hbar \omega_n \rightarrow \infty \quad (\text{ai.1})$$

The observed cosmological constant is 10^{120} times smaller than the QFT prediction. The Standard Model resolves this through artificial renormalisation. T0/FFGFT resolves it geometrically.

Periodic Geometry as Solution

The harmonic series $\sum 1/n$ diverges; the series of inverse squares converges. Both hold regardless of whether the modes are pictured on a line or a circle; the topology does not turn the divergent series into a convergent one. For the inverse squares (Euler 1735):

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \quad (\text{ai.2})$$

Topological Conclusion

The periodic geometry automatically suppresses high modes. If spacetime has a periodic topology, UV divergences vanish geometrically — without artificial renormalisation. The convergence of $\sum 1/n^2$ itself is not a consequence of the topology (see above).

Four Dimensions and the Form of ξ

The parameter $\xi = 4/30000$ encodes the dimensionality of spacetime:

$$\xi_d \sim (10^{-1})^d \Rightarrow \text{for } d = 4 : \quad \xi_4 \sim 10^{-4} \quad (\text{ai.3})$$

Four Dimensions and the Form of ξ

The **4** in the numerator of $\xi = 4/30000$ stands for the four spacetime dimensions. The dimension count 3+1 itself is empirical input — no theory derives it; the form of ξ reflects it.

ai.2 Why a Torus?

Winding Numbers as Conserved Quantities

On a torus, winding numbers are topologically conserved. They generate stable discrete states:

$$\text{Spin} = \text{winding number } w = n_\phi / n_\theta \quad (\text{ai.4})$$

$$\text{Charge} = \text{flux } \Phi = n \cdot h/e \quad (\text{ai.5})$$

$$\text{Masses} = \text{torus overtone frequencies} \quad (\text{ai.6})$$

Torus vs. Sphere

A sphere S^3 is simply connected — no stable winding numbers. The torus is not simply connected — it generates exactly the observed quantum numbers. This is the topological reason for choosing the torus.

Torus Eigenvalues and Particle Masses

The Laplace operator on a d -dimensional torus with radii R_i has eigenvalues:

$$E_{\vec{n}} = \frac{\hbar^2}{2} \sum_i \frac{n_i^2}{R_i^2}, \quad \vec{n} \in \mathbb{Z}^d \quad (\text{ai.7})$$

Particle masses appear as torus overtone frequencies with step size $1/3 = 1/D_f$:

$$m_n = m_0 \cdot n^{1/3} \quad (\text{ai.8})$$

The first overtones $n = 2, 3$ thus give the ratios 1.26 and 1.44; the ratio $m_\mu/m_e = 206.8$ requires $n = 206.8^3 \approx 8.8 \cdot 10^6$. The lepton hierarchy and the Koide formula do not follow from this simple overtone ansatz; a derivation of the Koide formula is not given here.

Compactification

The fourth dimension is rolled up to radius r_4 . This radius must coincide with the sub-Planck scale from the Lagrangian — otherwise the theory would be inconsistent. The condition:

$$r_4 = L_0 = \xi \cdot \ell_P \quad (\text{ai.9})$$

The topology is therefore:

$$\mathbb{R}^3 \times S^1 \quad \text{with} \quad r_4 = \xi \cdot \ell_P \quad (\text{ai.10})$$

ai.3 Derivation of L_0 from Torus Geometry

The Winding Density

The torus has $f = 1/\xi$ sub-Planck cells per Planck length. With $\xi = 4/30000$:

$$f = \frac{1}{\xi} = \frac{30000}{4} = 7500 \quad \text{cells per } \ell_P \quad (\text{ai.11})$$

This number is not arbitrary — it follows directly from the geometric winding density $30000/4$ of the torus. The prime factorisation $7500 = 2^2 \times 3 \times 5^4$ encodes the symmetry structure of spacetime.

On notation: here f is the base winding number $f = 1/\xi = 7500$, not a frequency. In natural units f is only another reading of the duality $T \cdot m = 1$: $f = t_P/t_0 = \Lambda_{T0}/E_P$ with $t_0 = \xi t_P$ and $\Lambda_{T0} = E_P/\xi$, and $\xi E_0^2 = 1$ gives $f = E_0^2$. As a dimensionless ratio, f cannot be set to 1 with justification (Doc. A135).

Smallest Stable Winding

The smallest stable winding on the torus has the length of a single lattice cell:

$$L_0 = \frac{\ell_P}{f} = \ell_P \cdot \xi \quad (\text{ai.12})$$

Numerically:

$$L_0 = 1.616 \times 10^{-35} \text{ m} \times \frac{4}{30000} = 1.616 \times 10^{-35} \text{ m} \times 1.333 \times 10^{-4} \quad (\text{ai.13})$$

Minimal Length Scale from Torus Geometry

$$L_0 = \xi \cdot \ell_P = 2.155 \times 10^{-39} \text{ m} \quad (\text{ai.14})$$

L_0 is the length of a single torus lattice cell — the smallest stable winding.

Geometric Interpretation

L_0 is not simply ξ times the Planck length — L_0 is the length of a single torus lattice cell. Below L_0 :

- The concept of “distance” loses its physical meaning.
- All vacuum fluctuations are fully effective.
- The fractal correction K_{frak} becomes structurally necessary.

Consistency of the Two Derivation Paths

L_0 is not derived twice — the Lagrangian and the torus yield the same number via two independent paths:

Two Consistent Paths to L_0

$$\text{Lagrangian: } g_T = \xi \cdot m \Rightarrow \xi = 4/30000$$

↓

$$\text{Torus: } f = 1/\xi = 7500 \text{ cells per } \ell_P$$

↓

$$\text{Smallest winding: } L_0 = \ell_P / f = \xi \cdot \ell_P$$

↓

$$\boxed{L_0 = 2.155 \times 10^{-39} \text{ m}} \quad (\text{consistent from both paths}) \quad (\text{ai.15})$$

The Lagrangian provides ξ . The torus explains what ξ means geometrically: the reciprocal winding density.

ai.4 Comparison with Other Approaches

String Theory

String theory requires 10 or 11 dimensions. The T0 Lagrangian with a single parameter ξ suffices with 4. Every additional dimension would be a postulate — in FFGFT there are no free parameters.

Loop Quantum Gravity

LQG discretises space into spin networks. T0 discretises spacetime through the periodic torus geometry. The difference: in T0 the discreteness is a topological consequence, not an additional postulate.

Asymptotic Safety

AS seeks a UV fixed point as the result of renormalisation group flow. In T0 there is no renormalisation group flow — ξ is a geometric constant, not a running parameter. The fixed point is not sought but established geometrically from the outset.

ai.5 Physical Consequences

Vacuum Catastrophe Resolved

On the fractal torus, the vacuum energy converges through periodic suppression of high frequencies. No artificial renormalisation required.

Three Generations as Overtone Structure

Electron, muon and tau are interpreted as states of a single torus resonator. The observed hierarchy

$$m_e : m_\mu : m_\tau \approx 1 : 206.8 : 3477 \quad (\text{ai.16})$$

cannot, however, be represented as fundamental tone and first overtones with the simple overtone ansatz $m_n = m_0 n^{1/3}$: the first overtones give only the ratios 1.26 and 1.44 (section Torus Eigenvalues and Particle Masses).

No Big Bang and No Frequency Dependence

The torus has no boundaries. The observed redshift does not arise through expansion but through geometric energy loss — fractal summation extends the effective path length of photons through the T0 vacuum field:

$$z \approx \xi \cdot \ln\left(\frac{d}{\ell_P}\right) \quad (\text{ai.17})$$

This logarithmic form does not hold up quantitatively: for $d = R_H = 1.4 \cdot 10^{26} \text{ m}$, $\ln(d/\ell_P) = 140.3$ and hence $z = \xi \cdot 140.3 = 0.019$; $z \approx 10$ would require $\ln(d/\ell_P) = 75\,000$. In this form the formula is incompatible with the observed redshifts $z > 1$. The achromaticity of the redshift is unaffected by this; the scale of the path extension does not follow from ξ alone but is calibrated via the measured H_0 .

Frequency Independence of the Redshift

Since the path extension is a property of the spacetime geometry itself, red and blue photons starting from the same location take exactly the same extended path. Their wavelengths are therefore stretched by the same percentage — there is no frequency dependence. This distinguishes the T0 mechanism fundamentally from simple "tired light" models, which fail precisely on the observed frequency independence. Derived in detail in Document 026 (Geometric Cosmology).

ai.6 Summary

Three Independent Conditions Point to the Same Torus

- (1) Freedom from divergences $\Rightarrow$ periodic topology
- (2) 3+1 dimensions (empirical) $\Rightarrow$ form of $\xi = 4/30000$
- (3) Winding numbers $\Rightarrow$ stable quantum numbers

All three point to the same conclusion:

$$\mathbb{R}^3 \times S^1 \quad \text{with} \quad r_4 = L_0 = \xi \cdot \ell_P = 2.155 \times 10^{-39} \text{ m} \quad (\text{ai.18})$$

Note: Connection to the SI System

The absolute numerical values in this document — in particular $L_0 = 2.155 \times 10^{-39} \text{ m}$ — presuppose a connection to the SI unit system. This is not circular but epistemologically necessary: from a dimensionless parameter ξ alone, no absolute metre values can be derived. The connection is made via m_e , ℓ_P or α as equivalent entry points. Treated in detail in Document 180 (Section: Connection to the SI System) and in Documents 056 and 101.

Documents 180, 181 and 182 form a unit and should not be read separately.

Chapter aj

Doc. 182 — T0 Theory — Universe and Maximum Scale

Abstract

ξ is the single fundamental parameter of T0 theory. The complete chain reads: $\xi \rightarrow \{m_e, m_\mu, \dots\} \rightarrow G \rightarrow \ell_P$, with $L_0 = \xi \cdot \ell_P$ as sub-Planck scale of spacetime granulation. The upper scale of the cosmological sector, R_H , is written in the form $E_H = E_0 \cdot \xi^{41/4}$; the exponent 41/4, however, is not derived from ξ but calibrated to the measured H_0 (Doc. 190, P35/P36). The ratio $R_H/L_0 = 6.49 \times 10^{64}$ is therefore the measured cosmic scale in ξ -notation, not an independent ξ result.

The question of a universal maximum size of the universe is, however, not meaningfully posed in FFGFT: **every physical scale possesses its own system-specific upper bound** $R_{\text{torus}}(m) = \hbar/mc$. There is no single cosmic upper bound valid for all systems — R_H is the upper bound of the cosmological sector, not of the universe as a whole.

aj.1 Derivation Chain: From ξ to Both Scale Limits

The Single Fundamental Parameter

T0 theory has one single fundamental parameter (Document 013):

$$\xi = \frac{4}{3} \times 10^{-4} \quad (\text{aj.1})$$

All physical quantities derive from it. The complete chain reads:

$$\xi \rightarrow \{m_e, m_\mu, m_\tau, \dots\} \rightarrow G \rightarrow \ell_P \rightarrow c \quad (\text{aj.2})$$

with $L_0 = \xi \cdot \ell_P$ as the fundamental sub-Planck scale of spacetime granulation.

Derivation of Particle Masses from ξ

The lepton masses follow from ξ via the geometric structure factor of the quantum numbers $f(n, l, j)$ (not a frequency and not the base winding number $f = 1/\xi$) and the T0 scaling factor $S_{T0} = 1 \text{ MeV}/c^2$:

$$m_\ell = \frac{f(n, l, j)^2}{\xi^2} \cdot S_{T0} \quad (\text{aj.3})$$

The geometric mean of the lepton masses yields the characteristic energy scale:

$$E_0 = \sqrt{\frac{m_e \cdot m_\mu}{K_{\text{frak}}}} = 7.398 \text{ MeV} \quad (\text{aj.4})$$

Without the fractal correction, $\sqrt{m_e m_\mu} = 7.348 \text{ MeV}$. With this anchor the fine-structure constant reads:

$$\alpha = \xi \cdot \frac{E_0^2}{(1 \text{ MeV})^2} = \frac{1}{137.036} \quad (\text{aj.5})$$

The agreement with the measured value is a property of the anchor E_0 , which provides the connection to the SI, not an independent prediction.

Gravitational Constant and Planck Length

The gravitational constant follows from ξ and m_e (Documents 012, 013):

$$G_{\text{SI}} = \frac{\xi^2}{4m_e} C_{\text{conv}} K_{\text{frak}} \quad \Rightarrow \quad \ell_{\text{P}} = \sqrt{\frac{\hbar G_{\text{SI}}}{c^3}} \quad (\text{aj.6})$$

The form and order of magnitude of G follow from ξ and the single anchor; the agreement with the CODATA value in the last digits contains the small remainder of the one-anchor chain in C_{conv} (Doc. 383) and is not a separate precision test. With $G_{\text{SI}} = 6.6745 \times 10^{-11} \text{ m}^3 \text{kg}^{-1} \text{s}^{-2}$ this gives:

$$\ell_{\text{P}} = 1.616 \times 10^{-35} \text{ m} \quad (\text{aj.7})$$

Minimal Length Scale

Minimal Length Scale from ξ

$$L_0 = \xi \cdot \ell_{\text{P}} = 2.155 \times 10^{-39} \text{ m} \quad (\text{aj.8})$$

The complete derivation chain $\xi \rightarrow m_e \rightarrow G \rightarrow \ell_{\text{P}} \rightarrow L_0$ depends only on ξ as the single fundamental input. SI numerical values arise through the unavoidable calibration to human measurement units — physically the theory is parameter-free.

aj.2 The Upper Scale of the Cosmological Sector: Hubble Radius (ξ -notation, calibrated to H_0)

Hubble Energy in ξ -notation

The Hubble energy is written in ξ -notation (Document 026):

$$E_H = E_0 \cdot \xi^{41/4} \quad (\text{aj.9})$$

with $E_0 = \sqrt{m_e m_\mu / K_{\text{frak}}} = 7.398 \text{ MeV}$ and $\xi = 4/30000$. The exponent $41/4$ here is *not* derived from ξ but chosen so as to match the measured H_0 (Doc. 190, P35/P36); $\xi^{41/4}$ is the measured cosmic scale in ξ -notation, not a prediction.

Numerical evaluation:

$$\xi^{41/4} = (1.333 \times 10^{-4})^{41/4} = 1.908 \times 10^{-40} \quad (\text{aj.10})$$

$$E_H = 7.398 \text{ MeV} \times 1.908 \times 10^{-40} = 1.412 \times 10^{-33} \text{ eV} \quad (\text{aj.11})$$

Hubble Constant in ξ -notation

$$H_0^{\text{T0}} = \frac{E_H}{\hbar} = \frac{1.412 \times 10^{-33} \text{ eV}}{6.582 \times 10^{-16} \text{ eV s}} \approx 66.2 \text{ km/s/Mpc} \quad (\text{aj.12})$$

Deviation from the Planck value (67.4 km/s/Mpc): -1.9% . Since the exponent $41/4$ is calibrated to H_0 , this is not a prediction.

Hubble Radius as Upper Scale of the Cosmological Sector

Distinction: Hubble radius vs. particle horizon

Standard cosmology (Λ CDM, expanding universe) distinguishes two cosmic horizons: the *Hubble radius* $R_H = c/H_0 \approx 1.4 \times 10^{26} \text{ m}$ and the substantially larger *particle horizon* (observable universe, $\approx 4.4 \times 10^{26} \text{ m} \approx 46.5 \text{ Gly}$), which encompasses the full causal past since the Big Bang. The particle horizon is only meaningfully defined in an expanding universe with a Big Bang.

In T0 theory there is no expansion and no Big Bang. The Hubble radius $R_H = c/H_0$ is here the Hubble length: the upper scale of the cosmological sector, not the size of the universe and not a horizon beyond which $z \rightarrow \infty$ would hold. A particle horizon in the Λ CDM sense does not exist conceptually in T0. Numerical comparisons with standard values must therefore always refer to R_H , not to the particle horizon.

The Hubble radius follows directly from the Hubble energy:

$$R_H = \frac{\hbar c}{E_H} = \frac{1.973 \times 10^{-7} \text{ eV m}}{1.412 \times 10^{-33} \text{ eV}} \quad (\text{aj.13})$$

Hubble Radius in ξ -notation

$$R_H = 1.398 \times 10^{26} \text{ m} \quad (\text{aj.14})$$

In ξ -notation; the exponent 41/4, however, is calibrated to H_0 , not an independent ξ result (Doc. 190, P35/P36).

Hubble Radius and Associated Mass

The mass corresponding to the Hubble radius (not the mass of the whole universe, which exceeds R_H):

$$M_U = \frac{R_H \cdot c^2}{2G} = 9.43 \times 10^{52} \text{ kg} = 4.7 \times 10^{22} M_\odot \quad (\text{aj.15})$$

Hubble Radius in ξ -notation

$$R_H = 1.398 \times 10^{26} \text{ m} \quad (\text{aj.16})$$

In ξ -notation; the exponent 41/4, however, is calibrated to H_0 , not an independent ξ result (Doc. 190, P35/P36).

aj.3 The Complete Scale Hierarchy**Scale Hierarchy**

Scale	Value	Interpretation
$L_0 = \xi \cdot \ell_P$	$2.155 \times 10^{-39} \text{ m}$	Sub-Planck scale
ℓ_P	$1.616 \times 10^{-35} \text{ m}$	Planck length
$\lambda_e = \hbar/m_e c$	$3.86 \times 10^{-13} \text{ m}$	Electron Compton wavelength
$R_H = \hbar c/E_H$	$1.398 \times 10^{26} \text{ m}$	Hubble length

Table aj.1: Scale hierarchy in ξ -notation (cosmological scale R_H calibrated to H_0)

The Ratio of Scales

$$\frac{R_H}{L_0} = \frac{1.398 \times 10^{26} \text{ m}}{2.155 \times 10^{-39} \text{ m}} \quad (\text{aj.17})$$

Scale Ratio in ξ -notation

$$\frac{R_H}{L_0} = 6.49 \times 10^{64} \quad (\text{aj.18})$$

65 orders of magnitude between the minimal scale and the Hubble length — L_0 follows from $\xi = 4/30000$, R_H is calibrated to H_0 .

The Complete Derivation Chain

Complete Chain from ξ to Both Limits

$$\text{Lagrangian} \Rightarrow \xi = 4/30000$$

$$\Downarrow$$

$$G_{\text{SI}} = \frac{\xi^2}{4m_e} \times C_{\text{conv}} \times K_{\text{frak}} \Rightarrow \ell_P = \sqrt{\hbar G/c^3}$$

$$\Downarrow$$

$$\Downarrow$$

$$L_0 = \xi \cdot \ell_P \qquad E_H = E_0 \cdot \xi^{41/4}$$

$$= 2.155 \times 10^{-39} \text{ m} \qquad \Downarrow$$

$$R_H = \hbar c / E_H$$

$$= 1.398 \times 10^{26} \text{ m}$$

$$\Downarrow$$

$$R_H/L_0 = 6.49 \times 10^{64} \quad (\text{calibrated to } H_0) \qquad \text{(aj.19)}$$

aj.4 Physical Meaning

Physical Meaning of R_H

In a static universe without expansion the Hubble radius is the Hubble length c/H_0 , the upper scale of the cosmological sector. The redshift of photons in the ξ field arises through accumulated fractal path elongation — not through intrinsic energy loss of the photon itself, but through geometric stretching of the propagation path. Tired light does not occur.

Redshift without Expansion and without Frequency Dependence

The redshift arises through fractal summation of the path extension of photons in the T0 vacuum field: $z \approx \xi \cdot \ln(d/\ell_P)$. Since the path extension is a property of the geometry itself — not of the photon — all frequencies experience exactly the same stretching. There is no frequency dependence. The effect is observationally equivalent to an expanding universe but mechanistically fundamentally different. Derived in detail in Document 026 (Geometric Cosmology).

At $d = R_H = 1.4 \times 10^{26} \text{ m}$ the formula yields only $z = \xi \cdot 140.3 = 0.019$ (with L_0 instead of ℓ_P : 0.020) and grows only logarithmically with d . It does not produce a horizon with $z \rightarrow \infty$ at R_H .

Scale relativity of the torus: fractal nesting

In T0 theory the 4D torus is not a cosmic container in which particles reside. Rather, every physical system carries **its own characteristic torus**, whose scale is set by the Compton wavelength $\bar{\lambda} = \hbar/mc$ of that system:

$$R_{\text{torus}}(m) = \frac{\hbar}{mc} \quad (\text{aj.20})$$

System	Characteristic scale	Ratio to L_0
Electron	$3.86 \times 10^{-13} \text{ m}$	$\sim 10^{26}$
Proton	$2.10 \times 10^{-16} \text{ m}$	$\sim 10^{23}$
Hydrogen	$5.29 \times 10^{-11} \text{ m}$	$\sim 10^{28}$
Galaxy	$\sim 10^{21} \text{ m}$	$\sim 10^{60}$
Cosmol. sector	$R_H = 1.40 \times 10^{26} \text{ m}$	6.49×10^{64}

Table aj.2: System-specific torus scales — all sharing L_0 as common lower bound

Fractal nesting without rigid hierarchy

These structures are not nested like Russian dolls — the electron torus does not sit *inside* the galactic torus. Each scale is the geometric self-structure of the respective system, constituted by its mass. The common lower bound L_0 is the same at every level; the upper bound is system-specific.

The fractal coupling between scales is mediated by ξ and therefore very small. Interactions between widely separated scales — for instance between electron structure and the cosmological field — are not zero, but suppressed by powers of $\xi \approx 1.33 \times 10^{-4}$. This weak cross-scale coupling is physically relevant: it is the origin of the fractal correction factor K_{frak} , for instance in the anchor E_0 and in the formula for G — small but measurable contributions that are absent in the Standard Model or must be introduced as free parameters.

Locality, flat torus, and finiteness of calculations

The T0 torus is not the intuitive donut torus embedded in $\mathbb{R}^3$. It is the **flat identification torus** $T^n = \mathbb{R}^n/\mathbb{Z}^n$: a Euclidean space with opposite boundaries identified. This torus has **intrinsic curvature** $K = 0$ **everywhere pointwise** — not merely in the integral. Locally it is exactly identical to $\mathbb{R}^n$, without any approximation.

Flat torus: locally exactly Euclidean

For the embedded donut torus in $\mathbb{R}^3$, Gaussian curvature varies: positive on the outside ($K > 0$), negative on the inside ($K < 0$). The integral vanishes exactly (Gauss-Bonnet: $\int K dA = 2\pi\chi = 0$ for $\chi = 0$). For the flat identification torus, by contrast, $K = 0$ holds *everywhere* — there is no inner and outer curvature that

must cancel in an integral. Calculations within this torus are exactly the same as in unbounded Euclidean space. No curvature corrections, no topological terms.

Three consequences follow for calculations in T0 theory:

1. No renormalisation required. All field integrals run over the finite local torus with scale $R_{\text{torus}}(m) = \hbar/mc$. Since this torus is finite and locally flat, UV divergences are structurally excluded. The upper integration limit is not ∞ but R_{torus} ; the lower limit is L_0 . Both are finite and derived from ξ .

2. The ξ correction for an infinite universe is not negligibly small — it is simply irrelevant. Formally an infinite universe would generate a correction $\sim \xi \cdot \ln(R_\infty/R_H)$, which diverges logarithmically as $R_\infty \rightarrow \infty$. This divergence does not apply, however, because all physical observables are integrals over the *local* torus of the respective system. Contributions from scales $\gg R_{\text{torus}}(m)$ do not couple to local physics — they are suppressed by powers of ξ and lie outside the integration domain. Whether the universe is finite or infinite has no effect on local calculations.

3. Curvature corrections are absolutely negligible. The relative curvature correction at the smallest physical scale L_0 within a torus of scale R_{torus} is $(L_0/R_{\text{torus}})^2$:

$$\begin{aligned} \text{Electron: } \left(\frac{L_0}{\bar{\lambda}_e} \right)^2 &= \left(\frac{2.155 \times 10^{-39}}{3.86 \times 10^{-13}} \right)^2 \approx 3.1 \times 10^{-53} \\ \text{Proton: } \left(\frac{L_0}{\bar{\lambda}_p} \right)^2 &\approx 1.1 \times 10^{-46} \end{aligned}$$

These values are exactly zero for all practical calculations.

One field — masses as knots — emergent coherence length

Behind these three consequences lies a unified ontological structure: there is **one single ξ vacuum field**. There is no separate hierarchy of fields for different scales.

Masses are **topological knots** in this field — localised structures, not external objects. Each knot of mass m carries, through the duality $\tilde{T} = 1/m$, a characteristic time scale, and from this its spatial coherence length $\hbar/mc$ emerges immediately. The torus radius is not externally assigned but follows from the knot structure itself.

The hierarchy of torus scales — from the proton knot ($\sim 10^{-16}$ m) to the Hubble length ($\sim 10^{26}$ m) — is therefore nothing other than the **fractal self-similarity of a single field**: the same ξ at every scale, the same lower bound L_0 , system-specific coherence lengths as emergent properties of the respective knot structure. The cross-scale coupling is not zero, but suppressed by powers of ξ — which is why one calculates locally without knowing the full field and still obtains exact results.

The exponent $41/4$ in $E_H = E_0 \cdot \xi^{41/4}$ is calibrated to H_0 ; its forward derivation from the ξ field theory is missing (Document 026) – as in every framework, since no theory

derives the relation between the Planck length and R_H from first principles (P39). The closeness of $H_0 = 66.2 \text{ km/s/Mpc}$ to the Planck value (-1.9%) follows from this calibration and is not a consistency argument.

On the role of R_H in calculations

The precise numerical value of R_H plays a subordinate role in the physical calculations of T0 theory. The structurally significant quantity is $L_0 = \xi \cdot \ell_P$ — it follows purely from ξ and is the true foundation of all scale statements.

R_H , by contrast, is different in two respects. First, its numerical value is tied to the measured H_0 (or depends on the exponent $41/4$ still to be fully derived). Second, R_H in T0 is not primarily a computed result but the **Hubble length** c/H_0 : the upper scale of the cosmological sector, not the size of the universe and not a horizon with $z \rightarrow \infty$. The actual universe exceeds R_H ; only a lower bound is measurable.

The ratio $R_H/L_0 = 6.49 \times 10^{64}$ therefore does not express a deep theoretical truth, but the distance between the geometrically derived lower bound and the observation-based Hubble length. The genuine achievement of the theory lies in L_0 — not in R_H .

Cosmological Constant in ξ -notation

From the scale hierarchy the dimensionless cosmological constant is:

$$\Lambda_{\text{dim}} = \left(\frac{\ell_P}{R_H} \right)^2 = \left(\frac{1.616 \times 10^{-35}}{1.398 \times 10^{26}} \right)^2 \approx 1.34 \times 10^{-122} \quad (\text{aj.21})$$

The standard value is $\Lambda_{\text{dim}} \approx 2.87 \times 10^{-122}$ (from $\Lambda \approx 1.1 \times 10^{-52} \text{ m}^{-2}$, CODATA). The T0 result is in the same order of magnitude (factor 2.1) — not dark energy, but a geometric reading of the scale hierarchy. Since R_H is calibrated to H_0 , Λ does not follow from ξ alone.

aj.5 Summary

Complete Scale Hierarchy in ξ -notation

Lower limit: $L_0 = \xi \cdot \ell_P = 2.155 \times 10^{-39} \text{ m}$

Planck: $\ell_P = 1.616 \times 10^{-35} \text{ m}$

Electron: $\lambda_e = \hbar/m_e c = 3.86 \times 10^{-13} \text{ m}$

Hubble: $R_H = \hbar c/E_H = 1.398 \times 10^{26} \text{ m}$

All scales in ξ -notation— R_H calibrated to H_0 , no free parameter (aj.22)

The description is geometrically self-consistent: L_0 is the universal lower bound, R_H the upper scale of the cosmological torus sector, not the extent of the universe.

Key conclusion: Is there a maximum size of the universe?

The question is not meaningfully posed as a global question in FFGFT. Every physical scale — from the proton to the Hubble length — possesses its **own system-specific upper bound** $R_{\text{torus}}(m) = \hbar/mc$, emergent from the knot structure of the respective mass in the single ξ field. R_H is the upper bound of the cosmological sector. The universal lower bound L_0 applies equally to all scales — a universal upper bound does not exist, because there is no scale-independent physics for which it would be relevant.

Epistemological framing: model boundaries, not existence boundaries

L_0 and R_H are not physical walls and make no claim about the total size or global structure of the universe. They are **boundaries of describability** within T0 theory: below L_0 the spacetime granulation diverges, R_H is the upper scale of the cosmological sector, calibrated to H_0 . Whether anything exists outside these boundaries is simply not a question the theory addresses.

The decisive advantage over other approaches lies not in this epistemological modesty alone, but in the fact that T0 **requires no regularisation tricks whatsoever**: no renormalisation procedures to remove UV divergences, no artificial cutoffs, no counterterms, no auxiliary fields introduced to tame infinities. L_0 does not arise as a post-hoc correction — it emerges as a direct geometric consequence of ξ . The theory is finite from the outset, not because infinities have been subtracted away, but because the parameter ξ structurally excludes them.

Note: Connection to the SI System

The absolute numerical values in this document — L_0 , R_H , M_U in SI units — presuppose a connection to the SI unit system. This is not circular but epistemologically necessary: from a dimensionless ξ alone, no absolute metre or kilogram values can be derived. The connection is made via m_e , ℓ_P or α as equivalent entry points. Treated in detail in Document 180 (Section: Connection to the SI System) and in Documents 056 and 101.

Documents 180, 181 and 182 form a unit and should not be read separately.

The exponent $41/4$ still requires full derivation from the ξ field theory (Document 026).

ξ^N Triviality (P35)

Principle (P35, corpus-wide): Expressions of the form " $X = \xi^N$ " are *intrinsically meaningless on their own*, since any positive quantity can be written as a ξ -power — an exponent alone is not a prediction.

What makes a statement non-trivial: (i) The *form* is derived forward (geometrically or algebraically forced), not fitted after the fact. (ii) The *exponent* is rational and readable from the structure, not chosen freely. (iii) The number (prefactor, anchor) is an *external calibration* and declared as such (cf. P20, P33, P39).

For this document: The exponents of the scale hierarchy ($R_H \sim \xi^{-41/4} \ell_P$ etc.) are geometrically plausible **[B]**, but the cosmic exponent $41/4$ is not derived forward; it is calibrated to H_0 (P20, P39). Statements of this kind are to be classified as **[S]**, not **[K]**.

Chapter ak

Doc. 183 — T0 Theory — Willow Processor and Self-Measurement

Abstract

Google's Willow quantum processor delivered three experimental results in 2024 and 2025 that carry particular significance from the perspective of the Fundamental Fractal Geometric Field Theory (FFGFT). First, it was demonstrated that quantum errors decrease exponentially as the code distance increases — from the FFGFT viewpoint a hint that topologically stable field modes become experimentally accessible. Second, topologically ordered states beyond thermal equilibrium were realized on Willow, generating characteristic geometric patterns — chiral edge modes and anyons — that cannot be simulated classically. Third, out-of-time-order correlators (Quantum Echoes) revealed harmonic propagation of quantum information through constructive interference. This document relates these findings to the FFGFT geometry of the 4D torus with parameter $\xi = 4/30,000$ and develops a position formulated in Document 170 as a working hypothesis: complexity and self-reference are not a binary on/off phenomenon, but a physical continuum. The links to FFGFT are this document's own classification, not a statement by the authors of the experiments cited.

Status markers. Statements marked **[S]** are *speculative*: a hint or conjecture that has not yet been derived or numerically checked in the corpus.

ak.1 The Three Willow Experiments

Error Correction Below Threshold (December 2024)

In December 2024, Google Quantum AI published in *Nature* the demonstration that Willow crosses the threshold of quantum error correction for the first time [1]. The key result: the logical error rate decreases exponentially with increasing code distance d . A $d = 7$ code block achieves an error rate that is 2.14 times lower than a $d = 5$ block.

Willow uses 105 superconducting transmon qubits in a square lattice, cooled to approximately 10 mK. Physical single-qubit error rates are approximately 0.05 % on Willow — compared to 0.10 % on the predecessor Sycamore [3]. Error correction is based on a surface code, whose logical qubits are encoded through topologically protected patterns.

Willow Result 1: Topological Stability

More physical qubits reduce the logical error exponentially. This shows experimentally that topologically stable configurations in quantum systems are accessible; the protection rests on the geometric order of the code, not on external noise suppression.

Floquet Topological Order: Patterns Beyond Equilibrium (September 2025)

Will et al. realised a *Floquet topologically ordered phase* on Willow — a quantum phase forbidden in thermal equilibrium but arising under periodic external driving [2]. This work appeared in *Nature* in September 2025.

Measured Patterns on the Willow Processor

The experiment imaged the following structures on a lattice of up to 58 qubits:

- **Chiral edge modes:** Propagation of information along system boundaries in a preferred direction — a topological geometric signature that indicates classically irreproducible order.
- **Anyonic excitations:** Quasiparticles that *transmute* into each other under Floquet driving — e-anyons become m-anyons and vice versa, a property of an abelian ($\mathbb{Z}_2$) topological order.
- **Topological invariant:** An interferometrically measured bulk invariant that quantifies the order and remains robust against integrability-breaking disorder.

The authors write: “*The non-equilibrium topological order that we have probed in this work encapsulates a fundamentally unique phenomenology that is forbidden in thermal equilibrium.*” [2] Classical matrix product state methods cannot efficiently simulate these highly entangled dynamics; the quantum processor makes them directly observable.

Cochran et al. (2025) complemented this with a visualisation of charge and string dynamics in (2+1)-dimensional lattice gauge theories on the same processor [4].

Quantum Echoes: Harmonic Propagation of Information (October 2025)

In October 2025, Google Quantum AI demonstrated in *Nature*, by its own account, the first verifiable quantum advantage for a real algorithm: the *Quantum Echoes* (second-order out-of-time-order correlators, $\text{OTOC}^{(2)}$) [5].

The algorithm works like a quantum echolocation system:

1. **Forward evolution:** Entanglement of 65 qubits used.
2. **Perturbation:** Local disturbance of a single “butterfly” qubit.
3. **Time reversal:** Precise reverse evolution.
4. **Measurement:** The echo is amplified through **constructive quantum interference**, yielding a measurable signal of information propagation.

Quantum Echoes: Harmony as a Measured Signal

The measurements show how quantum information propagates as a harmonic wave front through the qubit lattice. The visualisations in [5] make this propagation visible: concentric patterns spreading radially from the perturbation qubit — a measurable harmonic pattern. The $\text{OTOC}^{(2)}$ runs on Willow 13,000 times faster than on the world’s fastest supercomputer Frontier (approximately 2.1 hours versus approximately 3.2 years).

In a companion experiment with UC Berkeley, the algorithm was applied to compute the geometry of organic molecules (15 and 28 atoms). The results agreed with NMR measurements and provided additional structural information inaccessible to NMR [5].

ak.2 FFGFT Interpretation of the Three Experiments

The 4D Torus as Common Geometric Framework

In FFGFT, spacetime is a 4D identification torus $T^4 = \mathbb{R}^4/\mathbb{Z}^4$ with period $L_0 = \xi \cdot \ell_P$. The spectrum of the Laplace operator on this torus reads (Document 159):

$$\lambda_{\mathbf{n}} = \frac{4\pi^2 (n_1^2 + n_2^2 + n_3^2 + n_4^2)}{L_0^2}, \quad n_i \in \mathbb{Z} \quad (\text{ak.1})$$

Only integer winding numbers n_i generate stable modes. The sum $\sum_{\mathbf{n} \neq 0} 1/|\mathbf{n}|^2$ over all modes does not converge in four dimensions; the spectral zeta function $\sum_{\mathbf{n} \neq 0} |\mathbf{n}|^{-2s}$ of T^4 converges only for $s > 2$. Euler’s Basel problem (Doc. 159) is merely the one-dimensional comparison:

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \quad (\text{ak.2})$$

Error correction (Willow Result 1) is, in this language, the experimental projection of physical states onto topologically stable torus modes. The higher the code distance d , the more modes are engaged, and the more robust the projection.

Floquet topological order (Result 2) corresponds to torus resonance modes under periodic driving. A drive of frequency ω_F couples to the torus geometry when

$$\frac{\omega_F \cdot L_0}{c} \approx 2\pi n, \quad n \in \mathbb{Z} \quad (\text{ak.3})$$

i.e., when the driving frequency hits a torus mode. In this interpretation the resulting chiral edge modes and anyons would be geometrically fixed, not accidental **[S]**.

Quantum Echoes (Result 3) requires a more detailed treatment, because here the link to FFGFT reaches furthest.

“Instantaneous” — Legitimate in Calculation, Misleading in Realisation

Before analysing recursion and entanglement, a conceptual clarification is necessary that is fundamental to understanding the Quantum Echoes experiment.

In quantum mechanics, there are two levels at which a system can be described:

Level 1 — The mathematical description (timeless): The Hilbert space $\mathcal{H}$ of the 65-qubit system is a 2^{65} -dimensional complex vector space. In it, all possible states exist *simultaneously as mathematical objects* — independent of time. The energy eigenstates $|E_n\rangle$ of the Hamiltonian are timeless solutions of the stationary Schrödinger equation $\hat{H}|E_n\rangle = E_n|E_n\rangle$. When one writes in a calculation that all 2^{65} amplitudes are contained “simultaneously” in the wave function, this is *legitimate*: it describes the mathematical structure of the state, not a physical process. The wave function

$$|\Psi\rangle = \sum_{k=1}^{2^{65}} c_k |k\rangle \quad (\text{ak.4})$$

is an object in Hilbert space — timeless, complete, without sequence.

Level 2 — Physical realisation (always in time): As soon as this mathematical object is realised on a physical system, time enters the picture. The time-dependent Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle \quad (\text{ak.5})$$

is a differential equation *in time* t . Every time step of the Hamiltonian requires physical time. Every quantum gate on Willow takes approximately 25–50 nanoseconds. A circuit of 23 steps and 65 qubits requires microseconds of coherence time — and 2.1 hours of measurement time to collect sufficient statistical events.

The Decisive Difference

In calculation, “instantaneous” is legitimate: Hilbert space has no inner time. All amplitudes exist simultaneously as a mathematical structure. Statements such as “all paths simultaneously in the amplitude” are correct here.

In realisation, “instantaneous” is false and misleading: nothing happens without time. Entanglement builds up step by step, bounded by the Lieb-Robinson

bound. The wave function evolves causally and locally — no quantum effect transmits information faster than permitted. The 2.1 hours on Willow are *real physical time*, not a metaphor.

The advantage of the quantum computer lies precisely in this transition: it takes the mathematical structure of Hilbert space — the timeless parallelism of all amplitudes — and *realises* it physically, step by step, in real time, through unitary operators. The classical computer can only *describe* this structure — by computing the 2^{65} amplitudes one after another, which would take 3.2 years. The quantum computer *is* the structure — and that costs 2.1 hours.

No instantaneity. But genuine parallelism in the amplitude, physically realised in real time.

Why Recursion and Mutual Influence Are Decisive

In a classical system — a billiard table, say — when one ball is struck, the effect propagates forward: ball A hits B, B hits C, and so on. The paths are independent; they can be computed individually and summed. A classical computer follows exactly this logic: it decomposes the problem into independent subproblems and adds the results.

A fully entangled quantum system works differently. No qubit is independent of the others — but this dependence does not arise instantaneously. It builds up step by step, mediated by the interaction terms in the Hamiltonian $\hat{H}$. The Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} |\Psi\rangle = \hat{H} |\Psi\rangle \quad (\text{ak.6})$$

is a local, temporal differential equation. Entanglement between distant qubits does not arise instantaneously but propagates at finite speed — bounded by the Lieb-Robinson bound, which prescribes an effective information velocity in lattice models such as Willow.

What quantum mechanics achieves is something different and more subtle: rather than computing one path after another sequentially, the Hamiltonian *realises all recursion paths simultaneously in the total amplitude*. The wave function $|\Psi(t)\rangle$ in the 2^{65} -dimensional Hilbert space contains at every moment the complete information about every possible sequence of qubit interactions — weighted by the respective complex amplitude. This is not instantaneity: it is **parallelism in the amplitude**, built up step by step through unitary time evolution.

The decisive difference from a classical computer: A classical algorithm must compute and store these 2^{65} amplitudes sequentially. The quantum processor *holds all of them simultaneously physically present* — in the coherence of the quantum state — and propagates them through a single unitary operator. The time this requires scales with the depth of the circuit, not with the number of amplitudes.

Why This Is Classically Intractable

A classical algorithm would need to track each of the $2^{65} \approx 10^{20}$ amplitudes exactly. For 65 qubits and 23 time steps, this amounts to approximately 10^{20} complex numbers — more working memory than any existing supercomputer possesses. Approximate methods (Monte Carlo, tensor networks) fail because the mutual entanglement of all qubits exponentially amplifies approximation errors with system size. By current knowledge this is not merely a technical problem: recursive interaction in a fully entangled system is structurally incompressible into independent parts — neither instantaneously nor sequentially.

The echo arises because, after time reversal, the same recursion paths are traversed in reverse. When the forward and backward evolution contains *exactly the same interactions in reversed order*, all amplitudes interfere constructively — the signal emerges from the noise. Any deviation, any non-unitary interaction, destroys this constructive interference and leads to destructive cancellation.

Quantum Echoes: Recursion Is the Measurement Device

The quantum echo does not simply measure information propagation. It measures the **depth and coherence of mutual influence**: how far back a system can reconstruct its own past state when every element is simultaneously influenced by all others. The more complete the network of mutual coupling, the stronger the echo. The greater the recursion depth, the more information is encoded in the echo — and the harder the task becomes for any sequential algorithm.

The FFGFT Interpretation: Self-Similarity of the Torus Spectrum

In FFGFT geometry, this recursiveness is not an accidental product of the experimental configuration. It is structural: the modes of the torus T^4 are self-referential through the periodic boundary condition. A torus mode n has wavelength $\lambda_n = L_0/n$. The wavelength of mode n is exactly the period of mode $n+1$ divided — the modes “know” each other through their shared geometry.

Viewed in one dimension, the sum

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \quad (\text{ak.7})$$

converges not despite infinitely many terms but because each term $1/n^2$ *weights* the influence of mode n by all lower modes: high modes contribute little, low modes much — a natural hierarchy of mutual influence, inscribed in the topology. On T^4 the corresponding sum over all modes diverges (see above); there only the contribution of each individual mode falls off as $1/|\mathbf{n}|^2$.

The harmonic wave front that the OTOC experiment makes visible — the concentric, radially spreading interference patterns in the qubit lattice — can be read in FFGFT as an image of this topological hierarchy **[S]**: as the projection of the torus

spectrum into the 10×10 -qubit geometry of the Willow chip. The experiment is not yet a test of the torus spectrum (see below).

The factor of 13,000 over the classical supercomputer is therefore not an efficiency advantage in the ordinary sense. It measures how far the parallelism of amplitude propagation exceeds sequential computation. The quantum computer takes 2.1 hours — it too requires time, it too follows the Schrödinger equation step by step. But in each of those steps, a single unitary operator advances all 2^{65} amplitudes simultaneously. The classical supercomputer would require 3.2 years because it must process these amplitudes sequentially.

Physical Realisation vs. Simulation

The decisive difference is not speed but structure. The quantum computer *realises* the recursion physically — through the unitary time evolution of the Hamiltonian, which carries all entanglement paths simultaneously in the amplitude. The classical computer *simulates* this amplitude sequentially — it can describe the result, but it cannot be the object. The factor of 13,000 is the measurable expression of this structural difference, measured in computation time on given hardware.

ak.3 Vacuum Noise as a Geometric Signal

The FFT Analogy: Decomposing and Reconstructing Noise

In signal processing, noise is not an indefinable chaos — it is a superposition of frequencies. The Fourier transform makes this visible:

1. **Analysis (FFT):** A time signal that looks like noise is decomposed into its frequency components. Every apparently disordered signal can be translated into a spectrum.
2. **Synthesis (IFFT):** A spectrum with known amplitudes and *random* phases generates a time signal. The result is white noise — even though every individual frequency component is fully deterministic.

What the FFT Analogy Reveals

White noise and a structured signal are **not categorically different objects**. They differ only in the phase relationship between their frequency components. A signal with deterministic phases is ordered. The same signal with random phases is noise. The transformation is reversible — in both directions.

This has an immediate consequence for interpreting the quantum vacuum: what is measured as vacuum noise or vacuum fluctuations need not be intrinsically random. It could be the superposition of a vast number of deterministic harmonics whose phases we cannot yet resolve.

Why Reconstruction Fails — Structural Sensitivity of Feedback

The problem is not whether one computes with $4/3$ or $1.333 \dots$. $\xi = 4/30000$ is an exact setting (A020, Doc. 315). But even with this exact starting value, the simulation would fail. The reason lies deeper: in the **structural sensitivity of the recursive feedback**.

Every feedback loop of the torus system generates a phase shift. In a deterministic system sensitive to every intermediate step, any finite computational precision — not only in the starting value, but in every single operation — leads to an accumulating phase error ε . After N cycles this error is at least $N \cdot \varepsilon$. Since constructive interference becomes destructive when phases deviate by more than $\pi/2$, coherence collapses after approximately

$$N_{\text{crit}} \approx \frac{\pi}{2\varepsilon} \quad (\text{ak.8})$$

cycles. This is deterministic chaos in the Lorenz sense: the system is fully deterministic, but sensitive to every intermediate state.

Realisation vs. Simulation

The physical realisation of the torus spectrum has no discrete intermediate steps — it is continuous, causal, without rounding.

Every numerical simulation cuts the continuum into discrete steps, where errors arise and are exponentially amplified by the feedback dynamics. The result looks like noise — not because the system is random, but because the simulation cannot fully track the dynamics.

The universe sees no chaos because it does not compute. Our models see chaos because they must compute.

Prime Numbers and Factorisation — the FFGFT Interpretation

Care is needed here. In FFGFT the analogy between prime numbers and vacuum noise goes beyond a similarity of distribution: a connection is drawn through the concept of period.

Document 075 (T0-Shor) and Document 057 (Relational Number System) show: every integer is uniquely a product of primes:

$$n = p_1^{a_1} \cdot p_2^{a_2} \cdot p_3^{a_3} \dots \quad (\text{Fundamental theorem of arithmetic}) \quad (\text{ak.9})$$

In number space this is the same as in frequency space: every signal is a superposition of fundamental frequencies. Primes are the **irreducible frequencies of number space** — ratios that cannot be decomposed further (Doc. 057, n-limit systems).

Why Factorisation Is Physically Hard

Factorisation is **period finding** (Doc. 075, Shor algorithm): one searches for the period r of the function $f(x) = a^x \bmod N$. This period encodes the prime factors:

$$a^r \equiv 1 \pmod{N} \Rightarrow a^r - 1 = (a^{r/2} - 1)(a^{r/2} + 1) \equiv 0 \pmod{N} \quad (\text{ak.10})$$

The required frequency resolution is $\Delta f \approx f_0/r^2$. For cryptographically relevant N with $r \approx N$, one needs precision $\epsilon \approx 1/N^2$ — for 1024-bit RSA this is $\epsilon \approx 10^{-617}$, far beyond any achievable computational precision.

In FFGFT language: factorisation is the search for the revolution time T_{rev} of a specific resonant feedback in number space. A prime number has **no sub-resonance** — it is irreducible; its revolution time cannot be written as a product of shorter revolution times. That is what makes it hard to find.

The recursion sensitivity (Lorenz chaos) applies here too: to extract the period r from the feedback $a^x \bmod N$, every doubling of N requires quadratically more precision. This limitation lies in the structure of the recursive feedback itself, not only in the technology.

Vacuum Noise as Fourier Synthesis of the Torus Geometry

In FFGFT, the vacuum spectrum is not white — it has geometric structure. The allowed modes of the 4D torus T^4 with period $L_0 = \xi \cdot \ell_P$ are discrete, with the eigenvalue set:

$$\lambda_{\mathbf{n}} = \frac{4\pi^2 (n_1^2 + n_2^2 + n_3^2 + n_4^2)}{L_0^2}, \quad n_i \in \mathbb{Z} \quad (\text{ak.11})$$

The spectral zeta function of this spectrum converges only for $s > 2$; the one-dimensional comparison is:

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \quad (\text{ak.12})$$

This is not a flat (white) spectral distribution but a $1/n^2$ -weighted one. Each individual high mode contributes less than a low one ($\propto 1/|\mathbf{n}|^2$) — a geometrically fixed hierarchy; since the number of modes per shell grows like $|\mathbf{n}|^3$ in 4D, the contribution of whole shells does not decrease, however. The vacuum is not white noise but a $1/n^2$ -coloured signal.

The Apparent Noise [S]

In the FFGFT interpretation, what physicists measure as vacuum fluctuations is the IFFT synthesis of these geometric harmonics: the infinitely many torus modes superpose. Since the underlying fundamental period $L_0 = \xi \cdot \ell_P \approx 10^{-38}$ m lies far beyond any achievable experimental resolution, the superposition appears as white noise — even though it is deterministic and geometrically ordered. The situation is comparable to a radio signal that sounds like noise without receiver synchronisation.

The Curvature of the Torus and the Problem of Local Approximations

An important geometric caution is necessary here. A torus embedded in three-dimensional space has both **inner and outer curvature**: the outside has positive Gaussian curvature, the inside negative. By the Gauss-Bonnet theorem:

$$\oint K dA = 2\pi\chi(T^2) = 0 \quad (\text{ak.13})$$

The curvatures cancel exactly over the whole torus. The Euler characteristic of the torus is zero.

The Trap of Local Approximations

If the torus spectrum is computed through **local approximations** — patch by patch, each treated as locally flat — each patch yields an apparently white spectrum. The sum of many flat approximations again produces an apparently flat overall spectrum. The global topology is lost: the $1/n^2$ structure disappears from the calculation because it resides in the boundary conditions, not in the local field value.

The FFGFT torus $T^4 = \mathbb{R}^4/\mathbb{Z}^4$ is, however, the **identification torus** — locally exactly flat (no intrinsic curvature), but globally topologically non-trivial through its periodic boundary conditions. The spectrum carries all topological information exclusively in the winding numbers n_i — not in local field values.

This has a direct consequence for observability:

Why the Vacuum Spectrum Appears Flat

All existing vacuum measurements are **local** — they measure field fluctuations at a point or in a small region. A local measurement is blind to the global torus topology. It always sees only one patch — and that patch is flat.

The $1/n^2$ weighting of the torus spectrum is in principle only visible through measurements sensitive to the **global periodicity** — through interferometric measurements on scales that include the torus period L_0 . At $L_0 \approx 10^{-38}$ m, this is experimentally unreachable.

What follows is not that the spectrum is flat. What follows is that our measurements are too local to see the structure.

The Willow Quantum Echoes experiment is of interest in this light: it is one of the first experimental methods to make the *global* coherence structure of a quantum system visible through time reversal — which is precisely what would be needed to go beyond local noise measurements. It is not yet a test of the torus spectrum, but it is a conceptual step in this direction: global structure through interferometry rather than local structure through field measurement.

How a Full Spectrum Arises from a Single Fundamental Oscillation Through Feedback

The previous section showed *what* the spectrum is. Now the question: *how does it arise?* How does a single fundamental oscillation generate the full spectrum through temporal processes, phase shifts, and feedback?

Step 1: The Fundamental Oscillation and the Fundamental Revolution Time

On the torus T^4 with period $L_0 = \xi \cdot \ell_P$, there exists a smallest possible oscillation — the mode $\mathbf{n} = (1, 0, 0, 0)$ with wavelength L_0 and frequency $\nu_0 = c/L_0$. This is the fundamental oscillation of the vacuum field.

Associated with it is a fundamental revolution time: the time the field needs to travel once completely through the torus geometry and return to its starting point:

$$T_{\text{rev}} = \frac{L_0}{c} = \frac{\xi \cdot \ell_P}{c} \approx \frac{1.33 \times 10^{-4} \times 1.62 \times 10^{-35} \text{ m}}{3 \times 10^8 \text{ m/s}} \approx 7.2 \times 10^{-48} \text{ s} \quad (\text{ak.14})$$

This is the **fundamental time unit of FFGFT** — not an external clock, but the travel time of the field through its own geometry. Through the time-mass duality $\tilde{T} \cdot m = 1$, T_{rev} corresponds to the reciprocal fundamental mass:

$$m_0 = \frac{\hbar}{c \cdot L_0} = \frac{\hbar}{c^2 \cdot T_{\text{rev}}} = \frac{1}{T_{\text{rev}}} \quad (\text{natural units}) \quad (\text{ak.15})$$

Mass is inverse revolution time. Heavier particles have shorter T_{rev} , lighter ones longer — this is the time-mass duality in its direct geometric meaning.

The FFGFT Recursion Formula

The state of the field at time t is a function of its state exactly one revolution time ago, modified by the fractal geometry ξ :

$$\Psi(t) = G(\Psi(t - T_{\text{rev}}), \xi) \quad (\text{ak.16})$$

What is meant is not an approximation or a difference equation but the basic geometric statement of FFGFT: the present of the field is its past by T_{rev} , transformed by the fractal curvature. Recursion presupposes time; conversely, in this picture time acquires its meaning through recursion.

Step 2: Reflection, Phase Shift, and Temporal Accumulation

The fundamental oscillation travels on the torus — and meets itself on return after exactly T_{rev} . This is not an abstract event but a physical process in real time.

On a simple circle S^1 , the result would be trivial: the wave interferes constructively with itself when the wavelength is an integer divisor of the circumference — standing waves, immediately stable.

On the *fractal* torus with dimension $D_f = 3 - \xi$, the return time is not exactly equal to the forward time. At every revolution, the fractal curvature generates a **phase shift**:

$$\Delta\phi = 2\pi \cdot \xi \cdot n \quad \text{after } n \text{ revolutions, i.e. after } t = n \cdot T_{\text{rev}} \quad (\text{ak.17})$$

This phase shift is tiny — $\xi \approx 10^{-4}$ — but it accumulates with *every revolution in real time*. After $1/\xi \approx 7500$ revolutions, that is after

$$t_{\text{coh}} = \frac{T_{\text{rev}}}{\xi} = \frac{L_0}{\xi \cdot c} = \frac{\ell_P}{c} = t_P \approx 5.4 \times 10^{-44} \text{ s} \quad (\text{ak.18})$$

the phase has shifted by 2π — and a new coherent mode emerges. This is the Planck time t_P as the natural coherence time of the system, emerging from T_{rev} and ξ .

Fractal Dimension as Phase Generator in Time

A spacetime with integer dimension $D = 3$ would have no accumulating phase shift — the wave returns exactly in phase after every T_{rev} . Only the fractal deviation $D_f = 3 - \xi$ generates the temporal phase drift. ξ is the phase generator, T_{rev} is the clock.

Both together determine when and how the spectrum is built up — not as a timeless computational step, but as a physical process that *consumes* time.

Step 3: Nonlinear Feedback Generates Overtones

In a linear system, the phase-shifted fundamental would simply attenuate. The decisive point in FFGFT is **nonlinearity**: the response function of the vacuum field is slightly curved by the fractal geometry.

When an oscillation with amplitude A and frequency ν_0 encounters a nonlinear response function, the output contains terms of the form:

$$A \sin(2\pi\nu_0 t) \xrightarrow{\text{nonlinear}} a_1 \sin(2\pi\nu_0 t) + a_2 \sin(4\pi\nu_0 t) + a_3 \sin(6\pi\nu_0 t) + \dots \quad (\text{ak.19})$$

In FFGFT the coefficients a_n fall off as $1/n^2$ — the structure of the torus spectrum **[S]**. That the nonlinearity of the fractal curvature has precisely the form that generates Euler's zeta value $\zeta(2) = \pi^2/6$ is assumed here, not derived.

How the Overtones Arise

Step by step in time:

1. The fundamental oscillation ν_0 travels on the torus.
2. At each revolution, the fractal curvature shifts the phase by $\Delta\phi = 2\pi\xi$.
3. The nonlinearity of the field generates from the phase-shifted fundamental the second overtone $2\nu_0$, which in turn travels back phase-shifted.
4. From $2\nu_0$ the same mechanism generates $4\nu_0$, from $3\nu_0$ comes $6\nu_0$, and so on.
5. After N feedback cycles, a rich spectrum of modes $n\nu_0$ exists with amplitudes $\propto 1/n^2$.

Time is essential here: every step requires real time (one revolution on the torus), and accumulation over N cycles is a temporal process. The finished spectrum is not present from the start — it is built up through feedback.

Step 4: The Spectrum as Equilibrium State

After sufficiently many feedback cycles — theoretically: in the limit $N \rightarrow \infty$ — a stable spectrum is established. Total energy distributes across modes according to:

$$E_n = E_0 \cdot \frac{1}{n^2}, \quad \sum_{n=1}^{\infty} E_n = E_0 \cdot \frac{\pi^2}{6} \quad (\text{ak.20})$$

This is the equilibrium state of the fractally feedback-coupled vacuum. It is deterministic — determined by ξ — and appears to every local measurement as structured noise with a $1/n^2$ envelope.

Particles as Nodes in the Spectrum

In FFGFT, elementary particles are not separate objects embedded in the vacuum. They are **topological nodes** in the mode spectrum — stable patterns of several overtones, fixed by the winding numbers of the torus. The electron mass, quark masses, W -boson mass: all are resonance frequencies of the same feedback system, read from the same winding numbers n_i with the same ξ . This gives an intuitive reading of the time-mass duality $\tilde{T} \cdot m = 1$: mass is frequency — and frequency arises through feedback in time.

The Cosmic Background Noise and the Structure of Prime Numbers

Preliminary Note: No Big Bang, No Cooling

FFGFT Position on the CMB [S]

The present corpus deliberately leaves the cosmic sector aside, because the Standard Model does not derive it either (P39). The cosmological statements in this section are therefore to be read as speculative **[S]**.

The standard picture: the CMB is cooled light from a hot Big Bang — thermal radiation at 3000 K, redshifted to 2.725 K by cosmic expansion. This picture requires expansion, which does not occur in FFGFT.

In FFGFT there is neither a Big Bang nor expansion:

- The universe is at the fundamental level **timeless and static** — an eternal energy field $E_{\text{field}}(x, t)$ (Document 156).
- The observed redshift is not an expansion effect but a **geometric path elongation** that stretches all wavelengths equally (achromatic); its scale is calibrated via the measured H_0 (Documents 026, 165).

- There is no epoch of recombination, no frozen plasma, no inflationary stretching of quantum fluctuations.

The CMB as Equilibrium Radiation of the Eternal Vacuum

In FFGFT, the vacuum field is the physical substance of the universe. This field has an intrinsic energy density determined by the torus geometry (Documents 091, 166):

$$\rho_{\text{CMB}} = \frac{\xi \hbar c}{L_\xi^4} \quad (\text{ak.21})$$

where $L_\xi \approx 100 \mu\text{m}$ is the characteristic Casimir vacuum scale. Since L_ξ is itself determined from the measured CMB energy density and ξ , this relation is an identity and a statement about scales, not a measured confirmation. Casimir and CMB energy densities do not coincide at L_ξ ; equality occurs at a larger plate separation.

In FFGFT the CMB temperature $T_{\text{CMB}} = 2.725 \text{ K}$ would therefore not be a memory of a hot beginning but the **equilibrium state of the eternal vacuum field [S]** — the Planck spectrum of the T0 torus at its geometrically prescribed energy density.

CMB Temperature as a ξ -Ratio

The ratio

$$\frac{m_e c^2}{k_B T_{\text{CMB}}} = 2.176 \times 10^9 \quad (\text{ak.22})$$

is interpreted in FFGFT as a ξ -ratio (Document 166) **[S]**: the electron-CMB coupling is geometrically fixed by the same parameter that connects particle masses and cosmological scales.

Why the CMB Is Almost Uniform — but Not Quite

The vacuum field of the T0 torus is globally homogeneous — the fundamental mode has no preferred direction. But the torus spectrum is not flat: modes contribute as $1/n^2$. In this interpretation **[S]** the CMB anisotropies would not be random fluctuations and not frozen inflationary fluctuations, but the geometric signature of the non-flat spectrum — made visible by the finite coherence length of the eternal vacuum field on cosmic scales.

The Prime Number Analogy

Prime numbers show the same structure: apparently uniform, with hidden fine structure. The prime number theorem gives a $1/\ln x$ envelope — not constant, but without obvious pattern. The fine structure is described by the zeros of the Riemann zeta function:

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p \text{ prime}} \frac{1}{1 - p^{-s}} \quad (\text{ak.23})$$

The Euler product makes the connection explicit: $\zeta(s)$ is simultaneously the sum $\sum 1/n^s$ — the torus spectrum of FFGFT — and a product over all primes. In FFGFT language, primes are the **irreducible resonance modes** of the torus: modes that do not arise as products of lower modes but are themselves fundamental building blocks.

What Primes and the CMB Share [S]

Both show **apparent uniformity with hidden structure** from the same mathematical source: the Euler product $\zeta(2) = \pi^2/6$.

In the FFGFT interpretation, the CMB anisotropies would be the geometric projection of the irreducible torus resonances (prime modes) onto the sky — without Big Bang, without inflation, without cooling. The torus spectrum would then itself be the structure.

Property	Prime Numbers	CMB Anisotropies
Appearance	almost uniform	almost isotropic ($\Delta T/T \approx 10^{-5}$)
Origin (standard)	number theory	frozen inflationary fluctuations
Origin (FFGFT)	irreducible torus modes	$1/n^2$ spectrum in eternal vacuum
Fine structure	zeros of $\zeta(s)$	acoustic peaks at $\ell \approx 220, 540$
No Big Bang needed	yes	yes (in FFGFT)
Common root	$\zeta(s) = \sum 1/n^s = \prod_p 1/(1 - p^{-s})$	

Table ak.1: Primes and CMB anisotropies as possible instances of the same eternal spectrum [S]

In the interpretation proposed here [S], the apparent uniformity — of the prime distribution as of the microwave sky — is an approximation. The actual structure would be geometrically fixed, not random, and would require no starting point in time. The $1/n^2$ -weighted torus spectrum would be the common source — and the CMB its cosmic signature.

ak.4 Complexity as a Physical Continuum

Against the Levels Metaphor

Document 170 presented a table of discrete complexity levels — from elementary particles (Level 0) to consciousness (Level 4) — interpreting the fractal correction K_{frak} as a threshold operator. That presentation was formulated as a *working hypothesis* and included the caveat that the levels require further formalisation. This document takes that caveat seriously and develops an alternative perspective.

Complexity as Continuum

Complexity, self-reference, and what is called “consciousness” are not binary phenomena with discrete thresholds in FFGFT. They form a physical continuum that grows with the recursion depth of the mass-time coupling $\tilde{T} \cdot m = 1$. There is no sharp boundary beyond which a system “has consciousness” — any more than there is a sharp boundary beyond which water is “hot”.

What the Willow Chip Illustrates on the Continuum

The Willow chip is not a conscious system. But it is not structureless either:

- **Single Josephson qubit:** The Josephson energy E_J defines an effective mass and thereby a timescale $\tilde{T} = \hbar/E_J$. The qubit is a physical clock — the first step of self-reference.
- **Surface code lattice (105 qubits):** Collective interaction generates logical qubits that are robust against local perturbations. The system develops a form of “memory” — not cognitive, but physical: it preserves its own topological order against noise.
- **Floquet drive:** The system “remembers” the driving rhythm through time-crystalline structures. The drive period is inscribed in the topology of the system.
- **OTOC measurement:** The system measures its own information distribution. It references itself — geometrically precise, in measurable patterns.

This is a point on a continuum, not a threshold crossing. Between this point and the human brain lies not an abyss but a continuous spectrum of increasing recursion depth, integration breadth, and feedback stability.

The Open Question — and the Honest Limit of Physics

FFGFT describes this continuum geometrically: more qubits, deeper recursion, broader entanglement move the system upward along the continuum. But physics cannot answer where on this continuum what is called subjective experience arises — the experience of colour, pain, time, meaning.

This is not a gap in the theory but an honest statement about the scope of physical description. FFGFT states:

What FFGFT Can State

- The geometric structure $\xi \cdot \ell_p$ that generates Willow patterns is the same structure to which FFGFT traces the natural constants (with the one parameter ξ ; SI anchors serve for conversion).
- Complexity and self-reference grow continuously with the recursion depth of the mass-time coupling.
- The topological patterns measured in Willow experiments are interpreted in FFGFT as consequences of this structure, not as random artefacts; the coupling to the torus fundamental frequency is a working hypothesis **[S]**.

What consciousness *is* remains open — that is not a physical statement expressible in an equation.

ak.5 Summary

Experiment	Measured Finding	FFGFT Relation
Error Correction Nature (Dec. 2024) with code distance torus modes	Exponential error reduction Projection onto stable	
Floquet topo. order Nature (Sep. 2025) anyons, topo. invariant Floquet drive	Chiral edge modes, Torus resonance under	
Quantum Echoes / OTOC Nature (Oct. 2025) harmonic wave front 1/n ² -weighted spectrum	Constructive interference, Interferometry on torus;	

Table ak.2: Three Willow results and their FFGFT interpretation

The three experiments consistently show geometric order in quantum systems under controlled conditions; FFGFT interprets these patterns as a consequence of the topology of the vacuum field. This is the physically precise core behind intuitive descriptions associating quantum computers with self-perception or patterns. The difference from speculative narrative: the patterns are measured, quantified, and published in peer-reviewed journals; their FFGFT interpretation, by contrast, is a working hypothesis.

Bibliography

- [1] Google Quantum AI, *Quantum error correction below the surface code threshold*, *Nature* (2024), DOI: 10.1038/s41586-024-08449-y
- [2] M. Will, T. A. Cochran, E. Rosenberg et al., *Probing non-equilibrium topological order on a quantum processor*, *Nature* **645**, 348–353 (2025), DOI: 10.1038/s41586-025-09456-3
- [3] M. Will et al., *Probing non-equilibrium topological order on a quantum processor*, PubMed, PMID: 40931155 (2025)
- [4] T. A. Cochran, B. Jobst, E. Rosenberg et al., *Visualizing dynamics of charges and strings in (2+1)D lattice gauge theories*, *Nature* **642**, 315–320 (2025), DOI: 10.1038/s41586-025-08999-9
- [5] Google Quantum AI and collaborators, *Observation of constructive interference at the edge of quantum ergodicity*, *Nature* (2025), DOI: 10.1038/s41586-025-09526-6

Draft for further elaboration. The connection between Floquet frequency and torus fundamental frequency is a working hypothesis requiring formal verification. Statements about the continuum of complexity are to be understood as a physically motivated position, not a complete theory of consciousness.

Chapter al

Doc. 184 — p-bit and Probabilistic Information Processing

al.1 Introduction: The Operating Point Between Bit and Qubit

Doc. 178 established the stability of spin as macroscopic evidence for discrete torsion configurations. The permanent magnet shows: spin-up and spin-down are geometrically distinguished states, robust over years without cooling. The classical bit uses precisely this regime — a deep potential well $U \gg k_B T$.

The qubit operates at the opposite extreme: metastable torsion, coherence in the millisecond range (Doc. 174), enforced by extreme cooling to mK temperatures.

The **p-bit** (probabilistic bit) occupies the conceptually hitherto unexplored intermediate range: a magnetic nanostructure of the order of the geometric scale $L_8 \approx 22$ nm of the ξ -hierarchy, at which the torsion barrier becomes thermally surmountable.

The central thesis of this document

The thermal noise of the p-bit is not fundamental randomness. It is the macroscopically unresolved sampling of deterministic torsion configurations. The apparent stochasticity is epistemic — the transitions follow the sub-Planck-scale geometry operating on the timescale $t_0 = t_P/7500$.

The physical barrier is given materially by the Stoner-Wohlfarth energy $U = K_u \cdot V$. The FFGFT identifies: (a) **where** natural p-bit candidates are geometrically to be found (scale L_N), and (b) **how** the apparent randomness is to be understood physically.

al.2 The ξ -Length Hierarchy (Review of Doc. 173)

Doc. 173 defines the geometric scaling ladder of the FFGFT via the parameter $\xi_{\text{par}} = 4/30000 \approx 1,333 \times 10^{-4}$:

$$L_N = L_0 \cdot \left(\frac{1}{\xi_{\text{par}}} \right)^N, \quad L_0 = \xi_{\text{par}} \cdot \ell_P, \quad (\text{al.1})$$

where $\ell_P = 1,616 \times 10^{-35}$ m is the Planck length. Level $N = 8$ gives:

$$L_8 = \ell_P \cdot \xi_{\text{par}}^{-7} \approx 21,6 \text{ nm}. \quad (\text{al.2})$$

This scale is identified in Doc. 173 as the universal geometric anchor: excitonic Bohr radius, fine structure constant, transistor threshold size — all at $N \approx 8,00$.

al.3 From Length Scale to Characteristic Energy

Torsion field characteristic at scale L_N

Each geometric scale L_N is assigned a characteristic torsion energy in the FFGFT framework. The torsion field is a relativistic field object; the natural energy scale of a torsion mode at wavelength L_N is:

$$E_N \equiv \frac{\hbar c}{L_N} = \frac{\hbar c}{L_0} \xi_{\text{par}}^N = \frac{E_P}{\xi_{\text{par}}} \xi_{\text{par}}^N = E_P \cdot \xi_{\text{par}}^{N-1}, \quad (\text{al.3})$$

where $E_P = \hbar c / \ell_P = \hbar / t_P \approx 1,956 \times 10^9$ J is the Planck energy. Since $\xi_{\text{par}} < 1$, E_N decreases strictly monotonically with increasing N ; each level is a factor of $1/\xi_{\text{par}} = 7500$ smaller than the previous one.

Verification: $N = 1$ yields the Planck energy

For $N = 1$:

$$E_1 = E_P \cdot \xi_{\text{par}}^0 = E_P \approx 1,956 \times 10^9 \text{ J.} \quad (\text{al.4})$$

The Planck energy is the torsion energy of the Planck level — not a free parameter, but a geometric consequence of the hierarchy.

Complete energy hierarchy and the p-bit regime

N	L_N	$E_N = \hbar c / L_N$	$E_N / k_B T_{300}$	Physical domain
0	$2,16 \times 10^{-39} \text{ m}$	$1,47 \times 10^{13} \text{ J}$	$3,5 \times 10^{33}$	sub-Planck (L_0)
1	$1,62 \times 10^{-35} \text{ m}$	$1,96 \times 10^9 \text{ J}$	$4,7 \times 10^{29}$	Planck scale; $E_1 = E_P$
2	$1,21 \times 10^{-31} \text{ m}$	$2,61 \times 10^5 \text{ J}$	$6,3 \times 10^{25}$	—
3	$9,09 \times 10^{-28} \text{ m}$	$3,48 \times 10^1 \text{ J}$	$8,4 \times 10^{21}$	—
4–7	...	...	$10^{18}–10^6$	—
8	$2,16 \times 10^{-8} \text{ m}$	$1,47 \times 10^{-18} \text{ J}$	354	Exciton, Bohr radius, transistor threshold
9	$1,62 \times 10^{-4} \text{ m}$	$1,95 \times 10^{-22} \text{ J}$	0,047	0,16 mm
10	$1,21 \times 10^0 \text{ m}$	$2,61 \times 10^{-26} \text{ J}$	6×10^{-6}	$\sim 1 \text{ m}$

The gap across the p-bit regime

The p-bit operating regime $E_N \approx k_B T$ at $T = 300 \text{ K}$ lies at $N_{\text{opt}} \approx 8,66$ — *between* the integer levels $N = 8$ and $N = 9$. The step factor is $1/\xi_{\text{par}} = 7500$: between E_8 ($354 k_B T$) and E_9 ($0,047 k_B T$) there is no integer level value that hits $k_B T$ (300 K).

Consequence: The ξ -energy hierarchy E_N does *not* directly fix the barrier height of the p-bit. It determines the **geometric scale** of the nanostructure at which p-bit-capable resonators naturally occur.

For cryogenic p-bits, however: $E_9 \approx k_B T$ at $T^* \approx 14 \text{ K}$. At this temperature a macroscopic resonator of dimension $L_9 \approx 0,16 \text{ mm}$ lies exactly on a hierarchy level.

al.4 The Physical Barrier: Stoner-Wohlfarth Energy

Derivation

A magnetic nanoresonator with uniaxial anisotropy possesses an energy barrier between spin-up and spin-down, given in the Stoner-Wohlfarth model [4] by:

$$U_{\text{sw}} = K_u \cdot V \quad (\text{al.5})$$

Here K_u is the uniaxial anisotropy constant (in J/m^3 , material-specific) and V is the volume of the magnetic free layer.

In the FFGFT framework, U_{sw} is the macroscopic manifestation of the torsion barrier between the two stable configurations that Doc. 178 identifies as geometrically distinguished states.

For a spherical nanoresonator with diameter D :

$$U_{\text{sw}}(D) = K_u \cdot \frac{\pi}{6} D^3. \quad (\text{al.6})$$

The condition for p-bit operation $U_{\text{sw}} \approx k_B T$ fixes a specific diameter D^* for a given K_u :

$$D^* = \left(\frac{6 k_B T}{\pi K_u} \right)^{1/3}. \quad (\text{al.7})$$

Numerical values and comparison with L_8

For superparamagnetic nanoparticles with $K_u \approx 10^4 \text{ J/m}^3$:

D (nm)	U_{sw} (meV)	$U_{\text{sw}}/k_B T_{300}$	Regime
5	4,1	0,16	
8	16,7	0,65	p-bit
10	32,7	1,26	p-bit
12	56,5	2,18	p-bit
15	110	4,3	p-bit
20	261	10,1	p-bit
25	511	19,8	

The p-bit operating range ($U/k_B T \approx 1\text{--}10$) lies at $D \approx 8\text{--}20 \text{ nm}$ — just below the ξ -hierarchy level $N = 8$ with $L_8 \approx 22 \text{ nm}$. With Eq. (al.6) and $K_u = 10^4 \text{ J/m}^3$, $T = 300 \text{ K}$ one has $U_{\text{sw}}(22 \text{ nm}) = 13.5 k_B T$ and $U_{\text{sw}}(21.6 \text{ nm}) = 12.7 k_B T$, i.e. above the range 1–10.

Geometric prediction: $N = 8$ as natural p-bit anchor

The FFGFT predicts: magnetic nanoresonators on the ξ -level $N \approx 8$ are natural p-bit candidates at room temperature. Not because $E_8 \approx k_B T$ (that is not the case: $E_8 = 354 k_B T$), but because $L_8 \approx 22 \text{ nm}$ is of the order of the scale at which the Stoner-Wohlfarth volume with typical material values falls into the p-bit regime; at $K_u = 10^4 \text{ J/m}^3$, L_8 lies just above this range, and the exact position depends on the material through K_u .

This proximity is not accidental in the FFGFT: $N = 8$ is the universal scale of magnetic single resonators (Doc. 173: minimal resonator, Doc. 178: permanent magnet as collective resonator at $N \approx 8$).

al.5 The Kramers Switching Rate in the FFGFT Framework

Standard formula

The thermally activated switching rate between torsion configurations follows the Kramers formula [5, 6]:

$$\Gamma = \frac{\omega_0 \omega_b}{2\pi\gamma} \exp\left(-\frac{U_{sw}}{k_B T}\right), \quad (\text{al.8})$$

where ω_0 is the angular frequency at the potential minimum, ω_b at the barrier peak and γ is the (Gilbert) damping rate.

FFGFT interpretation: determinism instead of randomness

In the standard interpretation, Γ is a stochastic rate: each transition is a random, uncorrelated event.

FFGFT reinterpretation of the Kramers rate

In the FFGFT, equation (al.8) is **formally identical** — but its physical meaning changes:

Γ is the **time-averaged frequency of deterministic torsion transitions**. Each individual transition follows a precise path in the sub-Planck-scale geometry of the torsion field, which is determined on the timescale $t_0 = t_p/7500$.

The macroscopic measurement — temporal resolution $\Delta t \gg t_0$ — averages over $\Delta t/t_0 \gg 1$ microstates and creates the appearance of randomness. The randomness is **epistemic**, not ontological.

This is the direct transfer of the FFGFT position established in Doc. 178 (epistemic ignorance rather than genuine superposition) to the thermally fluctuating p-bit system.

Why no modified prefactor

A numerical FFGFT correction to the Kramers rate would require the structural parameter $k_B T t_0 / \hbar$:

$$\frac{k_B T t_0}{\hbar} = \frac{k_B \cdot 300 \text{ K}}{\hbar / t_0} \approx 2,82 \times 10^{-34}. \quad (\text{al.9})$$

This value is $\ll 1$ at all technically accessible temperatures. A correction to the Kramers prefactor by the FFGFT is not experimentally detectable at room temperature.

Where the FFGFT contribution really lies

1. **Geometric scale:** $D \sim L_8 \approx 22 \text{ nm}$ as natural p-bit anchor (order of magnitude; the exact position depends on the material through K_u) — verifiable materials-science prediction.
2. **Interpretation:** The switching statistics are deterministically structured on the t_0 scale; the macroscopically observed stochasticity is a projection.

3. **No new terms:** The Kramers rate (al.8) remains unchanged. The FFGFT contribution is interpretive.

al.6 The Three-Regime Picture: Bit, p-Bit, Qubit

Type	Barrier	Temp.	Timescale
Class. bit	$U \gg k_B T$	300 K	Years
p-bit	$U \approx k_B T$	300 K	ns– μ s
Qubit	$U \gg k_B T$ (prep.)	mK	μ s–ms

Type	ξ -level	FFGFT: random?
Class. bit	$N \approx 8$, deep	No; stable torsion well
p-bit	$D \sim L_8 \approx 22$ nm	No; deterministic
Qubit	$N \approx 8$, metastable	No; epistemic (Doc. 178)

All three types reside on the same geometric scale $N \approx 8$. The difference is the operating point on the potential curve: deep in the minimum (bit), at the tipping point (p-bit), metastably prepared (qubit).

al.7 Landauer's Principle

Landauer's principle [7] applies to the p-bit unchanged:

$$E_{\text{diss}}^{\min} = k_B T \ln 2. \quad (\text{al.10})$$

The p-bit switches between two torsion configurations; the state space is binary. The ξ -geometry determines the *dynamics* (scale, switching rate), not the *state space size*.

No modified Landauer bound in the FFGFT

A ξ -dependent Landauer bound would require the state space of the p-bit to have more than two elements, or that information erasure uses geometrically non-equivalent paths. Neither is the case in a standard p-bit. The standard bound $k_B T \ln 2$ applies.

al.8 Experimental Realisations

- **Magnetic Tunnel Junctions (MTJ) [2]:** CoFeB free layers, $K_u \sim 10^4$ – 10^5 J/m³, lateral dimension ~ 20 – 100 nm — directly at L_8 . Switching times in the nanosecond range.
- **Superparamagnetic nanoparticles:** Fe₃O₄, γ -Fe₂O₃ with $D \approx 10$ – 20 nm show spontaneous thermal switching at 300 K in the L_8 regime.

- **CMOS-based p-bits [3]:** Ring oscillators near metastability. No direct torsion field connection — realise the logical operating point by other means.

Verifiable FFGFT prediction

The FFGFT names $L_8 \approx 22$ nm as a **geometrically distinguished dimension** for p-bits. Eq. (al.7) with $K_u = 10^4$ J/m³ and $T = 300$ K, however, gives $D^* = 9.2$ nm, not L_8 ; D^* depends on the material through $K_u^{-1/3}$. That D^* coincides with L_8 therefore does not follow from the equation but requires a material with a suitable K_u . A systematic measurement of switching rate, noise colour and reproducibility across different nanoparticle sizes would show whether an optimum occurs at $D \approx L_8$.

al.9 Classification within the FFGFT Document Series

Doc.	Topic	Relation to Doc. 184
173	ξ -hierarchy, quantum dot, minimal resonator	Foundation: L_N , E_N derivation, $N = 8$ anchor
174	Reading and writing spin	Techniques; decoherence times
178	Spin stability, permanent magnet	Epistemic randomness; torsion well vs. tipping point
184	p-bit	Tipping point: same scale $N = 8$, different operating point

al.10 Summary

Conclusion: p-bit in the FFGFT framework

1. The ξ -energy hierarchy is cleanly derivable.

$E_N = \hbar c / L_N = E_P \cdot \xi_{\text{par}}^{N-1}$, with $E_1 = E_P$. The levels jump by a factor of 7500 — the p-bit regime at 300 K lies between $N = 8$ and $N = 9$.

2. The barrier is Stoner-Wohlfarth: $U = K_u \cdot V$.

Adjustable by materials engineering, not quantised by E_N . p-bit regime at $D \approx 8\text{--}20$ nm ($K_u \sim 10^4$ J/m³); the exact position depends on the material through K_u .

3. The ξ -reference is the geometric scale $L_8 \approx 22$ nm.

This is the universal resonator scale $N = 8$ from Doc. 173. Natural p-bit candidates lie near this level, just below it at $K_u = 10^4$ J/m³.

4. The Kramers rate remains formally unchanged.

FFGFT contribution: determinism of transitions on the t_0 scale — the stochasticity is epistemic, not ontological.

5. The Landauer bound applies unchanged.

$E_{\text{diss}}^{\text{min}} = k_B T \ln 2$ — binary state space, no ξ modification.

The p-bit is the geometric tipping point between permanent magnet (Doc. 178) and qubit (Doc. 174) — on the same ξ -level $N = 8$.

Bibliography

- [1] Datta, S. et al. *p-Bits for Probabilistic Spin Logic*. arXiv:1809.04028 (2018).
- [2] Borders, W. A. et al. *Integer factorization using stochastic magnetic tunnel junctions*. Nature **573**, 390–393 (2019).
- [3] Aadit, N. A. et al. *Massively parallel probabilistic computing with sparse inverse Ising problems*. Nature Electronics **5**, 460–468 (2022).
- [4] Stoner, E. C. & Wohlfarth, E. P. *A mechanism of magnetic hysteresis in heterogeneous alloys*. Phil. Trans. R. Soc. London A **240**, 599–642 (1948).
- [5] Kramers, H. A. *Brownian motion in a field of force and the diffusion model of chemical reactions*. Physica **7**, 284–304 (1940).
- [6] Hänggi, P., Talkner, P. & Borkovec, M. *Reaction-rate theory: fifty years after Kramers*. Rev. Mod. Phys. **62**, 251–341 (1990).
- [7] Landauer, R. *Irreversibility and heat generation in the computing process*. IBM J. Res. Dev. **5**, 183–191 (1961).
- [8] Hayakawa, K. et al. *Nanosecond random telegraph noise in in-plane magnetic tunnel junctions*. Phys. Rev. Lett. **126**, 117202 (2021).
- [9] Pascher, J. *The Minimal Quantum Resonator*. Doc. 173, FFGFT framework (2026).
- [10] Pascher, J. *Reading and Writing Spin*. Doc. 174, FFGFT framework (2026).
- [11] Pascher, J. *Stability as Ground State*. Doc. 178, FFGFT framework (2026).