

The T0 Energy Field Model: Mathematical Formulation

Contents

.1	Units Convention	3
	T0 natural units (Heaviside-Lorentz)	3
	Dimensions in natural units	3
.2	Time-Energy Duality	3
	Fundamental relation	3
	Intrinsic time field	4
.3	Universal Field Equation	4
	Wave equation	4
	With sources	4
.4	Lagrangian Density	4
	Universal Lagrangian density	4
	Euler-Lagrange equation	4
.5	Characteristic Energy	4
	Definition	4
	Numerical values	5
	Usage	5
.6	The Parameter ξ	5
	Definition	5
	Geometric components	5
.7	Fine Structure Constant	5
	In T0 units	5
	Reconstruction of SI value	6
	Numerical calculation	6
	Dimension check	6
.8	Gravitational Constant	6
	T0 formula	6
	Fundamental relation	6
.9	Characteristic Lengths	7
	T0 characteristic length	7
	Derivation	7
	Time scale	7
.10	Scale Hierarchy	7
	Planck length as reference	7
	Scale ratio	7
.11	Particles as Field Excitations	8
	Classification by energy	8
	Antiparticles	8
.12	Lepton Masses	8
.13	Anomalous Magnetic Moments	8
.14	Three Field Geometries	8

	Type 1: Localized spherical	8
	Type 2: Localized non-spherical	8
	Type 3: Extended homogeneous	9
.15	Mathematical Identities	9
	Energy field normalization	9
	Duality consistency	9
.16	Dimensional Analysis Verifications	9
	Field equation	9
	Characteristic length	9
	Lagrangian density	9
.17	Formula Reference	10
	Fundamental equations	10
	Derived constants	10
	Characteristic scales	10
	Prediction formulas	10
.18	Numerical Values	10
	Fundamental constants (in natural units)	10
	T0 parameters	11
	Lepton energies	11
	Energy ratios	11
.19	Calculation Examples	11
	Fine structure constant	11
	Characteristic length (electron)	12
0.1	Symbol Reference	12
0.2	Unit Conversions	12
	Natural $\rightarrow$ SI	12
	Standard natural units	12
0.3	Relations to Other Documents	13

Correction notes

Corrections from the calculation review (30 Sept. 2026). The following places are corrected or annotated in the text; each carries a dated note there with the previous value:

- E_0 : The two values are the same E_0 , bare ($\sqrt{m_e m_\mu} = 7.348$ MeV) and fractally corrected ($E_0/\sqrt{K_{\text{frak}}} = 7.398$ MeV, Doc. 190, R72). The corrected value serves as the anchor for α_i from the masses alone, $\alpha = 1/137.059$ (+0.017 %).
- α : agreement 0.0005 % instead of 0.03 %.
- Anomalous magnetic moments: the short formula $a_\ell = \frac{\xi}{2\pi}(E_\ell/E_e)^2$ has been removed; recalculated it gave $a_\mu = 0.907$ instead of 1.166×10^{-3} . Authoritative are Docs. 018 and 158.
- Dimension of the Lagrangian density: $[E^4]$, not $[E^2]$.
- G_{SI} formula: $\frac{\xi^2}{4m_e} \times C_{\text{dim}} \times C_{\text{conv}} \rightarrow \frac{\xi^2}{4m_e} \times C_{\text{conv}} \times K_{\text{frak}}$ – with C_{dim} but without the compensating factor E_{char} , G would be too small by a factor of $E_{\text{char}} = 28.8$. Likewise: $\xi^2/(4m_e)$ has $[E^{-1}]$, not $[E^{-2}]$.

Abstract

The T0 model describes physical phenomena through a universal energy field $E_{\text{field}}(x, t)$ with the parameter $\xi = \frac{4}{3} \times 10^{-4}$. The field equation is $\square E_{\text{field}} = 0$, the Lagrangian density $\mathcal{L} = \xi(\partial E)^2$. The model uses standard natural units with $\hbar = c = 1$.

Fundamental quantities:

- Characteristic energy: $E_0 = \sqrt{m_e \cdot m_\mu} = 7.348 \text{ MeV (bare)}$; fractally corrected $E_0/\sqrt{K_{\text{frak}}} = 7.398 \text{ MeV}$ (anchor for α)
- Fine structure constant: $\alpha = \xi(E_0/1 \text{ MeV})^2 \approx 1/137$
- Gravitational constant: $G = \xi^2/(4m_e) \times \text{factors}$

Predictions: Lepton masses with 2% accuracy, fine structure constant $\alpha \approx 1/137$ (from the masses alone $1/137.059, +0.017 \%$).

Detailed derivations: See Document 011 (fine structure), 012 (gravitation), 018 (g-2 geometric), 019 (Lagrangian).

.1 Units Convention

T0 natural units (Heaviside-Lorentz)

T0 uses Heaviside-Lorentz natural units:

$$\hbar = c = 4\pi\epsilon_0 = 1 \quad (1)$$

In this system:

- Fine structure constant: $\alpha = e^2 = 1$ (by convention in T0)
- Energy = Mass: $E = m$
- Length = Time = Energy⁻¹: $[L] = [T] = [E^{-1}]$

Note: In SI units, $\alpha \approx 1/137$. The choice $\alpha = 1$ in T0 is a unit convention that simplifies formulas.

Dimensions in natural units

$$[E] = E \quad (2)$$

$$[m] = E \quad (3)$$

$$[t] = E^{-1} \quad (4)$$

$$[L] = E^{-1} \quad (5)$$

$$[G] = E^{-2} \quad (6)$$

$$[\partial_\mu] = E \quad (7)$$

.2 Time-Energy Duality

Fundamental relation

$$T_{\text{field}}(x, t) \cdot E_{\text{field}}(x, t) = 1 \quad (8)$$

with $[T_{\text{field}}] = E^{-1}$ and $[E_{\text{field}}] = E$.

Intrinsic time field

$$T_{\text{field}}(x, t) = \frac{1}{E_{\text{field}}(x, t)} \quad (9)$$

.3 Universal Field Equation

Wave equation

$$\square E_{\text{field}} = 0 \quad (10)$$

with d'Alembert operator:

$$\square = \nabla^2 - \frac{\partial^2}{\partial t^2} \quad (11)$$

With sources

$$\nabla^2 E_{\text{field}} = 4\pi G\rho \cdot E_{\text{field}} \quad (12)$$

Dimension check: $[E^3] = [E^{-2}][E^4][E] = [E^3] \checkmark$

.4 Lagrangian Density

Universal Lagrangian density

$$\mathcal{L} = \xi \cdot (\partial_\mu E_{\text{field}})(\partial^\mu E_{\text{field}}) \quad (13)$$

with $\xi = \frac{4}{3} \times 10^{-4}$.

Euler-Lagrange equation

$$\frac{\partial \mathcal{L}}{\partial E} - \partial_\mu \frac{\partial \mathcal{L}}{\partial (\partial_\mu E)} = 0 \quad (14)$$

yields:

$$\square E_{\text{field}} = 0 \quad (15)$$

.5 Characteristic Energy

Definition

The characteristic energy E_0 is the geometric mean of electron and muon mass (derivation in Document 011):

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (16)$$

Numerical values

From experimental masses:

$$E_0 = \sqrt{0.511 \times 105.66} \quad (17)$$

$$= \sqrt{53.99} \quad (18)$$

$$= 7.348 \text{ MeV} \quad (19)$$

Fractally corrected value (Doc. 190, R72):

$$E_0^{T0} = \frac{E_0}{\sqrt{K_{\text{frak}}}} = \frac{7.348}{\sqrt{1 - 100\xi}} = 7.397 \approx 7.398 \text{ MeV} \quad (20)$$

Both values are the same E_0 , bare and fractally corrected: absolute values carry K_{frak} , ratios do not. The square-root factor follows because E_0 enters $\alpha = \xi E_0^2$ quadratically. The corrected value 7.398 MeV serves as the anchor for α .

(Corrected on 30 Sept. 2026: previously "Theoretical T0 value: $E_0^{T0} = 7.398 \text{ MeV}$; Deviation: 0.7 % (within geometric corrections)". The relation $7.398/7.348 = 1.00682 \approx 1/\sqrt{K_{\text{frak}}} = 1.00673$ (0.008 %) was not stated.)

Usage

E_0 serves as energy scale for:

- Fine structure constant: $\alpha = \xi(E_0/1 \text{ MeV})^2$
- Normalization of electromagnetic effects

.6 The Parameter ξ

Definition

$$\xi = \frac{4}{3} \times 10^{-4} = 1.3333 \times 10^{-4} \quad (21)$$

Dimensionless: $[\xi] = 1$.

Geometric components

$$\xi = G_3 \times S_{\text{ratio}} \quad (22)$$

where:

- $G_3 = \frac{4}{3}$: Geometric factor (sphere-cube ratio)
- $S_{\text{ratio}} = 10^{-4}$: Scale ratio

.7 Fine Structure Constant

In T0 units

In T0 natural units: $\alpha = 1$ (by convention)

Reconstruction of SI value

Important: The SI value can be reconstructed from T0 parameters!

$$\alpha_{\text{SI}} = \xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2 \quad (23)$$

Numerical calculation

With $\xi = \frac{4}{3} \times 10^{-4}$ and $E_0 = 7.398 \text{ MeV}$:

$$\alpha_{\text{SI}} = 1.3333 \times 10^{-4} \times (7.398)^2 \quad (24)$$

$$= 1.3333 \times 10^{-4} \times 54.73 \quad (25)$$

$$= 7.297 \times 10^{-3} \quad (26)$$

$$= \frac{1}{137.04} \quad (27)$$

Experimental: $\alpha_{\text{exp}} = \frac{1}{137.036}$

Agreement: 0.0005 % (with the anchor $E_0 = 7.398 \text{ MeV}$). From the masses alone, $\alpha = \xi E_0^2 / K_{\text{frak}}$ with $E_0 = 7.348 \text{ MeV}$ gives $\alpha = 1/137.059$ (+0.017 %).

(Corrected on 30 Sept. 2026: previously "Agreement: 0.03%"; $1/137.035$ against $1/137.036$ is 0.0005 %.)

Dimension check

$$[\alpha_{\text{SI}}] = [\xi] \times \left[\frac{E}{E} \right]^2 = 1 \times 1 = 1 \quad \checkmark \quad (28)$$

.8 Gravitational Constant

T0 formula

The gravitational constant is derived from ξ and m_e (derivation in Document 012):

$$G = \frac{\xi^2}{4m_e} \times C_{\text{conv}} \times K_{\text{frak}} \quad (29)$$

(Corrected on 30 Sept. 2026: previously $\frac{\xi^2}{4m_e} \times C_{\text{dim}} \times C_{\text{conv}} - C_{\text{dim}} = 1/E_{\text{char}} = 3.472 \times 10^{-2}$ is cancelled again by the factor $E_{\text{char}} = 28.8$ in the step chain of Doc. 012; in the formula alone the product would give 2.35×10^{-12} instead of 6.674×10^{-11} .)

where:

- C_{dim} : Dimension correction
- C_{conv} : SI conversion factor

Fundamental relation

In natural units:

$$\xi = 2\sqrt{G \cdot m_e} \quad (30)$$

Solved for G :

$$G_{\text{nat}} = \frac{\xi^2}{4m_e} \quad (31)$$

Dimension: $[G] = [E^{-2}]$ in natural units; the right-hand side $\xi^2/(4m_e)$, however, has $[E^{-1}]$ – the missing factor $1/E_{\text{char}}$ and the SI conversion reside in C_{conv} (Doc. 012). *(Corrected on 30 Sept. 2026: previously without this remark, so that $\xi^2/(4m_e)$ appeared as $[E^{-2}]$.)*

.9 Characteristic Lengths

T0 characteristic length

$$r_0 = 2GE \quad (32)$$

Dimension: $[r_0] = [E^{-2}][E] = [E^{-1}] = [L] \checkmark$

Derivation

For spherically symmetric point source $\rho(r) = E_0 \delta^3(\vec{r})$:

Solution of $\nabla^2 E = 4\pi G \rho E$:

$$E(r) = E_0 \left(1 - \frac{r_0}{r}\right) \quad (33)$$

with $r_0 = 2GE_0$.

Time scale

$$t_0 = \frac{r_0}{c} = r_0 = 2GE \quad (34)$$

(since $c = 1$)

.10 Scale Hierarchy

Planck length as reference

$$\ell_P = \sqrt{G} = 1 \quad (\text{in nat. units}) \quad (35)$$

Scale ratio

$$\xi_{\text{ratio}} = \frac{\ell_P}{r_0} = \frac{\sqrt{G}}{2GE} = \frac{1}{2\sqrt{G} \cdot E} \quad (36)$$

For $E \sim 1 \text{ GeV}$:

$$\frac{r_0}{\ell_P} = 2\sqrt{G}E = \frac{2E}{E_P} \sim 1.6 \times 10^{-19} \quad (\text{sub-Planck}) \quad (37)$$

(Corrected on 29 Sept. 2026: previously $r_0/\ell_P \sim 10^7$ – with $r_0 = 2GE$ and $\ell_P = \sqrt{G}$ one has $r_0/\ell_P = 2E/E_P$, i.e. 1.6×10^{-19} for $E = 1 \text{ GeV}$.)

Particle	Energy [MeV]
Electron	0.511
Muon	105.658
Tau	1776.86

.11 Particles as Field Excitations

Classification by energy

Antiparticles

Negative field excitations: $E_{\text{field}} < 0$

.12 Lepton Masses

The T0 model predicts lepton masses (derivation in Document 003):

Lepton	T0 [MeV]	Exp [MeV]	Δ [%]
Electron	0.507	0.511	0.87
Muon	103.5	105.7	2.09
Tau	1815	1777	2.16

.13 Anomalous Magnetic Moments

Anomalous magnetic moments are treated in dedicated documents: Doc. 018 (with its revisions) and Doc. 158.

(Corrected on 30 Sept. 2026: previously this section gave the short formula $a_\ell = \frac{\xi}{2\pi}(E_\ell/E_e)^2$ with $a_\mu = 1.166 \times 10^{-3}$, $a_e = 2.122 \times 10^{-5}$ and $a_\tau = 1.28 \times 10^{-3}$. Recalculated, the formula gives $a_\mu = 2.122 \times 10^{-5} \times 42\,753 = 0.907$ and $a_\tau = 256.6$; a_e lies a factor of 55 below the measured value 1.15965×10^{-3} . The formula has been removed, as have its repetitions in the dimensional analysis, formula reference and calculation examples.)

.14 Three Field Geometries

Type 1: Localized spherical

$$E(r) = E_0 \left(1 - \frac{\beta}{r}\right), \quad \beta = r_0 \quad (38)$$

Application: Individual particles (electron, muon, tau)

Type 2: Localized non-spherical

$$E(\vec{r}) = E_0 \left(1 - \frac{\beta_{ij} r_i r_j}{r^3}\right) \quad (39)$$

Application: Bound systems

Type 3: Extended homogeneous

Effective parameter:

$$\xi_{\text{eff}} = \frac{\xi}{2} = \frac{2}{3} \times 10^{-4} \quad (40)$$

Application: Cosmology (see Document 026)

.15 Mathematical Identities

Energy field normalization

$$E_{\text{field}}(\vec{r}, t) = E_0 \cdot f(\vec{r}, t) \cdot e^{i\phi(\vec{r}, t)} \quad (41)$$

with:

- E_0 : Characteristic energy
- $f(\vec{r}, t)$: Normalized profile
- $\phi(\vec{r}, t)$: Phase

Duality consistency

Time-mass (Document 003): $T \cdot m = 1$

Time-energy (this document): $T \cdot E = 1$

In natural units ($c = 1$):

$$E = mc^2 = m \quad \Rightarrow \quad T \cdot m = T \cdot E \quad (42)$$

.16 Dimensional Analysis Verifications

Field equation

$$[\nabla^2 E] = [L^{-2}][E] = [E^2][E] = [E^3] \quad (43)$$

$$[4\pi G \rho E] = [E^{-2}][E^4][E] = [E^3] \quad \checkmark \quad (44)$$

Characteristic length

$$[r_0] = [2GE] = [E^{-2}][E] = [E^{-1}] = [L] \quad \checkmark \quad (45)$$

Lagrangian density

$$[\mathcal{L}] = [\xi][(\partial E)^2] = [1][E^4] = [E^4] \quad (\text{correct for Lagrangian density}) \quad (46)$$

(Corrected on 30 Sept. 2026: previously $[1][E^2] = [E^2]$; $[\partial_\mu E] = E^2$, hence $[(\partial E)^2] = E^4$, as in the symbol table.)

.17 Formula Reference

Fundamental equations

$$\text{Duality: } T_{\text{field}} \cdot E_{\text{field}} = 1 \quad (47)$$

$$\text{Wave equation: } \square E_{\text{field}} = 0 \quad (48)$$

$$\text{With sources: } \nabla^2 E = 4\pi G \rho E \quad (49)$$

$$\text{Lagrangian density: } \mathcal{L} = \xi(\partial E)^2 \quad (50)$$

Derived constants

$$\text{Characteristic energy: } E_0 = \sqrt{m_e \cdot m_\mu} = 7.348 \text{ MeV (bare; corrected 7.398)} \quad (51)$$

$$\text{Fine structure constant: } \alpha = \xi(E_0/1 \text{ MeV})^2 \approx 1/137 \quad (52)$$

$$\text{Gravitational constant: } G = \frac{\xi^2}{4m_e} \times \text{factors} \quad (53)$$

Characteristic scales

$$\text{T0 length: } r_0 = 2GE \quad (54)$$

$$\text{T0 time: } t_0 = 2GE \quad (55)$$

$$\text{Planck length: } \ell_P = \sqrt{G} = 1 \quad (56)$$

$$\text{Scale ratio: } \xi_{\text{ratio}} = \frac{1}{2\sqrt{GE}} \quad (57)$$

Prediction formulas

$$\text{Parameter: } \xi = \frac{4}{3} \times 10^{-4} \quad (58)$$

$$\text{Effective parameter: } \xi_{\text{eff}} = \frac{\xi}{2} \quad (59)$$

.18 Numerical Values

Fundamental constants (in natural units)

$$\hbar = 1 \quad (60)$$

$$c = 1 \quad (61)$$

$$\alpha = \frac{1}{137.036} \approx 7.297 \times 10^{-3} \quad (62)$$

$$G = 1 \text{ (numerically, dimension } [E^{-2}]) \quad (63)$$

T0 parameters

$$\xi = \frac{4}{3} \times 10^{-4} = 1.3333 \times 10^{-4} \quad (64)$$

$$\xi^2 = 1.7778 \times 10^{-8} \quad (65)$$

$$\frac{\xi}{2\pi} = 2.1221 \times 10^{-5} \quad (66)$$

$$\xi_{\text{eff}} = 6.6667 \times 10^{-5} \quad (67)$$

$$E_0 = 7.348 \text{ MeV (from exp. masses)} \quad (68)$$

$$E_0^{\text{T0}} = 7.398 \text{ MeV (fractally corrected, } E_0/\sqrt{K_{\text{frak}}}) \quad (69)$$

Lepton energies

$$E_e = 0.511 \text{ MeV} \quad (70)$$

$$E_\mu = 105.658 \text{ MeV} \quad (71)$$

$$E_\tau = 1776.86 \text{ MeV} \quad (72)$$

Energy ratios

$$\frac{E_\mu}{E_e} = 206.768 \quad (73)$$

$$\frac{E_\tau}{E_e} = 3477.2 \quad (74)$$

$$\frac{E_\tau}{E_\mu} = 16.817 \quad (75)$$

.19 Calculation Examples

Fine structure constant

Given:

- $\xi = 1.3333 \times 10^{-4}$
- $E_0 = 7.398 \text{ MeV}$

Calculation:

$$\left(\frac{E_0}{1 \text{ MeV}} \right)^2 = (7.398)^2 = 54.73 \quad (76)$$

$$\alpha = 1.3333 \times 10^{-4} \times 54.73 \quad (77)$$

$$= 7.297 \times 10^{-3} \quad (78)$$

$$= \frac{1}{137.04} \quad (79)$$

Experimental: $\alpha_{\text{exp}} = \frac{1}{137.036}$

Deviation: 0.0005 % (corrected on 30 Sept. 2026: previously 0.03 %)

Characteristic length (electron)

Given:

- $E_e = 0.511 \text{ MeV} = 0.511 \times 1.6 \times 10^{-13} \text{ J} = 8.2 \times 10^{-14} \text{ J}$
- $G = 6.674 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$
- $c = 3 \times 10^8 \text{ m/s}$

Conversion to SI units:

$$r_0 = \frac{2GE}{c^4} \approx 1.35 \times 10^{-57} \text{ m} \quad (80)$$

Planck comparison:

$$\frac{r_0}{\ell_P} = \frac{1.35 \times 10^{-57}}{1.6 \times 10^{-35}} \approx 8.4 \times 10^{-23} \quad (81)$$

(Corrected on 29 Sept. 2026: previously $r_0 = 2GE \approx 10^{-28} \text{ m}$ and $r_0/\ell_P \approx 10^7$ – with the SI values given, the formula reads $r_0 = 2GE/c^4$ and yields $1.35 \times 10^{-57} \text{ m}$; r_0 thus lies far below the Planck length.)

0.1 Symbol Reference

Symbol	Meaning	Dimension
ξ	Fundamental parameter	1
E_0	Characteristic energy	E
E_{field}	Universal energy field	E
T_{field}	Intrinsic time field	E^{-1}
r_0	T0 characteristic length	$L = E^{-1}$
t_0	T0 characteristic time	$T = E^{-1}$
ℓ_P	Planck length	$L = E^{-1}$
G	Gravitational constant	E^{-2}
α	Fine structure constant	1
E_e, E_μ, E_τ	Lepton energies	E
m_e, m_μ, m_τ	Lepton masses (= E in nat. units)	E
$\mathcal{L}$	Lagrangian density	E^4
$\square$	d'Alembert operator	E^2
ξ_{eff}	Effective parameter ($\xi/2$)	1

0.2 Unit Conversions

Natural $\rightarrow$ SI

$$1 \text{ (Energy)} = 1 \text{ GeV} = 1.6 \times 10^{-10} \text{ J} \quad (82)$$

$$1 \text{ (Length)} = \frac{\hbar c}{1 \text{ GeV}} = 0.197 \text{ fm} \quad (83)$$

$$1 \text{ (Time)} = \frac{\hbar}{1 \text{ GeV}} = 6.58 \times 10^{-25} \text{ s} \quad (84)$$

Standard natural units

In standard convention ($\hbar = c = 1$):

- $\alpha = \frac{e^2}{4\pi\epsilon_0} \approx \frac{1}{137}$ (dimensionless)
- All quantities in powers of energy
- Physical predictions identical to other conventions

0.3 Relations to Other Documents

- **Document 003:** Time-mass duality, foundations, origin of ξ
- **Document 011:** Fine structure constant (geometric derivation)
- **Document 012:** Gravitational constant (systematic derivation)
- **Document 018:** Geometric g-2 formulation (fractal geometry)
- **Document 019:** Lagrangian formulation (quantum field theory)
- **Document 026:** Cosmology ($\xi_{\text{eff}} = \xi/2$)
All formulations are based on $\xi = \frac{4}{3} \times 10^{-4}$.