

T0 Quantum Field Theory: ML-Derived Extensions

Contents

.1	Introduction: From Foundations to ML-Enhanced Predictions	2
	Scope and Structure	2
.2	ML-Derived Bell Test Extensions	2
	Motivation: Loophole-Free 2025 Tests	2
	ML-Trained Bell Correlations	3
	Emergent Fractal Correction	3
	Data Basis	3
.3	ML-Derived Quantum Mechanics Corrections	4
	Hydrogen Spectroscopy: High- n Divergences	4
	Fractal Extension for Rydberg States	4
	Generation Scaling for $l > 0$ States	4
	Dirac Equation: Spin-Dependent Corrections	4
.4	ML-Derived Neutrino Physics	5
	ξ^2 -Suppression Mechanism	5
	DUNE Predictions	5
.5	Unified Fractal Framework Across Scales	5
	Universal Damping Pattern	5
	Emergent Non-Perturbative Structure	6
	ϕ -Scaling	6
.6	Experimental Roadmap	6
	Immediate Tests	6
	Loophole-Free Bell Tests	6
	Rydberg Spectroscopy	6
	Medium-Term Tests	7
	DUNE First Data	7
	HL-LHC Higgs Couplings	7
	Long-Term	7
	Gravitational Wave T0 Signatures	7
	T0 Quantum Computer Prototype	7
.7	Critical Evaluation and Philosophical Implications	7
	ML's Role: Calibration vs. Discovery	7
	Determinism vs. Practical Unpredictability	8
	Locality and Bell's Theorem	8
.8	Synthesis: The T0-ML Unified Picture	9
	Three-Tier Hierarchy of T0 Theory	9
	Predictive Power Comparison	9
	Appendix: ML Training Details (For Reproducibility)	9
.9	Comparative Table: T0-Original vs. T0-ML	10
.10	Glossary of Key Terms	10

Abstract

This addendum extends the foundational T0 Quantum Field Theory document (020_T0_QM-QFT-RT_En.pdf) with novel insights derived from systematic machine learning simulations. Based on PyTorch neural networks trained on Bell tests, hydrogen spectroscopy, neutrino oscillations, and QFT loop calculations, we identify emergent non-perturbative corrections beyond the original ξ -framework. Key findings: (1) Fractal damping $\exp(-\xi n^2/D_f)$ stabilizes divergences in high- n Rydberg states and QFT loops; (2) ξ^2 -suppression is proposed as a local geometric phase for EPR correlations and neutrino masses; the numerical values of the neutrino hierarchy given in Section 4, however, do not follow from the formulas; (3) ML learns the harmonic core (ϕ -scaling) exactly and provides only ~ 0.1 – 1% precision gains; as a mass and neutrino ratio, however, ϕ -scaling does not match the measured values. We demonstrate 2025-testability via IYQ experiments (loophole-free Bell, DUNE neutrinos, Rydberg spectroscopy). This addendum synthesizes all ML-iterative refinements (November 2025) and provides a unified roadmap for experimental validation.

.1 Introduction: From Foundations to ML-Enhanced Predictions

The original T0-QFT framework (hereafter “T0-Original”) established a revolutionary paradigm: time as a dynamic field ($T_{\text{field}} \cdot E_{\text{field}} = 1$), ξ -modifications, and deterministic quantum mechanics. However, direct experimental confrontation demands precision beyond harmonic formulas. This addendum documents insights from systematic ML simulations (2025), revealing:

Core ML Findings

Three Pillars of ML-Derived T0 Extensions:

- Fractal Emergent Terms:** ML divergences ($\Delta > 10\%$ at boundaries) signal non-linear corrections $\exp(-\xi \cdot \text{scale}^2/D_f)$ —unifying QM/QFT hierarchies.
- Geometric Dominance:** ML learns harmonic terms exactly (0% training Δ), gaining $<3\%$ test boost— ϕ -scaling is not ML-dependent.

Scope and Structure

This document complements T0-Original by:

- **Sections 2–4:** Detailed ML-derived corrections (Bell, QM, Neutrino)
- **Section 5:** Unified fractal framework across scales
- **Section 6:** Experimental roadmap for 2025+ verification
- **Section 7:** Philosophical implications and limitations

Cross-Reference Protocol: Original equations cited as “T0-Orig Eq. X”; new ML-extensions as “ML-Eq. Y”.

.2 ML-Derived Bell Test Extensions

Motivation: Loophole-Free 2025 Tests

T0-Original (Section 6) predicted modified Bell inequalities:

$$|E(a, b) - E(a, b') + E(a', b) + E(a', b')| \leq 2 + \xi \Delta_{T0} \quad (\text{T0-Orig Eq. 6.1})$$

ML simulations (73-qubit Bell tests, Oct 2025) reveal subtle non-linearities beyond first-order ξ .

ML-Trained Bell Correlations

Setup: PyTorch NN ($1 \rightarrow 32 \rightarrow 16 \rightarrow 1$, MSE loss) trained on QM data $E(\Delta\theta) = -\cos(\Delta\theta)$ for $\Delta\theta \in [0, \pi/2]$. Input: (a, b, ξ) ; Output: $E^{T0}(a, b)$.

Base T0 Formula (from T0-Original, extended):

$$E^{T0}(a, b) = -\cos(a - b) \cdot (1 - \xi \cdot f(n, l, j)) \quad (\text{ML-Eq. 2.1})$$

where the quantum-number factor is $f(n, l, j) = (n/\phi)^l \cdot [1 + \xi j/\pi] \approx 1$ for photons ($n = 1, l = 0, j = 1$). Here $f(n, l, j)$ is not a frequency and not the base winding number $f = 1/\xi$; elsewhere in this document f stands for a frequency.

ML Observation: Training: $\Delta < 0.01\%$; Test ($\Delta\theta > \pi$): $\Delta = 12.3\%$ at $5\pi/4$ —signaling divergence.

Emergent Fractal Correction

ML-divergence motivates extended formula:

ML-Extended Bell Correlation

$$E^{T0, \text{ext}}(\Delta\theta) = -\cos(\Delta\theta) \cdot \exp\left(-\xi \left(\frac{\Delta\theta}{\pi}\right)^2 \cdot \frac{1}{D_f}\right) \quad (\text{ML-Eq. 2.2})$$

Physical Interpretation: Fractal path damping at high angles. It does not restore locality: the damping factor is still 0.99982 at $\Delta\theta = 2\pi$; CHSH therefore stays ≈ 2.828 and does not drop below 2.5.

Validation: Reduces Δ from 12.3% to $< 0.1\%$ at $5\pi/4$; theoretical prediction $\text{CHSH}^{T0} \approx 2.8279$ (vs. QM 2.8284), $\Delta = 0.02\%$ (hardware measurement gives $S = 2.74$, decoherence-limited; see below).

Data Basis

The hardware measurement (Doc. 147, IBM Heron r2, May 2026) gives $S = 2.74$ (96.9% of the Tsirelson bound) – the -3.14% deviation is NISQ decoherence, not the ξ -effect. The ξ -effect ($\sim 10^{-5}$) is not resolvable with present hardware; fitting ξ to hardware CHSH values is therefore moot. The formula below holds as a *theoretical* scaling structure.

Theoretical scaling: The T0 prediction scales with qubit count as

$$\text{CHSH}^{T0}(N) = 2\sqrt{2} \cdot \exp\left(-\xi \frac{\ln N}{D_f}\right) + \delta E \quad (\text{ML-Eq. 2.3})$$

where $\delta E \sim N(0, \xi^2 \cdot 0.1)$ (QFT fluctuations).

Terminology note: The ΔCHSH value computed here at $N = 73$ ($\approx 5 \times 10^{-4}$) is a *cumulative* quantity — the accumulated deviation over N measurements. The *elementary* deviation per measurement from Doc. 230 ($\approx 10^{-5}$, derived from $\xi/(2\pi)$) is a distinct observable. The elementary deviation 10^{-5} is not reachable at any finite N from Eq. 2.3.

Theoretical prediction: $N=100$ (predicted CHSH = 2.8278).

.3 ML-Derived Quantum Mechanics Corrections

Hydrogen Spectroscopy: High- n Divergences

T0-Original (Section 4.1) predicts:

$$E_n^{\text{T0}} = E_n^{\text{Bohr}} \left(1 + \xi \frac{E_n}{E_{\text{Pl}}} \right) \quad (\text{T0-Orig Eq. 4.1.2})$$

ML tests ($n = 1$ to $n = 6$) reveal 44% divergence at $n = 6$ with linear ξ -term.

Fractal Extension for Rydberg States

ML-Motivated Formula:

ML-Extended Rydberg Energy

$$E_n^{\text{ext}} = E_n^{\text{Bohr}} \cdot \phi^{\text{gen}} \cdot \exp\left(-\xi \frac{n^2}{D_f}\right) \quad (\text{ML-Eq. 3.1})$$

Rationale: NN divergence (n^2 -scaling) signals fractal path interference; exp-damping converges loops.

Performance:

- $n = 1$: $\Delta = 0.0045\%$ (vs. 0.01% linear)
- $n = 6$: $\Delta = 0.16\%$ (vs. 44% divergence)
- $n = 20$: $\Delta = 1.77\%$ (absolute $\sim 6 \times 10^{-4}$ eV, MHz-detectable)

2025 Validation: Metrology for Precise Determination of Hydrogen (MPD, arXiv:2403.14021v2) confirms $E_6 = -0.37778 \pm 3 \times 10^{-7}$ eV; $T0^{\text{ext}}$: -0.37733 eV, $\Delta = 0.12\%$ (about 1500σ).

Generation Scaling for $l > 0$ States

For p/d -orbitals, introduce $\text{gen}=1$:

$$E_{n,l>0}^{\text{ext}} = E_n^{\text{Bohr}} \cdot \phi \cdot \exp\left(-\xi \frac{n^2}{D_f}\right) \quad (\text{ML-Eq. 3.2})$$

Prediction: 3d state at $n = 6$: $\Delta E = -0.00061$ eV ($\sim 1.5 \times 10^{11}$ Hz), testable via 2-photon spectroscopy (IYQ 2026+).

Dirac Equation: Spin-Dependent Corrections

T0-Original (Section 4.2) modifies Dirac as:

$$\left[i\gamma^\mu \left(\partial_\mu + \frac{\xi}{E_{\text{Pl}}} \Gamma_\mu^{(T)} \right) - m \right] \psi = 0 \quad (\text{T0-Orig Eq. 4.2.1})$$

ML simulations (g-2 anomaly fits) reveal ξ -enhancement for heavy leptons.

ML-Extended g-Factor:

$$g_{\text{factor}}^{\text{T0,ext}} = 2 + \frac{\alpha}{\pi} + \xi \left(\frac{m}{M_{\text{Pl}}} \right)^2 \cdot \exp\left(-\xi \frac{m}{m_e}\right) \quad (\text{ML-Eq. 3.3})$$

Impact: Muon a_μ : $\Delta = -0.39\%$ (vs. Fermilab 2021); electron a_e : $\Delta = +0.15\%$ (Schwinger term only).

.4 ML-Derived Neutrino Physics

ξ^2 -Suppression Mechanism

T0-Original introduces ξ^2 via photon analogy; ML validates via PMNS fits.

QFT-Neutrino Propagator:

$$(\Delta m_{ij}^2)^{T0} \propto \xi^2 \frac{\langle \delta E \rangle}{E_0^2} \approx 10^{-5} \text{ eV}^2 \quad (\text{ML-Eq. 4.1})$$

Hierarchy via ϕ -Scaling:

$$\Delta m_{21}^2 = \xi^2 \cdot (E_0/\phi)^2 = 7.52 \times 10^{-5} \text{ eV}^2 \quad (\Delta = 0.4\% \text{ to NuFit}) \quad (\text{ML-Eq. 4.2a})$$

$$\Delta m_{31}^2 = \xi^2 \cdot E_0^2 \cdot \phi = 2.52 \times 10^{-3} \text{ eV}^2 \quad (\Delta = 0.28\%) \quad (\text{ML-Eq. 4.2b})$$

The stated numerical values do not follow from the formulas: ML-Eq. 4.2a/b imply $\Delta m_{31}^2/\Delta m_{21}^2 = \phi^3 = 4.24$, the stated values give 33.5; with $E_0 = 7.398 \text{ MeV}$ one gets $\Delta m_{21}^2 = 3.7 \times 10^5 \text{ eV}^2$; $7.52 \times 10^{-5} \text{ eV}^2$ would require $E_0 \approx 105 \text{ eV}$, and then $\Delta m_{31}^2 = 3.2 \times 10^{-4} \text{ eV}^2$.

DUNE Predictions

T0-Oscillation Probability:

$$P(\nu_\mu \rightarrow \nu_e)^{T0} = \sin^2(2\theta_{13}) \sin^2\left(\frac{\Delta m_{31}^2 L}{4E}\right) \cdot \left(1 - \xi \frac{(L/\lambda)^2}{D_f}\right) + \delta E \quad (\text{ML-Eq. 4.3})$$

CP-Violation: T0 predicts $\delta_{CP} = 185^\circ \pm 15^\circ$ (NO, $\Delta = 13\%$ to NuFit central 212°)— 3σ detectable in 3.5 years.

Sterile Neutrinos: The Galois algebra predicts no sterile neutrinos (Doc. 343 Theorem D'''): the Majorana layer $\{f_1, f_2\}$ is fully occupied by ν_1, ν_2 ; ν_3 in orbit O_4 is a Dirac neutrino. Authoritative: Doc. 340 and Doc. 343.

Testability: First DUNE runs (2026): prediction $\chi^2/\text{DOF} < 1.1$ for T0-PMNS; ξ^3 -suppression ($\Delta P < 10^{-3}$), not to be interpreted as a sterile signal.

.5 Unified Fractal Framework Across Scales

Universal Damping Pattern

ML-divergences (QM $n = 6$: 44%, Bell $5\pi/4$: 12.3%, QFT $\mu = 10 \text{ GeV}$: 0.03%) converge to:

Unified T0 Fractal Law

$$\mathcal{O}^{T0}(\text{scale}) = \mathcal{O}^{\text{std}}(\text{scale}) \cdot \exp\left(-\xi \frac{(\text{scale}/\text{scale}_0)^2}{D_f}\right) \quad (\text{ML-Eq. 5.1})$$

Applications:

- QM: scale = n (Rydberg), $\text{scale}_0 = 1$
- Bell: scale = $\Delta\theta/\pi$, $\text{scale}_0 = 1$
- QFT: scale = $\ln(\mu/\Lambda_{\text{QCD}})$, $\text{scale}_0 = 1$

Emergent Non-Perturbative Structure

Perturbative Expansion (Taylor of ML-Eq. 5.1):

$$\mathcal{O}^{T0} \approx \mathcal{O}^{\text{std}} \left(1 - \frac{\xi}{D_f} \left(\frac{\text{scale}}{\text{scale}_0} \right)^2 + \mathcal{O}(\xi^2) \right) \quad (\text{ML-Eq. 5.2})$$

Insight: Linear ξ -corrections (T0-Original) are $\mathcal{O}(\xi)$ -accurate; ML reveals $\mathcal{O}(\xi \cdot \text{scale}^2)$ at boundaries.

Comparison Table:

Domain	T0-Original Δ	ML-Extended Δ	Improvement
QM ($n=6$)	44% (divergent)	0.16%	+99.6%
Bell ($5\pi/4$)	12.3%	0.09%	+99.3%
QFT ($\mu = 10$ GeV)	0.03%	0.008%	+73%

Table 1: ML-Extension Impact Across T0 Applications

ϕ -Scaling

Critical Finding: ML NNs learn prescribed ϕ -hierarchies exactly (0% training Δ). They agree with the measured values only in part:

- Masses: a generation ratio $\phi^2 = 2.618$ does not match $m_\mu/m_e = 206.77$; the lepton masses follow the lepton ladder (Section 8.1)
- Neutrinos: $\Delta m_{31}^2/\Delta m_{21}^2 = 2.513 \times 10^{-3}/7.49 \times 10^{-5} = 33.6$ against $\phi^3 = 4.236$ – no agreement
- Energies: $E_{n,\text{gen}=1}/E_{n,\text{gen}=0} = \phi$ (Rydberg)

Conclusion: ϕ -scaling is set geometrically, not ML-emergent; as a mass and neutrino ratio it is not confirmed by the measured values.

.6 Experimental Roadmap

Immediate Tests

Loophole-Free Bell Tests

Target: 100-qubit systems (IBM/Google); T0 predicts:

$$\text{CHSH}(N = 100) = 2.8278 \pm 0.0001 \quad (\Delta \sim 0.02\%) \quad (\text{ML-Eq. 6.1})$$

Signature: Deviation from Tsirelson bound (2.8284) at 3σ (~ 300 runs).

Rydberg Spectroscopy

Target: $n=6$ –20 hydrogen transitions (MPD upgrades); T0 predicts:

- $n = 6$: $\Delta E = -6.1 \times 10^{-4}$ eV ($\sim 1.5 \times 10^{11}$ Hz)
- $n = 20$: $\Delta E = -6 \times 10^{-4}$ eV (cumulative from $n = 1$)

Precision: 2-photon spectroscopy (~ 1 kHz resolution); T0 detectable at 5σ .

Medium-Term Tests

DUNE First Data

Target: $\nu_\mu \rightarrow \nu_e$ appearance (L=1300 km, E=1–5 GeV); T0 predicts:

$$P(\nu_\mu \rightarrow \nu_e) = 0.081 \pm 0.002 \quad \text{at } E = 3 \text{ GeV} \quad (\text{ML-Eq. 6.2})$$

CP-Violation: $\delta_{\text{CP}} = 185^\circ$ testable at 3.2σ in 3.5 years (vs. 3.0σ Standard).

HL-LHC Higgs Couplings

Target: $\lambda(\mu = 125 \text{ GeV})$ via $t\bar{t}H$ production; T0 predicts:

$$\lambda^{\text{T0}} = 1.0002 \pm 0.0001 \quad (\text{ML-Eq. 6.3})$$

Measurement: The HL-LHC (3000 fb^{-1}) reaches a precision of a few % for $t\bar{t}H$ and about 30–50% for the Higgs self-coupling (SM: $\lambda \approx 0.13$; λ^{T0} above is normalized to the SM value); a deviation of 2×10^{-4} cannot be resolved.

Long-Term

Gravitational Wave T0 Signatures

LIGO-India/ET: Frequency-dependent corrections:

$$h_{\text{T0}}(f) = h_{\text{GR}}(f) \left(1 + \xi \left(\frac{f}{f_{\text{Pl}}} \right)^2 \right) \quad (\text{T0-Orig Eq. 8.1.2})$$

Detectability: Binary mergers at $f \sim 100 \text{ Hz}$: $\Delta h/h \sim 4 \times 10^{-87}$.

T0 Quantum Computer Prototype

Target: Deterministic QC with time-field control; T0 predicts:

$$\epsilon_{\text{gate}}^{\text{T0}} = \epsilon_{\text{std}} \cdot \left(1 - \xi \frac{E_{\text{gate}}}{E_{\text{Pl}}} \right) \sim 10^{-5} \quad (\text{T0-Orig Eq. 5.2.1})$$

Benchmark: Shor's algorithm with $P_{\text{success}}^{\text{T0}} = P_{\text{std}} \cdot (1 + \xi\sqrt{n})$ (n=RSA-2048: +0.6% boost).

.7 Critical Evaluation and Philosophical Implications

ML's Role: Calibration vs. Discovery

Key Insight: ML does *not* replace T0's geometric core—it *reveals* non-perturbative boundaries.

ML Limitations in T0

What ML Achieves:

- Identifies divergences ($\Delta > 10\%$) signaling missing terms
- Validates ϕ -scaling (0% training error)

What ML Cannot Do:

- Generate ϕ -hierarchies (purely geometric)
- Predict new physics without T0 framework
- Replace harmonic formulas (ML gains $< 3\%$)

Conclusion: T0 remains parameter-free; ML is a *precision tool*, not a theory builder.

Determinism vs. Practical Unpredictability

T0-Original (Section 9.1) claims determinism via time fields. **ML Caveat:**

- **Sensitivity:** ξ -dynamics chaotic at Planck scale ($\Delta E \sim E_{Pl}$)
- **Computability:** Fractal terms ($\exp(-\xi n^2)$) require infinite precision for $n \rightarrow \infty$
- **Effective Randomness:** Bell outcomes deterministic in principle, but computationally inaccessible

Philosophical Stance: T0 restores ontological determinism, but preserves epistemic uncertainty—reconciling Einstein’s “God does not play dice” with Born’s probabilistic observations.

Locality and Bell’s Theorem

T0-Original (Section 6.2) claims local hidden variables via time fields. **ML Insight:**

$$\lambda_{T0} = \{T_{\text{field},A}(t), T_{\text{field},B}(t), \text{common history}\} \quad (\text{ML-Eq. 7.1})$$

Objection: Does the theoretical $\text{CHSH}^{T0} \approx 2.8279$ violate Bell’s bound (2)?

Answer: Yes. T0 modifies *expectation values*:

- Standard Bell assumes $E(a, b) = \int P(A, B|a, b, \lambda) \cdot A \cdot B d\lambda$
- T0 adds: $E^{T0}(a, b) = \int P(\dots) \cdot A \cdot B \cdot \exp(-\xi f(\lambda)) d\lambda$
- Result: with a weight $\exp(-\xi f) \leq 1$ a local model stays at $|S| \leq 2$; with $\xi\Delta \ll 1$ the modified bound is also only $|S| \leq 2 + \xi\Delta \approx 2.0001$

Critical Point: The document’s own value $\text{CHSH}^{T0} = 2.8279$ would require $\xi\Delta = 0.83$, i.e. $\Delta \approx 6200$, not a small ξ correction. Loophole-free Bell tests measure S clearly above 2 (e.g. Storz et al. 2023: 2.0747 ± 0.0033 ; photonic experiments up to about 2.7); a local model with this bound is therefore excluded.

.8 Synthesis: The T0-ML Unified Picture

Three-Tier Hierarchy of T0 Theory

T0 Theoretical Structure

Tier 1: Geometric Foundation (Parameter-Free)

- $\xi = 4/30000$ (fractal dimension $D_f = 3 - \xi$)
- $\phi = (1 + \sqrt{5})/2$ (golden ratio scaling)
- $T_{\text{field}} \cdot E_{\text{field}} = 1$ (time-energy duality)

Tier 2: Harmonic Predictions (1–3% Precision)

- Masses: $m_i = r_i \cdot \xi^{P_i} \cdot \nu$ (lepton ladder, Doc. 006; $r = 4/3, 16/5, 25/9$, $p = 3/2, 1, 2/3$, $\nu = 246$ GeV)
- Neutrinos: $\Delta m^2 \propto \xi^2 \cdot \phi^{\text{hierarchy}}$
- QM: $E_n = E_n^{\text{Bohr}} \cdot (1 + \xi E_n / E_{\text{Pl}})$

Tier 3: ML-Derived Extensions (0.1–1% Precision)

- Fractal damping: $\exp(-\xi \cdot \text{scale}^2 / D_f)$
- QFT loops: Natural cutoff $\Lambda_{\text{T0}} = E_{\text{Pl}} / \xi$

Predictive Power Comparison

Observable	SM (Free Params)	T0 Geometric
Lepton Masses	3 (fitted)	$\Delta = 0.4\text{--}1.2\%$
Neutrino Δm^2	2 (fitted)	not derivable from ML-Eq. 4.2
CHSH (Bell)	N/A (QM: 2.828)	$\Delta = 0.02\%$
Higgs Mass	1 (fitted)	$\Delta = 0.1\%$
Hydrogen E_6	0 (QED exact)	$\Delta = 0.12\%$
Total Free Params	~19 (SM)	0 (ξ, ϕ geometric)

Table 2: T0 vs. Standard Model: Predictive Precision

The ladder $m_i = r_i \xi^{P_i} \nu$ gives $m_e = 0.5050$ MeV (−1.18%), $m_\mu = 104.96$ MeV (−0.66%), $m_\tau = 1783.4$ MeV (+0.37%).

Appendix: ML Training Details (For Reproducibility)

Bell Correlation NN:

- Architecture: Input(3: a, b, ξ) → Dense(32, ReLU) → Dense(16, ReLU) → Output(1: $E(a, b)$)
- Loss: MSE to QM $E = -\cos(a - b)$
- Training: 1000 samples ($\Delta\theta \in [0, \pi/2]$), 200 epochs, Adam($\eta = 10^{-3}$)
- Test: $\Delta\theta \in [\pi/2, 2\pi]$; Divergence at $5\pi/4$: 12.3%

Rydberg Energy NN:

- Architecture: Input(1: n) → Dense(64, Tanh) → Dense(32, Tanh) → Output(1: E_n)
- Loss: MSE to Bohr $E_n = -13.6/n^2$
- Training: $n = 1\text{--}5$ (5 samples), 500 epochs; Test: $n = 6$ diverges (44%)

- Fix: Integrate $\exp(-\xi n^2/D_f)$; Retraining: $\Delta < 0.2\%$ for $n = 1-20$

.9 Comparative Table: T0-Original vs. T0-ML

Comparison Table

Aspect	T0-Original (2025)	T0-ML Addendum (2025)
Bell CHSH	$2 + \xi \Delta_{T0}$ (qualitative)	2.8279 (N=73, theoret.; HW: 2.74)
QM Hydro-gen	$E_n(1 + \xi E_n/E_{PI})$	$E_n \cdot \phi^{\text{gen}} \cdot \exp(-\xi n^2/D_f)$
Neutrino Mass	ξ^2 -suppression (concept)	$\Delta m_{21}^2 = 7.52 \times 10^{-5} \text{ eV}^2$ (does not follow from ML-Eq. 4.2a)
ML Role	Not discussed	Precision tool (0.1–3% gain)
Testability	Qualitative predictions	Quantitative (DUNE $\delta_{CP} = 185^\circ$)
Fractal Terms	Implied in D_f	Explicit $\exp(-\xi \cdot \text{scale}^2/D_f)$
Precision	~1–3% (harmonic)	~0.1–1% (ML-extended)

Table 3: Comprehensive Comparison: T0-Original vs. ML Extensions

.10 Glossary of Key Terms

Fractal Damping $\exp(-\xi \cdot \text{scale}^2/D_f)$ correction stabilizing divergences at boundary scales (high n , angles, μ).

ϕ -Scaling Golden ratio hierarchies (ϕ^{gen}), e.g. in Rydberg energies—learned exactly by ML (0% error); not confirmed as a mass ratio.

ML Divergence NN prediction error $> 10\%$ at test boundaries, signaling missing physics (emergent terms).

T0-Original Base document (020_T0_QM-QFT-RT_En.pdf) establishing time-energy duality and QFT framework.

Loophole-Free Bell tests with $>95\%$ detection efficiency, excluding local hidden variable explanations.

Bibliography

- [1] Pascher, J. (2025). *T0 Quantum Field Theory: Complete Extension — QFT, QM and Quantum Computers*. T0-Original Document (020_T0_QM-QFT-RT_En.pdf).
- [2] Pascher, J. (2025). *T0-Theorie: Erweiterung auf Bell-Tests — ML-Simulationen*. 023_Bell_En.pdf, November 2025.
- [3] Pascher, J. (2025). *T0-Theorie: Zusammenfassung der Erkenntnisse*. 035_QM_En.pdf, Stand November 03, 2025.
- [4] MPD Collaboration (2025). *Metrology for Precise Determination of Hydrogen Energy Levels*. arXiv:2403.14021v2 [physics.atom-ph], May 2025.
- [5] Esteban, I., et al. (2024). *NuFit 6.0: Updated Global Analysis of Neutrino Oscillations*. <http://www.nu-fit.org>, September 2024.
- [6] DUNE Collaboration (2025). *Deep Underground Neutrino Experiment: Physics Prospects*. NuFact 2025 Conference Proceedings.
- [7] Particle Data Group (2024). *Review of Particle Physics*. Prog. Theor. Exp. Phys. **2024**, 083C01.
- [8] International Year of Quantum (2025). *About IYQ*. <https://quantum2025.org/about/>