

T0 Model: Field-Theoretic Derivation of the β -Parameter

Contents

.1	Introduction and Motivation	1
.2	Natural Units Framework	2
.3	Fundamental Structure of the T0 Model	2
	Time-Mass Duality	2
	Fundamental Field Equation	2
.4	Geometric Derivation of the β -Parameter	3
	Spherically Symmetric Point Source	3
	Solution of the Field Equation	3
	Determination of Integration Constants	3
	The Characteristic Length Scale	4
	Definition of the β -Parameter	4
.5	Physical Interpretation of the β -Parameter	4
	Dimensional Analysis	4
	Connection to Classical Physics	4
	Limiting Cases and Application Domains	4
.6	Comparison with Established Theories	5
	Connection to General Relativity	5
	Differences from the Standard Model	5
.7	Experimental Predictions	5
	Time Dilation Effects	5
	Spectroscopic Tests	5
.8	Mathematical Consistency	5
	Conservation Laws	5
	Solution Stability	6

.1 Introduction and Motivation

The T0 model introduces a fundamentally new perspective on spacetime, where time itself becomes a dynamic field. At the center of this theory lies the dimensionless β -parameter, which characterizes the strength of the time field and establishes a direct connection between gravitational and electromagnetic interactions.

This work focuses exclusively on the mathematically rigorous derivation of the β -parameter from the fundamental field equations of the T0 model, avoiding the complexity of additional scaling parameters.

Central Result

The β -parameter is derived as:

$$\beta = \frac{Gm}{r} = \frac{r_s}{2r} \quad (1)$$

where G is the gravitational constant, m is the source mass, r is the distance from the source, and $r_s = 2Gm$ is the Schwarzschild radius.

.2 Natural Units Framework

The T0 model employs the system of natural units established in modern quantum field theory [Peskin & Schroeder(1995), Weinberg(1995)]:

- $\hbar = 1$ (reduced Planck constant)
- $c = 1$ (speed of light)

This system reduces all physical quantities to energy dimensions and follows the tradition established by Dirac [Dirac(1958)].

Dimensions in Natural Units

- Length: $[L] = [E^{-1}]$
- Time: $[T] = [E^{-1}]$
- Mass: $[M] = [E]$
- The β -parameter: $[\beta] = [1]$ (dimensionless)

.3 Fundamental Structure of the T0 Model**Time-Mass Duality**

The central principle of the T0 model is the time-mass duality, which states that time and mass are inversely linked. This relationship differs fundamentally from the conventional treatment in general relativity [Einstein(1915), Misner et al.(1973)].

Theory	Time	Mass	Reference
Einstein GR	$dt' = \sqrt{g_{00}}dt$	$m_0 = \text{const}$	[Einstein(1915), Misner et al.(1973)]
Special Relativity	$t' = \gamma t$	$m_0 = \text{const}$	[Einstein(1905)]
T0 Model	$T(x) = \frac{1}{m(x)}$	$m(x) = \text{dynamic}$	This work

Table 1: Comparison of time-mass treatment in different theories

Fundamental Field Equation

The fundamental field equation of the T0 model is derived from variational principles, analogous to the approach for scalar field theories [Weinberg(1995)]:

$$\nabla^2 m(x) = 4\pi G \rho(x) \cdot m(x) \quad (2)$$

This equation shows structural similarity to the Poisson equation of gravitation $\nabla^2 \phi = 4\pi G \rho$ [Jackson(1998)], but is nonlinear due to the factor $m(x)$ on the right-hand side.

The time field follows directly from the inverse relationship:

$$T(x) = \frac{1}{m(x)} \quad (3)$$

.4 Geometric Derivation of the β -Parameter

Spherically Symmetric Point Source

For a point mass source, we use the established methodology for solving Einstein's field equations [[Schwarzschild\(1916\)](#), [Misner et al.\(1973\)](#)]. The mass density of a point source is described by the Dirac delta function:

$$\rho(\vec{x}) = m_0 \cdot \delta^3(\vec{x}) \quad (4)$$

where m_0 is the mass of the point source.

Solution of the Field Equation

Outside the source ($r > 0$), where $\rho = 0$, the field equation reduces to:

$$\nabla^2 m(r) = 0 \quad (5)$$

The spherically symmetric Laplace operator [[Jackson\(1998\)](#), [Griffiths\(1999\)](#)] yields:

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{dm}{dr} \right) = 0 \quad (6)$$

The general solution to this equation is:

$$m(r) = \frac{C_1}{r} + C_2 \quad (7)$$

Determination of Integration Constants

Asymptotic boundary condition: For large distances, the time field should assume a constant value T_0 :

$$\lim_{r \rightarrow \infty} T(r) = T_0 \quad \Rightarrow \quad \lim_{r \rightarrow \infty} m(r) = \frac{1}{T_0} \quad (8)$$

This gives us: $C_2 = \frac{1}{T_0}$

Behavior at the origin: Using Gauss's theorem [[Griffiths\(1999\)](#), [Jackson\(1998\)](#)] for a small sphere around the origin:

$$\oint_S \nabla m \cdot d\vec{S} = 4\pi G \int_V \rho(r) m(r) dV \quad (9)$$

For a small radius ϵ :

$$4\pi\epsilon^2 \left. \frac{dm}{dr} \right|_{r=\epsilon} = 4\pi G m_0 \cdot m(\epsilon) \quad (10)$$

With $\frac{dm}{dr} = -\frac{C_1}{r^2}$ and $m(\epsilon) \approx \frac{1}{T_0}$ for small ϵ :

$$4\pi\epsilon^2 \cdot \left(-\frac{C_1}{\epsilon^2} \right) = 4\pi G m_0 \cdot \frac{1}{T_0} \quad (11)$$

This yields: $-C_1 = \frac{Gm_0}{T_0}$, hence $C_1 = -\frac{Gm_0}{T_0}$.

The Characteristic Length Scale

The complete solution reads:

$$m(r) = \frac{1}{T_0} \left(1 - \frac{Gm_0}{r} \right) \quad (12)$$

The corresponding time field is:

$$T(r) = \frac{T_0}{1 - \frac{Gm_0}{r}} \quad (13)$$

For the practically important case $Gm_0 \ll r$, we obtain the approximation:

$$T(r) \approx T_0 \left(1 + \frac{Gm_0}{r} \right) \quad (14)$$

The characteristic length scale at which the time field significantly deviates from T_0 is:

$$r_0 = Gm_0 \quad (15)$$

This scale is equal to half the Schwarzschild radius $r_s = 2GM/c^2 = 2Gm$ in geometric units [[Misner et al.\(1973\)](#), [Carroll\(2004\)](#)].

Definition of the β -Parameter

The dimensionless β -parameter is defined as the ratio of the characteristic length scale to the actual distance:

$$\beta = \frac{r_0}{r} = \frac{Gm_0}{r} \quad (16)$$

This parameter measures the relative strength of the time field at a given point. In units of the Schwarzschild radius, $\beta = r_s/(2r)$. The form $\beta = 2Gm/r = r_s/r$ that is sometimes used does not follow from the derivation; it is twice as large.

.5 Physical Interpretation of the β -Parameter

Dimensional Analysis

The dimensionlessness of the β -parameter in natural units:

$$[\beta] = \frac{[G][m]}{[r]} = \frac{[E^{-2}][E]}{[E^{-1}]} = [1] \quad (17)$$

Connection to Classical Physics

The β -parameter shows direct connections to established physical concepts:

- **Gravitational potential:** β is proportional to the Newtonian potential $\Phi = -Gm/r$
- **Schwarzschild radius:** $\beta = r_s/(2r)$ in geometric units
- **Escape velocity:** β is related to v_{esc}^2/c^2

Limiting Cases and Application Domains

For the hydrogen atom, $\beta = Gm_p/(a_0 c^2) = 2.3 \times 10^{-44}$; the frequently quoted value 10^{-39} is the ratio of gravitational to Coulomb force, not β .

Physical System	Typical β -Value	Regime
Hydrogen atom	$\sim 2 \times 10^{-44}$	Quantum mechanics
Earth (surface)	$\sim 10^{-9}$	Weak gravitation
Sun (surface)	$\sim 10^{-6}$	Stellar physics
Neutron star	~ 0.1	Strong gravitation
Schwarzschild horizon	$\beta = 1/2$	Limiting case

Table 2: Typical β -values for various physical systems

.6 Comparison with Established Theories

Connection to General Relativity

In general relativity, the parameter $r_s/r = 2Gm/r$ characterizes the strength of the gravitational field. The T0 parameter $\beta = Gm/r = r_s/(2r)$ is half as large; it is exactly the first-order term in $\sqrt{g_{00}} = \sqrt{1 - 2Gm/r} \approx 1 - Gm/r$.

Differences from the Standard Model

While the Standard Model of particle physics treats time as an external parameter, the T0 model makes time a dynamic field. The β -parameter quantifies this dynamics and represents a measurable deviation from standard physics.

.7 Experimental Predictions

Time Dilation Effects

The T0 model predicts a modified time dilation:

$$\frac{dt}{dt_0} = 1 - \beta = 1 - \frac{Gm}{r} \quad (18)$$

This relationship is identical to the gravitational time dilation of GR in first order, $dt/dt_0 = \sqrt{1 - 2Gm/r} \approx 1 - Gm/r$, but offers a fundamentally different theoretical foundation. With the form $\beta = 2Gm/r$, $1 - \beta$ would be twice as large and would not agree with GR to first order.

Spectroscopic Tests

The β -parameter could be tested through high-precision spectroscopy:

- Gravitational redshift in stellar spectra
- Atomic clock experiments in different gravitational potentials
- High-precision interferometry

.8 Mathematical Consistency

Conservation Laws

The derivation of the β -parameter respects fundamental conservation laws:

- **Energy conservation:** Guaranteed by the Lagrangian formulation

- **Momentum conservation:** From spatial translation invariance
- **Dimensional consistency:** Verified in all derivation steps

Solution Stability

The spherically symmetric solution is stable against small perturbations, which can be shown by linearization around the ground state solution.

Bibliography

- [Carroll(2004)] Carroll, S. M. *Spacetime and Geometry: An Introduction to General Relativity*. Addison-Wesley, San Francisco, CA (2004).
- [Dirac(1958)] Dirac, P. A. M. *The Principles of Quantum Mechanics*. Oxford University Press, Oxford, 4th edition (1958).
- [Einstein(1905)] Einstein, A. Zur Elektrodynamik bewegter Körper. *Annalen der Physik*, **17**, 891–921 (1905).
- [Einstein(1915)] Einstein, A. Die Feldgleichungen der Gravitation. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften*, 844–847 (1915).
- [Griffiths(1999)] Griffiths, D. J. *Introduction to Electrodynamics*. Prentice Hall, Upper Saddle River, NJ, 3rd edition (1999).
- [Jackson(1998)] Jackson, J. D. *Classical Electrodynamics*. John Wiley & Sons, New York, 3rd edition (1998).
- [Misner et al.(1973)] Misner, C. W., Thorne, K. S., and Wheeler, J. A. *Gravitation*. W. H. Freeman and Company, New York (1973).
- [Peskin & Schroeder(1995)] Peskin, M. E. and Schroeder, D. V. *An Introduction to Quantum Field Theory*. Addison-Wesley, Reading, MA (1995).
- [Schwarzschild(1916)] Schwarzschild, K. Über das Gravitationsfeld eines Massenpunktes nach der Einsteinschen Theorie. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften*, 189–196 (1916).
- [Weinberg(1995)] Weinberg, S. *The Quantum Theory of Fields, Volume I: Foundations*. Cambridge University Press, Cambridge (1995).