

Justification of the 4D Torus Geometry

and Derivation of L_0 as the Smallest Stable Winding

Johann Pascher · March 2026

<https://github.com/jpascher/T0-Time-Mass-Duality>

March 2026

Contents

.1	Why a Fourth Dimension?	2
	The Divergence Problem of Quantum Field Theory	2
	Periodic Geometry as Solution	2
	Four Dimensions and the Form of ξ	2
.2	Why a Torus?	3
	Winding Numbers as Conserved Quantities	3
	Torus Eigenvalues and Particle Masses	3
	Compactification	3
.3	Derivation of L_0 from Torus Geometry	3
	The Winding Density	3
	Smallest Stable Winding	4
	Geometric Interpretation	4
	Consistency of the Two Derivation Paths	4
.4	Comparison with Other Approaches	5
	String Theory	5
	Loop Quantum Gravity	5
	Asymptotic Safety	5
.5	Physical Consequences	5
	Vacuum Catastrophe Resolved	5
	Three Generations as Overtone Structure	5
	No Big Bang and No Frequency Dependence	5
.6	Summary	6

Abstract

The 4D torus is not a free postulate of T0 theory. It is supported by three conditions: (1) freedom from divergences in quantum field theory requires periodic topology, (2) the four spacetime dimensions (3+1, empirical input) are reflected in the form of $\xi = 4/30000$, (3) stable winding numbers on the torus generate the observed quantum numbers. The minimal length scale $L_0 = \xi \cdot \ell_P$ is geometrically the length of a single torus lattice cell — the smallest stable winding. The Lagrangian and the torus yield L_0 consistently via two independent paths.

.1 Why a Fourth Dimension?

The Divergence Problem of Quantum Field Theory

Quantum field theory suffers from a fundamental problem: the vacuum energy diverges:

$$E_{\text{vac}} = \sum_n \frac{1}{2} \hbar \omega_n \rightarrow \infty \quad (1)$$

The observed cosmological constant is 10^{120} times smaller than the QFT prediction. The Standard Model resolves this through artificial renormalisation. T0/FFGFT resolves it geometrically.

Periodic Geometry as Solution

The harmonic series $\sum 1/n$ diverges; the series of inverse squares converges. Both hold regardless of whether the modes are pictured on a line or a circle; the topology does not turn the divergent series into a convergent one. For the inverse squares (Euler 1735):

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \quad (2)$$

Topological Conclusion

The periodic geometry automatically suppresses high modes. If spacetime has a periodic topology, UV divergences vanish geometrically — without artificial renormalisation. The convergence of $\sum 1/n^2$ itself is not a consequence of the topology (see above).

Four Dimensions and the Form of ξ

The parameter $\xi = 4/30000$ encodes the dimensionality of spacetime:

$$\xi_d \sim (10^{-1})^d \quad \Rightarrow \quad \text{for } d = 4 : \quad \xi_4 \sim 10^{-4} \quad (3)$$

Four Dimensions and the Form of ξ

The **4** in the numerator of $\xi = 4/30000$ stands for the four spacetime dimensions. The dimension count 3+1 itself is empirical input — no theory derives it; the form of ξ reflects it.

.2 Why a Torus?

Winding Numbers as Conserved Quantities

On a torus, winding numbers are topologically conserved. They generate stable discrete states:

$$\text{Spin} = \text{winding number } w = n_\phi / n_\theta \quad (4)$$

$$\text{Charge} = \text{flux } \Phi = n \cdot h/e \quad (5)$$

$$\text{Masses} = \text{torus overtone frequencies} \quad (6)$$

Torus vs. Sphere

A sphere S^3 is simply connected — no stable winding numbers. The torus is not simply connected — it generates exactly the observed quantum numbers. This is the topological reason for choosing the torus.

Torus Eigenvalues and Particle Masses

The Laplace operator on a d -dimensional torus with radii R_i has eigenvalues:

$$E_{\vec{n}} = \frac{\hbar^2}{2} \sum_i \frac{n_i^2}{R_i^2}, \quad \vec{n} \in \mathbb{Z}^d \quad (7)$$

Particle masses appear as torus overtone frequencies with step size $1/3 = 1/D_f$:

$$m_n = m_0 \cdot n^{1/3} \quad (8)$$

The first overtones $n = 2, 3$ thus give the ratios 1.26 and 1.44; the ratio $m_\mu/m_e = 206.8$ requires $n = 206.8^3 \approx 8.8 \cdot 10^6$. The lepton hierarchy and the Koide formula do not follow from this simple overtone ansatz; a derivation of the Koide formula is not given here.

Compactification

The fourth dimension is rolled up to radius r_4 . This radius must coincide with the sub-Planck scale from the Lagrangian — otherwise the theory would be inconsistent. The condition:

$$r_4 = L_0 = \xi \cdot \ell_P \quad (9)$$

The topology is therefore:

$$\mathbb{R}^3 \times S^1 \quad \text{with} \quad r_4 = \xi \cdot \ell_P \quad (10)$$

.3 Derivation of L_0 from Torus Geometry

The Winding Density

The torus has $f = 1/\xi$ sub-Planck cells per Planck length. With $\xi = 4/30000$:

$$f = \frac{1}{\xi} = \frac{30000}{4} = 7500 \quad \text{cells per } \ell_P \quad (11)$$

This number is not arbitrary — it follows directly from the geometric winding density $30000/4$ of the torus. The prime factorisation $7500 = 2^2 \times 3 \times 5^4$ encodes the symmetry structure of spacetime.

On notation: here f is the base winding number $f = 1/\xi = 7500$, not a frequency. In natural units f is only another reading of the duality $T \cdot m = 1$: $f = t_P/t_0 = \Lambda_{T0}/E_P$ with $t_0 = \xi t_P$ and $\Lambda_{T0} = E_P/\xi$, and $\xi E_0^2 = 1$ gives $f = E_0^2$. As a dimensionless ratio, f cannot be set to 1 with justification (Doc. A135).

Smallest Stable Winding

The smallest stable winding on the torus has the length of a single lattice cell:

$$L_0 = \frac{\ell_P}{f} = \ell_P \cdot \xi \quad (12)$$

Numerically:

$$L_0 = 1.616 \times 10^{-35} \text{ m} \times \frac{4}{30000} = 1.616 \times 10^{-35} \text{ m} \times 1.333 \times 10^{-4} \quad (13)$$

Minimal Length Scale from Torus Geometry

$$L_0 = \xi \cdot \ell_P = 2.155 \times 10^{-39} \text{ m} \quad (14)$$

L_0 is the length of a single torus lattice cell — the smallest stable winding.

Geometric Interpretation

L_0 is not simply ξ times the Planck length — L_0 is the length of a single torus lattice cell. Below L_0 :

- The concept of "distance" loses its physical meaning.
- All vacuum fluctuations are fully effective.
- The fractal correction K_{frak} becomes structurally necessary.

Consistency of the Two Derivation Paths

L_0 is not derived twice — the Lagrangian and the torus yield the same number via two independent paths:

Two Consistent Paths to L_0

$$\begin{array}{lcl}
 \text{Lagrangian:} & g_T = \xi \cdot m \Rightarrow \xi = 4/30000 & \\
 \Downarrow & & \\
 \text{Torus:} & f = 1/\xi = 7500 \text{ cells per } \ell_P & \\
 \Downarrow & & \\
 \text{Smallest winding:} & L_0 = \ell_P / f = \xi \cdot \ell_P & \\
 \Downarrow & & \\
 & L_0 = 2.155 \times 10^{-39} \text{ m} & \text{(consistent from both paths)} \quad (15)
 \end{array}$$

The Lagrangian provides ξ . The torus explains what ξ means geometrically: the reciprocal winding density.

.4 Comparison with Other Approaches

String Theory

String theory requires 10 or 11 dimensions. The T0 Lagrangian with a single parameter ξ suffices with 4. Every additional dimension would be a postulate — in FFGFT there are no free parameters.

Loop Quantum Gravity

LQG discretises space into spin networks. T0 discretises spacetime through the periodic torus geometry. The difference: in T0 the discreteness is a topological consequence, not an additional postulate.

Asymptotic Safety

AS seeks a UV fixed point as the result of renormalisation group flow. In T0 there is no renormalisation group flow — ξ is a geometric constant, not a running parameter. The fixed point is not sought but established geometrically from the outset.

.5 Physical Consequences

Vacuum Catastrophe Resolved

On the fractal torus, the vacuum energy converges through periodic suppression of high frequencies. No artificial renormalisation required.

Three Generations as Overtone Structure

Electron, muon and tau are interpreted as states of a single torus resonator. The observed hierarchy

$$m_e : m_\mu : m_\tau \approx 1 : 206.8 : 3477 \quad (16)$$

cannot, however, be represented as fundamental tone and first overtones with the simple overtone ansatz $m_n = m_0 n^{1/3}$: the first overtones give only the ratios 1.26 and 1.44 (section Torus Eigenvalues and Particle Masses).

No Big Bang and No Frequency Dependence

The torus has no boundaries. The observed redshift does not arise through expansion but through geometric energy loss — fractal summation extends the effective path length of photons through the T0 vacuum field:

$$z \approx \xi \cdot \ln\left(\frac{d}{\ell_P}\right) \quad (17)$$

This logarithmic form does not hold up quantitatively: for $d = R_H = 1.4 \cdot 10^{26}$ m, $\ln(d/\ell_P) = 140.3$ and hence $z = \xi \cdot 140.3 = 0.019$; $z \approx 10$ would require $\ln(d/\ell_P) = 75\,000$. In this form the formula is incompatible with the observed redshifts $z > 1$. The achromaticity of the redshift is unaffected by this; the scale of the path extension does not follow from ξ alone but is calibrated via the measured H_0 .

Frequency Independence of the Redshift

Since the path extension is a property of the spacetime geometry itself, red and blue photons starting from the same location take exactly the same extended path. Their wavelengths are therefore stretched by the same percentage — there is no frequency dependence. This distinguishes the T0 mechanism fundamentally from simple “tired light” models, which fail precisely on the observed frequency independence. Derived in detail in Document 026 (Geometric Cosmology).

.6 Summary

Three Independent Conditions Point to the Same Torus

- (1) Freedom from divergences $\Rightarrow$ periodic topology
- (2) 3+1 dimensions (empirical) $\Rightarrow$ form of $\xi = 4/30000$
- (3) Winding numbers $\Rightarrow$ stable quantum numbers

All three point to the same conclusion:

$$\mathbb{R}^3 \times S^1 \quad \text{with} \quad r_4 = L_0 = \xi \cdot \ell_P = 2.155 \times 10^{-39} \text{ m} \quad (18)$$

Note: Connection to the SI System

The absolute numerical values in this document — in particular $L_0 = 2.155 \times 10^{-39} \text{ m}$ — presuppose a connection to the SI unit system. This is not circular but epistemologically necessary: from a dimensionless parameter ξ alone, no absolute metre values can be derived. The connection is made via m_e , ℓ_P or α as equivalent entry points. Treated in detail in Document 180 (Section: Connection to the SI System) and in Documents 056 and 101.

Documents 180, 181 and 182 form a unit and should not be read separately.