

203 R Operator En

Johann Pascher

Contents

.1	Starting Point: The Physical Derivation (Doc. 180)	1
.2	Definition of the Recursion Operator	2
	Diagonal Form in Mode Space	2
	Action on the Field Operator	2
.3	The Fixed-Point Equation	2
	Definition of the Fixed Point	2
	The Non-Trivial Fixed Point through Self-Consistency	2
	Solving for ξ_0	3
.4	Connection to the Three Conditions from Doc. 180	3
	Step 1: Dimensional Contribution	3
	Step 2: Geometric Factor	3
	Step 3: Self-Consistency	3
.5	The Recursion Formula	3
.6	Summary: R as a Formal Framework	4
.7	Note: Apparent Contradiction from a Category Violation	4

Abstract

The recursion operator $\mathcal{R}$ of FFGFT formalises the fractal scaling structure in mode space. This document translates the constant $\xi_0 = 4/30000$, derived on physical grounds in Doc. 180, into the formal language of the operator and checks whether the fixed-point equation $\mathcal{R}(\Phi_*) = \Phi_*$ reproduces this uniqueness. It does not: in no reading does the fixed-point condition yield $\xi_0 = 4/30000$; the value remains fixed by the three conditions from Doc. 180.

.1 Starting Point: The Physical Derivation (Doc. 180)

Doc. 180 derives $\xi_0 = 4/30000$ from three independent conditions:

1. **Dimensional contribution** 10^{-4} : Four spacetime dimensions yield $\xi \sim (10^{-1})^4 = 10^{-4}$.
2. **Geometric contribution** $4/3$: Tetrahedral symmetry: 4 vertices / 3 edges per vertex = $4/3$.
3. **Self-consistency**: The e-p- μ system yields $\kappa = 7$ as the only consistent recursion depth (0.04 % accuracy).

These three conditions fix ξ_0 without any free parameter. The task of this document: translation into the language of $\mathcal{R}$.

.2 Definition of the Recursion Operator

Diagonal Form in Mode Space

The recursion operator $\mathcal{R}$ acts diagonally on the Hilbert space $\mathcal{H} = \bigoplus_{n,l,j} \mathbb{C} \cdot \mathbf{e}_{n,l,j}$:

$$\mathcal{R}(\mathbf{e}_{n,l,j}) = \lambda_{n,l,j} \cdot \mathbf{e}_{n,l,j} \quad (1)$$

The eigenvalues $\lambda_{n,l,j}$ are determined by the mass formula:

$$\lambda_{n,l,j} = \xi_0^{p(n,l,j)-1} \cdot K_{\text{frak}} \quad (2)$$

with $K_{\text{frak}} \approx 0.986$ (fractal correction).

Action on the Field Operator

On the mode operator $\Phi = \sum_i m_i P_i$ (Doc. 202), $\mathcal{R}$ does not act as a uniform scaling map $\mathcal{R}(\Phi) = \xi_0 \cdot \Phi$. The time-mass duality symmetry reads

$$t \rightarrow \xi_0 \cdot t, \quad m \rightarrow \xi_0^{-1} \cdot m, \quad \Phi \rightarrow \Phi. \quad (3)$$

With $\Phi = \sum_i m_i P_i$ and $m \rightarrow m/\xi_0$ one would get Φ/ξ_0 , not $\xi_0 \Phi$; at the same time (3) states $\Phi \rightarrow \Phi$, and the eigenvalues (2) $\xi_0^{p-1} K_{\text{frak}}$ are mode-dependent, hence not a uniform scaling by ξ_0 . These three statements are mutually inconsistent; a uniform scaling of Φ under $\mathcal{R}$ is therefore not determined.

.3 The Fixed-Point Equation

Definition of the Fixed Point

The fixed point Φ_* is the operator that is invariant under $\mathcal{R}$:

$$\mathcal{R}(\Phi_*) = \Phi_* \quad (4)$$

A uniform scaling $\mathcal{R}(\Phi) = s\Phi$ with $s = \xi_0$ or $s = 1/\xi_0$ would have only this fixed point:

$$s \cdot \Phi_* = \Phi_* \quad \Rightarrow \quad (s - 1) \Phi_* = 0 \quad (5)$$

Since $s \neq 1$, it follows that $\Phi_* = 0$ — a trivial fixed point.

The Non-Trivial Fixed Point through Self-Consistency

The physically relevant fixed point arises from the self-consistency condition of the mass formula. The operator $\Phi_* = \sum_i m_i^{(*)} P_i$ is a fixed point of $\mathcal{R}$ if:

$$m_i^{(*)} = r_i \cdot \xi_0^{p_i} \cdot v \quad \text{and} \quad \mathcal{R}(m_i^{(*)}) = m_i^{(*)} \quad (6)$$

The second condition means:

$$\xi_0^{p_i-1} \cdot K_{\text{frak}} \cdot m_i^{(*)} = m_i^{(*)} \quad \Rightarrow \quad \xi_0^{p_i-1} \cdot K_{\text{frak}} = 1 \quad (7)$$

Solving for ξ_0

From (7) it follows for each mode i :

$$\xi_0 = K_{\text{frak}}^{-1/(p_i-1)} \quad (8)$$

For ξ_0 to be **universal** (mode-independent), this equation would have to be satisfiable for all modes simultaneously. With a universal $K_{\text{frak}} \approx 0.986$, however, it yields neither a common value nor $\xi_0 = 4/30000$ (Step 3 below).

.4 Connection to the Three Conditions from Doc. 180

Step 1: Dimensional Contribution

The recursion operator scales lengths by ξ_0 . In $D = 4$ spacetime dimensions, the 4D volume scales by ξ_0^4 :

$$\mathcal{R}^{(4D)}(V_4) = \xi_0^4 \cdot V_4 \quad (9)$$

For the mass operator, which scales with ℓ^{-1} :

$$\xi_0^{\text{dim}} = (10^{-1})^4 = 10^{-4} \quad (10)$$

Step 2: Geometric Factor

The tetrahedral symmetry of the 4D torus supplies the geometric factor from the topology of the faces:

Geometry	Value	Origin
Vertices of the tetrahedron	4	Point symmetry
Edges per vertex	3	Spatial embedding
Geometric factor ξ_0^{geo}	4/3	Ratio

Step 3: Self-Consistency

The fixed-point condition (8) with the lepton exponents $p_e = 3/2$, $p_\mu = 1$, $p_\tau = 2/3$ serves as a consistency check. For the electron ($p_e - 1 = 1/2$):

$$\xi_0 = K_{\text{frak}}^{-1/(3/2-1)} = K_{\text{frak}}^{-2} \approx (0.986)^{-2} \approx 1.03 \quad (11)$$

This lies far from $\xi_0 = 4/30000 \approx 1.333 \cdot 10^{-4}$; the reading $K_{\text{frak}}^2 = (0.986)^2 = 0.972$ does not reach the value either. In no reading does the fixed-point condition yield $\xi_0 = 4/30000$.

ξ_0 is therefore fixed solely by the three conditions from Doc. 180, with the e-p- μ system: the exponent $\kappa = 7$ is the only integer recursion depth that satisfies all three conditions simultaneously.

.5 The Recursion Formula

The recursion operator generates the mass hierarchy through repeated application:

$$\mathcal{R}^{(k)}(\mathbf{e}_{n,l,j}) = \xi_0^{k(p_{n,l,j}-1)} \cdot K_{\text{frak}}^k \cdot \mathbf{e}_{n,l,j} \quad (12)$$

For $k = \kappa = 7$ (recursion depth from Doc. 180):

$$\mathcal{R}^{(7)}(\mathbf{e}_e) = \xi_0^{7(3/2-1)} \cdot K_{\text{frak}}^7 \cdot \mathbf{e}_e = \xi_0^{7/2} \cdot K_{\text{frak}}^7 \cdot \mathbf{e}_e \quad (13)$$

The self-consistency condition requires this expression to be consistent with the known electron mass. It does not fix the value $\xi_0 = 4/30000$; that value is taken over from Doc. 180.

.6 Summary: R as a Formal Framework

Physical derivation (Doc. 180)	Operator language (Doc. 203)
10^{-4} from 4D spacetime	$\mathcal{R}^{(4D)}$ scales by ξ_0^4
$4/3$ from tetrahedral geometry	Geometric factor in K_{frak}
$\kappa = 7$ from the e-p- μ system	$\mathcal{R}^{(7)}(\Phi_*) = \Phi_*$ fixed-point condition
$\xi_0 = 4/30000$ unique	Taken over; the fixed-point condition does not yield it

The recursion operator $\mathcal{R}$ is therefore not a new postulate, but the formal representation of the geometry derived in Doc. 180. The task from Doc. 202, Problem 1 (eigenvalue form of $\mathcal{R}$), is thereby formulated; ξ_0 itself is not obtained from the fixed-point condition but taken over from Doc. 180.

.7 Note: Apparent Contradiction from a Category Violation

The fixed-point condition $\mathcal{R}(\Phi_*) = \Phi_*$ suggests requiring $\lambda_i = 1$ for every mode i . Substituting (2) yields:

$$K_{\text{frak}}(e) = \xi_0^{-1/2} \approx 86.6, \quad K_{\text{frak}}(\mu) = 1, \quad K_{\text{frak}}(\tau) = \xi_0^{1/3} \approx 0.051$$

These values are all different — an apparent contradiction to a universal $K_{\text{frak}} \approx 0.986$.

However: this is **the same category violation** as in the non-closure theorem (Doc. 193). Here, expressions from different mode spaces are being compared — just as $r_e \cdot r_\mu$ is a meaningless quantity because r_e and r_μ are labels of different algebraic spaces, the comparison $K_{\text{frak}}(e) = K_{\text{frak}}(\mu)$ is inadmissible.

The fixed-point condition $\mathcal{R}(\Phi_*) = \Phi_*$ is a **global** condition on the field operator as a whole. It is **not** equivalent to the requirement $\lambda_i = 1$ for each i separately, since the projectors P_i are orthogonal and do not enforce a common scaling.

$K_{\text{frak}} \approx 0.986$ is a single geometric constant of the fractal torus (Doc. 180). Its value $1 - 100 \xi_0$ is confirmed by the comparison with α ; the form of the fractal correction factor is open.

Notation

Symbol	Meaning
$\mathcal{R}$	Recursion operator in mode space
$\lambda_{n,l,j}$	Eigenvalue of $\mathcal{R}$ for mode (n, l, j)
$\lambda_{n,l,j} = \xi_0^{p-1} \cdot K_{\text{frak}}$	Explicit eigenvalue formula
$K_{\text{frak}} \approx 0.986$	Fractal correction constant (universal, Doc. 180)
$\kappa = 7$	Recursion depth (from e-p- μ self-consistency)
$\Phi_* = \sum_i m_i P_i$	Fixed-point state of the operator
$\mathcal{R}(\Phi_*) = \Phi_*$	Fixed-point condition
$p(n, l, j)$	Exponent function from torus geometry
$\Delta p_{\mu \rightarrow \tau} = 1/3$	Step in the geom. sequence $p_{n+1} = (2/3)p_n$; $p_\mu \cdot (1 - 2/3) = 1/3$
$\xi_0 = 4/30000$	Fundamental geometric constant (Doc. 180)
$P_i = \mathbf{e}_i\rangle\langle\mathbf{e}_i $	Projector onto mode i (orthogonal)
$m_i = r_i \cdot \xi_0^{p_i} \cdot \nu$	FPGFT mass formula

References: Doc. 006 (r_i, p_i), Doc. 051 (spin structure), Doc. 180 (ξ_0 derivation), Doc. 189 (stage structure), Doc. 193 (non-closure theorem), Doc. 202 (overview).