

Detailed Field-Geometric Spectral Theory

Full Derivation of Particle Spectra, Mass Hierarchies
and Gauge Groups from the $T^4/\mathbb{Z}_3$ Geometry

Contents

.1	Starting Point: the Sub-Planck Geometry	2
	Why a sub-Planck level?	2
	The D_4 lattice and its symmetry	2
.2	The Geometry Parameter ξ and its Origin	2
	Definition and numerical value	2
	Connection to Higgs geometry	3
	The D_4 projection coefficient η_{D4}	3
.3	Charged Leptons: Derivation from Winding Numbers	3
	Why winding numbers?	3
	The exponent ladder	3
	Step-by-step mass calculation	4
	Parameter-free mass ratios	4
.4	Neutrino Sector: Fixed-Point Localisation on the Orbifold	4
	Structural difference from charged leptons	5
	The neutrino mass formula	5
	Step-by-step neutrino mass calculation	5
	Mass-squared differences and oscillation dynamics	5
	Solar mass-squared difference Δm_{21}^2	5
	Atmospheric mass-squared difference Δm_{32}^2	5
	Cosmological mass sum	6
.5	Symmetry Breaking and Gauge Structure	6
	$SU(3)_c$: colour charges from Z_3 fibration	6
	Electroweak bosons ($W^\pm, Z^0$)	7
.6	Complete Overview of All Spectra	7
.7	Cross-references and Open Bridges	7

List of Symbols

Symbol	Meaning	Value / Unit
ξ	FFGFT geometry parameter	$4/30000 = 1.3\bar{3} \times 10^{-4}$
v	Higgs vacuum expectation value	246.0 GeV
m_e	electron mass (neutrino sector ref.)	0.511 MeV
r_i	topological winding ratio	n_ϕ^2/n_θ
p_i	exponent in mass formula	dimensionless
η_{D4}	D_4 projection coefficient	81/640
F_{D4}	D_4 volume element	27/1 600 000
[K]	derivable from ξ	status marker
[B]	algebraically proved	status marker
[S]	sketched / plausible	status marker

Summary

The FFGFT derives all particle masses of the leptonic sector from the single dimensionless geometry parameter $\xi = 4/30000$ and the topology of the $T^4/\mathbb{Z}_3$ orbifold. This document follows the derivation explicitly in every step: from the sub-Planck lattice, through the D_4 projection and topological winding numbers, to the measurable masses. The mathematical Hilbert-space foundation is provided in Doc. 322.

Status markers

All statements carry FFGFT status markers:

[K] derivable from ξ **[B]** algebraically proved **[S]** sketched/plausible **[SETZUNG]** axiomatic setting

.1 Starting Point: the Sub-Planck Geometry

Why a sub-Planck level?

The Standard Model describes particle masses through Yukawa couplings – free parameters measured experimentally but not explained within the framework itself. FFGFT goes one level deeper: masses do not originate from free couplings but from the geometry of space-time on scales below the Planck length ℓ_P .

[SETZUNG] At the sub-Planck scale t_0 space-time has a discrete packing structure. The most efficient regular lattice in four dimensions is the D_4 lattice – the 4D analogue of the densest sphere packing in 3D.

The D_4 lattice and its symmetry

Properties of the D_4 lattice:

- **4 orthogonal dimensions** with identical scaling.
- **Automorphism group** $\text{Aut}(D_4) \cong W(F_4)$ with $|W(F_4)| = 1152$ elements.
- **24 nearest neighbours** per lattice point.
- The torus $T^4/\mathbb{Z}_3$ is the natural compactification, with the $\mathbb{Z}_3$ identification reflecting the three quark colour charges.

[K] The fundamental FFGFT relation $\tilde{T} \cdot m = 1$ follows from the duality between torus radius $R \propto \tilde{T}$ and mass $m \propto 1/R$: on a compact torus the smallest mode spacing is inverse to the radius.

.2 The Geometry Parameter ξ and its Origin

Definition and numerical value

[K]

$$\xi = \frac{4}{30000} = \frac{1}{7500} = 1.3\bar{3} \times 10^{-4}. \quad (1)$$

Numerator 4 equals the number of orthogonal dimensions of the D_4 lattice – topologically fixed, not freely chosen.

Denominator 30000 equals the phase volume of the fractal T^4 compactification: product of the $\mathbb{Z}_3$ orbifold order (3) and the number of fundamental Fourier modes on the torus (10^4 , the fourth power of the first non-trivial mode).

Connection to Higgs geometry

The Higgs mass provides a consistency check of ξ (Doc. 354, A190, Doc. 385). With the Standard Model self-coupling $\lambda_h = m_h^2/(2v^2) = 0.129$ one has $\xi_{\text{EFT}} = \lambda_h^2 v^2 / (16\pi^3 m_h^2) = m_h^2 / (64\pi^3 v^2) \approx 1.30 \cdot 10^{-4}$, about 2.3 % below $4/30000$. The Higgs geometry is thus a consistency check of ξ to within a few per cent, not an exact derivation and not a second independent route to the same value. Inserting $\lambda_h = m_h/v$ does not hold: it would give $1/(16\pi^3) = 2.016 \cdot 10^{-3}$, 15 times $4/30000$, and an iteration $\xi_{n+1} = \xi_n(1 - 100\xi_n)$ does not lead from there to $4/30000$ but towards 0 (its only fixed point).

The D_4 projection coefficient η_{D_4}

The projection coefficient used is

$$\eta_{D_4} = \frac{81}{640} = 0.1265625; \quad (2)$$

[S] its origin is open: a counting of the Lie-algebra modes ($N_c^2 \cdot 9$ over $N_g \cdot 8 \cdot 8$ with $N_g = 80$) gives $81/5120$, one eighth of this value. The D_4 volume element:

$$F_{D_4} = \xi \cdot \eta_{D_4} = \frac{27}{1\,600\,000} = 1.6875 \times 10^{-5}. \quad (3)$$

Numerical check: $4 \cdot 81 = 324$, $30000 \cdot 640 = 19\,200\,000$, $324/19\,200\,000 = 27/1\,600\,000$. ✓

.3 Charged Leptons: Derivation from Winding Numbers

Why winding numbers?

Field modes on a compact torus T^2 are standing waves. The periodicity condition $\psi(x+2\pi R) = \psi(x)$ permits only integer winding numbers $n \in \mathbb{Z}$. On the T^2 torus two independent circles carry winding numbers n_θ and n_ϕ .

[K] The three generations of charged leptons correspond to the three topologically distinguished (n_θ, n_ϕ) states forming the simplest rational ratios $r_i = n_\phi^2/n_\theta$:

Table 1: FFGFT quantum numbers of the charged leptons

Particle	n_θ	n_ϕ	$r_i = n_\phi^2/n_\theta$	p_i
Electron e	3	2	$4/3$	$3/2$
Muon μ	5	4	$16/5$	1
Tau τ	9	5	$25/9$	$2/3$

The exponent ladder

[K] The exponents p_i form a geometric sequence with ratio $2/3$ per generation, $p_g = \frac{3}{2} \left(\frac{2}{3}\right)^{g-1}$:

$$p_e = \frac{3}{2}, \quad p_\mu = 1, \quad p_\tau = \frac{2}{3}. \quad (4)$$

The steps $3/2 \rightarrow 1 \rightarrow 2/3$ are $-1/2$ and $-1/3$; the constant quantity is the ratio $2/3$. The general mass formula:

$$m_i = r_i \cdot \xi^{P_i} \cdot v, \quad v = 246.0 \text{ GeV}. \quad (5)$$

Step-by-step mass calculation

Pre-computed ξ powers:

$$\xi^{3/2} = 1.5396 \times 10^{-6}, \quad \xi^1 = 1.33 \times 10^{-4}, \quad \xi^{2/3} = 2.6099 \times 10^{-3}. \quad (6)$$

[K]

$$m_e = \frac{4}{3} \cdot 1.5396 \times 10^{-6} \cdot 246.0 \times 10^3 \text{ MeV} = \mathbf{0.50499 \text{ MeV}} \quad (\text{PDG: } 0.511 \text{ MeV}, -1.18 \%), \quad (7)$$

$$m_\mu = \frac{16}{5} \cdot 1.33 \times 10^{-4} \cdot 246.0 \times 10^3 \text{ MeV} = \mathbf{104.96 \text{ MeV}} \quad (\text{PDG: } 105.66 \text{ MeV}), \quad (8)$$

$$m_\tau = \frac{25}{9} \cdot 2.6099 \times 10^{-3} \cdot 246.0 \times 10^3 \text{ MeV} = \mathbf{1783.4 \text{ MeV}} \quad (\text{PDG: } 1776.86 \text{ MeV}). \quad (9)$$

Parameter-free mass ratios

[K] The VEV v cancels in ratios:

$$m_\mu/m_e = \frac{12}{5} \cdot \xi^{-1/2} = 207.85 \quad (\text{PDG: } 206.77, +0.52 \%), \quad (10)$$

$$m_\tau/m_\mu = \frac{125}{144} \cdot \xi^{-1/3} = 16.99 \quad (\text{PDG: } 16.82, +1.04 \%), \quad (11)$$

$$m_\tau/m_e = \frac{25}{12} \cdot \xi^{-5/6} = 3532 \quad (\text{PDG: } 3477, +1.57 \%). \quad (12)$$

Table 2: Lepton masses and ratios: FFGFT vs. PDG

Quantity	Formula	Theory	PDG	Dev.	Status
m_e	$\frac{4}{3}\xi^{3/2}v$	0.50499 MeV	0.511 MeV	-1.18 %	[K]
m_μ	$\frac{16}{5}\xi^1v$	104.96 MeV	105.66 MeV	-0.66 %	[K]
m_τ	$\frac{25}{9}\xi^{2/3}v$	1783.4 MeV	1776.86 MeV	+0.37 %	[K]
m_μ/m_e	$\frac{12}{5}\xi^{-1/2}$	207.85	206.77	+0.52 %	[K]
m_τ/m_μ	$\frac{125}{144}\xi^{-1/3}$	16.99	16.82	+1.04 %	[K]

.4 Neutrino Sector: Fixed-Point Localisation on the Orbifold

This section derives neutrino masses from orbifold fixed points (F_2, F_5, F_7 , exponents $\{9/4, 2, 9/5\}$) and achieves Δm_{atm}^2 with a deviation of +16.0 % (with $K_{\text{frak}} + 12.8 \%$). Doc. 339/340 derives both mass differences from the $GF(27)^*$ Frobenius orbit structure and closes the atmospheric discrepancy at 1.0 % deviation [K]. The fixed-point derivation is a development step; the valid derivation is Doc. 340.

According to Doc. 343 (Theorem D'''), the quadratic character mod 13 (self-inverseness of the Frobenius orbit under $k \mapsto k^{-1}$) determines Majorana/Dirac for the Galois masses in Doc. 340: v_1 (orbit O_1) and v_2 (orbit O_3) are Majorana, v_3 (orbit O_4) is a **Dirac neutrino [B]**. The fixed-point assignment F_2, F_5, F_7 in this section provides no Majorana/Dirac character; this follows only from the Galois algebra of Doc. 340/343.

Structural difference from charged leptons

Charged leptons arise as winding modes on the T^2 torus and couple to the Higgs VEV v . Neutrinos are fundamentally different: they are localised at the *geometric fixed points* of the $T^4/\mathbb{Z}_3$ orbifold – points x_0 satisfying $\tau(x_0) = x_0$. This topological distinction produces a double ξ -suppression compared to the charged leptons. For the formal Hilbert-space treatment of the fixed-point boundary conditions see Doc. 322, Eq. (15).

The neutrino mass formula

[K]

$$m_{\nu_i} = m_e \cdot \xi^{p_i}, \quad m_e = 0.511 \text{ MeV}. \quad (13)$$

Table 3: Fixed-point localisation of neutrino flavours (Doc. 322)

Fixed pt.	Position	Neutrino	Exponent p_i	Mass
F_2	$\frac{2\pi R}{3}(1, 1, 1, 0)$	ν_1	$9/4$	0.976 meV
F_5	$\frac{4\pi R}{3}(1, 1, 1, 0)$	ν_2	2	9.084 meV
F_7	$\frac{4\pi R}{3}(1, 1, 1, 1)$	ν_3	$9/5$	54.11 meV

Step-by-step neutrino mass calculation

[K] Pre-computed ξ powers and masses:

$$\xi^{9/4} = 1.9103 \times 10^{-9}, \quad m_{\nu_1} = 511\,000 \text{ eV} \cdot 1.9103 \times 10^{-9} = \mathbf{0.976 \text{ meV}}, \quad (14)$$

$$\xi^2 = 1.7 \times 10^{-8}, \quad m_{\nu_2} = 511\,000 \text{ eV} \cdot 1.7 \times 10^{-8} = \mathbf{9.084 \text{ meV}}, \quad (15)$$

$$\xi^{9/5} = 1.059 \times 10^{-7}, \quad m_{\nu_3} = 511\,000 \text{ eV} \cdot 1.059 \times 10^{-7} = \mathbf{54.11 \text{ meV}}. \quad (16)$$

Mass-squared differences and oscillation dynamics

Solar mass-squared difference Δm_{21}^2

$$\Delta m_{21}^2 = (9.084 \text{ meV})^2 - (0.976 \text{ meV})^2 = 82.52 - 0.95 = 81.57 \text{ meV}^2 = \mathbf{8.16 \times 10^{-5} \text{ eV}^2}. \quad (17)$$

PDG 2023: $7.53 \times 10^{-5} \text{ eV}^2$. Deviation: +8.3 %.

Atmospheric mass-squared difference Δm_{32}^2

$$\Delta m_{32}^2 = (54.11 \text{ meV})^2 - (9.084 \text{ meV})^2 = 2927.9 - 82.5 = 2845.4 \text{ meV}^2 = \mathbf{2.845 \times 10^{-3} \text{ eV}^2}. \quad (18)$$

PDG (NuFIT 5.3): $2.453 \times 10^{-3} \text{ eV}^2$. Deviation: +16.0 %.

Open point: atmospheric mass-squared difference

Δm_{21}^2 agrees to +8.3 %, within the typical FFGFT range. Δm_{32}^2 deviates by +16.0 % – about twice as much and with the same sign: m_{ν_3} is too large. Equal sign and comparable magnitude point to a common cause rather than two independent problems.

The fixed-point exponents p_i are secured [K]; the source of the discrepancy is an open point. Three candidates were examined:

- **K_{frak} correction at fixed point F_7** (Doc. 315): F_7 is the only fixed point in the full (1, 1, 1, 1) direction and thus carries a mass-circle component. With $K_{\text{frak}} = 1 - 100\xi = 74/75$ one obtains $m_3 = 53.39$ meV and $\Delta m_{32}^2 = 2.768 \times 10^{-3} \text{ eV}^2$, i.e. +12.8 % – an improvement, but not sufficient.
- **Mixing term F_5 – F_7** : the fixed points are not fully orthogonal. An off-diagonal term in the mass matrix on the orbifold would shift both eigenvalues and would carry the right sign. This requires the complete mass matrix, not worked out here – the most promising candidate.
- **A different exponent** – unlikely: the exponent required for the PDG value is $p_3 = 1.8081$, only 0.45 % above $9/5 = 1.8000$. No rational fraction with small numerators lies closer ($16/9 \rightarrow +74\%$, $11/6 \rightarrow -37.5\%$, $20/11 \rightarrow -17.1\%$). The exponent $9/5$ is the best rational candidate and is grounded in the D_4 coordination.

The deviation therefore most likely does not stem from a wrong exponent but from an effect correcting $m_{\nu_i} = m_e \xi^{p_i}$ multiplicatively. Status: [S].

Cosmological mass sum

$$\sum_i m_{\nu_i} = 0.976 + 9.084 + 54.11 \text{ meV} = 64.17 \text{ meV} = 0.0642 \text{ eV}. \quad (19)$$

Planck 2018 limit: $\sum m_\nu < 0.12 \text{ eV}$. ✓

Table 4: Neutrino spectrum: FFGFT prediction vs. experiment

Quantity	Formula	Prediction	Experiment	Status
m_{ν_1}	$m_e \xi^{9/4}$	0.976 meV	$< 0.8 \text{ eV}$	[K]
m_{ν_2}	$m_e \xi^2$	9.084 meV	$\sim 8.7 \text{ meV}$	[K]
m_{ν_3}	$m_e \xi^{9/5}$	54.11 meV	$\sim 50 \text{ meV}$	[K]
Δm_{21}^2	$m_{\nu_2}^2 - m_{\nu_1}^2$	$8.16 \times 10^{-5} \text{ eV}^2$	$7.53 \times 10^{-5} \text{ eV}^2$	[K] +8.3 %
Δm_{32}^2	$m_{\nu_3}^2 - m_{\nu_2}^2$	$2.845 \times 10^{-3} \text{ eV}^2$	$2.453 \times 10^{-3} \text{ eV}^2$	[S] +16.0 %
$\sum m_\nu$	$\sum m_{\nu_i}$	0.0642 eV	$< 0.12 \text{ eV}$	[K]

.5 Symmetry Breaking and Gauge Structure

$SU(3)_c$: colour charges from $\mathbb{Z}_3$ fibration

[K] The $\mathbb{Z}_3$ symmetry generates three discrete complex-conjugate phase levels $\omega = e^{2\pi i/3}$, corresponding to the three colour charges Red, Green, Blue. The eight generators of $\mathfrak{su}(3)$ follow from the eight real degrees of freedom of the $\mathbb{Z}_3$ projection. **[B]** Full algebraic derivation of the Gell-Mann matrices from orbifold modes is in Doc. 321.

Electroweak bosons ($W^\pm, Z^0$)

[B]

$$m_W = \frac{1}{2} g v \approx 80.38 \text{ GeV}, \quad m_Z = m_W / \cos \theta_W \approx 91.19 \text{ GeV}. \quad (20)$$

The numerical values insert the measured g and θ_W and are inputs, not a prediction: with $g = 0.6517$, $v = 246 \text{ GeV}$ one gets $m_W = 80.16 \text{ GeV}$, 80.38 GeV requires $g = 0.6535$; $m_Z = m_W / \cos \theta_W$ with $\cos \theta_W = m_W / m_Z$ is circular.

.6 Complete Overview of All Spectra

Table 5: Systematic overview of all FFGFT spectra, Doc. 320

Group	Particle	Prediction	PDG	Status
Leptons	Electron e	0.50499 MeV	0.511 MeV	[K]
	Muon μ	104.96 MeV	105.66 MeV	[K]
	Tau τ	1783.4 MeV	1776.86 MeV	[K]
Neutrinos	ν_1 (F_2)	0.976 meV	$< 0.8 \text{ eV}$	[K]
	ν_2 (F_5)	9.084 meV	$\sim 8.7 \text{ meV}$	[K]
	ν_3 (F_7)	54.11 meV	$\sim 50 \text{ meV}$	[K]
Oscillations	Δm_{21}^2	$8.16 \times 10^{-5} \text{ eV}^2$	$7.53 \times 10^{-5} \text{ eV}^2$	[K] +8.3 %
	Δm_{32}^2	$2.845 \times 10^{-3} \text{ eV}^2$	$2.453 \times 10^{-3} \text{ eV}^2$	[S] +16.0 %
	$\sum m_\nu$	0.0642 eV	$< 0.12 \text{ eV}$	[K]
Bosons	γ, g	0	0	[K]
	$W^\pm$	80.38 GeV (input)	80.38 GeV	input
	Z^0	91.19 GeV (input)	91.19 GeV	input

The $W^\pm$ and Z^0 values insert the measured g and θ_W and are inputs; θ_W is not derived from ξ in this document (RG running from ξ : Doc. 323). For the neutrino rows the derivation in Doc. 340 is authoritative (see the neutrino-sector section).

.7 Cross-references and Open Bridges

Table 6: Derivation chain Doc. 320 and cross-references

Step	Doc.	Status
$\xi = 4/30000$ from D_4 lattice	Docs. 001, 205	[K]
Higgs consistency check of ξ ($\approx 2.3\%$)	Doc. 354, A190	[K]
K_{frac} interface factor	Docs. 012, 133	[K]
$\tilde{T} \cdot m = 1$	Doc. 003	[SETZUNG]
Winding numbers (n_θ, n_ϕ)	Doc. 317	[K]
Lepton masses m_e, m_μ, m_τ	Docs. 006, 317	[K]
Hilbert-space embedding, operator $\hat{F}$	Doc. 322	[K/S]
Fixed-point positions F_2, F_5, F_7	Doc. 322	[K]
Δm_{21}^2 (+8 %)	this Doc.	[K]
Δm_{32}^2 (+16.0 %)	this Doc.	[S] open
$SU(3)_c$ from $\mathbb{Z}_3$	Doc. 321	[B]
Weinberg angle θ_W from ξ	Doc. 323	[K]
Quark sector (hadrons)	Docs. 318, 319	open bridge