

Frobenius Separation of Massive and Massless Sectors in $GF(27)^*$

Algebraic Structure of the $\tilde{T} \cdot m = 1$ Transition
and Connection to 8 Gluons

Johann Pascher
ORCID 0009-0000-6518-4064

29 August 2026

Contents

Frobenius Separation of Massive and Massless Sectors	2
.1 Starting Point	1
.2 Decomposition of $\mathbb{Z}_{26}$ under the Frobenius	1
Basic structure [B]	1
Orbits of ϕ on $\mathbb{Z}_{13}$ [B]	1
Complete orbit table in $\mathbb{Z}_{26}$ [B]	1
.3 Physical Identification [K]	2
Massive sector: fixed points $= \mathbb{Z}_2$	2
Massless $SU(3)$ sector: eight gluons	2
$U(1)$ sector: photon from the fixed field $GF(3)$	2
.4 Algebraic Necessity [B]	2
.5 Complete Picture	2
.6 Scales of the Duality and Effective Masslessness [K]	3
The two poles of $\tilde{T} \cdot m = 1$	3
What "massless" means in FFGFT	3
Conclusion	3
.7 Spin Assignment from the Frobenius Structure [K]	3
Derivation	3
Compatibility with the FFGFT corpus	4
Result	4
.8 Context: Connection to HLV and the Hilbert Space Doubling [K]	4
The fork from Doc. 307	4
FFGFT, HLV and the doubling $3 \rightarrow 6$	4
What the Frobenius decomposition contributes to the bridge [K]	5
.9 Extension: Two Pictures, One Structure [K]	5
Unrolled picture and conformal symmetry	5
Accessible calculations per picture	6
.10 Note on Authorship	6
.11 Notation	6

Frobenius Separation of Massive and Massless Sectors in $GF(27)^*$

Abstract

The multiplicative group $GF(27)^* \cong \mathbb{Z}_{26}$ decomposes under the Frobenius automorphism $\phi : x \mapsto x^3$ into exactly two types: two fixed points $\{+1, -1\}$ and eight three-element orbits. Via the decomposition $\mathbb{Z}_{26} \cong \mathbb{Z}_2 \times \mathbb{Z}_{13}$ (CRT), four $\mathbb{Z}_{13}$ three-orbits times two $\mathbb{Z}_2$ parities yield the eight gluons of the adjoint $SU(3)$ representation; the fixed field $GF(3)$ of the Frobenius supplies the $U(1)$ sector (photon). The fixed points $\{+1, -1\}$ correspond to the massive $\mathbb{Z}_2$ sector (particle/antiparticle). All results are algebraically forced. Status: **[K]**.

.1 Starting Point

In Doc. 338

$$\left(\frac{m_\mu}{m_e}\right)^2 = 43200 = |GF(9)^*|^2 \cdot 5^2 \cdot |GF(27)| = 64 \cdot 25 \cdot 27$$

is established as an algebraically exact relation **[B]**. The fundamental relation $\tilde{T} \cdot m = 1$ describes a duality: massive sector ($m > 0$, compact torus S_m^1) and massless sector ($m \rightarrow 0$, $\tilde{T} \rightarrow \infty$). In Doc. 285 the $3'$ extension is identified as a kinetic (massless) window. This note asks: what algebraic structure separates both sectors within $GF(3) \subset GF(9) \subset GF(27)$?

.2 Decomposition of $\mathbb{Z}_{26}$ under the Frobenius

Basic structure **[B]**

$GF(27)^* \cong \mathbb{Z}_{26}$, cyclic of order $3^3 - 1 = 26 = 2 \cdot 13$. Since $\gcd(2, 13) = 1$:

$$\mathbb{Z}_{26} \cong \mathbb{Z}_2 \times \mathbb{Z}_{13}.$$

The Frobenius $\phi : x \mapsto x^3$ acts:

- On $\mathbb{Z}_2$: $3 \equiv 1 \pmod{2}$ — ϕ is *trivial*.
- On $\mathbb{Z}_{13}$: $\text{ord}(3, \mathbb{Z}_{13}) = 3$ (since $3^3 = 27 \equiv 1 \pmod{13}$) — ϕ has order 3.

Orbits of ϕ on $\mathbb{Z}_{13}$ **[B]**

Type	Elements
Fixed point	$\{0\}$
Three-orbit 1	$\{1, 3, 9\}$
Three-orbit 2	$\{2, 5, 6\}$
Three-orbit 3	$\{4, 10, 12\}$
Three-orbit 4	$\{7, 8, 11\}$

Complete orbit table in $\mathbb{Z}_{26}$ **[B]**

$\mathbb{Z}_2$ parity	$\mathbb{Z}_{13}$ orbit	$\mathbb{Z}_{26}$ orbit
0	$\{0\}$	$\{0\}$ (fixed point)
1	$\{0\}$	$\{13\}$ (fixed point)
0	$\{1, 3, 9\}$	$\{14, 16, 22\}$
1	$\{1, 3, 9\}$	$\{1, 3, 9\}$
0	$\{2, 5, 6\}$	$\{2, 6, 18\}$
1	$\{2, 5, 6\}$	$\{5, 15, 19\}$
0	$\{4, 10, 12\}$	$\{4, 10, 12\}$
1	$\{4, 10, 12\}$	$\{17, 23, 25\}$
0	$\{7, 8, 11\}$	$\{8, 20, 24\}$
1	$\{7, 8, 11\}$	$\{7, 11, 21\}$

Check: $2 + 8 \times 3 = 26 = |GF(27)^*|$. ✓

.3 Physical Identification [K]

Massive sector: fixed points $= \mathbb{Z}_2$

The two fixed points $\{+1, -1\}$ form the massive sector: every massive particle has an antiparticle ($\mathbb{Z}_2$ structure), consistent with the compact torus sector.

Massless $SU(3)$ sector: eight gluons

$$8 \text{ gluons} = 4 \mathbb{Z}_{13}\text{-orbits} \times 2 \mathbb{Z}_2\text{-parities} = |GF(9)^*|.$$

The three elements of each $\mathbb{Z}_{13}$ orbit correspond to the three colours $\{R, G, B\}$; the $\mathbb{Z}_2$ parity distinguishes colour from anticolour. This is the adjoint representation of $SU(3)$, derived from Galois structure alone.

$U(1)$ sector: photon from the fixed field $GF(3)$

The fixed field of the Frobenius on $GF(27)$ is $GF(3)$ (Fermat: $x^3 = x$ for $x \in GF(3)$). The projection $GF(27) \rightarrow GF(3)$ gives:

Structure	Sector
$GF(3)^* = \{1, 2\} \cong \mathbb{Z}_2$	Photon: $U(1)$, own antiparticle
$27 - 3 = 24$ remaining elements	$SU(3)$ sector (quarks, gluons)

.4 Algebraic Necessity [B]

$$\text{Non-fixed elements in } GF(27)^* = 26 - 2 = 24, \quad (1)$$

$$24 = 8 \cdot 3 = |GF(9)^*| \cdot |\text{colours}| = (3^2 - 1) \cdot 3 = 3^3 - 3, \quad (2)$$

$$8 \text{ gluons} = \frac{26-2}{3} = 8. \quad (3)$$

All three relations follow solely from $|GF(27)| = 3^3$ and the order 3 of the Frobenius on $\mathbb{Z}_{13}$.

.5 Complete Picture

Galois structure	Physical sector	Symmetry
$GF(3)^* = \mathbb{Z}_2$	Photon	$U(1)$
Fixed points $\{+1, -1\} \subset GF(27)^*$	Massive particle/antiparticle	$\mathbb{Z}_2$
4×2 three-orbits in $GF(27)^*$	8 gluons	$SU(3)_{\text{adj}}$

The relation $8 \times 3 = 24 = \text{Kissing}(D_4)$ is algebraically forced ($3^3 - 3 = 24$). Whether $\text{Kissing}(D_4)$ has a deeper geometric connection to this Galois structure is subject to further investigation.

Verification script: `pruef_339_frobenius.py` (6/6 assertions).

.6 Scales of the Duality and Effective Masslessness [K]

The two poles of $\tilde{T} \cdot m = 1$

The fundamental relation $\tilde{T} \cdot m = 1$ has two excluded edges (Doc. 306), both finite and quantified:

Pole	Length	Time	Energy per cell
Dense (derived)	$L_0 = \xi \ell_P$	$T_0 = \xi t_P$	$E_{\max} = E_P/\xi \approx 9.2 \times 10^{22} \text{ GeV}$
Thin (calibrated)	$R_H = \hbar/E_H$	$T_H = \hbar/E_H$	$E_H = E_0 \xi^{41/4} \approx 1.4 \times 10^{-33} \text{ eV}$

At both ends the reciprocity closes exactly: $T_0 \cdot E_{\max} = t_P \cdot E_P = 1$ and $T_H \cdot E_H = 1$ (natural units). There is no infinity — neither above nor below. The dense pole is derived from ξ [B]; the thin pole is calibrated to the measured H_0 (Doc. 182, P35/P36) [S].

What “massless” means in FFGFT

$\tilde{T} \cdot m = 1$ has no solution at $m = 0$ — FFGFT knows no empty state, only a thinnest one (Doc. 306). The minimal rest mass per cell is:

$$m_0 = \frac{E_H}{c^2} \approx 1.4 \times 10^{-33} \text{ eV}/c^2.$$

Gluons and photons are modes in the limit $m \rightarrow m_0$, not at $m = 0$. The time circle S_m^1 opens to the time axis $\mathbb{R}_t$ in this limit (type-III pullback, Doc. 332, R96).

The deviation of the dispersion relation from the massless formula $\omega = |\mathbf{k}|$ is:

$$\frac{\delta\omega}{\omega} = \frac{m_0^2}{2k^2} \approx \begin{cases} 2.5 \times 10^{-67} & k = 2 \text{ eV (visible light)} \\ 1.7 \times 10^{-59} & k = 2.4 \times 10^{-4} \text{ eV (CMB)} \\ 9.8 \times 10^{-85} & k = 1 \text{ GeV (QCD)} \end{cases}$$

The experimental upper bound on the photon rest mass is $m_\gamma < 10^{-18} \text{ eV}/c^2$ (PDG). The FFGFT minimum mass m_0 lies 10^{15} times below this — beyond any conceivable measurement.

Conclusion

“Massless” in FFGFT is not an algebraic exactness ($m \neq 0$) but a structural statement: the rest mass lies below the Hubble energy scale E_H/c^2 and is indistinguishable from zero for all physical purposes. Both poles of the duality are finite and quantified — there is no infinity, but both limits lie far beyond any experimental reach.

.7 Spin Assignment from the Frobenius Structure [K]

Derivation

The derivation proceeds in four steps.

Step 1 [K]: The Frobenius map $\phi : x \mapsto x^3$ on $GF(27)^*$ corresponds in the type-III pullback picture (Doc. 332, R96) to the conformal scaling of the wave vector:

$$k \mapsto 3k, \quad \omega(3k) = 3\omega(k).$$

This follows directly from the massless dispersion relation $\omega = c|k|$. Massive modes do not satisfy this relation: for $m > 0$, $\omega^2 = c^2 k^2 + (mc^2/\hbar)^2$, so $\omega(3k) \neq 3\omega(k)$. Fixed points transform trivially: $\phi(x) = x$ corresponds to $k \mapsto k$.

Step 2 [B, standard result]: In the representation theory of the Poincaré group (Wigner 1939), massless fields are classified by their behaviour under scalings $k \mapsto \lambda k$:

- *Spin-1 fields* (vector bosons A_μ): scale linearly, $A_\mu \rightarrow \lambda A_\mu$, i.e. $\omega(\lambda k) = \lambda \omega(k)$.
- *Massive fields* (rest mass): invariant in the non-relativistic limit ($k \rightarrow 0$, $\omega \rightarrow mc^2/\hbar$).

Step 3 [K]: The 8 three-orbits of $GF(27)^*$ carry the conformal scaling $k \mapsto 3k$ and therefore correspond to massless spin-1 fields. The 2 fixed points $\{+1, -1\}$ transform trivially and correspond to the massive sector.

Step 4 [K]: The fixed field $GF(3)^* = \{1, 2\} \cong \mathbb{Z}_2$ is conformally invariant (fixed under the projection $GF(27) \rightarrow GF(3)$) and carries a single spin-1 singlet — the photon.

Compatibility with the FFGFT corpus

In FFGFT, spin arises from winding numbers (n_θ, n_ϕ) on the spin bundle $L^2(T^3, S)$ (Doc. 231), not from an external Poincaré representation. The Wigner classification in Step 2 is therefore a standard result [B] that is consistent with the FFGFT spin-bundle framework but does not replace it. The explicit connection — that the three helicity states of a spin-1 field on T^3 under $\mathbb{Z}_3$ reproduce exactly the three-orbits of $GF(27)^*$ — is structurally plausible and remains as an open bridge to Doc. 231.

Result

Frobenius type	Sector	Spin	Physical object
Fixed points $\{+1, -1\}$	Massive, $k \mapsto k$	–	Leptons, quarks
8 three-orbits	Massless, $k \mapsto 3k$	1	8 gluons
Fixed field $GF(3)^*$	Massless, $k \mapsto k$	1	Photon (singlet)

Note: The spin of the massive fixed points (leptons: $\frac{1}{2}$, quarks: $\frac{1}{2}$) does not follow from the Frobenius structure but from the winding numbers on $L^2(T^3, S)$ (Doc. 231).

.8 Context: Connection to HLV and the Hilbert Space Doubling [K]

The fork from Doc. 307

Doc. 307 poses the fundamental question: is time a parameter *beside* the state space (unrolled picture) or a compact dimension *inside* it (rolled-up picture)? Both descriptions use the same operator formalism; the difference lies in where time is located.

The rolled-up picture ($\tilde{T} \cdot m = 1$, $L^2(T^4) \otimes \mathbb{C}^3$) is the massive side: discrete modes, winding numbers (n_θ, n_ϕ) , lepton masses from $\xi = 4/30000$. The unrolled picture ($L^2(\mathbb{R}_t)$) is the massless side — and exactly there, as the Frobenius decomposition of $GF(27)^*$ shows algebraically, the gluons live.

FFGFT, HLV and the doubling $3 \rightarrow 6$

In Doc. 285 (FFGFT-HLV dimensional bridge) the doubling $\mathbf{6} = \mathbf{3} \oplus \mathbf{3}'$ is specified from the Hilbert space perspective:

- **FFGFT sector (3, compact torus S_m^1):** $\mathbb{Z}_3$ -crystallographic, 3-fold symmetry, lives natively in 3D/4D. Protected invariant: $\theta = 2/9$ (Koide), $r = \sqrt{2}$ ($Q = 2/3$). This is the massive sector.
- **HLV sector (3', type-III pullback to $\mathbb{R}_t$):** A_5 -icosahedral, 5-fold symmetry, requires 6D embedding. The unrolling referred to here is the lossless mathematical type (isometry $L^2(S_m^1) \cong \text{Per}_{R_m}(\mathbb{R}_t)$, Doc. 332 R96, type III), *not* the measurement act (type I, lossy, Doc. 333). Protected invariant: φ (golden ratio), $r = 2/\sqrt{5}$. This is the massless candidate.

The 6D splitting is forced by the crystallographic restriction: 5-fold symmetry is forbidden in 3D ($2 \cos 72^\circ = 1/\varphi \notin \mathbb{Z}$) but exists in 6D. The **3'** sector is geometrically present in FFGFT but physically empty without dynamics.

What the Frobenius decomposition contributes to the bridge [K]

The Frobenius decomposition of $GF(27)^*$ provides a new algebraic precision that Doc. 285 did not yet have:

Structure	Hilbert space picture	Galois picture
Massive sector (3)	$L^2(S_m^1)$, rolled-up	Fixed points $\{+1, -1\}$ of ϕ
Massless sector (3')	$L^2(\mathbb{R}_t)$, type-III pullback (R96)	8 three-orbits of ϕ
Koide $\theta = 2/9$	$\mathbb{Z}_3$ circulant	$GF(9)^*$ structure (Doc. 338)
Golden ratio φ	A_5 icosahedron	not anchored in $GF(27)^*$

The golden ratio φ does not appear in the $\mathbb{Z}_{13}$ picture: $\mathbb{Z}_{13}$ carries no 5-fold symmetry (since $5 \nmid 12$). This is consistent with φ being the HLV invariant, not the FFGFT invariant.

The Frobenius structure of $GF(27)^*$ confirms algebraically what Doc. 285 shows geometrically: FFGFT is the $\mathbb{Z}_3$ -crystallographic core (massive fixed points), HLV would be the A_5 sector (massless **3'** orbits).

.9 Extension: Two Pictures, One Structure [K]

Unrolled picture and conformal symmetry

The Frobenius map $\phi : x \mapsto x^3$ on $GF(27)^*$ corresponds in the unrolled picture ($\mathbb{R}_t$ instead of S_m^1) to a conformal scaling of the wave vector:

$$k \mapsto 3k, \quad \omega \mapsto 3\omega.$$

This is consistent with the massless dispersion relation $\omega = c|k|$:

$$\omega(3k) = c \cdot |3k| = 3 \cdot c \cdot |k| = 3\omega(k). \quad \checkmark$$

For massive modes, $\omega^2 = c^2 k^2 + (mc^2/\hbar)^2$ gives $\omega(3k) \neq 3\omega(k)$ — massive modes are *not* conformally invariant. The Frobenius structure thus explains the separation algebraically:

Frobenius type	Physical sector	Symmetry
Fixed points $\{+1, -1\}$	Massive, $\omega(3k) \neq 3\omega(k)$	no conformal inv.
Three-orbits (8 total)	Massless, $\omega(3k) = 3\omega(k)$	conformal inv.

Accessible calculations per picture

Content	Picture	Status
Lepton masses, winding numbers (n_θ, n_ϕ)	Compact	[B]
Galois identity $43200 = 64 \cdot 25 \cdot 27$	Compact	[B]
$K_{\text{frak}}^{-36} \approx 16/\pi^2$	Compact	[B]
Gluon dispersion $\omega = c k $	Unrolled	[K]
Frobenius = conformal scaling $k \mapsto 3k$	Unrolled	[K]
Hilbert space isometry $L^2(S_m^1) \cong \text{Per}_{R_m}(\mathbb{R}_t)$	Both	[B]
$\xi = 4/30000$, mass ratios	Both	[B]

The conformal dimension of gluons ($\Delta = 1$, spin-1 vector bosons) remains to be derived from the FFGFT field theory — noted as an open point.

Declared 29 August 2026

.10 Note on Authorship

This document was developed on 29 August 2026 in collaboration with the AI assistant Claude (Anthropic, Sonnet 4.6). The algebraic computations (Frobenius orbits, $\mathbb{Z}_2 \times \mathbb{Z}_{13}$ decomposition, conformal symmetry test) were carried out by the assistant and verified by the author. The physical interpretation and all status markers (**[B]**, **[K]**) are the author's responsibility.

.11 Notation

[ASSUMPTION]	Axiom or declared external input – not derived
[K]	Core – derived from ξ and the assumptions, numerically verified
[B]	Bridge – algebraically proved
[S]	Sketch – plausible, not yet fully worked out