

T0 Deterministic Quantum Computing: Complete Analysis of Important Algorithms

Contents

.1	Introduction: T0 Quantum Computing	1
	Motivation and Scope	1
	Analysis Structure	2
.2	Algorithm 1: Deutsch Algorithm	2
	Problem Statement	2
	Standard Quantum Mechanics Implementation	2
	Algorithm Steps	2
	Mathematical Analysis	3
	T0 Energy Field Implementation	3
	T0 Gate Operations with Updated ξ	3
	Mathematical Analysis with Updated ξ	3
	Result Comparison	4
	T0-Specific Predictions with Updated ξ	4
.3	Algorithm 2: Bell State Generation	4
	Standard QM Bell States	4
	T0 Energy Field Bell States with Updated ξ	5
.4	Algorithm 3: Grover Search	5
	T0 Energy Field Grover with Updated ξ	5
.5	Algorithm 4: Shor Factorization	5
	T0 Energy Field Shor with Updated ξ	5
	T0 Quantum Fourier Transform with ξ Corrections	6
	T0-Specific Corrections with Updated ξ	6
.6	Experimental Distinction with Updated ξ	6
	Universal Distinction Tests	6
	Repeatability Test	6
	ξ -Parameter Precision Tests with Updated Value	6
	Step-by-Step Calculation	6
	Ideal ξ Parameter from Measurement Error Analysis	7
	NOT Gate Error Analysis	7
	Comparison with Higgs Calculation	7
	Information Structure in Energy Field Amplitudes	7
	Experimental Roadmap	8
	Phase I - Precision Validation	8
	Phase II - Information Layer Access	8
	Phase III - Higgs Coupling Detection	8

Abstract

This comprehensive document presents a complete analysis of important quantum computing algorithms within the T0 energy field formulation. We systematically examine four fundamental quantum algorithms: Deutsch, Bell states, Grover, and Shor, demonstrating

that the T0 approach reproduces all standard quantum mechanical results while offering fundamentally different physical interpretations. The T0 formulation replaces probabilistic amplitudes with deterministic energy field configurations, leading to single-measurement predictability and novel experimental signatures. **This updated version uses the numerical simulation parameter $\xi_{\text{num}} = 1.0 \times 10^{-5}$ (physical: $\xi = 4/30000$, Dok. 009) and shows that energy field amplitude deviations are information carriers rather than computational errors.** Our analysis demonstrates that deterministic quantum computing is not only theoretically possible but offers practical advantages including perfect repeatability, spatial energy field structure, and systematic ξ -parameter corrections measurable at the ppm level.

.1 Introduction: T0 Quantum Computing

Motivation and Scope

Standard quantum mechanics has achieved experimental successes, yet its probabilistic foundation creates fundamental interpretational problems. The measurement problem, wavefunction collapse, and the quantum-classical boundary remain unresolved after nearly a century of development.

The T0 theoretical framework offers a radical alternative: deterministic quantum mechanics based on energy field dynamics. This work presents the first comprehensive analysis of how important quantum computing algorithms function within the T0 formulation.

Core T0 Principles with Updated ξ Parameter

Fundamental T0 Relations:

$$T(x, t) \cdot m(x, t) = 1 \quad (\text{time-mass duality}) \quad (1)$$

$$\partial^2 E(x, t) = 0 \quad (\text{universal field equation}) \quad (2)$$

$$\xi_{\text{num}} = 1.0 \times 10^{-5} \quad (\text{simulation parameter;}) \quad (3)$$

$$\text{physical: } \xi = 4/30000, \text{ Dok. 009}) \quad (4)$$

Quantum State Representation:

$$\text{Standard QM: } |\psi\rangle = \sum_i c_i |i\rangle \quad \rightarrow \quad \text{T0: } \{E(x, t)_i(x, t)\} \quad (5)$$

Updated ξ -Parameter Justification: Note on ξ values: The physical T0 parameter is $\xi_{\text{phys}} = 4/30000 \approx 1.333 \times 10^{-4}$ (Doc. 009). The Higgs matching (consistency check) via the Higgs sector (Doc. 097) gives $\xi \approx \lambda_h^2 v^2 / (16\pi^3 m_h^2) \approx 1.30 \times 10^{-4}$ (2.2% deviation with the inputs of the appendix). This is a consistency check, not a derivation (Doc. 385). This document uses the **numerical simulation parameter** $\xi_{\text{num}} = 1.0 \times 10^{-5}$, which optimizes numerical stability in the algorithm number space. All concrete values below (gate amplitudes, ppm shifts) are calculated with ξ_{num} .

Analysis Structure

We examine four quantum algorithms of increasing complexity:

1. **Deutsch Algorithm:** Single-qubit oracle problem (deterministic result)
2. **Bell States:** Two-qubit entanglement generation (correlation without superposition)
3. **Grover Algorithm:** Database search (deterministic amplification)

4. **Shor Algorithm:** Integer factorization (deterministic period finding)

For each algorithm we provide:

- Complete mathematical analysis in both formulations
- Algorithmic result comparisons
- Physical interpretation differences
- T0-specific predictions and experimental tests

.2 Algorithm 1: Deutsch Algorithm

Problem Statement

The Deutsch algorithm determines whether a black-box function $f : \{0, 1\} \rightarrow \{0, 1\}$ is constant or balanced, using only one function evaluation.

Classical Complexity: 2 evaluations required

Quantum Advantage: 1 evaluation sufficient

Standard Quantum Mechanics Implementation

Algorithm Steps

1. Initialization: $|\psi_0\rangle = |0\rangle$
2. Hadamard: $|\psi_1\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$
3. Oracle: $|\psi_2\rangle = U_f|\psi_1\rangle$ where $U_f|x\rangle = (-1)^{f(x)}|x\rangle$
4. Hadamard: $|\psi_3\rangle = H|\psi_2\rangle$
5. Measurement: $0 \rightarrow \text{constant}, 1 \rightarrow \text{balanced}$

Mathematical Analysis

Constant function ($f(0) = f(1) = 0$):

$$|\psi_0\rangle = |0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (6)$$

$$|\psi_1\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (7)$$

$$|\psi_2\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad (\text{no phase change}) \quad (8)$$

$$|\psi_3\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \rightarrow P(0) = 1.0 \quad (9)$$

Balanced function ($f(0) = 0, f(1) = 1$):

$$|\psi_2\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix} \quad (\text{phase flip at } |1\rangle) \quad (10)$$

$$|\psi_3\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \rightarrow P(1) = 1.0 \quad (11)$$

T0 Energy Field Implementation

T0 Gate Operations with Updated ξ

T0 Qubit State: $\{E(x, t)_0(x, t), E(x, t)_1(x, t)\}$
T0 Hadamard Gate with $\xi = 1.0 \times 10^{-5}$:

$$H_{T0} : \begin{cases} E(x, t)_0 \rightarrow \frac{E(x, t)_0 + E(x, t)_1}{2} \times (1 + \xi) \\ E(x, t)_1 \rightarrow \frac{E(x, t)_0 - E(x, t)_1}{2} \times (1 + \xi) \end{cases} \quad (12)$$

T0 Oracle Operation:

$$U_f^{T0} : \begin{cases} \text{Constant} : & E(x, t)_0 \rightarrow +E(x, t)_0, & E(x, t)_1 \rightarrow +E(x, t)_1 \\ \text{Balanced} : & E(x, t)_0 \rightarrow +E(x, t)_0, & E(x, t)_1 \rightarrow -E(x, t)_1 \end{cases} \quad (13)$$

Mathematical Analysis with Updated ξ

Constant function:

$$\text{Start} : \{E(x, t)_0, E(x, t)_1\} = \{1.000000, 0.000000\} \quad (14)$$

$$\text{After } H_{T0} : \{E(x, t)_0, E(x, t)_1\} = \{0.500005, 0.500005\} \quad (15)$$

$$\text{After Oracle} : \{E(x, t)_0, E(x, t)_1\} = \{0.500005, 0.500005\} \quad (16)$$

$$\text{After } H_{T0} : \{E(x, t)_0, E(x, t)_1\} = \{0.500010, 0.000000\} \quad (17)$$

T0 Measurement: $|E(x, t)_0| > |E(x, t)_1| \rightarrow \text{Result: 0 (constant)}$

Balanced function:

$$\text{After Oracle} : \{E(x, t)_0, E(x, t)_1\} = \{0.500005, -0.500005\} \quad (18)$$

$$\text{After } H_{T0} : \{E(x, t)_0, E(x, t)_1\} = \{0.000000, 0.500010\} \quad (19)$$

T0 Measurement: $|E(x, t)_1| > |E(x, t)_0| \rightarrow \text{Result: 1 (balanced)}$

Result Comparison

Function Type	Standard QM	T0 Approach	Agreement
Constant	0	0	✓
Balanced	1	1	✓

Table 1: Deutsch Algorithm: Perfect Result Agreement with Updated ξ

T0-Specific Predictions with Updated ξ

- Deterministic Repeatability:** Identical results for identical conditions
- Spatial Energy Structure:** $E(x, t)(x, t)$ has measurable spatial extent with characteristic scale $\sim \lambda\sqrt{1 + \xi}$
- Minimal Measurement Errors:** Gate operations deviate only by $\xi \times 100\% = 0.001\%$ from ideal values
- Information Enhancement:** 51× more physical information per qubit compared to standard QM

.3 Algorithm 2: Bell State Generation

Standard QM Bell States

Generation Protocol:

1. Initialization: $|00\rangle$
2. Hadamard on qubit 1: $\frac{1}{\sqrt{2}}(|00\rangle + |10\rangle)$
3. CNOT(1 $\rightarrow$ 2): $\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ (Bell state)

Mathematical Calculation:

$$|00\rangle \rightarrow \frac{1}{\sqrt{2}}(|00\rangle + |10\rangle) \quad (20)$$

$$\rightarrow \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) \quad (21)$$

Correlation Properties:

- $P(00) = P(11) = 0.5$
- $P(01) = P(10) = 0.0$
- Perfect correlation: Measurement of one qubit determines the other

T0 Energy Field Bell States with Updated ξ

T0 Two-Qubit State: $\{E(x, t)_{00}, E(x, t)_{01}, E(x, t)_{10}, E(x, t)_{11}\}$

T0 Hadamard on Qubit 1 with $\xi = 1.0 \times 10^{-5}$:

$$E(x, t)_{00} \rightarrow \frac{E(x, t)_{00} + E(x, t)_{10}}{2} \times (1 + \xi) \quad (22)$$

$$E(x, t)_{10} \rightarrow \frac{E(x, t)_{00} - E(x, t)_{10}}{2} \times (1 + \xi) \quad (23)$$

$$E(x, t)_{01} \rightarrow \frac{E(x, t)_{01} + E(x, t)_{11}}{2} \times (1 + \xi) \quad (24)$$

$$E(x, t)_{11} \rightarrow \frac{E(x, t)_{01} - E(x, t)_{11}}{2} \times (1 + \xi) \quad (25)$$

T0 CNOT Gate: Energy transfer from $|10\rangle$ to $|11\rangle$

$$\text{T0-CNOT : } E(x, t)_{10} \rightarrow 0, \quad E(x, t)_{11} \rightarrow E(x, t)_{11} + E(x, t)_{10} \times (1 + \xi) \quad (26)$$

Mathematical Calculation with Updated ξ :

$$\text{Start : } \{1.000000, 0.000000, 0.000000, 0.000000\} \quad (27)$$

$$\text{After H : } \{0.500005, 0.000000, 0.500005, 0.000000\} \quad (28)$$

$$\text{After CNOT : } \{0.500005, 0.000000, 0.000000, 0.500010\} \quad (29)$$

T0 Correlations with Minimal Errors:

$$P(00) = 0.499995 \approx 0.5 \quad (\text{Error: } 0.001\%) \quad (30)$$

$$P(11) = 0.500005 \approx 0.5 \quad (\text{Error: } 0.001\%) \quad (31)$$

$$P(01) = P(10) = 0.000000 \quad (\text{exact}) \quad (32)$$

.4 Algorithm 3: Grover Search

T0 Energy Field Grover with Updated ξ

T0 Concept: Deterministic energy field focusing instead of probabilistic amplification

T0 Operations with $\xi = 1.0 \times 10^{-5}$:

1. Uniform energy distribution: $\{0.25, 0.25, 0.25, 0.25\}$
2. T0 Oracle: Energy inversion for marked element with ξ -correction
3. T0 Diffusion: Energy rebalancing toward inverted element

Mathematical Calculation with Updated ξ :

$$\text{Start : } \{0.250000, 0.250000, 0.250000, 0.250000\} \quad (33)$$

$$\text{After T0 Oracle : } \{0.250000, 0.250000, 0.250000, -0.250003\} \quad (34)$$

$$\text{After T0 Diffusion : } \{-0.000001, -0.000001, -0.000001, 0.500004\} \quad (35)$$

T0 Measurement: $|E(x, t)_{11}| = 0.500004$ is maximum $\rightarrow$ Result: $|11\rangle$

Search Accuracy: 99.999% (error significantly less than 0.001%)

.5 Algorithm 4: Shor Factorization

T0 Energy Field Shor with Updated ξ

T0 Approach: Period finding through energy field resonance with minimal systematic errors

T0 Quantum Fourier Transform with ξ Corrections

T0 Resonance Transformation: $E(x, t)(x, t) \rightarrow E(x, t)(\omega, t)$ via resonance analysis

$$\frac{\partial^2 E(x, t)}{\partial t^2} = -\omega^2 E(x, t) \quad \text{with } \omega = \frac{2\pi k}{N} \times (1 + \xi) \quad (36)$$

T0-Specific Corrections with Updated ξ

$$\omega_{T0} = \omega_{\text{standard}} \times (1 + \xi) = \omega \times 1.00001 \quad (37)$$

Measurable Frequency Shift: 10 ppm (calculated with $\xi = 1.0 \times 10^{-5}$)

.6 Experimental Distinction with Updated ξ

Universal Distinction Tests

Repeatability Test

Protocol: Execute each algorithm 1000 times under identical conditions

Predictions:

- **Standard QM:** Results consistent within statistical error bounds
- **T0:** Perfect repeatability with 0.001% systematic precision

ξ -Parameter Precision Tests with Updated Value

Protocol: High-precision measurements searching for systematic deviations

Predictions:

- **Standard QM:** No systematic corrections predicted
- **T0:** 10 ppm systematic shifts in gate operations
- **Detection Threshold:** Requires precision better than 1 ppm

Higgs matching of the ξ parameter (consistency check, not a derivation, Doc. 385):

$$\xi = \frac{\lambda_h^2 v^2}{16\pi^3 m_h^2} \quad (38)$$

$$m_h = 125.25 \pm 0.17 \text{ GeV} \quad (\text{Higgs boson mass}) \quad (39)$$

$$v = 246.22 \text{ GeV} \quad (\text{vacuum expectation value}) \quad (40)$$

$$\lambda_h = \frac{m_h^2}{2v^2} = 0.129383 \quad (\text{Higgs self-coupling}) \quad (41)$$

Step-by-Step Calculation

$$\lambda_h^2 = (0.129383)^2 = 0.01674 \quad (42)$$

$$v^2 = (246.22 \times 10^9)^2 = 6.062 \times 10^{22} \text{ eV}^2 \quad (43)$$

$$16\pi^3 = 496.10 \quad (44)$$

$$m_h^2 = (125.25 \times 10^9)^2 = 1.569 \times 10^{22} \text{ eV}^2 \quad (45)$$

Higgs result (consistency check):

$$\xi_{\text{QFT}} = 1.304 \times 10^{-4} \quad (46)$$

This is 2.2 % below 4/30000.

Ideal ξ Parameter from Measurement Error Analysis

To determine the ideal ξ value, we analyze acceptable measurement errors in quantum gate operations.

NOT Gate Error Analysis

The NOT gate operation in T0 formulation:

$$|0\rangle \rightarrow |1\rangle \times (1 + \xi) \quad (47)$$

For ideal output amplitude 1.0, the measurement error is:

$$\text{Error} = \frac{|(1 + \xi) - 1|}{1} = |\xi| \quad (48)$$

With acceptable error threshold of 0.001%:

$$|\xi| = 0.001\% = 1.0 \times 10^{-5} \quad (49)$$

Ideal ξ parameter: $\xi_{\text{ideal}} = 1.0 \times 10^{-5}$

Source	ξ Value	Agreement
Measurement error requirement	1.000×10^{-5}	Reference
Higgs matching (Doc. 097, $16\pi^3$)	1.304×10^{-4}	4/30000 (2.2 %)
Adopted value	1.0×10^{-5}	Ideal

Table 2: ξ Parameter Source Comparison**Comparison with Higgs Calculation**

The Higgs matching (Doc. 097) yields $\xi_{\text{QFT}} \approx 1.30 \times 10^{-4}$, consistent with the physical $\xi = 4/30000 \approx 1.33 \times 10^{-4}$ (Doc. 009) to 2.2 % (consistency check). The simulation parameter $\xi_{\text{num}} = 10^{-5}$ is a separate numerical choice for algorithm stability.

Information Structure in Energy Field Amplitudes

The energy field amplitude deviations encode specific physical information:

Hadamard Gate Analysis:

$$\text{Ideal QM amplitude: } \pm \frac{1}{\sqrt{2}} = \pm 0.7071067812 \quad (50)$$

$$\text{T0 energy field amplitude: } \pm 0.5 \times (1 + \xi) = \pm 0.5000050000 \quad (51)$$

$$\text{Deviation: } 29.3\% \text{ (information carrier, not error)} \quad (52)$$

This 29.3% deviation contains:

1. **Spatial scaling information:** Field extent factor $\sqrt{1 + \xi} = 1.000005$
 2. **Energy density information:** Density ratio $(1 + \xi/2) = 1.000005$
 3. **ξ -parameter information:** Direct measure of $\xi_{\text{num}} = 1.0 \times 10^{-5}$
 4. **Vacuum structure information:** Connection to electroweak symmetry breaking
- Total information enhancement:** 51 bits per qubit (compared to 1 bit in standard QM)

Experimental Roadmap**Phase I - Precision Validation**

Goal: Verification of 0.001% systematic errors in quantum gates **Methods:**

- High-precision amplitude measurements
- Statistical vs. deterministic behavior tests
- Gate fidelity analysis beyond standard error bounds

Expected timeframe: 1-2 years with existing quantum hardware

Phase II - Information Layer Access

Goal: Demonstration of access to enhanced information layers **Methods:**

- Spatial field mapping with nanometer resolution
- Time-resolved field evolution measurements
- Multi-modal information extraction protocols

Expected timeframe: 3-5 years with specialized equipment

Phase III - Higgs Coupling Detection**Goal:** Direct measurement of ξ parameter effects **Methods:**

- Quantum field correlation measurements
- Vacuum structure probes

Expected timeframe: 5-10 years with next-generation technology