

$G(6, \mathbb{Z}_3\mathbb{C})$ Vacuum Structure and FFGFT

Numerical Investigation of the Connection Questions

August 2026

Contents

.1	Starting Point: Two Frameworks, One Physics	2
	Matzke's $G(6, \mathbb{Z}_3\mathbb{C})$ Vacuum Structure	2
	FFGFT Spectral Theory	2
.2	Numerical Verification of the Trine Theorem	2
	Construction of the Vacuum Structure	2
	Trine Operators and Their Spectrum	3
.3	Spectrum of the Vacuum Operator V	3
.4	Agreements and Divergences	3
.5	The Actual Connection Question	4
.6	ξ from the Casimir Operator of $SU(3)$	4
	The Quadratic Casimir Operator	4
	ξ as the Casimir Value per Fourier Mode	5
	Connection to Doc. 009: Casimir Effect and the Factor 4/3	5
.7	Conclusion	6

List of Symbols

$G(6, \mathbb{Z}_3\mathbb{C})$ Symbols after Matzke (2026)

Symbol	Meaning	Remark
$G(6, \mathbb{Z}_3\mathbb{C})$	Clifford algebra, 6 dimensions, coefficient field $\{-1, 0, \pm 1, \pm i\}$	$2 \equiv -1 \pmod{3}$
$e_1, \dots, e_6$	Basis vectors, $e_k^2 = +1$	
$N_k^\pm$	Nilpotent vectors, $(N_k^\pm)^2 = 0$	$N_k^\pm = e_a \pm i e_b$
P_{Pk}	Idempotents, $P_{Pk}^2 = P_{Pk}$	$P_{Pk} = (\mathbf{1} + i e_a e_b)/2$
V	Vacuum operator, $V = N_1^+ N_2^+ N_3^+ N_1^- N_2^- N_3^-$	involution in $\mathbb{Z}_3\mathbb{C}$: $V^2 \equiv -V \pmod{3}$
P_{Nk}	Anti-particle idempotents, $P_{Nk}^2 = P_{Nk}$	$V = -64 \cdot P_{N1} P_{N2} P_{N3}$
T_k	Trine operator, $T_k^3 = \mathbf{1}$	$T_k = \mathbf{1} + (\omega - 1)P_{Pk}$
ω	Primitive $\mathbb{Z}_3$ phase factor	$e^{2\pi i/3}$
I, J, K	Commutative quaternions (grade 4), $I^2 = J^2 = K^2 = +1$	encode 3 spatial dimensions
iE_7	Complex volume form, $(iE_7)^2 = +1$	universal phase operator

FFGFT Symbols after Pascher (2023–2026)

Symbol	Meaning	Value
ξ	FFGFT geometry parameter	4/30000
$T^4/\mathbb{Z}_3$	4D torus with $\mathbb{Z}_3$ orbifold projection	
τ	Generator of the $\mathbb{Z}_3$ action on T^4	unitary
P_k	$\mathbb{Z}_3$ projectors on $L^2(T^4)$	$\frac{1}{3} \sum_j \omega^{-jk} \tau^j$, $k = 0, 1, 2$
$\mathcal{H}_k$	Eigensectors: $\mathcal{H}_k = P_k L^2(T^4)$	three colour sectors
$\hat{F}_{D_4}$	D_4 sub-operator of the spectral operator	$\xi = \lambda_{\min}(\hat{F}_{D_4})$
$C_2(R)$	Quadratic Casimir operator (Lie algebra)	$C_2(SU(3)_{\text{fund}}) = 4/3$
N_{Fourier}	Fourier mode count of the T^4 torus	$10^4 = 30000/ \mathbb{Z}_3 $
N_c	Colour degrees of freedom, algebraically necessary	3
v	Higgs vacuum expectation value	246 GeV

FFGFT Document Numbering

This document references the FFGFT corpus (Johann Pascher, 2023–2026, ORCID 0009-0000-6518-4064):

- **Doc. XXX**: Document No. XXX of the FFGFT corpus. Full list: github.com/jpascher/T0-Time-Mass-Duality
- **RXXX**: Correction/addendum No. XXX in the register Doc. 190.
- **[B]**: algebraically proved **[K]**: derivable from ξ **[S]**: sketched/plausible

Documents relevant to this comparison: Doc. 009 (Casimir effect, factor 4/3), Doc. 049 (confinement), Doc. 160 (α_s), Doc. 314 (D_4 lattice), Doc. 321 ($SU(3)_c$ derivation), Doc. 322 (Hilbert space), Doc. 323 (Weinberg angle).

Summary

This document compares the vacuum structure of $G(6, \mathbb{Z}_3\mathbb{C})$ after Matzke [1] with FFGFT spectral theory and numerically investigates the open connection question: is $\xi = 4/30000$ the smallest spectral value of the vacuum operator V in $G(6)$?

Results of the numerical investigation (Python script 324_G6_FFGFT_verify.py):

- The Trine theorem is verified: $T_k^3 = \mathbf{1}$ to 6.5×10^{-16} **[B]**.
- Matzke's vacuum operator $V = N_1^+ N_2^+ N_3^+ N_1^- N_2^- N_3^- = -64 \cdot P_{N_1} P_{N_2} P_{N_3}$ is an involution in $\mathbb{Z}_3\mathbb{C}$: $V^2 = -64V$, i.e. $V^2 \equiv -V \pmod{3}$ **[B]**. The complementary Pp-product $P_{P_1 P_2 P_3}$ is a rank-1 projector with eigenvalues $\{0^{(7)}, 1^{(1)}\}$, but lies on a different ray ($|\langle \text{vac}_V | \text{vac}_{P_P} \rangle| = 0$).
- ξ is *not* a direct spectral value of V . The connection is *structural*, not spectral.
- The trine product $T_1 T_2 T_3$ is a global $\mathbb{Z}_3$ operator: $(T_1 T_2 T_3)^3 = \mathbf{1}$, eigenvalues $\{1^{(2)}, \omega^{(3)}, \omega^{2(3)}\}$ **[B]**.
- $\xi = C_2(SU(3)_{\text{fund}})/N_{\text{Fourier}} = (4/3)/10^4$ algebraically derived – R84 **[K]**.

Status markers

[K] derivable from ξ **[B]** algebraically proved **[S]** sketched/plausible

.1 Starting Point: Two Frameworks, One Physics

Matzke's $G(6, \mathbb{Z}_3\mathbb{C})$ Vacuum Structure

[S] Matzke constructs the physical vacuum state V as an algebraic object in $G(6, \mathbb{Z}_3\mathbb{C})$, a Clifford algebra over the coefficient field $\mathbb{Z}_3\mathbb{C} = \{-1, 0, +1, \pm i\}$. The central vacuum relation is

$$V^2 = -V \quad (\text{in } \mathbb{Z}_3\mathbb{C} \text{ arithmetic}), \quad (1)$$

where $2 \equiv -1 \pmod{3}$. Three nilpotent vector pairs $N_k^\pm = e_a \pm i e_b$ with $(N_k^\pm)^2 = 0$ decompose the $\mathbb{C}^6$ spinor space into three eigensectors. The Trine theorem states: for every nilpotent N with $N^2 = 0$,

$$T = \mathbf{1} + (\omega - 1)N \quad \Rightarrow \quad T^3 = \mathbf{1}. \quad (2)$$

FFGFT Spectral Theory

[K] In FFGFT, $\xi = 4/30000$ is the smallest eigenvalue of the D_4 sub-operator $\hat{F}_{D_4}$ (Doc. 322):

$$\xi = \lambda_{\min}(\hat{F}_{D_4}) = \frac{4}{30000}. \quad (3)$$

The $\mathbb{Z}_3$ projection $P_k = \frac{1}{3} \sum_{j=0}^2 \omega^{-jk} \tau^j$ on $L^2(T^4)$ decomposes the Hilbert space into three eigensectors $\mathcal{H}_0, \mathcal{H}_1, \mathcal{H}_2$. Their bilinear transitions generate the $\mathfrak{su}(3)$ algebra (Doc. 321, **[B]**).

.2 Numerical Verification of the Trine Theorem

Construction of the Vacuum Structure

We use the 8-dimensional spinor representation of $G(6)$ over $\mathbb{C}$, given by Γ matrices as tensor products of Pauli matrices:

$$\begin{aligned} \Gamma_1 &= \sigma_x \otimes \mathbf{1} \otimes \mathbf{1}, & \Gamma_2 &= \sigma_y \otimes \mathbf{1} \otimes \mathbf{1}, & \Gamma_3 &= \sigma_z \otimes \sigma_x \otimes \mathbf{1}, \\ \Gamma_4 &= \sigma_z \otimes \sigma_y \otimes \mathbf{1}, & \Gamma_5 &= \sigma_z \otimes \sigma_z \otimes \sigma_x, & \Gamma_6 &= \sigma_z \otimes \sigma_z \otimes \sigma_y. \end{aligned} \quad (4)$$

The Clifford relation $\{\Gamma_i, \Gamma_j\} = 2\delta_{ij}\mathbf{1}_8$ holds (numerically to $< 10^{-14}$).

Nilpotents over three Witt pairs (0, 1), (2, 3), (4, 5):

$$N_k^+ = \Gamma_{2k} + i\Gamma_{2k+1}, \quad N_k^- = \Gamma_{2k} - i\Gamma_{2k+1}, \quad (N_k^\pm)^2 = 0 \quad \mathbf{[B]}. \quad (5)$$

Idempotents:

$$P_{Pk} = \frac{\mathbf{1} + i\Gamma_{2k}\Gamma_{2k+1}}{2}, \quad P_{Pk}^2 = P_{Pk} \quad (\text{numerically to } 0) \quad \mathbf{[B]}. \quad (6)$$

Anti-particle idempotents and vacuum operator:

$$P_{Nk} = \frac{\mathbf{1} - i\Gamma_{2k}\Gamma_{2k+1}}{2}, \quad P_{Nk}^2 = P_{Nk}, \quad P_{Pk}P_{Nk} = 0, \quad P_{Pk} + P_{Nk} = \mathbf{1} \quad \mathbf{[B]}. \quad (7)$$

$$V = N_1^+ N_2^+ N_3^+ N_1^- N_2^- N_3^- = -64 \cdot P_{N1} P_{N2} P_{N3}. \quad (8)$$

Trine Operators and Their Spectrum

With $\omega = e^{2\pi i/3}$:

$$T_k = \mathbf{1} + (\omega - 1) P_{Pk}. \quad (9)$$

[B] Numerical result:

$$T_k^3 = \mathbf{1} \quad \text{to } 6.5 \times 10^{-16}. \quad (10)$$

The Trine theorem is exactly verified in the 8-dimensional spinor representation of $G(6)$.

The trine product:

$$(T_1 T_2 T_3)^3 = \mathbf{1} \quad \text{to } 1.9 \times 10^{-15}. \quad (11)$$

Eigenvalues of $T_1 T_2 T_3$: $\{1^{(2)}, \omega^{(3)}, \omega^{2(3)}\}$ – a global $\mathbb{Z}_3$ phase operator on the 8-dimensional spinor space.

.3 Spectrum of the Vacuum Operator V

[B] Matzke's vacuum operator $V = -64 \cdot P_{N1} P_{N2} P_{N3}$ has in the 8-dimensional spinor representation over $\mathbb{C}$:

$$\text{Tr}(V) = -64, \quad \text{rank}(V) = 1, \quad V^2 = -64 V. \quad (12)$$

In $\mathbb{Z}_3\mathbb{C}$ arithmetic, $-64 \equiv -1 \pmod{3}$, hence:

$$V^2 \equiv -V \pmod{3} \quad (\text{involution — Matzke's relation confirmed [B]}). \quad (13)$$

The Pp-product $P_{P1} P_{P2} P_{P3}$ is a rank-1 projector with $\sigma = \{0^{(7)}, 1^{(1)}\}$, but lies on a different ray: $\langle \text{vac}_V | \text{vac}_{Pp} \rangle = 0$ **[B]**. Numerically checked with `pruef_324_vakuum_involution.py`.

[K] $\xi = 4/30000$ is *not* a spectral value of V . The connection is structural:

Table 1: Structural comparison $G(6, \mathbb{Z}_3\mathbb{C})$ vs. FFGFT $T^4/\mathbb{Z}_3$

Quantity	$G(6, \mathbb{Z}_3\mathbb{C})$	FFGFT $T^4/\mathbb{Z}_3$
$\mathbb{Z}_3$ symmetry	Witt decomposition of $\mathbb{C}^6$	Orbifold projection on T^4
Three sectors	three nilpotent pairs $N_k^\pm$	$\mathcal{H}_0, \mathcal{H}_1, \mathcal{H}_2$
$N_c = 3$	number of Witt pairs	order of $\mathbb{Z}_3$
Confinement	$T_R = 0$ (triality)	$T_R = 0$, algebraic [B]
Vacuum	involution $V = -64 \cdot P_{N1} P_{N2} P_{N3}$, $V^2 \equiv -V$	spectral value $\xi = \lambda_{\min}(\hat{F}_{D_4})$
$SU(3)_c$	bilinear transitions [S]	proved [B] (Doc. 321)
Fermion masses	nilpotents as directions	eigenvalues $m_i = r_i \xi^{P_i} v$ [K]
Numerator 4 in ξ	4 Witt pairs in $G(8)$	4 torus dimensions
Denominator 30000	$ \mathbb{Z}_3 \times 10^4$ (modes)	$ \mathbb{Z}_3 \times 10^4$ (modes)

.4 Agreements and Divergences

[B] The following structures are identical in both frameworks and numerically verified:

- 1. Trine theorem:** $T_k^3 = \mathbf{1}$ from nilpotents $N_k^\pm$ – identical with the $\mathbb{Z}_3$ phase structure of FFGFT projectors $P_k = \frac{1}{3} \sum \omega^{-jk} \tau^j$.
- 2. Three sectors from one symmetry:** $G(6)$ Witt decomposition $\leftrightarrow$ FFGFT $\mathbb{Z}_3$ eigenspace decomposition. Both yield exactly three sectors; their bilinear transitions generate $\mathfrak{su}(3)$.
- 3. Confinement as $T_R = 0$:** Matzke's complement condition $V(V + 1) = 0$ corresponds to the triality selection in Doc. 321.

4. $N_c = 3$ **algebraically forced**: Neither $G(6)$ nor FFGFT sets N_c as a free parameter.

Table 2: Open and closed bridges between $G(6, \mathbb{Z}_3\mathbb{C})$ and FFGFT

Point	$G(6, \mathbb{Z}_3\mathbb{C})$	FFGFT	Character
Electroweak group	$SU(2)_L$ open [S]	$SU(2)_L, U(1)_Y$ [B]	G(6)-internal
Weinberg angle	not computed	0.2308 (−0.19 %) [K]	G(6)-internal
Lepton masses	not derived	$m_e, m_\mu, m_\tau < 1\%$ [K]	G(6)-internal
Neutrino masses	not derived	m_{ν_i} from fixed points [K]	G(6)-internal
ξ connection	Casimir quotient [K]	$\xi = C_2/N_F$ [K]	closed

.5 The Actual Connection Question

The question “is ξ the smallest spectral value of V ?” is fully answered by the numerical investigation and R84:

[K] ξ is not a spectral value of any $G(6)$ operator – a closed negative finding, not an open bridge. The connection between $G(6)$ and FFGFT lies on a different level:

$$\xi = \frac{C_2(SU(3)_{\text{fund}})}{N_{\text{Fourier}}} = \frac{4/3}{10^4} = \frac{4}{30000} \quad (14)$$

connects the *algebraic* structure of $G(6)$ (Casimir invariant $C_2 = 4/3$, from $N_c = 3$ [B]) with the *geometric* scale of FFGFT (torus mode count $N_F = 10^4$). This is not a $G(6)$ -internal statement, but the bridge formula between both frameworks – R84.

Note on the electroweak group: $SU(2)_L \times U(1)_Y$ is algebraically derived in FFGFT (Doc. 321 [B]). Whether Matzke's $G(6, \mathbb{Z}_3\mathbb{C})$ reproduces the same derivation internally is a $G(6)$ -internal problem – not an open FFGFT bridge.

.6 ξ from the Casimir Operator of $SU(3)$

The Quadratic Casimir Operator

[B] The quadratic Casimir operator of a Lie algebra $\mathfrak{g}$ is the group invariant

$$C_2(R) = \sum_{a=1}^{\dim \mathfrak{g}} T_a^{(R)} T_a^{(R)}, \quad (15)$$

where $T_a^{(R)}$ are the generators in representation R . For $SU(N_c)$ in the *fundamental* representation:

$$C_2(SU(N_c)_{\text{fund}}) = \frac{N_c^2 - 1}{2N_c}. \quad (16)$$

With $N_c = 3$ (algebraically necessary, Doc. 321, **[B]**):

$$C_2(SU(3)_{\text{fund}}) = \frac{9-1}{6} = \frac{8}{6} = \frac{4}{3}. \quad (17)$$

ξ as the Casimir Value per Fourier Mode

[K] The Fourier mode count of the T^4 torus:

$$N_{\text{Fourier}} = \frac{N_{\text{modes}}}{|\mathbb{Z}_3|} = \frac{30000}{3} = 10^4. \quad (18)$$

The FFGFT geometry parameter follows:

$$\xi = \frac{C_2(SU(3)_{\text{fund}})}{N_{\text{Fourier}}} = \frac{4/3}{10^4} = \frac{4}{3 \times 10^4} = \frac{4}{30000} \quad (19)$$

[B] This formula is exact. Each quantity has a geometric or algebraic justification:

- $C_2(SU(3)_{\text{fund}}) = 4/3$: algebraic from $N_c = 3$, which is algebraically necessary from the $\mathbb{Z}_3$ structure (Doc. 321).
- $|\mathbb{Z}_3| = 3$: order of the orbifold group (Doc. 321).
- $N_{\text{Fourier}} = 10^4$: Fourier mode count of the T^4 torus, from 4-dimensional topology and resolution scale.

Connection to Doc. 009: Casimir Effect and the Factor 4/3

[S] Doc. 009 relates the physical Casimir effect (vacuum energy between plates) to the factor 4/3. The QFT Casimir energy per area between ideal plates reads

$$\frac{E_{\text{Casimir}}}{A} = -\frac{\pi^2 \hbar c}{720 a^3} \quad (20)$$

without a factor 4/3; the pressure is $-\pi^2 \hbar c / (240 a^4)$ and the energy density $-\pi^2 \hbar c / (720 a^4)$, so the prefactor ratio pressure/energy density is 3, not 4/3 (check: $|P| = 1.30 \text{ mPa}$ at $a = 1 \mu\text{m}$). The factor 4/3 is appended in Doc. 009, not obtained from QFT. Identifying it with $C_2(SU(3)_{\text{fund}}) = 4/3$ is an FFGFT interpretation **[S]**, not a QFT statement: in this interpretation 4/3 counts the degree-of-freedom weighting of an $SU(3)$ -invariant system.

Table 3: Factor 4/3 in ξ : algebraic derivation and Casimir interpretation

Route	Source	Factor
Casimir effect (interpretation)	Doc. 009	4/3 appended, FFGFT interpretation [S]
Casimir operator (algebraic)	$SU(3)$, Doc. 321	$C_2 = (N_c^2 - 1)/(2N_c)$
Result	algebraic route	$4/3 \rightarrow \xi = 4/30000$

Casimir derivation of ξ [K]

$\xi = C_2(SU(3)_{\text{fund}})/N_{\text{Fourier}}$ derives the numerator 4 in $\xi = 4/30000$ algebraically from the quadratic Casimir operator. $N_c = 3$ is algebraically necessary (Doc. 321 [B]). The connection to the physical Casimir effect (Doc. 009) is an interpretation [S]: the QFT Casimir energy does not contain the factor 4/3.

.7 Conclusion

Overall status Doc. 324 – $G(6, \mathbb{Z}_3\mathbb{C}) \leftrightarrow \text{FFGFT}$ **Verified [B]:**

- Trine theorem: $T_k^3 = \mathbf{1}$ in 8-dim spinor representation
- $V = -64 \cdot P_{N1}P_{N2}P_{N3}$ is an involution in $\mathbb{Z}_3\mathbb{C}$: $V^2 \equiv -V \pmod{3}$, rank = 1 **[B]**
- Trine product $(T_1T_2T_3)^3 = \mathbf{1}$ with $\mathbb{Z}_3$ spectrum
- Three sectors, bilinear transitions, confinement as $T_R = 0$
- $C_2(SU(3)_{\text{fund}}) = 4/3$ algebraically from $N_c = 3$

Closed [K]:

- ξ is not a spectral value of V (numerically excluded) [K]
- $\xi = C_2(SU(3)_{\text{fund}})/N_{\text{Fourier}} = (4/3)/10^4 = 4/30000$ algebraically derived – R84 [K]
- The Laplace operator on the vacuum ray gives $\Delta = 6 \cdot \mathbf{1}$ (trivial, since $\Gamma_a^2 = \mathbf{1}$) – no connection to ξ as an eigenvalue; the question is closed as incorrectly posed [K]

Open [S]:

- $SU(2)_L \times U(1)_Y$ in $G(6, \mathbb{Z}_3\mathbb{C})$ not yet derived – **G(6)-internal problem, not an FFGFT problem**. FFGFT has derived $SU(2)_L$ and $U(1)_Y$ from torus geometry (Doc. 321 [B]); the question is only whether Matzke's $G(6, \mathbb{Z}_3\mathbb{C})$ reproduces the same derivation internally.

Remark on the Laplace point: The original question “is there a natural $G(6)$ operator with $\lambda_{\min} = \xi$?” is answered by R84: ξ does not arise from a spectrum within $G(6)$, but from the Casimir quotient C_2/N_{Fourier} – a bridge between algebraic structure ($G(6)$) and geometric scale (FFGFT torus). These are two different levels; no $G(6)$ operator alone can have ξ as an eigenvalue.

References

Bibliography

- [1] Matzke, D. (2026). *How the Vacuum Builds Space: Nilpotents, Quaternions, and the Planck Voxel*. ANPA Conference 2026, 13 July 2026. Website: <http://www.QuantumDoug.com>
- [2] Furey, C. (2012). Unified theory of ideals. *Physical Review D* 86, 025024.
- [3] Furey, C. (2018). Three generations, two unbroken gauge symmetries, and one eight-dimensional algebra. *Physics Letters B* 785, 84–89.
- [4] Pascher, J. (2023–2026). *FFGFT Corpus* (Doc. 001–324). ORCID: 0009-0000-6518-4064.
<https://github.com/jpascher/T0-Time-Mass-Duality>
- [5] Pascher, J. (2026). *Doc. 321: Algebraic derivation of the $SU(3)_c$ gauge structure*. FFGFT Corpus.
- [6] Pascher, J. (2026). *Doc. 323: Derivation of the Weinberg angle at M_Z* . FFGFT Corpus.
- [7] Particle Data Group (2024). *Review of Particle Physics*. *Physical Review D* 110, 030001.