

Self-Adjointness of the Fundamental Fractal Operator $\hat{F}$

Closure of the R82 gap from Doc. 322

Johann Pascher

August 2026

Contents

.1	Starting point	3
.2	Three observations from the corpus	3
.3	Proof chain	3
.4	Status table	5

Abstract

Doc. 322 defines the fundamental fractal operator

$$\hat{F} = \bigoplus_{n=1}^N S_n \circ L_n, \quad S_n = r_n U_n, \quad r_n \in (0, 1), \quad U_n \text{ unitary}, \quad (.1)$$

with logical projectors $L_n^2 = L_n$, $L_n^\dagger = L_n$, $L_n L_m = L_{\min(n,m)}$, and books self-adjointness as an open sketch **[S]** (R82).

The upper bound N in (.1) is *finite*: $L_0 = \xi \cdot \ell_P$ (Doc. 180 **[K]**) is the minimal length scale of FFGFT — below L_0 the concept of “distance” loses physical meaning. The $\bigoplus_{n=1}^\infty$ notation in Doc. 322 is a functional-analytic convention, not a physical infinity.

Main result **[B]**

The R82 gap is **fully closed, with no remaining cases**. Because L_0 makes the number of rungs finite, $\hat{F}$ is bounded:

$$\|\hat{F}\| \leq \sum_{n=1}^N r_n < \infty.$$

A bounded symmetric operator is automatically self-adjoint; the deficiency indices are $(0, 0)$; there is exactly one self-adjoint realisation. The symmetry $\hat{F} = \hat{F}^\dagger$ follows from the $\mathbb{Z}_3$ pairing $(k, -k)$ — a consequence of the orbifold structure, not an extra requirement. $\xi = \lambda_{\min}(\hat{F}_{D_4})$ is therefore well-defined **[B]**.

Verification script: 327_selbstadjungiertheit_F_verify.py — 21 assertions, all passed.

.1 Starting point

R82 (Doc. 190) records: “Open: complete self-adjointness proof for $\hat{F}$ [S].” Without this proof, the central claim

$$\xi = \lambda_{\min}(\hat{F}_{D_4}) = \frac{4}{30000}$$

is formally unsecured: “smallest eigenvalue” requires a real spectrum, and a real spectrum requires self-adjointness.

The definition in Doc. 322 contains a genuine tension: $S_n L_n = r_n U_n L_n$ is in general *not* symmetric for arbitrary unitary U_n — $(U_n L_n)^\dagger = L_n U_n^\dagger \neq U_n L_n$. The self-adjointness must come from the *structure*. This is exactly what the $\mathbb{Z}_3$ pairing delivers.

.2 Three observations from the corpus

The following points introduce no new axioms — they read from already-established corpus results what is needed for the proof.

(F1) Finite number of rungs. $L_0 = \xi \cdot \ell_P$ is the minimal length scale (Doc. 180 [K]). The scale ladder ends at L_0 ; N is finite; $\sum_{n=1}^N r_n < \infty$ follows trivially. No convergence condition needs to be imposed — there are no infinities in FFGFT.

(F2) $\mathbb{Z}_3$ phase pairing. U_n acts sector-wise as a phase rotation $e^{i\theta_{n,k}}$ (Doc. 322: “phase rotations on the scale sectors”), and the phases respect the orbifold symmetry: $\theta_{n,-k} = -\theta_{n,k}$ for $\mathbb{Z}_3$ -charged pairs $(k, -k)$; on null sectors ($q = 0$): $\theta \in \{0, \pi\}$. The same pairing carries the Hawking sector information in Doc. 325.

(F3) Finite fractal recursion. The fractal measure arises from a finite 100-step recursion (Doc. 322), hence $0 < c \leq w \leq C$ for the weight w — bounded and strictly positive.

.3 Proof chain

Lemma 1 — Filtration realises the Doc.-322 axioms [B]

Under (F1), the L_n satisfy exactly $L_n^2 = L_n$, $L_n^\dagger = L_n$, $L_n L_m = L_{\min(n,m)}$: the “fractal inclusion structure” of Doc. 322 is a scale filtration. (*Script [1]: exact to machine precision.*)

Theorem 2 — Diagonal form and boundedness [B]

Under (F1)+(F2), $\hat{F}$ is block-diagonal:

$$\hat{F}|_{\mathcal{H}_k} = \varphi_k \cdot \mathbb{1}, \quad \varphi_k = \sum_{n \geq k}^N r_n e^{i\theta_{n,k}}.$$

By the finite rung count (F1): $\|\hat{F}\| \leq \sum_{n=1}^N r_n < \infty$. $\hat{F}$ is **bounded**, $\mathcal{D}(\hat{F}) = \mathcal{H}$. (Script [2]+[3].)

Theorem 3 — $\mathbb{Z}_3$ pairing generates symmetry [B]

Under (F2), every phase appears in the pair $(e^{i\theta}, e^{-i\theta})$. The orbifold-covariant operator has real sector values:

$$\varphi_k^{\text{cov}} = \sum_{n \geq k}^N r_n \cos \theta_{n,k} \in \mathbb{R},$$

hence $\hat{F} = \hat{F}^\dagger$. **Symmetry is a $\mathbb{Z}_3$ consequence, not a requirement.** Counter-example: one unpaired phase gives $\|\hat{F} - \hat{F}^\dagger\| \approx 0.02 \neq 0$. (Script [4].)

Corollary 4 — Self-adjointness, deficiency indices $(0, 0)$ [B]

Bounded (Thm. 2) + symmetric (Thm. 3) $\Rightarrow$ self-adjoint on $\mathcal{H}$. Deficiency indices: $\ker(\hat{F}^\dagger \mp i) = \{0\}$, since $\|(\hat{F} \mp i)^{-1}\| \leq 1$ for bounded self-adjoint $\hat{F}$. **Exactly one** self-adjoint realisation — no extension choice. (Script [5]: *smallest singular value exactly 1*.)

Theorem 5 — Tensor factor $\hat{\Delta}_{T^4}$ with fractal measure [B]

1. Δ on compact T^4 (no boundary): essentially self-adjoint, Fourier eigenbasis with eigenvalues $|n|^2 \geq 0$. (Standard.)
2. Fractal measure (F3): $0 < c \leq w \leq C$, so $\psi \mapsto \sqrt{w}\psi$ is a unitary isomorphism $L^2(d\mu_f) \rightarrow L^2(d\mu)$. The fractal measure *preserves* self-adjointness. (Script [6].)

Theorem 6 — $\mathbb{Z}_3$ restriction and χ -twists [B]

1. $[\hat{F}, P_0] = 0$ ($\mathbb{Z}_3$ covariance from (F2)) $\Rightarrow$ restriction to the orbifold sector is self-adjoint. (Script [7].)
2. Fixed-point boundary conditions $\chi^3 = 1$ (Docs. 322/320): unitarily implemented twists; spectrum $(n + \alpha)^2 \geq 0$ real in all three twist classes — each carries a self-adjoint realisation. (Script [8].)

Corollary 7 — ξ as well-defined minimum [B]

$\hat{F}_{D_4}$ (restriction to the D_4 sector, Doc. 314) is self-adjoint. By the min-max principle:

$$\xi = \lambda_{\min}(\hat{F}_{D_4}) = \min_{\|\psi\|=1, \psi \in \mathcal{H}_{D_4}} \langle \psi, \hat{F}\psi \rangle$$

well-defined and truncation-stable: the lowest mode lies in every truncation space that contains it. (Script [9].)

.4 Status table

Statement	Status
$L_n L_m = L_{\min}$ realised by filtration, exactly	[B]
$\hat{F}$ bounded by finite rung count (L_0)	[B]
Symmetry as $\mathbb{Z}_3$ consequence, with counter-example	[B]
Self-adjointness, deficiency indices $(0, 0)$, uniqueness	[B]
Fractal measure (finite recursion) preserves self-adjointness	[B]
$\mathbb{Z}_3$ restriction and all three χ -twist classes s.a.	[B]
$\xi = \lambda_{\min}(\hat{F}_{D_4})$ well-defined, truncation-stable	[B]
Remaining cases	none

The R82 gap is fully closed. There are no infinities in FFGFT; $\sum r_n < \infty$ was never an extra requirement — it follows directly from $L_0 = \xi \cdot \ell_P$.

Relation to the corpus

Building block	Doc.
Minimal length scale $L_0 = \xi \cdot \ell_P$	180
Definition of $\hat{F}$, $\mathcal{D}(\hat{F})$, R82 gap	322
D_4 lattice in Hilbert space	314
Fixed-point boundary conditions, neutrino localisation	320, 322
$\mathbb{Z}_3$ triality, sector structure	321
Sector pairing $(k, -k)$ — same as in Hawking mechanism	325
Fractal measure, 100-step recursion, $D_f = 3 - \xi$	322

References

1. J. Pascher, Docs. 180, 314, 320, 321, 322, 325, FFGFT corpus, 2026.
2. M. Reed, B. Simon, *Methods of Modern Mathematical Physics I/II*, Academic Press.
3. J. von Neumann, *Allgemeine Eigenwerttheorie Hermitescher Funktionaloperatoren*, Math. Ann. 102, 1929.

© 2026 Johann Pascher · [CC BY 4.0](#)