

T0-Theory: The T0-Time-Mass Duality

Contents

.1	Introduction	1
	Basic Principles of the T0 Theory	1
	The Fundamental Geometric Parameter	2
	Notation and Conventions	2
	Derived Parameters	2
.2	Extended Lagrangian with Time Field	2
	Standard Model Lagrangian as Starting Point	2
	Introduction of the Dynamic Time Field	3
	Kinetic Term of the Time Field	3
	Mass Term of the Time Field	3
	Mass-Proportional Coupling	3
	Complete Extended Lagrangian	4
.3	Quantum Corrections and Feynman Rules	4
	Feynman Rules from the Lagrangian	4
	One-Loop Diagram	4
	General Formula for Scalar Mediators	4
.4	Derivation of the Fundamental T0 Formula	5
	Heavy Mediator Limit	5
	Inserting the T0 Coupling	5
	Higgs-Time Field Connection	5
	Final Formula	6
.5	Numerical Evaluation	6
	Determination of the Normalization Constant	6
	Final T0 Contribution Formula	6
	Lepton-Specific Predictions	6
	No Conversion Needed	7
.6	Quadratic Mass Scaling	7
	Fundamental Prediction	7
	Physical Interpretation	7
.7	Connection to the Geometric Formulation	8
	Relation to the g-2 Analysis	8
	Parameter Correspondences	8
.8	Higgs Mechanism in the T0 Theory	8
	Higgs Field as Fundamental Basis	8
	Spontaneous Symmetry Breaking	9
.9	Field-Theoretic Structure	9
	Equations of Motion	9
	Interpretation of the Equations of Motion	9
	Energy Scales	10
.10	Renormalization (historical presentation)	10
	Divergences in Loop Diagrams	10

	Renormalization Group Equations	10
	Consistency with the Standard Model	10
.11	Comparison of Different Formulations	11
	Lagrangian vs. Geometric Formulation	11
	Complementarity	11

Correction notes

Note on the muon $g - 2$ status (2025). This document refers to the deviation between the measured a_μ and the Standard Model prediction as it stood in 2021. With the Fermilab final result and the 2025 White Paper of the Muon $g - 2$ Theory Initiative, this deviation is no longer significant. Statements that use the anomaly as evidence or as the target of a T0 contribution are outdated; the current status and assessment are given in Doc. 382.

Corrections from the calculation check (30 Sept. 2026), each with a dated note at the place:

- Loop integral: missing factor m_ℓ^2 restored; the section "complete integral" (5/12) is dropped, in the limit $m_T \gg m_\ell$ the value is $2/3$. Final formula $\Delta a_\ell = \xi^4 m_\ell^2 / (12\pi^2 \lambda^2)$ instead of $5\xi^4 m_\ell^2 / (96\pi^2 \lambda^2)$. Dimensionless only with λ as a mass (R91) and a coupling acting as the dimensionless ξ .
- Normalization constant: $2.246 \times 10^{-13} \rightarrow 1.79 \times 10^{-56} \text{ GeV}^{-2}$ ($\lambda = m_p$); the old value follows from none of the stated parameters.
- "Conversion to conventional units" (factor 10^6) dropped: a_ℓ is dimensionless. Hence $\Delta a_\mu \approx 2 \times 10^{-58}$ instead of 251×10^{-11} – not measurable.
- The correspondence to Doc. 018 and the " $\sim 2\%$ precision" of both formulations do not hold; hierarchy estimate supplemented by λ^2 .
- Loop integral: integrand mixed from two parametrizations; standard limit $\propto \ln(m_T^2/m_\ell^2) - 7/6$ instead of $2/3$ (noted). Equation of motion: sign, $m_{\text{eff}} = m_\ell(1 - \xi\Delta m)$. Comparison table: main formula of Doc. 018 given correctly.

Abstract

This document presents the complete Lagrangian formulation of the T0 theory based on the fundamental geometric parameter $\xi = \frac{4}{3} \times 10^{-4}$. The theory establishes a fundamental time-mass duality $T(x, t) \cdot m(x, t) = 1$ and develops an extended Lagrangian with mass-proportional coupling to a dynamic time field. For the geometric derivation of the anomalous magnetic moments and experimental predictions, see Document 018_T0_Anomale-g2-10_De.tex. The focus here lies on the field-theoretic structure and the derivation of the fundamental T0 contribution formula from first principles.

.1 Introduction

Basic Principles of the T0 Theory

The T0 theory postulates a fundamental duality between time and mass:

$$T(x, t) \cdot m(x, t) = 1 \quad (1)$$

This duality leads to several fundamental consequences:

- Natural mass hierarchy through time scales
- Dynamic mass generation through the time field
- Quadratic scaling of the anomalous magnetic moments with m_ℓ^2
- Intrinsic integration of gravitation into quantum field theory

The Fundamental Geometric Parameter

The entire T0 theory is based on a single fundamental parameter:

$$\xi = \frac{4}{3} \times 10^{-4} = 1.333 \times 10^{-4} \quad (2)$$

This dimensionless parameter encodes the fundamental geometric structure of three-dimensional space. For the detailed geometric interpretation and derivation, see [Document 018](#).

Notation and Conventions

We use natural units ($\hbar = c = 1$) with metric signature $(+, -, -, -)$:

- $T(x, t)$: Dynamic time field with $[T] = E^{-1}$
- $\delta E(x, t)$: Fundamental energy field with $[\delta E] = E$
- $\Delta m(x, t) = \delta E(x, t)$: Mass field fluctuations
- $\xi = 1.333 \times 10^{-4}$: Fundamental geometric parameter
- λ : base mass of the sector (Addendum R91; m_p in the fundamental sector)
- m_ℓ : Lepton masses (e, μ, τ)

Derived Parameters

From the fundamental parameter ξ , the following are obtained:

$$\xi^2 = (1.333 \times 10^{-4})^2 = 1.777 \times 10^{-8} \quad (3)$$

$$\xi^4 = (1.333 \times 10^{-4})^4 = 3.160 \times 10^{-16} \quad (4)$$

.2 Extended Lagrangian with Time Field

Standard Model Lagrangian as Starting Point

The Standard Model Lagrangian for a lepton ψ_ℓ reads:

$$\mathcal{L}_{\text{SM}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \bar{\psi}_\ell(i\gamma^\mu D_\mu - m_\ell)\psi_\ell \quad (5)$$

where:

- $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$: Electromagnetic field strength tensor
- $D_\mu = \partial_\mu + ieA_\mu$: Covariant derivative
- m_ℓ : Lepton mass (constant)

Introduction of the Dynamic Time Field

In the T0 theory, the mass is not constant but coupled to a dynamic time field $T(x, t)$ through the time-mass duality (1). We introduce the mass field:

$$m(x, t) = m_0 + \Delta m(x, t) \quad (6)$$

where m_0 is the rest mass and $\Delta m(x, t)$ represents the dynamic mass fluctuations.

Kinetic Term of the Time Field

The time field itself requires a kinetic term:

$$\mathcal{L}_{\text{kin}}^{(T)} = \frac{1}{2}(\partial_\mu \Delta m)(\partial^\mu \Delta m) \quad (7)$$

This term describes the propagation of mass field fluctuations as dynamic degrees of freedom.

Mass Term of the Time Field

The time field has a characteristic mass m_T :

$$\mathcal{L}_{\text{mass}}^{(T)} = -\frac{1}{2}m_T^2 \Delta m^2 \quad (8)$$

The time field mass m_T is given through the Higgs-time field connection by:

$$m_T = \frac{\lambda}{\xi} \quad (9)$$

where λ is the base mass of the respective sector (m_p in the fundamental sector).
(Corrected on 30 Sept. 2026: previously "Higgs coupling parameter" – dimensionally λ must be a mass, see Addendum R91.)

Mass-Proportional Coupling

The fundamental new term in the T0 theory is the coupling of the lepton fields to the time field, proportional to the lepton mass:

$$\mathcal{L}_{\text{int}} = g_T^\ell \bar{\psi}_\ell \psi_\ell \Delta m \quad (10)$$

The coupling strength is given by:

$$\boxed{g_T^\ell = \xi m_\ell} \quad (11)$$

This mass-proportional coupling is the core of the T0 theory. It implies:

- Heavier leptons couple more strongly to the time field
- The coupling scales linearly with mass
- This leads to quadratic mass scaling in quantum corrections

Complete Extended Lagrangian

The complete T0 Lagrangian combines all terms:

$$\begin{aligned}
 \mathcal{L}_{T0} = & -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} \\
 & + \bar{\psi}_\ell(i\gamma^\mu D_\mu - m_\ell)\psi_\ell \\
 & + \frac{1}{2}(\partial_\mu \Delta m)(\partial^\mu \Delta m) \\
 & - \frac{1}{2}m_T^2 \Delta m^2 \\
 & + \xi m_\ell \bar{\psi}_\ell \psi_\ell \Delta m
 \end{aligned} \tag{12}$$

.3 Quantum Corrections and Feynman Rules

Feynman Rules from the Lagrangian

From the interaction term (10), the Feynman rules follow:

Vertex factor:


$$\bar{\psi}_\ell \psi_\ell \Delta m \longrightarrow -ig_T^\ell = -i\xi m_\ell \tag{13}$$

Time field propagator:

$$\Delta m(k) \longrightarrow \frac{i}{k^2 - m_T^2 + i\epsilon} \tag{14}$$

One-Loop Diagram

The fundamental one-loop diagram for the anomalous magnetic moment contribution has the structure:



General Formula for Scalar Mediators

For a scalar mediator with mass m_T and coupling g_T^ℓ , the general one-loop formula reads:

$$\Delta a_\ell = \frac{(g_T^\ell)^2}{8\pi^2} \int_0^1 dx \frac{m_\ell^2(1-x)(1-x^2)}{m_\ell^2 x^2 + m_T^2(1-x)} \tag{15}$$

This formula is standard in quantum field theory for scalar contributions to the anomalous magnetic moment. (Note of 30 Sept. 2026: numerator and denominator come from two different parametrizations. The standard form for a scalar mediator is $\Delta a_\ell = \frac{(g_T^\ell)^2}{8\pi^2} \int_0^1 dx \frac{m_\ell^2(1-x)^2(1+x)}{m_\ell^2(1-x)^2 + m_T^2 x}$; in the limit $m_T \gg m_\ell$ it gives $\frac{(g_T^\ell)^2 m_\ell^2}{8\pi^2 m_T^2} (\ln \frac{m_T^2}{m_\ell^2} - \frac{7}{6})$, not $\frac{2}{3}$ (numerically for $m_T/m_\ell = 1000$: 12.65 instead of 0.667). The factor $2/3$ in the following section, also in the note there, and hence the prefactor of the T0 formula rest on the mixed integrand; the m_ℓ^2 scaling up to the logarithm remains.)

.4 Derivation of the Fundamental T0 Formula

Heavy Mediator Limit

For $m_T \gg m_\ell$, the integral (15) can be simplified. In the denominator, the term $m_T^2(1-x)$ dominates:

$$\frac{m_\ell^2(1-x)(1-x^2)}{m_\ell^2 x^2 + m_T^2(1-x)} \approx \frac{m_\ell^2(1-x)(1-x^2)}{m_T^2(1-x)} = \frac{m_\ell^2(1-x^2)}{m_T^2} \quad (16)$$

This yields:

$$\begin{aligned} \Delta a_\ell &\approx \frac{(g_T^\ell)^2 m_\ell^2}{8\pi^2 m_T^2} \int_0^1 dx (1-x^2) \\ &= \frac{(g_T^\ell)^2 m_\ell^2}{8\pi^2 m_T^2} \left[x - \frac{x^3}{3} \right]_0^1 \\ &= \frac{(g_T^\ell)^2 m_\ell^2}{8\pi^2 m_T^2} \left(1 - \frac{1}{3} \right) \\ &= \frac{(g_T^\ell)^2 m_\ell^2}{8\pi^2 m_T^2} \cdot \frac{2}{3} \end{aligned} \quad (17)$$

(Corrected on 30 Sept. 2026: previously without the factor m_ℓ^2 – it appears in Eq. (16) and must not be lost when it is pulled in front of the integral.)

Inserting the T0 Coupling

With the mass-proportional coupling (11), we obtain:

$$\begin{aligned} \Delta a_\ell &= \frac{\xi^2 m_\ell^2}{8\pi^2 m_T^2} \cdot \frac{2}{3} \\ &= \frac{2\xi^2 m_\ell^2}{24\pi^2 m_T^2} \\ &= \frac{\xi^2 m_\ell^2}{12\pi^2 m_T^2} \end{aligned} \quad (18)$$

(Corrected on 30 Sept. 2026: previously first line $(\xi m_\ell)^2 / (8\pi^2 m_T^2) \cdot 2/3$ – there the coupling ξm_ℓ replaced the factor m_ℓ^2 lost in Eq. (17). With m_ℓ^2 from the loop integral, the result is dimensionless and quadratic in m_ℓ only if the coupling in the interaction term acts as the dimensionless ξ ; $g_T^\ell = \xi m_\ell$ would have the dimension of a mass and would give m_ℓ^4 with $[\Delta a] = E^2$. The mass proportionality thus resides in the loop factor m_ℓ^2 . The result of this line is unchanged.)

Higgs-Time Field Connection

With the time field mass (9), this becomes:

$$\begin{aligned} \Delta a_\ell &= \frac{\xi^2 m_\ell^2}{12\pi^2 (\lambda/\xi)^2} \\ &= \frac{\xi^2 m_\ell^2}{12\pi^2} \cdot \frac{\xi^2}{\lambda^2} \\ &= \frac{\xi^4 m_\ell^2}{12\pi^2 \lambda^2} \end{aligned} \quad (19)$$

Final Formula

Eq. (19) is already the final formula of the limit $m_T \gg m_\ell$:

$$\Delta a_\ell^{(T0)} = \frac{\xi^4}{12\pi^2\lambda^2} \cdot m_\ell^2 \quad (20)$$

(Corrected on 30 Sept. 2026: previously a section "Correction through the Complete Integral" with $\int_0^1 (1-x)(1-x^2) dx = 5/12$ and final formula $5\xi^4 m_\ell^2 / (96\pi^2 \lambda^2)$ – in the limit $m_T \gg m_\ell$ the factor $(1-x)$ cancels against the denominator (Eq. (16)); the integral is $\int_0^1 (1-x^2) dx = 2/3$, as computed above. $5/12$ would be the numerator without the denominator.)

Δa_ℓ is dimensionless exactly when λ is a mass – this is the reading of Addendum R91: $\lambda = M_{\text{base}}$, in the fundamental sector $M_{\text{base}} = m_p$.

.5 Numerical Evaluation

Determination of the Normalization Constant

The normalization constant reads:

$$C_{T0} = \frac{\xi^4}{12\pi^2\lambda^2} \quad (21)$$

With $\xi^4 = 3.160 \times 10^{-16}$ and $\lambda = m_p = 1.221 \times 10^{19} \text{ GeV}$ (Addendum R91) we obtain:

$$C_{T0} = \frac{3.160 \times 10^{-16}}{118.4 \times 1.491 \times 10^{38} \text{ GeV}^2} = 1.79 \times 10^{-56} \text{ GeV}^{-2} \quad (22)$$

(Corrected on 30 Sept. 2026: previously $C_{T0} \approx 2.246 \times 10^{-13} \text{ GeV}^{-2}$, supposedly from $\lambda_h \approx 0.13$, v , m_h and $\xi = \lambda_h^2 v^2 / (16\pi^3 m_h^2)$ – this value follows from none of the stated parameters: with $\lambda = 0.13$ the formula would give 1.6×10^{-16} (dimensionless), and $2.246 \times 10^{-13} \text{ GeV}^{-2}$ would require $\lambda = 2.7 \text{ MeV}$. The relation $\xi = \lambda_h^2 v^2 / (16\pi^3 m_h^2) = 1.32 \times 10^{-4}$ is independent of it.)

Final T0 Contribution Formula

The fully evaluated T0 contribution formula reads:

$$\Delta a_\ell^{(T0)} = 1.79 \times 10^{-56} \cdot m_\ell^2 [\text{GeV}^{-2}] \quad (23)$$

where m_ℓ is to be inserted in GeV.

Lepton-Specific Predictions

With the lepton masses:

- $m_e = 0.511 \text{ MeV} = 0.000511 \text{ GeV}$
- $m_\mu = 105.658 \text{ MeV} = 0.105658 \text{ GeV}$
- $m_\tau = 1776.86 \text{ MeV} = 1.77686 \text{ GeV}$

the T0 contributions are:

Electron:

$$\begin{aligned} \Delta a_e^{(T0)} &= 1.79 \times 10^{-56} \cdot (0.000511)^2 \\ &= 1.79 \times 10^{-56} \cdot 2.611 \times 10^{-7} \\ &= 4.7 \times 10^{-63} \end{aligned} \quad (24)$$

Muon:

$$\begin{aligned}\Delta a_\mu^{(T0)} &= 1.79 \times 10^{-56} \cdot (0.105658)^2 \\ &= 1.79 \times 10^{-56} \cdot 1.1164 \times 10^{-2} \\ &= 2.0 \times 10^{-58}\end{aligned}\tag{25}$$

Tau:

$$\begin{aligned}\Delta a_\tau^{(T0)} &= 1.79 \times 10^{-56} \cdot (1.77686)^2 \\ &= 1.79 \times 10^{-56} \cdot 3.1572 \\ &= 5.7 \times 10^{-56}\end{aligned}\tag{26}$$

(Corrected on 30 Sept. 2026: previously, with $C_{T0} = 2.246 \times 10^{-13}$, the values 5.86×10^{-20} , 2.51×10^{-15} and 7.09×10^{-13} .)

No Conversion Needed

The anomalous magnetic moment $a_\ell = (g_\ell - 2)/2$ is dimensionless; the values above are already the values to be compared with experiment (often quoted in units of 10^{-11}). The Lagrangian contribution of the time field is thus many orders of magnitude below any measurement precision.

(Corrected on 30 Sept. 2026: previously a section "Conversion to Conventional Units" that multiplied all values by 10^6 : $\Delta a_\mu = 2.51 \times 10^{-9} = 251 \times 10^{-11}$, $\Delta a_e = 5.86 \times 10^{-14}$, $\Delta a_\tau = 7.09 \times 10^{-7}$ – a dimensionless quantity does not change under a conversion, and a change MeV/GeV gives no factor either. 251×10^{-11} was the muon anomaly as of 2021; constant and factor produced this value only together. Since 2025 the anomaly is no longer significant (Doc. 382); a negligible Lagrangian contribution does not contradict this.)

.6 Quadratic Mass Scaling

Fundamental Prediction

The central prediction of the T0 theory is the quadratic mass scaling (23):

$$\Delta a_\ell^{(T0)} \propto m_\ell^2\tag{27}$$

This leads to natural hierarchies:

$$\frac{\Delta a_e^{(T0)}}{\Delta a_\mu^{(T0)}} = \left(\frac{m_e}{m_\mu}\right)^2 = \left(\frac{0.511}{105.658}\right)^2 = 2.34 \times 10^{-5}\tag{28}$$

$$\frac{\Delta a_\tau^{(T0)}}{\Delta a_\mu^{(T0)}} = \left(\frac{m_\tau}{m_\mu}\right)^2 = \left(\frac{1776.86}{105.658}\right)^2 = 282.8\tag{29}$$

Physical Interpretation

The quadratic mass scaling has deep physical significance:

Cause: The coupling is mass-proportional $g_T^\ell = \xi m_\ell$. In the one-loop contribution, $(g_T^\ell)^2 = \xi^2 m_\ell^2$ appears.

Consequences:

- Electron effects are smaller by $\sim 10^{-5}$ relative to the muon

- Muon effects are of order $\sim 10^{-58}$ and not measurable
- Tau effects are relatively the largest ($\sim 280\times$ larger than muon), in absolute terms also negligible

(Corrected on 30 Sept. 2026: previously "Muon effects are measurable ($\sim 10^{-9}$)" and "Tau effects are dominant" – consequence of the factor 10^6 and the constant 2.246×10^{-13} ; the ratios 2.34×10^{-5} and 282.8 remain valid.)

Comparison with QED: In the Standard Theory, the leading contribution scales as $\alpha/(2\pi)$, independent of the mass. The T0 theory adds a mass-dependent contribution.

.7 Connection to the Geometric Formulation

Relation to the g-2 Analysis

The T0 contributions $\Delta a_\ell^{(T0)}$ derived here are additional contributions to the Standard Model. They are **not** the same quantities as the values geometrically derived in Document 018: Doc. 018 obtains $\Delta a(\mu-e) = 4.37 \times 10^{-6}$ from $4\pi/f^p$, whereas here a contribution $\propto m_\ell^2$ of $\sim 10^{-58}$ results. The load-bearing g-2 line is the ratio line in Docs. 018 and 158. *(Corrected on 30 Sept. 2026: previously "They correspond to the values geometrically derived in Document 018" – the two formulations give contributions differing by many orders of magnitude, with different mass dependence.)*

The difference in notation:

- **Document 018:** Calculates directly a_ℓ (including SM + T0)
- **This document:** Calculates $\Delta a_\ell^{(T0)}$ (T0 contribution only)

The relation is:

$$a_\ell^{(\text{total})} = a_\ell^{(\text{SM})} + \Delta a_\ell^{(T0)} \quad (30)$$

Parameter Correspondences

The geometric parameters from Document 018 correspond to the Lagrangian parameters:

$$\xi = \frac{4}{3} \times 10^{-4} \quad (\text{identical in both formulations}) \quad (31)$$

$$f = 7500 \quad (\text{geometric sub-Planck factor}) \quad (32)$$

$$\varphi = \frac{1 + \sqrt{5}}{2} \quad (\text{pentagonal symmetry breaking}) \quad (33)$$

What the two formulations share is the parameter ξ ; there is no numerical connection between the contributions (see above). The projection factor k_{geom} belongs solely to the geometric 4D-3D projection in Doc. 018.

.8 Higgs Mechanism in the T0 Theory

Higgs Field as Fundamental Basis

In the T0 theory, the Higgs field is not an additional field but the fundamental basis of the time-mass duality:

$$T(x, t) \cdot m(x, t) = 1 \quad (34)$$

The universal parameter ξ follows directly from the Higgs parameters:

$$\xi = \frac{\lambda_h^2 v^2}{16\pi^3 m_h^2} \quad (35)$$

with:

- λ_h : Higgs self-coupling
- $v = 246$ GeV: Higgs vacuum expectation value
- $m_h = 125$ GeV: Higgs mass

Spontaneous Symmetry Breaking

The spontaneous symmetry breaking of the Higgs field generates:

- Lepton masses through Yukawa coupling
- The time field $T(x, t)$ as fluctuations around the VEV
- The time-mass duality as fundamental structure

In this picture, mass fluctuations Δm are directly connected to Higgs fluctuations:

$$\Delta m(x, t) = y_\ell \cdot \delta h(x, t) \quad (36)$$

where y_ℓ is the Yukawa coupling and δh is the Higgs fluctuation.

.9 Field-Theoretic Structure

Equations of Motion

From the Lagrangian (12), the Euler-Lagrange equations follow:

For the lepton field:

$$(i\gamma^\mu D_\mu - m_\ell + \xi m_\ell \Delta m)\psi_\ell = 0 \quad (37)$$

For the time field:

$$\square \Delta m + m_T^2 \Delta m = \xi m_\ell \bar{\psi}_\ell \psi_\ell \quad (38)$$

where $\square = \partial_\mu \partial^\mu$ is the d'Alembert operator.

Interpretation of the Equations of Motion

Equation (37) shows that the lepton has an effective mass:

$$m_{\text{eff}}(x, t) = m_\ell(1 - \xi \Delta m) \quad (39)$$

(Corrected on 30 Sept. 2026: previously $-\xi m_\ell \Delta m$ in the equation of motion and $m_{\text{eff}} = m_\ell(1 + \xi \Delta m)$ – from $\mathcal{L}_{\text{int}} = +\xi m_\ell \bar{\psi}_\ell \psi_\ell \Delta m$ follows $\partial \mathcal{L} / \partial \bar{\psi}_\ell = (i\gamma^\mu D_\mu - m_\ell + \xi m_\ell \Delta m)\psi_\ell$, hence $m_{\text{eff}} = m_\ell(1 - \xi \Delta m)$.)
The mass is modulated by the time field.

Equation (38) shows that the time field has a source:

$$\text{Source} = \xi m_\ell \bar{\psi}_\ell \psi_\ell = \xi m_\ell \rho_\ell \quad (40)$$

The lepton density ρ_ℓ generates time field fluctuations, proportional to the lepton mass.

Energy Scales

The characteristic energy scales in the theory are:

$$E_{\text{Lepton}} \sim m_\ell \quad (\text{lepton mass}) \quad (41)$$

$$E_{\text{Time}} \sim m_T = \frac{\lambda}{\xi} \quad (\text{time field mass}) \quad (42)$$

$$E_{\text{Coupling}} \sim \xi m_\ell \ll m_\ell \quad (\text{weak coupling}) \quad (43)$$

The hierarchy $E_{\text{Coupling}} \ll E_{\text{Lepton}} \ll E_{\text{Time}}$ justifies the perturbative treatment.

.10 Renormalization (historical presentation)

Note: This section uses the language of QFT renormalization (wavefunction renormalization, β -function, RG equations) as a descriptive tool. In the complete FFGFT formulation (Doc. 202), traditional renormalization is not required: the fractal torus provides a natural UV cutoff at E_p/ξ , and the scale structure follows from the recursion $\mathcal{R}$. The formulae used here are compatible with the FFGFT structure, but the terminology is closer to SM language than to native FFGFT language.

Divergences in Loop Diagrams

The one-loop integral (15) is finite for $m_T \gg m_\ell$. However, higher orders may contain divergences.

The renormalization of the T0 theory requires:

- Wave function renormalization for ψ_ℓ and Δm
- Mass renormalization for m_ℓ and m_T
- Coupling renormalization for g_T^ℓ

Renormalization Group Equations

The running coupling follows:

$$\mu \frac{dg_T^\ell}{d\mu} = \beta_{g_T}(g_T^\ell) \quad (44)$$

Due to the mass-proportional structure (11):

$$\beta_{g_T} = \xi \beta_m \quad (45)$$

where β_m is the anomalous mass dimension.

Consistency with the Standard Model

The T0 contributions are subdominant compared to the Standard Model contributions:

$$\frac{\Delta a_\ell^{(\text{T0})}}{a_\ell^{(\text{QED})}} \sim \frac{\xi^4 m_\ell^2}{6\pi\alpha\lambda^2} \ll 1 \quad (46)$$

(Corrected on 30 Sept. 2026: previously $\xi^4 m_\ell^2/\alpha$ – not dimensionless without λ^2 ; with $a^{(\text{QED})} \approx \alpha/(2\pi)$ and Eq. (20) the expression above follows.)

This guarantees consistency with experimental data and allows perturbative treatment.

.11 Comparison of Different Formulations

Lagrangian vs. Geometric Formulation

Aspect	Lagrangian (Doc. 019)	Geometric (Doc. 018)
Starting point	Time field $\Delta m(x, t)$	Torsion lattice, windings
Method	Field theory, loop integrals	Fractal geometry
Main formula	$\Delta a \propto \xi^4 m^2$	$a_e = S_3 / (f k_{\text{eff}})$, $\Delta a = 4\pi / f^p$
Prediction	T0 contribution Δa	Total value a
Parameters	ξ, λ, m_T	ξ, φ, f

Table 1: Comparison of formulations

Complementarity

The two formulations are complementary:

- **Lagrangian:** Provides the field-theoretic structure and Feynman rules
- **Geometric:** Provides the physical intuition and ratio predictions

The numerical predictions for a_ℓ come from the geometric formulation (Docs. 018, 158); the Lagrangian contribution of the time field is negligibly small. (*Corrected on 30 Sept. 2026: previously "Both lead to consistent numerical predictions with ~2 % precision" – see the section "Relation to the g-2 Analysis".*)

Bibliography

Bibliography

- [1] J. Pascher, *Anomalous Magnetic Moments in the FFGFT Theory: Geometric Derivation from the Time-Mass Duality*, [Document 018_T0_Anomale-g2-10_De.pdf](#), February 2026.
- [2] M. E. Peskin, D. V. Schroeder, *An Introduction to Quantum Field Theory*, Westview Press, 1995. Standard reference for Feynman rules and loop calculations.
- [3] Particle Data Group, *Review of Particle Physics*, Prog. Theor. Exp. Phys. 2024, 083C01 (2024). Experimental values for lepton masses and g-2.
- [4] P. W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, Phys. Rev. Lett. 13, 508 (1964). Original Higgs mechanism.
- [5] S. Weinberg, *The Quantum Theory of Fields, Volume I: Foundations*, Cambridge University Press, 1995. Comprehensive treatment of quantum field theory.

Addendum R91: Label correction $m_T = \lambda/\xi$ (7 September 2026)

The label λ as “Higgs coupling parameter” in the formula $m_T = \lambda/\xi$ is **misleading** (R91, Doc. 190-Archive-2).

Three reasons: (i) Dimensionally, λ must be a *mass*, since m_T is a mass and ξ is dimensionless; the Higgs quartic $\lambda_h \approx 0.129$ is dimensionless. (ii) λ_h runs under the RGE and is a low-energy parameter of the electroweak scale; a value measured there cannot be inserted at the Planck scale. (iii) Interpreted as a dimensionless factor in Planck units, λ_h would give range $7.75 \cdot L_0$ — misses L_0 .

Correct: $\lambda = M_{\text{basis}}$ is the basis mass of the respective sector; in the fundamental sector $M_{\text{basis}} = m_P$ (R89/R90, Doc. 180). The formula reads structurally $m_T = M_{\text{basis}}/\xi$, sector-dependent value, universal form.