

Universal Derivation of All Physical Constants from the Fine-Structure Constant and Planck Length

Contents

.1	Introduction and Basic Principle	3
	The Minimal Principle of Physics	3
	SI Base Definitions	3
.2	Derivation of Fundamental Constants	3
	Speed of Light c	3
	Vacuum Permittivity ϵ_0	4
	Reduced Planck Constant $\hbar$	4
	Gravitational Constant G	5
.3	Complete Planck Units	5
	Planck Time	5
	Planck Mass	5
	Planck Energy	5
	Planck Temperature	5
.4	Atomic and Molecular Constants	5
	Classical Electron Radius	5
	Compton Wavelength of the Electron	6
	Bohr Radius	6
	Rydberg Constant	6
.5	Thermodynamic Constants	6
	Stefan-Boltzmann Constant	6
	Wien's Displacement Law Constant	6
.6	Dimensional Analysis and Verification	6
	Consistency Check of the Fine-Structure Constant	6
	Consistency Check of the Gravitational Constant	7
	Consistency Check of $\hbar$	7
.7	The Characteristic Energy E_0 and T_0 Theory	7
	Definition of the Characteristic Energy	7
	Numerical Evaluation with Different Precision Levels	7
	Level 1: Rounded Standard Values	7
	Level 2: CODATA 2018 Precision Values	8
	Level 3: The Value $E_0 = 7.398 \text{ MeV}$	8
	Precise Fine-Structure Constant Calculation	8
	Comparison of Calculation Accuracy	8
	Detailed Calculation with $E_0 = 7.398 \text{ MeV}$	9
.8	On the Precision of the Value $E_0 = 7.398 \text{ MeV}$	9
	Why $E_0 = 7.398 \text{ MeV}$ Works Well	9
	The Mathematical Justification	9
.9	Comparison with Alternative Approaches	10
	Estimation with T_0 -Calculated Masses	10
	Correct Interpretation	10
.10	Dimensional Consistency of the E_0 Formula	10

	Correct Dimensionless Formulation	10
	Alternative Notation	10
.11	On the Circularity Objections	11
	The Apparent Circularity Objections	11
	Resolution of the Apparent Circularity	11
	The Structure of SI Definitions (since 2019)	11
	Corrected Hierarchy with Modern SI	11
	ℓ_P is Only ONE Possible Length Scale	12
	The Mathematics Works with ANY Length Scale	12
	The SI Reference is What Matters	12
	The Hierarchy in the T0 Interpretation	12
	Possible Independent Approaches	13
	Independent Measurement of ℓ_P	13
	Independent Measurement of α	13
	Formal Check of the Order of Definitions	13
	Definition Hierarchy	13
	Circularity Test	13
	The Philosophical Argument	14
	Reference Scales are Necessary	14
.12	Further Considerations	14
	Connection to the T0 Model	14

Correction notes

Corrections from the calculation review (30 Sept. 2026). The following places are corrected or annotated in the text; each carries a dated note there with the previous value:

- Speed of light: note – what carries is $c = \ell_P/t_P$ ($c = 1$); 299 792 458 m/s is the SI anchor, not the result of the dimensional analysis.
- α^{-1} with $E_0 = 7.398$ MeV (table, calculation, comparison box, alternative form): 137.038 $\rightarrow$ 137.035 (with the rounded 54.73: 137.036); deviation 0.0014 % $\rightarrow$ 0.0005 %.
- Table $G(L)$: $1.04 \times 10^{-10} \rightarrow 1.60 \times 10^{-10}$, $1.67 \times 10^{-11} \rightarrow 2.55 \times 10^{-11}$, $1.64 \times 10^{-10} \rightarrow 2.52 \times 10^{-10}$ (scaled with ℓ_P instead of 2.0×10^{-35} m).
- Table of calculation accuracy, upper rows: 139.15 $\rightarrow$ 138.91 (1.5 $\rightarrow$ 1.37 %), 139.07 $\rightarrow$ 138.89 (1.4 $\rightarrow$ 1.35 %).
- $E_0^{(2)} = \sqrt{0.5109989461 \times 105.6583745}$: 7.348566 $\rightarrow$ 7.347878 MeV; table row CODATA exact 138.89/1.35 % $\rightarrow$ 138.91/1.37 %.
- Interpretation of $E_0 = 7.398$ MeV: geometric mean of refined masses (QED, weak corrections) $\rightarrow$ fractally corrected mean $\sqrt{m_e m_\mu / K_{\text{frak}}}$ (R72).
- $\alpha = \xi E_0^2$ with $E_0 = \sqrt{m_e m_\mu} \rightarrow E_0 = \sqrt{m_e m_\mu / K_{\text{frak}}}$ (without K_{frak} : 138.91 instead of 137.06).
- Estimate with T0 masses: $\sqrt{0.511000 \times 105.658000} = 72.868 \rightarrow 7.348$ MeV; “obviously flawed” dropped. μ_0 : no longer exactly $4\pi \times 10^{-7}$ H/m but measured since SI 2019, $1.25663706127(20) \times 10^{-6}$ N/A²; note on the order of calculation (ε_0 from μ_0). Unit in the $\hbar$ check: C²·s/(kg·m³) $\rightarrow$ C²·s²/(kg·m³).

Abstract

This document shows how the physical constants treated here can be calculated in SI units from two experimental base quantities together with the SI definitions (and, for atomic quantities, the electron mass) - the dimensionless fine-structure constant $\alpha = 1/137.036$ and the Planck length $\ell_P = 1.616255 \times 10^{-35}$ m. Additionally, the value of the characteristic energy E_0 in T0 theory is discussed: the geometric mean of the CODATA electron and muon masses gives $E_0 = 7.3486$ MeV; the value $E_0 = 7.398$ MeV used for α is interpreted here as the geometric mean of corrected masses. Two common circularity objections are discussed. The calculation reduces the number of natural constants that must be entered independently to two experimental values plus SI conventions, and the T0 raw values reproduce mass ratios without correction to about 0.5 %.

Note on versions. The German version of this document is more extensive than this English one. Earlier revisions shortened or changed parts of the English version. Where the two differ, the more extensive German version applies.

Status markers. Statements marked **[S]** are *speculative*: a hint or conjecture that has not yet been derived or numerically checked in the corpus.

.1 Introduction and Basic Principle

The Minimal Principle of Physics

In modern physics, about 30 different natural constants appear to need independent experimental determination. This work shows that, together with the SI definitions, the constants treated here can be calculated from **two experimental values**:

Fundamental Input Data

- **Fine-structure constant:** $\alpha = \frac{1}{137.035999084}$ (dimensionless)
- **Planck length:** $\ell_P = 1.616255 \times 10^{-35}$ m

SI Base Definitions

Additionally, we use the modern SI base definitions (since 2019):

$$\mu_0 = 1.25663706127(20) \times 10^{-6} \text{ N/A}^2 \quad (\text{measured, CODATA 2022; exactly } 4\pi \times 10^{-7} \text{ H/m until 2019}) \quad (1)$$

$$e = 1.602176634 \times 10^{-19} \text{ C} \quad (\text{exact definition}) \quad (2)$$

$$k_B = 1.380649 \times 10^{-23} \text{ J/K} \quad (\text{exact definition}) \quad (3)$$

$$N_A = 6.02214076 \times 10^{23} \text{ mol}^{-1} \quad (\text{exact definition}) \quad (4)$$

.2 Derivation of Fundamental Constants

Speed of Light c

The speed of light follows from the relationship between Planck units. Since the Planck length is defined as:

$$\ell_P = \sqrt{\frac{\hbar G}{c^3}} \quad (5)$$

and all Planck units are interconnected through $\hbar$, G and c , dimensional analysis yields:

Speed of Light

$$c = 2.99792458 \times 10^8 \text{ m/s} \quad (6)$$

(Note of 30 Sept. 2026: What carries is the structure, $c = \ell_P/t_P$ as the ratio of length and time in the time-mass duality (Doc. 101, 145), in natural units $c = 1$. The number 299 792 458 m/s is the SI definition of the metre (1983) and serves as the conversion anchor; it is not the result of the dimensional analysis but expresses c in SI, see the section on the SI definitions.)

Vacuum Permittivity ϵ_0

From the Maxwell relation $\mu_0 \epsilon_0 = 1/c^2$ follows:

$$\epsilon_0 = \frac{1}{\mu_0 c^2} = \frac{1}{4\pi \times 10^{-7} \times (2.99792458 \times 10^8)^2} \quad (7)$$

Vacuum Permittivity

$$\epsilon_0 = 8.8541878128 \times 10^{-12} \text{ F/m} \quad (8)$$

(Note of 30 Sept. 2026: Since SI 2019, μ_0 is no longer exactly $4\pi \times 10^{-7} \text{ H/m}$ but measured: $\mu_0 = 2\alpha h/(e^2 c) = 1.25663706127(20) \times 10^{-6} \text{ N/A}^2$ (CODATA 2022); exact are h , e , c , k_B , N_A . The SI order is therefore reversed: $\epsilon_0 = e^2/(4\pi \hbar c \alpha)$ from the measured α , then $\mu_0 = 1/(\epsilon_0 c^2)$, as in the section on the SI definitions. The number $8.8541878128 \times 10^{-12} \text{ F/m}$ (CODATA 2018) differs from the $4\pi \times 10^{-7}$ value only by $\sim 10^{-10}$ relative.)

Reduced Planck Constant $\hbar$

The fine-structure constant is defined as:

$$\alpha = \frac{e^2}{4\pi \epsilon_0 \hbar c} \quad (9)$$

Solving for $\hbar$:

$$\hbar = \frac{e^2}{4\pi \epsilon_0 c \alpha} \quad (10)$$

Substituting known values:

$$\hbar = \frac{(1.602176634 \times 10^{-19})^2}{4\pi \times 8.8541878128 \times 10^{-12} \times 2.99792458 \times 10^8 \times \frac{1}{137.035999084}} \quad (11)$$

Reduced Planck Constant

$$\hbar = 1.054571817 \times 10^{-34} \text{ J}\cdot\text{s} \quad (12)$$

Gravitational Constant G

From the definition of the Planck length follows:

$$G = \frac{\ell_P^2 c^3}{\hbar} \quad (13)$$

Substituting calculated values:

$$G = \frac{(1.616255 \times 10^{-35})^2 \times (2.99792458 \times 10^8)^3}{1.054571817 \times 10^{-34}} \quad (14)$$

Gravitational Constant

$$G = 6.67430 \times 10^{-11} \text{ m}^3/(\text{kg} \cdot \text{s}^2) \quad (15)$$

.3 Complete Planck Units

With $\hbar$, c and G , all Planck units can be calculated:

Planck Time

$$t_P = \sqrt{\frac{\hbar G}{c^5}} = \frac{\ell_P}{c} = 5.391247 \times 10^{-44} \text{ s} \quad (16)$$

Planck Mass

$$m_P = \sqrt{\frac{\hbar c}{G}} = 2.176434 \times 10^{-8} \text{ kg} \quad (17)$$

Planck Energy

$$E_P = m_P c^2 = \sqrt{\frac{\hbar c^5}{G}} = 1.956082 \times 10^9 \text{ J} = 1.220890 \times 10^{19} \text{ GeV} \quad (18)$$

Planck Temperature

$$T_P = \frac{E_P}{k_B} = \frac{m_P c^2}{k_B} = 1.416784 \times 10^{32} \text{ K} \quad (19)$$

.4 Atomic and Molecular Constants

Classical Electron Radius

With the electron mass $m_e = 9.1093837015 \times 10^{-31} \text{ kg}$:

$$r_e = \frac{e^2}{4\pi\epsilon_0 m_e c^2} = \frac{\alpha \hbar}{m_e c} = 2.817940 \times 10^{-15} \text{ m} \quad (20)$$

Compton Wavelength of the Electron

$$\lambda_{C,e} = \frac{h}{m_e c} = \frac{2\pi\hbar}{m_e c} = 2.426310 \times 10^{-12} \text{ m} \quad (21)$$

Bohr Radius

$$a_0 = \frac{4\pi\epsilon_0\hbar^2}{m_e e^2} = \frac{\hbar}{m_e c \alpha} = 5.291772 \times 10^{-11} \text{ m} \quad (22)$$

Rydberg Constant

$$R_\infty = \frac{\alpha^2 m_e c}{2h} = \frac{\alpha^2 m_e c}{4\pi\hbar} = 1.097373 \times 10^7 \text{ m}^{-1} \quad (23)$$

.5 Thermodynamic Constants

Stefan-Boltzmann Constant

$$\sigma = \frac{2\pi^5 k_B^4}{15h^3 c^2} = \frac{2\pi^5 k_B^4}{15(2\pi\hbar)^3 c^2} = 5.670374419 \times 10^{-8} \text{ W/(m}^2 \cdot \text{K}^4) \quad (24)$$

Wien's Displacement Law Constant

$$b = \frac{hc}{k_B} \times \frac{1}{4.965114231} = 2.897771955 \times 10^{-3} \text{ m} \cdot \text{K} \quad (25)$$

.6 Dimensional Analysis and Verification

Consistency Check of the Fine-Structure Constant

$$[\alpha] = \frac{[e^2]}{[\epsilon_0][\hbar][c]} \quad (26)$$

$$= \frac{[\text{C}^2]}{[\text{F/m}][\text{J} \cdot \text{s}][\text{m/s}]} \quad (27)$$

$$= \frac{[\text{C}^2]}{[\text{C}^2 \cdot \text{s}^2 / (\text{kg} \cdot \text{m}^3)][\text{J} \cdot \text{s}][\text{m/s}]} \quad (28)$$

$$= \frac{[\text{C}^2]}{[\text{C}^2 / (\text{kg} \cdot \text{m}^2 / \text{s}^2)]} \quad (29)$$

$$= [1] \quad \checkmark \quad (30)$$

Consistency Check of the Gravitational Constant

$$[G] = \frac{[\ell_p^2][c^3]}{[\hbar]} \quad (31)$$

$$= \frac{[m^2][m^3/s^3]}{[J \cdot s]} \quad (32)$$

$$= \frac{[m^5/s^3]}{[kg \cdot m^2/s^2 \cdot s]} \quad (33)$$

$$= \frac{[m^5/s^3]}{[kg \cdot m^2/s^3]} \quad (34)$$

$$= [m^3/(kg \cdot s^2)] \quad \checkmark \quad (35)$$

Consistency Check of $\hbar$

$$[\hbar] = \frac{[e^2]}{[\epsilon_0][c][\alpha]} \quad (36)$$

$$= \frac{[C^2]}{[F/m][m/s][1]} \quad (37)$$

$$= \frac{[C^2]}{[C^2 \cdot s^2/(kg \cdot m^3)][m/s]} \quad (38)$$

$$= \frac{[C^2 \cdot kg \cdot m^3]}{[C^2 \cdot s \cdot m]} \quad (39)$$

$$= [kg \cdot m^2/s] = [J \cdot s] \quad \checkmark \quad (40)$$

.7 The Characteristic Energy E_0 and T0 Theory

Definition of the Characteristic Energy

Basic Definition

The fundamental definition of the characteristic energy is:

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (41)$$

This is **not a derivation** and **not a fit** – it is the mathematical definition of the geometric mean of two masses.

Numerical Evaluation with Different Precision Levels

Level 1: Rounded Standard Values

With the often cited rounded masses:

$$m_e = 0.511 \text{ MeV} \quad (42)$$

$$m_\mu = 105.658 \text{ MeV} \quad (43)$$

$$E_0^{(1)} = \sqrt{0.511 \times 105.658} = \sqrt{53.99} = 7.348 \text{ MeV} \quad (44)$$

Level 2: CODATA 2018 Precision Values

With the precise experimental masses:

$$m_e = 0.510,998,946,1 \text{ MeV} \quad (45)$$

$$m_\mu = 105.658,374,5 \text{ MeV} \quad (46)$$

$$E_0^{(2)} = \sqrt{0.5109989461 \times 105.6583745} = 7.347,878 \text{ MeV} \quad (47)$$

(Corrected on 30 Sept. 2026: previously $7.348,566 \text{ MeV} - \sqrt{0.5109989461 \times 105.6583745} = \sqrt{53.99125} = 7.347878 \text{ MeV}$.)

Level 3: The Value $E_0 = 7.398 \text{ MeV}$

Critical Question

Is $E_0 = 7.398 \text{ MeV}$ a fitted parameter?

Answer in this document: It is not treated as a free fit parameter.

$E_0 = 7.398 \text{ MeV}$ is the fractally corrected geometric mean $\sqrt{m_e m_\mu / K_{\text{frak}}} = 7.3479 / \sqrt{0.98667} = 7.397 \text{ MeV}$ (Doc. 190, R72); the geometric mean of the CODATA masses themselves is 7.3479 MeV (Level 2).

(Corrected on 30 Sept. 2026: previously interpreted as the geometric mean of refined mass values including QED and weak corrections, and 7.3486 MeV as the mean of the CODATA masses – $7.398/7.348 = 1.0068 = 1/\sqrt{K_{\text{frak}}}$, the difference is the fractal correction, not a radiative correction; the mean is 7.3479 MeV .)

Precise Fine-Structure Constant Calculation

The dimensionally correct formula:

$$\alpha = \xi \cdot \frac{E_0^2}{(1 \text{ MeV})^2} \quad (48)$$

where:

- $\xi = \frac{4}{3} \times 10^{-4} = 1.333\bar{3} \times 10^{-4}$ (exact)
- $(1 \text{ MeV})^2$ is the normalization energy for dimensionless calculation

Comparison of Calculation Accuracy

E_0 Value	Source	α_{T0}^{-1}	Deviation
7.348 MeV	Rounded masses	138.91	1.37%
7.347,878 MeV	CODATA exact	138.91	1.37%
7.398 MeV	Optimized	137.035	0.0005%
Experiment (CODATA):		137.035999084	Reference

Table 1: Comparison of calculation accuracy for different E_0 values

(Corrected on 30 Sept. 2026: previously $139.15/1.5\%$ and $139.07/1.4\%$ in the two upper rows, 137.038 and 0.0014% in the 7.398 MeV row – with $\xi = \frac{4}{3} \times 10^{-4}$ and $E_0 = 7.398 \text{ MeV}$ one

obtains $\alpha^{-1} = 137.035$, deviation 0.0005 %; with $E_0 = 7.348$ or 7.348566 MeV one obtains 138.91 or 138.89, deviation 1.37 or 1.35 %.)

(Corrected on 30 Sept. 2026: previously 7.348,566 MeV, 138.89, 1.35 % in the CODATA exact row – the geometric mean of the CODATA masses is 7.347878 MeV, hence $1/(\frac{4}{3} \times 10^{-4} \times 53.9913) = 138.91$, deviation 1.37 %; with K_{frak} one obtains $0.98667 \times 138.91 = 137.06$.)

Detailed Calculation with $E_0 = 7.398$ MeV

$$E_0^2 = (7.398)^2 = 54.7303 \text{ MeV}^2 \quad (49)$$

$$\frac{E_0^2}{(1 \text{ MeV})^2} = 54.7303 \quad (50)$$

$$\alpha = 1.333\bar{3} \times 10^{-4} \times 54.7303 \quad (51)$$

$$= 7.297 \times 10^{-3} \quad (52)$$

$$\alpha^{-1} = 137.035 \quad (53)$$

Comparison with Experiment

TO Prediction: $\alpha^{-1} = 137.035$

Experiment: $\alpha^{-1} = 137.035999084$

Relative Deviation: $\frac{|137.0353 - 137.0360|}{137.036} = 0.0005\%$

(Corrected on 30 Sept. 2026: previously $\alpha^{-1} = 137.038$ and relative deviation 0.0014 % – $1/(1.333\bar{3} \times 10^{-4} \times 54.7304) = 137.0353$.)

.8 On the Precision of the Value $E_0 = 7.398$ MeV

Why $E_0 = 7.398$ MeV Works Well

According to the interpretation of this document, the value $E_0 = 7.398$ MeV is meant to result from:

1. **Inclusion of QED corrections** in particle masses
2. **Incorporation of weak interaction effects**
3. **Geometric mean calculation** with full precision
4. **Consistency** with TO geometry $\xi = \frac{4}{3} \times 10^{-4}$

The Mathematical Justification

Geometric Interpretation

The geometric mean $E_0 = \sqrt{m_e \cdot m_\mu}$ is the natural energy scale between electron and muon.

On a logarithmic scale, E_0 lies precisely in the middle:

$$\log(E_0) = \frac{\log(m_e) + \log(m_\mu)}{2} \quad (54)$$

This is the **characteristic energy** of the first two lepton generations.

.9 Comparison with Alternative Approaches

Estimation with T0-Calculated Masses

If the particle masses themselves were calculated from T0 theory:

$$m_e^{T0} = 0.511,000 \text{ MeV} \quad (\text{theoretical}) \quad (55)$$

$$m_\mu^{T0} = 105.658,000 \text{ MeV} \quad (\text{theoretical}) \quad (56)$$

$$E_0^{T0} = \sqrt{0.511000 \times 105.658000} = 7.348 \text{ MeV} \quad (57)$$

Result: This calculation gives $E_0 = 7.348 \text{ MeV}$, the bare value; the approach is not flawed. *(Corrected on 30 Sept. 2026: previously 72.868 MeV and "obviously flawed (much too large)" – $0.511000 \times 105.658000 = 53.991$, $\sqrt{53.991} = 7.348$; the error was in the arithmetic, not in the approach.)*

Correct Interpretation

The correct approach is:

1. Use **experimental masses** as input
2. Calculate **geometric mean** exactly
3. Use **T0 geometry** ξ as theoretical parameter
4. Check **fine-structure constant** as output

.10 Dimensional Consistency of the E_0 Formula

Correct Dimensionless Formulation

The formula:

$$\alpha = \xi \cdot \frac{E_0^2}{(1 \text{ MeV})^2} \quad (58)$$

is dimensionally consistent:

$$[\alpha] = [\xi] \cdot \frac{[E_0^2]}{[(1 \text{ MeV})^2]} \quad (59)$$

$$= [1] \cdot \frac{[\text{Energy}^2]}{[\text{Energy}^2]} \quad (60)$$

$$= [1] \quad \checkmark \quad (61)$$

Alternative Notation

Equivalently can be written:

$$\frac{1}{\alpha} = \frac{(1 \text{ MeV})^2}{\xi \cdot E_0^2} = \frac{1}{\xi \cdot 54.73} = \frac{1}{1.333 \times 10^{-4} \times 54.73} = 137.036 \quad (62)$$

(Corrected on 30 Sept. 2026: previously 137.038 – with $\xi = \frac{4}{3} \times 10^{-4}$ and the rounded value 54.73 one obtains 137.036, with 54.7304 exactly 137.035.)

.11 On the Circularity Objections

The Apparent Circularity Objections

Common Criticisms

Objection 1: The Planck length ℓ_P is already defined via the gravitational constant G :

$$\ell_P = \sqrt{\frac{\hbar G}{c^3}} \quad (63)$$

Therefore, it is circular to derive G from ℓ_P .

Objection 2: The speed of light c is calculated from μ_0 and ε_0 :

$$c = \frac{1}{\sqrt{\mu_0 \varepsilon_0}} \quad (64)$$

But ε_0 is calculated from c - that is circular.

Resolution of the Apparent Circularity

The Structure of SI Definitions (since 2019)

Modern SI Base

Since the SI reform in 2019, the following quantities are **exactly defined**:

$$c = 299792458 \text{ m/s} \quad (\text{exact definition}) \quad (65)$$

$$e = 1.602176634 \times 10^{-19} \text{ C} \quad (\text{exact definition}) \quad (66)$$

$$\hbar = 1.054571817 \times 10^{-34} \text{ J}\cdot\text{s} \quad (\text{exact definition}) \quad (67)$$

$$k_B = 1.380649 \times 10^{-23} \text{ J/K} \quad (\text{exact definition}) \quad (68)$$

μ_0 and ε_0 are no longer fixed exactly but follow from the measured α : $\mu_0 = 2\alpha\hbar/(e^2c) = 1.25663706127(20) \times 10^{-6} \text{ N/A}^2$ (CODATA 2022)

Corrected Hierarchy with Modern SI

The actual derivation is therefore:

$$\text{Given (experimental): } \alpha, \ell_P \quad (69)$$

$$\text{Defined (SI 2019): } c, e, \hbar, k_B \quad (70)$$

$$\text{Calculated: } \varepsilon_0 = \frac{e^2}{4\pi\hbar c\alpha} \quad (71)$$

$$\mu_0 = \frac{1}{\varepsilon_0 c^2} \quad (72)$$

$$G = \frac{\ell_P^2 c^3}{\hbar} \quad (73)$$

Result: For c , ε_0 and μ_0 there is no circularity, since c and $\hbar$ are directly defined. For G it should be noted that, in practice, ℓ_P has so far been determined from the measured G .

ℓ_P is Only ONE Possible Length Scale

The Planck length is not the only fundamental length scale. One could equally well use:

$$L_1 = 2.5 \times 10^{-35} \text{ m} \quad (\text{arbitrarily chosen}) \quad (74)$$

$$L_2 = 1.0 \times 10^{-35} \text{ m} \quad (\text{round number}) \quad (75)$$

$$L_3 = \pi \times 10^{-35} \text{ m} \quad (\text{with } \pi) \quad (76)$$

$$L_4 = e \times 10^{-35} \text{ m} \quad (\text{with } e) \quad (77)$$

The Mathematics Works with ANY Length Scale

The general formula is:

$$G = \frac{L^2 \times c^3}{\hbar} \quad (78)$$

Crucial: Only with the specific length $\ell_P = 1.616255 \times 10^{-35} \text{ m}$ does one obtain the correct experimental value of G .

The SI Reference is What Matters

Length Scale L	Calculated G	Status
$2.5 \times 10^{-35} \text{ m}$	$1.60 \times 10^{-10} \text{ m}^3/(\text{kg}\cdot\text{s}^2)$	Wrong
$1.0 \times 10^{-35} \text{ m}$	$2.55 \times 10^{-11} \text{ m}^3/(\text{kg}\cdot\text{s}^2)$	Wrong
$\pi \times 10^{-35} \text{ m}$	$2.52 \times 10^{-10} \text{ m}^3/(\text{kg}\cdot\text{s}^2)$	Wrong
$\ell_P = 1.616 \times 10^{-35} \text{ m}$	$6.674 \times 10^{-11} \text{ m}^3/(\text{kg}\cdot\text{s}^2)$	Correct

Table 2: G-values for different length scales

(Corrected on 30 Sept. 2026: previously 1.04×10^{-10} , 1.67×10^{-11} and 1.64×10^{-10} – these values were scaled with $2.0 \times 10^{-35} \text{ m}$ instead of $\ell_P = 1.616 \times 10^{-35} \text{ m}$ and were too small by a factor of 0.65; $G(L) = G(\ell_P)^2$.)

The Hierarchy in the T0 Interpretation

Interpretation in the T0 Framework

In the T0 interpretation, ℓ_P and G are both expressions of the same underlying geometry.

The order in this interpretation:

1. Fundamental 3D space geometry $\rightarrow \xi = \frac{4}{3} \times 10^{-4}$
2. From this follows ℓ_P as natural scale
3. From this follows G as emergent property
4. SI units provide the reference to human measures

Possible Independent Approaches

Independent Measurement of ℓ_P

The Planck length could in principle become accessible independently of G through **[S]** (no such measurement exists so far):

1. **Quantum gravity experiments:** Direct measurement of the minimal length scale
2. **Black hole Hawking radiation:** ℓ_P determines the evaporation rate
3. **Cosmological observations:** ℓ_P influences quantum fluctuations of inflation (cosmic sector, deliberately set aside in the current corpus, P39)
4. **High-energy scattering experiments:** At Planck energies, ℓ_P becomes directly accessible

Independent Measurement of α

The fine-structure constant is measured through:

1. **Quantum Hall effect:** $\alpha = \frac{e^2}{h} \times \frac{R_K}{Z_0}$
 2. **Anomalous magnetic moment:** α from QED corrections
 3. **Atom interferometry:** α from recoil measurements
 4. **Spectroscopy:** α from hydrogen spectrum
- None of these methods uses G or ℓ_P .

Formal Check of the Order of Definitions

Definition Hierarchy

$$\textbf{Given: } \alpha \text{ (experimental), } \ell_P \text{ (experimental)} \quad (79)$$

$$\textbf{Defined: } \mu_0 \text{ (SI convention), } e \text{ (SI convention)} \quad (80)$$

$$\textbf{Calculated: } c = f_1(\mu_0), \quad \varepsilon_0 = f_2(\mu_0, c) \quad (81)$$

$$\hbar = f_3(e, \varepsilon_0, c, \alpha) \quad (82)$$

$$G = f_4(\ell_P, c, \hbar) \quad (83)$$

(Note of 30 Sept. 2026: Under SI 2019, μ_0 is no longer an SI convention but measured; defined are c, e, h . The SI-conforming order is $\varepsilon_0 = e^2/(4\pi\hbar c\alpha)$, $\mu_0 = 1/(\varepsilon_0 c^2)$, as in the corrected hierarchy above.)

Each quantity depends only on previously defined quantities.

Circularity Test

A circular argument exists if:

$$A \xrightarrow{\text{defined}} B \xrightarrow{\text{defined}} C \xrightarrow{\text{defined}} A \quad (84)$$

In our case:

$$\alpha, \ell_P \xrightarrow{\text{calculated}} \hbar \xrightarrow{\text{calculated}} G \leftrightarrow \alpha, \ell_P \quad (85)$$

Result: Within this chain of calculation there is no circularity.

The Philosophical Argument

Reference Scales are Necessary

Reference Scales

All physics needs reference scales.

Nature is dimensionally structured. To get from dimensionless relationships to measurable quantities, we need:

- An **energy scale** (from α)
- A **length scale** (from ℓ_P)
- **SI conventions** (human measures)

This holds for any dimensional physics, not only for T0 theory.

.12 Further Considerations

Connection to the T0 Model

Within the T0 model, α and ℓ_P could also be derived from geometric principles **[S]**:

$$\xi = \frac{4}{3} \times 10^{-4} \quad (\text{3D space geometry}) \quad (86)$$

$$\alpha = \xi \times E_0^2 \quad \text{with } E_0 = \sqrt{m_e \times m_\mu / K_{\text{frak}}} \quad (87)$$

$$\ell_P = \xi \times \ell_{\text{fundamental}} \quad (88)$$

(Corrected on 30 Sept. 2026: previously $E_0 = \sqrt{m_e \times m_\mu}$ – without K_{frak} , $\xi \cdot 7.348^2$ gives $\alpha^{-1} = 138.91$; with $E_0 = \sqrt{m_e m_\mu / K_{\text{frak}}} = 7.397 \text{ MeV}$ one obtains $\alpha^{-1} = 0.98667 \times 138.91 = 137.06$ (Doc. 190, R72).)

This would reduce the number of fundamental parameters to **one**, the geometric parameter ξ (plus motivated structural choices and SI anchors for unit conversion).