

Parameter System-Dependency in T0-Model:

Contents

.1	Introduction	1
	The Parameter Transfer Problem	1
	Scope and Methodology	2
.2	The ξ Parameter: Variable Across Mass Scales	2
.3	The Universal ξ -Field Framework	2
	Definition and Physical Meaning	2
	Connection to Higgs Physics	2
	ξ Values in the SI System	3
	ξ in T0-Natural Units	3
	Invariance of Ratios	3
.4	The Fine-Structure Constant α_{EM}	4
	The Mystification of $1/137$	4
	Electromagnetic Duality as the Key	4
	The Dual Nature of α : System-Dependent yet Invariant	4
	As an Invariant Ratio of Physical Quantities	4
	As a System-Dependent Numerical Value	5
	The System Dependency of α	5
	The Natural System of Units with $\alpha = 1$	5
	The Resolution of the "Mystery"	5
	The Anthropic Fallacy	6
	Sommerfeld's Harmonic Imprinting	6
	Physical Interpretation	6
	Historical Warning: The Eddington Saga	6
.5	The β_T Parameter	7
	Empirical vs. Theoretical Values	7
	Theoretical Foundation from Field Theory	7
	Circularity in SI Determination	7
.6	The Wien Constant α_W	7
	Mathematical vs. Conventional Values	7
.7	Parameter Comparison Table	8
.8	Yukawa Parameters: Variable and System-Dependent	8
	The Hierarchy of Yukawa Couplings	8
	Transformation Uncertainty	8
	Consistency Requirements	9
.9	Universal Warning: No Direct Parameter Transfer	9
	The Systematic Problem	9
	Required Transformation Protocol	9
	Experimental Prediction Guidelines	10
.10	The Circularity Resolution	10
	Apparent vs. Real Circularity	10
	System Consistency Verification	10

.11	Implications for T0-Model Testing	10
	System-Specific Predictions	10
	Transformation Validation	11

Correction notes

Corrections from the calculation check (30 Sept. 2026). The following places are corrected or annotated in the text; each carries a dated note with the previous value:

- $\alpha = r_e/\lambda_C \rightarrow r_e/(\lambda_C/2\pi)$ (otherwise $\alpha/2\pi$).
- "1/137 exclusively in the SI system" $\rightarrow$ in all systems not redefining α (Gaussian included).
- Rydberg constant: $m_e c \alpha^2 / (2\hbar) \rightarrow m_e c \alpha^2 / (2h)$.
- Wien constant: $\alpha_W^{\text{SI}} = 2.8977719 \dots$, $\alpha_W^{\text{nat}} = 1$, factor $2.898 \rightarrow b = 2.8977719 \times 10^{-3} \text{ m K}$, or $2\pi/x_W = 1.2655$ in natural units (no dimensionless factor); Higgs relation $m_h^2 = \lambda_h v^2 / 2 \rightarrow 2\lambda_h v^2$.

Abstract

This paper systematically analyzes the parameter dependency between SI units and T0-model natural units, revealing that fundamental parameters like α_{EM} , β_{T} , and Yukawa couplings have dramatically different numerical values in different unit systems. Through detailed calculations, we demonstrate that direct transfer of parameter values between systems leads to errors spanning multiple orders of magnitude. The analysis extends beyond specific parameters to establish universal transformation rules and provides critical warnings against naive parameter transfer. This work establishes that the apparent inconsistencies in T0-model parameters are actually systematic unit-system dependencies that require careful transformation protocols for experimental verification. *(Corrected on 29 Sept. 2026: ξ removed from the list – as the ratio of lengths r_0/ℓ_P it has the same value in both systems, see Section .3.)*

.1 Introduction

The Parameter Transfer Problem

The T0 model, formulated in natural units where $\hbar = c = G = k_B = \alpha_{\text{EM}} = \alpha_W = \beta_{\text{T}} = 1$, presents a fundamental challenge when compared with experimental data expressed in SI units. This paper demonstrates that the apparent inconsistencies between T0-model predictions and experimental observations are not physical contradictions but systematic unit-system dependencies.

The core insight is that parameters such as ξ , α_{EM} , and β_{T} represent fundamentally different quantities when expressed in different unit systems:

$$\xi_{\text{SI}} \neq \xi_{\text{nat}}, \quad \alpha_{\text{EM,SI}} \neq \alpha_{\text{EM,nat}}, \quad \beta_{\text{T,SI}} \neq \beta_{\text{T,nat}}$$

Scope and Methodology

This analysis covers:

- Systematic calculation of parameter ratios between SI and T0-natural units

- Demonstration of transformation invariance for dimensionless ratios
- Extension to variable parameters like ξ and Yukawa couplings
- Universal warnings against direct parameter transfer
- Guidelines for correct experimental comparison protocols

.2 The ξ Parameter: Variable Across Mass Scales

.3 The Universal ξ -Field Framework

The cornerstone of the T0-model is the universal geometric constant that serves as the fundamental parameter for all physical calculations.

The universal geometric constant:

$$\xi = \frac{4}{3} \times 10^{-4} = 1.3333... \times 10^{-4} \quad (1)$$

This dimensionless constant is used throughout T0 theory to connect quantum mechanical and gravitational phenomena. It establishes the characteristic strength of field interactions and provides the foundation for unified field descriptions.

For the detailed derivation and physical justification of this parameter, see the document "Parameter Derivation" (available at:).

This geometric constant determines a characteristic energy scale for the ξ -field:

$$E_\xi = \frac{1}{\xi} = \frac{3}{4 \times 10^{-4}} = 7500 \text{ (natural units)} \quad (2)$$

Definition and Physical Meaning

The parameter ξ is also the ratio of the Schwarzschild radius to the Planck length:

$$\xi = \frac{r_0}{\ell_P} = \frac{2Gm}{c^2 \ell_P} \quad (3)$$

(Corrected on 29 Sept. 2026: previously $\xi = 2Gm/\ell_P$ – the factor c^2 was missing; only $r_0 = 2Gm/c^2$ is a length, and only then is $\xi = r_0/\ell_P$ dimensionless. The same applies to the two occurrences in the running text.)

Crucial: The parameter ξ scales with the mass of the object under consideration according to $\xi(m) = 2Gm/(c^2 \ell_P)$. The Higgs mass defines the fundamental reference scale $\xi_0 = 1.33 \times 10^{-4}$, to which all other masses are normalized in the T0 model.

Connection to Higgs Physics

The T0 model establishes a fundamental connection between ξ and Higgs sector physics through the relationship derived in the complete field-theoretic framework

$$\xi = \frac{\lambda_h^2 v^2}{16\pi^3 m_h^2} \approx 1.33 \times 10^{-4} \quad (4)$$

where:

- $\lambda_h \approx 0.13$ (Higgs self-coupling)
- $v \approx 246$ GeV (Higgs VEV)

- $m_h \approx 125$ GeV (Higgs mass)

This represents the universal scale parameter that emerges from fundamental Standard Model physics, while the mass-dependent form $\xi = 2Gm/(c^2\ell_P)$ applies to specific objects.

ξ Values in the SI System

Using SI constants:

$$G = 6.674 \times 10^{-11} \text{ m}^3/(\text{kg} \cdot \text{s}^2) \quad (5)$$

$$\ell_P = 1.616 \times 10^{-35} \text{ m} \quad (6)$$

We calculate ξ_{SI} for various objects:

Object	Mass	ξ_{SI}
Electron	$9.109 \times 10^{-31} \text{ kg}$	8.37×10^{-23}
Proton	$1.673 \times 10^{-27} \text{ kg}$	1.54×10^{-19}
Human (70 kg)	$7.0 \times 10^1 \text{ kg}$	6.4×10^9
Earth	$5.972 \times 10^{24} \text{ kg}$	5.5×10^{32}
Sun	$1.989 \times 10^{30} \text{ kg}$	1.8×10^{38}
Planck mass	$2.176 \times 10^{-8} \text{ kg}$	2.0

Table 1: ξ values for different objects in SI units

The parameter ξ varies over about 60 orders of magnitude!

(Corrected on 29 Sept. 2026: table values recomputed with $\xi = 2Gm/(c^2\ell_P)$; previously electron 7.52×10^{-7} , proton 1.38×10^{-3} , human 6.4×10^6 , Earth 4.1×10^{28} as well as the statement of 46 orders of magnitude – these values had been computed with the formula lacking c^2 or inconsistently. Sun and Planck mass were already correct.)

ξ in T0-Natural Units

With $\xi = r_0/\ell_P = 2Gm/(c^2\ell_P)$, ξ is a pure ratio of lengths. In natural units $G = 1/E_P^2$ and $\ell_P = 1/E_P$, so again

$$\xi = \frac{2m}{E_P} = \frac{2m}{m_P}. \quad (7)$$

The numerical value is therefore the same in SI and in T0-natural units; no conversion factor between the systems arises for this quantity. The values in Table 1 hold unchanged in both systems.

(Corrected on 29 Sept. 2026: previously a conversion factor $\xi_{\text{nat}}/\xi_{\text{SI}} \approx 4100$ and a table of ξ_{nat} (electron 3.1×10^{-3} , proton 5.7, human 2.6×10^{10} , Sun 7.4×10^{41}) – they rested on the formula without c^2 ; a ratio of lengths has no conversion factor.)

Invariance of Ratios

Critical verification: Since the values themselves are already the same in both systems, this holds all the more for the ratios between different objects:

$$\frac{\xi_{\text{Sun}}}{\xi_e} = \frac{1.83 \times 10^{38}}{8.37 \times 10^{-23}} = \frac{m_{\text{Sun}}}{m_e} = 2.18 \times 10^{60} \quad (8)$$

Ratios are invariant under system transformation!

(Corrected on 29 Sept. 2026: previously $1.8 \times 10^{38} / 7.52 \times 10^{-7} = 2.4 \times 10^{44}$ in SI and $7.4 \times 10^{41} / 3.1 \times 10^{-3}$ in natural units – with the corrected formula the ratio equals the mass ratio.)

.4 The Fine-Structure Constant α_{EM}

The Mystification of 1/137

The fine-structure constant $\alpha_{EM} \approx 1/137$ has been declared one of the greatest mysteries of physics by prominent physicists:

- **Richard Feynman:** "It is one of the greatest damn mysteries of physics: a magic number that comes to us with no understanding whatsoever."
- **Wolfgang Pauli:** "When I die, I will ask God two questions: Why relativity? And why 137? I believe he will have an answer for the first one."
- **Max Born:** "If α were larger, molecules could not exist, and there would be no life."

Electromagnetic Duality as the Key

What all these statements overlook: The fine-structure constant possesses two mathematically equivalent representations that reveal its true nature:

$$\alpha_{EM} = \frac{e^2}{4\pi\epsilon_0\hbar c} \quad (\text{Standard form}) \quad (9)$$

$$\alpha_{EM} = \frac{e^2\mu_0 c}{4\pi\hbar} \quad (\text{Dual form}) \quad (10)$$

This equivalence is based on the Maxwell relation $c^2 = \frac{1}{\epsilon_0\mu_0}$ and reveals a fundamental electromagnetic duality:

$$\frac{1}{\epsilon_0 c} = \mu_0 c \quad (11)$$

The Dual Nature of α : System-Dependent yet Invariant

The fine-structure constant possesses a remarkable dual nature:

As an Invariant Ratio of Physical Quantities

Regardless of the chosen system of units, α remains constant as a **ratio** of fundamental lengths:

$$\alpha_{EM} = \frac{r_e}{\lambda_C/2\pi} = \frac{\text{Classical electron radius}}{\text{Reduced Compton wavelength}} \quad (12)$$

(Corrected on 30 Sept. 2026: previously $\alpha_{EM} = r_e/\lambda_C$ – with $\lambda_C = h/(m_e c)$ this gives $1.161 \times 10^{-3} = \alpha/(2\pi)$; the reduced Compton wavelength is meant, as in the following equation.)

Similarly, the inverse ratio:

$$\alpha_{EM}^{-1} = \frac{a_0}{\lambda_C/2\pi} = \frac{\text{Bohr radius}}{\text{Reduced Compton wavelength}} = 137.036... \quad (13)$$

These ratios are **system-of-units invariant** – they have the same numerical value in any consistent system of units, as the units cancel out in the ratio.

As a System-Dependent Numerical Value

Simultaneously, the numerical value of α depends on the choice of fundamental units:

- **SI system:** $\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} \approx 1/137$
- **Natural units:** $\alpha = 1$ (by suitable choice)
- **Gaussian units:** $\alpha = \frac{e^2}{\hbar c} \approx 1/137$

The System Dependency of α

The numerical value $\alpha_{\text{EM}} = 1/137$ holds **in all systems that do not redefine α by convention** (SI, Gaussian, Heaviside-Lorentz); only the T0 setting turns it into 1: *(Corrected on 30 Sept. 2026: previously "valid exclusively in the SI system" – contradicted the list above, according to which $\alpha = e^2/(\hbar c) \approx 1/137$ also holds in Gaussian units.)*

$$\text{SI system: } \alpha_{\text{EM}}^{\text{SI}} = \frac{e^2}{4\pi\epsilon_0\hbar c} \approx \frac{1}{137.036} \quad (14)$$

$$\text{Natural system of units: } \alpha_{\text{EM}}^{\text{nat}} = 1 \text{ (by suitable choice of units)} \quad (15)$$

Transformation factor:

$$\frac{\alpha_{\text{EM}}^{\text{nat}}}{\alpha_{\text{EM}}^{\text{SI}}} = 137.036 \quad (16)$$

The Natural System of Units with $\alpha = 1$

In a natural system of units that respects electromagnetic duality, we obtain:

- $\hbar_{\text{nat}} = 1$ (quantum mechanical scale)
- $c_{\text{nat}} = 1$ (relativistic scale)
- $\epsilon_{0,\text{nat}} = 1$ (electric constant)
- $\mu_{0,\text{nat}} = 1$ (magnetic constant)
- $e_{\text{nat}}^2 = 4\pi$ (elementary charge)

With these values, $\alpha = 1$ is verified in both the standard form and the dual form:

$$\alpha = \frac{4\pi}{4\pi \cdot 1 \cdot 1 \cdot 1} = 1 \quad (17)$$

The Resolution of the "Mystery"

The apparent mystification of $1/137$ arises from:

1. **Confusion of two aspects:** The invariance of the ratios is conflated with the system-dependency of the numerical representation.
2. **Treatment of the SI system as absolute:** The historically evolved SI units (meter, second, kilogram, ampere) force electromagnetic constants to take "unnatural" values.
3. **Forgetting the construction of unit systems:** All unit systems are human constructs. Nature knows no preferred units.
4. **Search for deeper meaning in conversion factors:** The number 137 has no deeper cosmic significance than, say, the factor 1609.344 between miles and meters.

The Anthropic Fallacy

Typical anthropic arguments claim:

- "If $\alpha_{\text{EM}} = 1/200 \rightarrow$ no atoms $\rightarrow$ no life"
- "If $\alpha_{\text{EM}} = 1/80 \rightarrow$ no stars $\rightarrow$ no life"
- "Therefore, $\alpha_{\text{EM}} = 1/137$ is 'fine-tuned' for life"

The problem: These arguments presuppose the SI system as absolute!

In natural units: $\alpha_{\text{EM}} = 1$ is perfectly natural and requires no fine-tuning whatsoever. The electromagnetic interaction has unit strength in the natural system of units, which respects the fundamental structure of quantum mechanics and relativity.

Sommerfeld's Harmonic Imprinting

An often overlooked historical aspect: In 1916, Arnold Sommerfeld actively searched for **harmonic ratios** in atomic spectra, guided by the philosophical conviction that nature follows musical principles.

His methodological approach:

1. **Expectation** of musical ratios in quantum transitions
2. **Calibration** of measurement systems to produce harmonic values
3. **Definition** of α_{EM} based on harmonic spectroscopic adjustments
4. **Attribution** of the resulting ratio to fundamental physics

The apparent "harmony" in $\alpha_{\text{EM}}^{-1} = 137 \approx (6/5)^{27}$ is therefore not a cosmic discovery, but the result of Sommerfeld's harmonic expectations embedded into the definition of the unit system.

Physical Interpretation

In natural units, $\alpha = 1$ represents the perfect balance between:

- **Electric field coupling** (via ϵ_0 with c^{-1})
- **Magnetic field coupling** (via μ_0 with c^{+1})
- **Quantum mechanical scale** (via $\hbar$)
- **Relativistic scale** (via c)

The electromagnetic duality $\frac{1}{\epsilon_0 c} = \mu_0 c$ ensures this perfect balance.

Historical Warning: The Eddington Saga

Arthur Eddington (1882-1944) attempted to "prove" $\alpha_{\text{EM}} = 1/137$ from first principles and developed elaborate numerological theories. The result was entirely speculative and wrong – a warning against mystifying system-dependent numbers.

However, modern analysis shows that the fine-structure constant is indeed derivable from fundamental electromagnetic vacuum constants and that $\alpha_{\text{EM}} = 1$ in natural units is not only possible but reveals the arbitrary nature of our choice of unit system.

.5 The β_T Parameter

Empirical vs. Theoretical Values

The β_T parameter shows the same system dependency:

$$\beta_{T,SI} \approx 0.008 \text{ (from astrophysical observations)} \quad (18)$$

$$\beta_{T,nat} = 1 \text{ (in T0-natural units)} \quad (19)$$

Transformation factor:

$$\frac{\beta_{T,nat}}{\beta_{T,SI}} = \frac{1}{0.008} = 125$$

Theoretical Foundation from Field Theory

The T0 model establishes $\beta_T = 1$ through the fundamental field-theoretic relationship [1]:

$$\beta_T = \frac{\lambda_h^2 v^2}{16\pi^3 m_h^2 \xi} = 1 \quad (20)$$

This relationship, combined with the Higgs-derived value of ξ , uniquely determines $\beta_T = 1$ in natural units, eliminating any free parameters from the theory.

Circularity in SI Determination

The SI value $\beta_{T,SI}$ is determined through:

$$z(\lambda) = z_0 \left(1 + \beta_T \ln \frac{\lambda}{\lambda_0} \right)$$

But this involves:

- Hubble constant $H_0 \rightarrow$ distance measurements
- Distance ladder $\rightarrow$ standard candles
- Photometry $\rightarrow$ Planck radiation law $\rightarrow$ fundamental constants

The determination is circular through cosmological parameters!

.6 The Wien Constant α_W

Mathematical vs. Conventional Values

Wien's displacement law gives:

$$\text{SI system: } \alpha_W^{SI} = 2.8977719... \times 10^{-3} \text{ m K} \quad (21)$$

$$\text{T0 system: } \alpha_W^{nat} = 2\pi/x_W = 1.2655 \quad (x_W = 4.965114) \quad (22)$$

Transformation factor:

$$\alpha_W^{SI} = \alpha_W^{nat} \cdot \frac{\hbar c}{k_B} = 1.2655 \times 2.2899 \times 10^{-3} \text{ m K}$$

(Corrected on 30 Sept. 2026: previously $\alpha_W^{SI} = 2.8977719 \dots$, $\alpha_W^{nat} = 1$ and factor 2.898 – Wien’s constant is $b = 2.8977719 \times 10^{-3} \text{ m K}$ (the factor 10^{-3} and the unit were missing); in $\hbar = c = k_B = 1$ it is fixed as $b = 2\pi/x_W = 2\pi/4.965114 = 1.2655$ and cannot additionally be set to 1; $1.2655 \cdot \hbar c/k_B = 2.8978 \times 10^{-3} \text{ m K}$. There is therefore no dimensionless conversion factor between the systems for b ; this also applies to the row α_W in Table 2 and to the statement on orders of magnitude below it.)

.7 Parameter Comparison Table

Parameter	SI Value	T0-nat Value	Ratio	Factor
ξ (electron)	8.4×10^{-23}	8.4×10^{-23}	1	10^0
α_{EM}	7.3×10^{-3}	1	137	$10^{2.1}$
β_T	0.008	1	125	$10^{2.1}$
α_W	$2.898 \times 10^{-3} \text{ m K}$	1.2655	$\hbar c/k_B$	–

Table 2: Systematic parameter differences between unit systems

(Corrected on 29 Sept. 2026: row ξ (electron) previously $7.5 \times 10^{-6} / 3.1 \times 10^{-2}$, ratio 4100 – $\xi = 2m_e/m_p$ is the same in both systems.)

α_{EM} , β_T and α_W show **0.5–2 orders of magnitude difference between systems**; ξ is the **same in both**. (Corrected on 29 Sept. 2026: previously “All parameters show 0.5-4 orders of magnitude” – the 4 orders came from the ξ row.)

.8 Yukawa Parameters: Variable and System-Dependent

The Hierarchy of Yukawa Couplings

In the Standard Model, Yukawa couplings vary dramatically:

Particle	y_i (SI system)
Electron	2.94×10^{-6}
Muon	6.09×10^{-4}
Tau	1.03×10^{-2}
Up quark	1.27×10^{-5}
Top quark	1.00
Bottom quark	2.25×10^{-2}

Table 3: Yukawa coupling hierarchy (5 orders of magnitude variation)

Transformation Uncertainty

The transformation of Yukawa parameters between systems requires careful consideration of the Higgs mechanism. The general form would be:

$$y_{i,nat} = y_{i,SI} \times T_{Yukawa}$$

where T_{Yukawa} depends on the transformation of Higgs vacuum expectation value and particle masses.

Consistency Requirements

The Higgs mechanism requires:

$$m_h^2 = 2\lambda_h v^2$$

(Corrected on 30 Sept. 2026: previously $m_h^2 = \lambda_h v^2/2$ – with $\lambda_h \approx 0.13$, $v = 246$ GeV this would give $m_h = 62.7$ GeV; $\lambda_h \approx 0.13$ belongs to the convention $m_h^2 = 2\lambda_h v^2$, $\sqrt{0.26} \cdot 246 = 125.4$ GeV.)

For transformation consistency:

$$T_m^2 = T_\lambda \times T_v^2$$

This gives:

$$y_{i,\text{nat}} = y_{i,\text{SI}} \times \sqrt{T_\lambda}$$

However, T_λ requires detailed specification of the T0-natural unit system transformation rules.

.9 Universal Warning: No Direct Parameter Transfer

The Systematic Problem

Warning

EVERY parameter symbol in T0-model documents may have different values than in SI system calculations!

Concrete danger zones:

$$G_{\text{nat}} = 1 \quad \text{vs.} \quad G_{\text{SI}} = 6.674 \times 10^{-11} \text{ m}^3/(\text{kg} \cdot \text{s}^2) \quad (23)$$

$$\alpha_{\text{EM,nat}} = 1 \quad \text{vs.} \quad \alpha_{\text{EM,SI}} = 1/137 \quad (24)$$

$$e_{\text{nat}} = 2\sqrt{\pi} \quad \text{vs.} \quad e_{\text{SI}} = 1.602 \times 10^{-19} \text{ C} \quad (25)$$

Direct transfer leads to errors of factors 10^2 to 10^{11} !

Required Transformation Protocol

For every parameter, explicitly specify:

1. **Which unit system** is being used
2. **How transformation occurs** between systems
3. **Which factors must be considered**
4. **Which consistency conditions** must be satisfied

Example of complete specification:

Parameter Specification Template

Parameter: Fine structure constant α_{EM}

SI value: $\alpha_{\text{EM,SI}} = 1/137.036$

T0 value: $\alpha_{\text{EM,nat}} = 1$

Transformation: $\alpha_{\text{EM,nat}} = \alpha_{\text{EM,SI}} \times 137.036$

Consistency: Dimensional analysis verified

Usage: Specify system before calculation

Experimental Prediction Guidelines

For QED calculations:

$$\text{WRONG: } \alpha_{\text{EM}} = 1 \text{ from T0-model directly in SI formulas} \quad (26)$$

$$\text{CORRECT: } \alpha_{\text{EM,SI}} = 1/137 \text{ with transformation to } \alpha_{\text{EM,nat}} = 1 \quad (27)$$

For gravitational calculations:

$$\text{WRONG: } G = 1 \text{ from T0-model directly in Newton's formulas} \quad (28)$$

$$\text{CORRECT: } G_{\text{SI}} = 6.674 \times 10^{-11} \text{ with transformation to } G_{\text{nat}} = 1 \quad (29)$$

.10 The Circularity Resolution

Apparent vs. Real Circularity

The circularity problem that seemed to plague T0-model parameter determination is resolved by recognizing:

1. **No real circularity exists** within each consistent system
2. **Both SI and T0 systems are internally consistent**
3. **The apparent contradiction** arose from comparing parameters across different systems
4. **Proper transformation** eliminates all apparent inconsistencies

System Consistency Verification

SI system consistency:

$$R_0 = \frac{m_e c (\alpha_{\text{EM,SI}})^2}{2h} \quad \checkmark \text{ (experimentally verified to 0.000001\%)}$$

(Corrected on 30 Sept. 2026: previously denominator $2\hbar$ – this gives $6.895 \times 10^7 \text{ m}^{-1}$; with $2h$ one gets $1.09737 \times 10^7 \text{ m}^{-1}$.)

T0 system consistency:

$$\text{All parameters} = 1 \quad \checkmark \text{ (by construction)}$$

Both systems work perfectly within their own frameworks!

.11 Implications for T0-Model Testing

System-Specific Predictions

Experimental tests must clearly specify which parameter system is used:

(Corrected on 29 Sept. 2026: row on galaxy rotation previously $\xi_{\text{nat}} \sim 10^{41}$ – ξ has the same value in both systems.)

Test Type	SI-based Prediction	T0-based Prediction
QED anomaly	$a_e \propto \alpha_{\text{EM,SI}} = 1/137$	$a_e \propto \alpha_{\text{EM,nat}} = 1$
Galaxy rotation	$v^2 \propto \xi \sim 10^{38}$	$v^2 \propto \xi \sim 10^{38}$ (same)
CMB temperature	$T \propto \beta_{T,\text{SI}} = 0.008$	$T \propto \beta_{T,\text{nat}} = 1$

Table 4: System-specific experimental predictions

Transformation Validation

The transformation factors can be validated by checking:

1. **Dimensional consistency** in both systems
2. **Known limits** are reproduced correctly
3. **Ratios remain invariant** between systems
4. **Internal consistency** of each system

Bibliography

- [1] J. Pascher. Derivation of Beta in the T0 Framework. T0-Theory Document Collection, 2025. URL: https://github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/pdf/093_DerivationVonBeta_En.pdf (verifizierter Link existiert).