

On the Mathematical Structure of the T0 Theory:

Why Numerical Ratios Must Not Be Directly Simplified

Johann Pascher

Original

Contents

.1	Circular Relationships and Fundamental Constants	3
	Resolution of the Circularity Problem	4
.2	The Role of Natural Units	4
.3	Foundation: The Single Geometric Constant	4
	The Universal Geometric Parameter	4
	Natural Units	5
.4	Building the Scale Hierarchy	5
	Step 1: Characteristic T0 Scales	5
	Step 2: Energy Scales from Geometry	5
.5	Deriving the Fine Structure Constant	6
	Origin of the Formula $\epsilon = \xi \cdot E_0^2$	6
	The Characteristic Energy E_0	6
	The Geometric Parameter ξ	6
	Numerical Verification and Fine Structure Constant	6
	From Fractal Geometry	7
	Fractal Dimension of Spacetime	7
	The Fine Structure Constant from Geometry	7
	Formula from ξ to α	7
.6	Lepton Mass Hierarchy from Pure Geometry	8
	Mechanism for Mass Generation	8
	Mass Calculations	8
	Electron Mass	8
	Muon Mass	8
	Tau Mass	8
	Mass Ratios	9
.7	Complete Hierarchy with Final Anomaly Formula	9
.8	Verification of Final Formula	9
	Complete Derivation Chain to Final Formula	9
	T0 Field Theory Verification of Final Formula	10
.9	The Fundamental Meaning of E_0 as Logarithmic Center	11
	The Central Geometric Definition	11
	Mathematical Properties	11
	Numerical Values	11
	Logarithmic Symmetry	11
.10	The Geometric Constant C	12
	Fundamental Relationship	12
	Tetrahedral Geometry	12
	Formula for α with Fractal Correction	12
.11	The Simplest Formula: The Geometric Mean	12
	The Fundamental Definition	12
	Direct Calculation	13

	The Complete Chain in One Line	13
	Why Is This So Simple?	13
	Logarithmic Centering	13
	Alternative Notations	13
	The Fine Structure Constant Directly	14
	Relation to the Other Derivations	14
	Meaning	14
.12	The Fundamental Dependence: $\alpha \sim \xi^{11/2}$	14
	Inserting the Mass Formulas	14
	Calculation of E_0	14
	Calculation of α	15
	What Does This Mean?	15
	1. Fundamental Connection	15
	2. Hierarchy Problem	15
	3. No Independence	15
	Numerical Verification	15
	The Units Problem	16
	Alternative View: Reduction to ξ	16
.13	Derivation of the Coefficients c_e and c_μ	16
	Starting Point: Mass Formulas	16
	Step 1: Quantum Numbers and Geometric Factors	16
	Step 2: Derivation of c_e (Electron)	17
	Step 3: Derivation of c_μ (Muon)	17
	Step 4: Physical Interpretation	17
	Step 5: Dimensional Analysis	17
	Step 6: Consistency Check	18
.14	Why Natural Units Are Necessary	18
	The Problem with Conventional Units	18
	Natural Units Simplify Physics	18
	Transformation to Natural Units	18
	The Coefficients in Natural Units	19
	Comparison of Representations	19
	Advantages of Natural Units	19
	Example: The Mass Formula	19
	Fundamental Interpretation	19
.15	The Formula from ξ to α	19
	Fundamental Relationship	19
	Coefficients	20
	Product of Coefficients	20
	Complete Formula	20
	Solving for α	20
.16	T0-Theory: Formulas and Values	20
	In T0-Theory	20
	Correct Assignment in Natural Units	20
	Mass Scaling Laws	20
	Geometric Constant	21
	Calculation of the Characteristic Energy	21
	Calculation of the Fine Structure Constant	21
	Numerical Values	21
	Interpretation	21

.17	Definitions	21
	Geometric Constant	21
	Mass Formulas	22
.18	Coefficients from T0-Theory	22
	Electron ($n=1, l=0, j=1/2$)	22
	Muon ($n=2, l=1, j=1/2$)	22
	Tauon ($n=3, l=2, j=1/2$)	22
.19	Mass Calculation	22
	Electron Mass	22
	Muon Mass	22
	Tauon Mass	23
.20	Characteristic Energy	23
.21	Fine Structure Constant	23
.22	Numerical Values	23
.23	The Formula from ξ to α (Detailed)	24
	From the Fundamental Relationship	24
	Inserting the Coefficients	24
	Complete Formula	24
	Solving for α	24
	Numerical Values	25
	With $\xi = 4/3 \times 10^{-4}$	25
	Symbol Explanation	25
	With Fractal Correction	25
	Final Fundamental Relationship	26
.24	Observation: α Cancels Out	26
	Equating the Formula Sets	26
	Solving for α	26
	For the Muon	26
	The Apparent Contradiction	26
	Resolution in the T0 Framework	27
	The Key Elements	27
	Meaning of the Different Representations	27
	Central Statement	27
	Why This Insight Is Important	27
.25	Why the Extended Form Is Important	28
	The Two Equivalent Representations	28
	The Apparent Contradiction	28
	Consequence	28
	Mathematical Structure	28
	Physical Interpretation	28
	Why Both Representations Are Important	29
	The Actual Statement of T0-Theory	29

Correction notes

Corrections from the calculation review (30 Sept. 2026). The following places are corrected or annotated in the text; each carries a dated note there with the previous value (the assessment of the muon anomaly is covered by the note on Doc. 382):

- Comparison with alternative theories: $138.93 \times 0.9862 = 137.036 \rightarrow 137.01$; Further connections: $K_{\text{frac}} = 0.9862$ comes from the older anchor $D_f = 2.94$ (annotated).
- Comparison with experiment: theoretical m_μ/m_e 206.768 (measured value) $\rightarrow \frac{12}{5}\xi^{-1/2} = 207.85$.
- Table 22.1.1: c_e mantissa = $e \rightarrow \sqrt{e}$; Section 19: the masses given are measured values (anchors), not reachable with a common anchor (annotated).
- Box 11.6.1: $\alpha = \frac{m_e m_\mu}{7500} \times 0.986 \rightarrow \frac{m_e m_\mu}{7500 \times 0.986}$ (K_{frac} acts on α^{-1}); boxed α formula (3.6.1, 9.3.1, 23.9.1) gives 2.4×10^{-9} , and K_{frac} belongs with α^{-1} (annotated).
- Mass ratios: $c_\mu/c_e = 3\sqrt{3}/(2\sqrt{\alpha}) \rightarrow \sqrt{3}/(2\sqrt{\alpha})$, giving $m_\mu/m_e \approx 878$.
- Section 11.3.2: $138.91 \times 0.986 = 137.04 \rightarrow 136.97$.
- Tau mass 4.2.3: $\frac{5}{4}\xi^{2/3} \cdot 246 \text{ GeV} = 1.777 \rightarrow 0.80 \text{ GeV}$.
- Chain 3.5.2: $3\pi \cdot 7500 \cdot \ln 10^4 / 2.973 \approx 137.036 \rightarrow 2.2 \times 10^5$.
- Sections 23.5/23.6: $\alpha \approx 7.292 \times 10^{-3}$, $\alpha^{-1} \approx 137.13 \rightarrow \alpha \approx 2.42 \times 10^{-9}$ ($0.5923^{0.4} = 0.8110$, $\xi^{2.2} = 2.984 \times 10^{-9}$).
- Section 23.8.1: $138.949 \times 0.9862 = 137.036 \rightarrow 138.911 \times 0.9862 = 136.99$.
- Sections 12.5 and 16.2.5: $\xi^{11/2} = 5.19 \times 10^{-22} \rightarrow 4.87 \times 10^{-22}$, $1.4 \times 10^{19} \rightarrow 1.5 \times 10^{19}$.
- Section 3.2.2: gravitational-geometric derivation $E_0^2 = 4\sqrt{2}m_\mu/\xi^4$ ($1.4 \times 10^9 \text{ MeV}$) removed $\rightarrow \sqrt{m_e m_\mu / K_{\text{frac}}}$.
- Section 13.7.1: "match $m_\mu/m_e \approx 207$ " $\rightarrow$ give $m_\mu/m_e \approx 878$; 14.1.1 and 18.1.1–18.3.1: the stated coefficient values do not follow from the formulas (annotated).
- Section 2.2: $E_0 = \frac{3}{4} \times 10^4 E_p$ contradicts $E_0 = 7.398 \text{ MeV}$ (annotated); hierarchy table: m_τ/m_μ from $y_\tau/y_\mu \sim 17 \rightarrow 5.73$; 14.3.1: $m^{\text{nat}} = m^{\text{SI}} G/(\hbar c) \rightarrow m^{\text{SI}} \sqrt{G/(\hbar c)}$, ξ not converted; 16.3.1: "Planck energy to electron volt" $\rightarrow$ to GeV; 20.1.1: $\sqrt{c_e c_\mu} \xi^{9/4} = 5.9 \times 10^{-8}$, not 7.346881 MeV (annotated); Table 22.1.1: $\xi^{5/2} = 3.098 \times 10^{-10} \rightarrow 2.0528 \times 10^{-10}$; $E_0^{\text{corr}} = K_{\text{frac}} E_0^{\text{bare}}$ contradicts R72 (annotated).

Note on the muon $g - 2$ status (2025). This document refers to the deviation between the measured a_μ and the Standard Model prediction as it stood in 2021. With the Fermilab final result and the 2025 White Paper of the Muon $g - 2$ Theory Initiative, this deviation is no longer significant. Statements that use the anomaly as evidence or as the target of a T0 contribution are outdated; the current status and assessment are given in Doc. 382.

On the Mathematical Structure of the T0-Theory: Why Numerical Ratios Must Not Be Directly Simplified

Note on versions. The German and English versions of this document differ in structure and extent; earlier revisions shortened, extended or changed parts of both. Where they

differ, the more extensive German version applies; sections contained only in this English version count as supplements.

Status markers. Statements marked **[S]** are *speculative*: a hint or conjecture that has not yet been derived or numerically checked in the corpus.

Introduction

In theoretical physics, the question often arises as to which mathematical operations are legitimate and which are not. A particularly interesting problem occurs in the T0-theory, where seemingly simple numerical ratios such as $\frac{2}{3}$ and $\frac{8}{5}$ possess a deeper structural significance that prohibits direct simplification.

The Fundamental Problem

The T0-theory postulates two equivalent representations for the lepton masses:

$$\textbf{Simple Form: } m_e = \frac{2}{3} \cdot \xi^{5/2}, \quad m_\mu = \frac{8}{5} \cdot \xi^2$$

$$\textbf{Extended Form: } m_e = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \cdot \xi^{5/2}, \quad m_\mu = \frac{9}{4\pi\alpha} \cdot \xi^2$$

At first glance, one might assume that the fractions $\frac{2}{3}$ and $\frac{8}{5}$ are simple rational numbers that could be simplified or reduced. However, this assumption would be incorrect.

Why Direct Simplification Is Not Allowed

Equating both representations leads to:

$$\frac{2}{3} = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}}, \quad \frac{8}{5} = \frac{9}{4\pi\alpha}$$

These equations show that the seemingly simple fractions are, in fact, composite expressions containing the constants (π, α) and geometric factors $(\sqrt{3})$.

Mathematical and Physical Consequences

1. **Structure Preservation:** Direct simplification would destroy the underlying geometric and physical structure.
2. **Information Loss:** The fractions encode information about spacetime geometry and electromagnetic coupling.
3. **Equivalence Principle:** Both representations are mathematically equivalent, but the extended form reveals the physical origin.

1 Circular Relationships and Fundamental Constants

In the T0-theory, seemingly circular relationships arise, which express how the constants are linked in the T0 framework:

$$\alpha = f(\xi)$$

$$\xi = g(\alpha)$$

This mutual dependence leads to an apparent chicken-and-egg problem: Which comes first, α or ξ ?

Resolution of the Circularity Problem

The solution lies in the realization that both constants are expressions of an underlying geometric structure:

α and ξ are not independent of each other but are emergent properties of the fractal spacetime geometry.

The apparent circularity dissolves when it is recognized that both constants originate from the same fundamental geometry.

.2 The Role of Natural Units

In natural units, we conventionally set $\alpha = 1$ for certain calculations. This is legitimate because:

- Fundamental physics should be independent of measurement units.
- Dimensionless ratios contain the actual physical statements.
- The choice $\alpha = 1$ represents a specific gauge.

However, this convention must not obscure the fact that α in the T0-theory has a specific numerical value determined by ξ .

The seemingly simple numerical ratios in the T0-theory are not arbitrarily chosen but represent complex physical relationships.

Directly simplifying these ratios would be mathematically possible but physically incorrect, as it would destroy the underlying structure of the theory. The extended form shows where these seemingly simple fractions come from in the T0 framework and makes their connection to π , α and geometric factors visible. The apparent circularity between α and ξ is an expression of their common geometric origin and not a logical problem of the theory.

.3 Foundation: The Single Geometric Constant

The Universal Geometric Parameter

1.1.1 The T0-theory begins with a single dimensionless constant motivated by the geometry of three-dimensional space:

Key Result

$$\xi = \frac{4}{3} \times 10^{-4} \quad (1)$$

1.1.2 This constant arises from:

- The tetrahedral packing density of 3D space: $\frac{4}{3}$
- The scale hierarchy between quantum and classical domains: 10^{-4} (motivated choice of the exponent)

Natural Units

1.2.1 We work in natural units where:

$$c = 1 \quad (\text{speed of light}) \quad (2)$$

$$\hbar = 1 \quad (\text{reduced Planck constant}) \quad (3)$$

$$G = 1 \quad (\text{gravitational constant, numerically}) \quad (4)$$

1.2.2 The Planck length serves as reference scale:

$$\ell_P = \sqrt{G} = 1 \quad (\text{in natural units}) \quad (5)$$

.4 Building the Scale Hierarchy**Step 1: Characteristic T0 Scales**

2.1.1 From ξ and the Planck reference, we derive the characteristic T0 scales:

$$r_0 = \xi \cdot \ell_P = \frac{4}{3} \times 10^{-4} \cdot \ell_P \quad (6)$$

$$t_0 = r_0 = \frac{4}{3} \times 10^{-4} \quad (\text{in units with } c = 1) \quad (7)$$

Step 2: Energy Scales from Geometry

2.2.1 The characteristic energy scale follows from dimensional analysis:

$$E_0 = \frac{1}{r_0} = \frac{3}{4} \times 10^4 \quad (\text{in Planck units}) \quad (8)$$

2.2.2 This yields the T0 energy hierarchy:

$$E_P = 1 \quad (\text{Planck energy}) \quad (9)$$

$$E_0 = \xi^{-1} E_P = \frac{3}{4} \times 10^4 E_P \quad (10)$$

(Note of 30 Sept. 2026: $E_0 = \xi^{-1} E_P = 7500 E_P = 9.2 \times 10^{22} \text{ GeV}$ contradicts $E_0 = 7.398 \text{ MeV}$ in Section 3; with this E_0 one would get $\xi E_0^2 = 1/\xi = 7500$, not α . Setting $E_0 = 1/r_0$ in Planck units is not compatible with the rest of the derivation.)

.5 Deriving the Fine Structure Constant

Origin of the Formula $\varepsilon = \xi \cdot E_0^2$

3.1.1 The fundamental formula of T0-theory for the coupling parameter ε is:

Key Result

$$\varepsilon = \xi \cdot E_0^2 \quad (11)$$

3.1.2 This relationship connects:

- ε – the T0 coupling parameter
- ξ – the geometric parameter from tetrahedral packing
- E_0 – the characteristic energy

The Characteristic Energy E_0

3.2.1 The characteristic energy E_0 is defined as the geometric mean of electron and muon masses:

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (12)$$

3.2.2 With the fractal correction (Doc. 190, R72) this becomes:

$$E_0 = \sqrt{\frac{m_e \cdot m_\mu}{K_{\text{frac}}}} \quad (13)$$

3.2.3 Bare and fractally corrected one obtains:

$$E_0 = 7.348 \text{ MeV (bare)}, \quad E_0 = 7.348 / \sqrt{0.98667} = 7.397 \approx 7.398 \text{ MeV (corrected)} \quad (14)$$

(Corrected on 30 Sept. 2026: previously 3.2.2 was the "gravitational-geometric" derivation $E_0^2 = 4\sqrt{2}m_\mu/\xi^4$ and "Both approaches consistently lead to $E_0 \approx 7.35$ to 7.398 MeV " – this formula gives $E_0 = 1.4 \times 10^9 \text{ MeV}$, is dimensionally inconsistent and not derived in the corpus; it has been removed as a second route (Docs. 011, 041, A130). The difference $7.348 \rightarrow 7.398$ is the factor $1/\sqrt{K_{\text{frac}}}$.)

The Geometric Parameter ξ

3.3.1 The parameter ξ is a fundamental geometric constant:

$$\xi = \frac{4}{3} \times 10^{-4} = 1.333... \times 10^{-4} \quad (15)$$

Numerical Verification and Fine Structure Constant

3.4.1 With the derived values, ε becomes:

$$\varepsilon = \xi \cdot E_0^2 \quad (16)$$

$$= (1.333 \times 10^{-4}) \times (7.398 \text{ MeV})^2 \quad (17)$$

$$= 7.297 \times 10^{-3} \quad (18)$$

$$= \frac{1}{137.036} \quad (19)$$

Comparison with the Fine Structure Constant

3.4.2 With $E_0 = 7.398$ MeV, the T0 coupling parameter ε reproduces the value of the fine structure constant $\alpha = 1/137.036$. The geometric mean $\sqrt{m_e m_\mu}$ alone gives $E_0 \approx 7.35$ MeV (Section 11.2); the difference is described below by the fractal correction factor K_{frac} .

From Fractal Geometry

Fractal Dimension of Spacetime

3.5.1 From topological considerations of 3D space with time:

$$D_f^{\text{eff}} \approx 2.973 \quad (20)$$

where $\delta = 0.06$ is the fractal correction.

The Fine Structure Constant from Geometry

3.5.2 The geometric approach gives:

Key Result

$$\alpha^{-1} = 3\pi \times \xi^{-1} \times \ln\left(\frac{\Lambda_{\text{UV}}}{\Lambda_{\text{IR}}}\right) \times D_f^{-1} \quad (21)$$

$$= 3\pi \times \frac{3}{4} \times 10^4 \times \ln(10^4) \times \frac{1}{2.973} \quad (22)$$

$$= \frac{9\pi}{4} \times 10^4 \times 9.21 \times 0.336 \quad (23)$$

$$\approx 2.2 \times 10^5 \quad (24)$$

(Corrected on 30 Sept. 2026: previously $= 9\pi \times 10^4 \times 9.21 \times 0.340 \approx 137.036 - 3\pi \times \frac{3}{4} \times 10^4 = \frac{9\pi}{4} \times 10^4$ and $1/2.973 = 0.336$; the chain gives 2.2×10^5 , not 137.)

Formula from ξ to α

3.6.1 The precise relationship is:

Key Result

$$\alpha = \left(\frac{27\sqrt{3}}{8\pi^2}\right)^{2/5} \cdot \xi^{11/5} \cdot K_{\text{frac}} \quad (25)$$

$$\text{with } K_{\text{frac}} = 0.9862 \quad (26)$$

(Note of 30 Sept. 2026: this formula gives numerically $\alpha \approx 2.4 \times 10^{-9}$, not $1/137$; moreover, according to Section 23.8, K_{frac} acts on α^{-1} ($\alpha^{-1} = 138.9 \cdot K_{\text{frac}}$), i.e. in α as $1/K_{\text{frac}}$, not as a factor.)

.6 Lepton Mass Hierarchy from Pure Geometry

Mechanism for Mass Generation

4.1.1 Masses arise from the coupling of the energy field to spacetime geometry:

$$m_\ell = r_\ell \cdot \xi^{p_\ell} \quad (27)$$

where r_ℓ are rational coefficients and p_ℓ are exponents.

Mass Calculations

Electron Mass

4.2.1 The electron mass calculation:

Key Result

$$m_e = \frac{2}{3} \xi^{5/2} \quad (28)$$

$$= \frac{2}{3} \left(\frac{4}{3} \times 10^{-4} \right)^{5/2} \quad (29)$$

$$= \frac{2}{3} \cdot \frac{32}{9\sqrt{3}} \times 10^{-10} \quad (30)$$

$$= \frac{64\sqrt{3}}{81} \times 10^{-10} \quad (31)$$

$$\approx 1.368 \times 10^{-10} \quad (\text{natural units}) \quad (32)$$

Muon Mass

4.2.2 The muon mass calculation:

Key Result

$$m_\mu = \frac{8}{5} \xi^2 \quad (33)$$

$$= \frac{8}{5} \left(\frac{4}{3} \times 10^{-4} \right)^2 \quad (34)$$

$$= \frac{128}{45} \times 10^{-8} \quad (35)$$

$$\approx 2.844 \times 10^{-8} \quad (\text{natural units}) \quad (36)$$

Tau Mass

4.2.3 The tau mass calculation:

Key Result

$$m_\tau = \frac{5}{4} \xi^{2/3} \cdot v_{\text{scale}} \quad (37)$$

$$= \frac{5}{4} \left(\frac{4}{3} \times 10^{-4} \right)^{2/3} \cdot v_{\text{scale}} \quad (38)$$

$$\approx 0.80 \text{ GeV} \quad (39)$$

with $v_{\text{scale}} = 246 \text{ GeV}$.

(Corrected on 30 Sept. 2026: previously $\approx 1.777 \text{ GeV} \approx 2.133 \times 10^{-4}$ (natural units) – $\frac{5}{4} \xi^{2/3} \cdot 246 \text{ GeV}$ gives 0.80 GeV ; 1.777 GeV is the measured tau mass.)

Mass Ratios**4.3.1** The electron to muon mass ratio:**Key Result**

$$\frac{m_e}{m_\mu} = \frac{\frac{64\sqrt{3}}{81} \times 10^{-10}}{\frac{128}{45} \times 10^{-8}} \quad (40)$$

$$= \frac{5\sqrt{3}}{18} \times 10^{-2} \quad (41)$$

$$\approx 4.811 \times 10^{-3} \quad (42)$$

.7 Complete Hierarchy with Final Anomaly Formula

6.1 The following table summarizes all derived quantities with the final anomaly formula:
(Corrected on 30 Sept. 2026: m_τ/m_μ from y_τ/y_μ previously $\sim 17 - \frac{5}{4} \xi^{2/3} / \frac{64}{15} \xi = 3.26 \times 10^{-3} / 5.69 \times 10^{-4} = 5.73$.)

.8 Verification of Final Formula**Complete Derivation Chain to Final Formula**

7.1.1 The complete derivation sequence:

1. **Start:** $\xi = \frac{4}{3} \times 10^{-4}$ (pure geometry)
2. **Reference:** $\ell_P = 1$ (natural units)
3. **Derivation:** $r_0 = \xi \ell_P$
4. **Energy:** $E_0 = r_0^{-1}$
5. **Fractal:** $D_f^{\text{eff}} \approx 2.973$ (cumulative topology over RG running)
6. **Fine structure:** $\alpha = f(\xi, D_f)$
7. **Yukawa:** $y_\ell = r_\ell \xi^{p_\ell}$ (geometry)
8. **Masses:** $m_\ell \propto y_\ell$

Quantity	Expression	Value
Fundamental		
ξ	$\frac{4}{3} \times 10^{-4}$	$1.333 \dots \times 10^{-4}$
D_f^{eff}	cumulative over RG running	2.973
Scales		
r_0/ℓ_P	ξ	$\frac{4}{3} \times 10^{-4}$
E_0/E_P	ξ^{-1}	$\frac{3}{4} \times 10^4$
Couplings		
α^{-1}	From Geometry	137.036
Yukawa Couplings		
y_e	$\frac{32}{9\sqrt{3}} \xi^{3/2}$	$\sim 10^{-6}$
y_μ	$\frac{64}{15} \xi$	$\sim 10^{-4}$
y_τ	$\frac{5}{4} \xi^{2/3}$	$\sim 10^{-3}$
Mass Ratios		
m_e/m_μ	$\frac{5\sqrt{3}}{18} \times 10^{-2}$	4.8×10^{-3}
m_τ/m_μ	From y_τ/y_μ	≈ 5.73

Table 1: Complete hierarchy with final quadratic anomaly formula

9. **Yukawa coupling:** $g_T^\ell = m_\ell \xi$

10. **One-loop calculation:** $\Delta a_\ell = \frac{(m_\ell \xi)^2}{8\pi^2} \cdot \frac{\xi^2}{\lambda^2}$

11. **Final formula:** $\Delta a_\ell = 251 \times 10^{-11} \times (m_\ell/m_\mu)^2$

T0 Field Theory Verification of Final Formula

7.2.1 The final formula follows from T0 field theory calculation:

- **Muon g-2 calculation:** $\frac{m_\mu^2 \xi^4}{8\pi^2 \lambda^2} = 251 \times 10^{-11}$ (T0 field theory prediction)
- **Electron prediction:** 5.87×10^{-15} (T0 prediction with the single parameter ξ)
- **Tau prediction:** 7.10×10^{-9} (testable in future experiments)
- **Quadratic scaling:** Follows from standard QFT one-loop calculation

.9 The Fundamental Meaning of E_0 as Logarithmic Center

The Central Geometric Definition

Fundamental Definition

8.1.1 The characteristic energy E_0 is the logarithmic center between electron and muon masses:

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (43)$$

This means:

$$\log(E_0) = \frac{\log(m_e) + \log(m_\mu)}{2} \quad (44)$$

Mathematical Properties

8.2.1 The fundamental relationships:

$$E_0^2 = m_e \cdot m_\mu \quad (45)$$

$$\frac{E_0}{m_e} = \sqrt{\frac{m_\mu}{m_e}} \quad (46)$$

$$\frac{m_\mu}{E_0} = \sqrt{\frac{m_\mu}{m_e}} \quad (47)$$

$$\frac{E_0}{m_e} \cdot \frac{m_\mu}{E_0} = \frac{m_\mu}{m_e} \quad (48)$$

Numerical Values

8.3.1 With T0-calculated masses:

$$m_e^{\text{T0}} = 0.5108082 \text{ MeV} \quad (49)$$


$$m_\mu^{\text{T0}} = 105.66913 \text{ MeV} \quad (50)$$

$$E_0^{\text{T0}} = \sqrt{0.5108082 \times 105.66913} \approx 7.346881 \text{ MeV} \quad (51)$$

Logarithmic Symmetry

8.4.1 The logarithmic symmetry (by definition of E_0):

$$\ln(E_0) - \ln(m_e) = \ln(m_\mu) - \ln(E_0) \quad (52)$$



.10 The Geometric Constant C

Fundamental Relationship

9.1.1 The fractal correction factor:

$$K_{\text{frac}} = 1 - \frac{D_f - 2}{C} = 1 - \frac{\gamma}{C} \quad (53)$$

where:

$$D_f^{\text{eff}} \approx 2.973 \quad (\text{effective fractal dimension}) \quad (54)$$

$$\gamma = D_f - 2 = 0.94 \quad (55)$$

$$C \approx 68.24 \quad (56)$$

Tetrahedral Geometry

Tetrahedral Numbers

9.2.1 Several tetrahedral combinations yield 72; whether this is related to $C \approx 68.24$ is open [S]:

$$6 \times 12 = 72 \quad (\text{edges} \times \text{rotations}) \quad (57)$$

$$4 \times 18 = 72 \quad (\text{faces} \times 18) \quad (58)$$

$$24 \times 3 = 72 \quad (\text{symmetries} \times \text{dimensions}) \quad (59)$$

Formula for α with Fractal Correction

9.3.1 The complete expression:

$$\alpha = \left(\frac{27\sqrt{3}}{8\pi^2} \right)^{2/5} \cdot \xi^{11/5} \cdot K_{\text{frac}} \quad \text{with} \quad K_{\text{frac}} = 0.9862 \quad (60)$$

(Note of 30 Sept. 2026: as in 3.6.1 – the formula gives $\alpha \approx 2.4 \times 10^{-9}$, and K_{frac} belongs with α^{-1} .)

.11 The Simplest Formula: The Geometric Mean

The Fundamental Definition

The Simplest Formula

11.1.1 The basic relation:

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (61)$$

This definition needs no further derivation, only the geometric mean.

Direct Calculation

11.2.1 Simple numerical evaluation:

$$E_0 = \sqrt{0.511 \text{ MeV} \times 105.658 \text{ MeV}} \quad (62)$$

$$= \sqrt{53.99 \text{ MeV}^2} \quad (63)$$

$$= 7.35 \text{ MeV} \quad (64)$$

The Complete Chain in One Line

11.3.1 The fundamental relationship:

$$\alpha^{-1} = \frac{7500}{m_e \cdot m_\mu} = \frac{7500}{E_0^2} \quad (65)$$

11.3.2 With numbers:

$$\alpha^{-1} = \frac{7500}{0.511 \times 105.658} \quad (66)$$

$$= \frac{7500}{53.99} \quad (67)$$

$$= 138.91 \quad (68)$$

(With fractal correction $\times 0.986 = 136.97$) (Corrected on 30 Sept. 2026: previously = 137.04
 $- 138.91 \times 0.986 = 136.97$.)


Why Is This So Simple?

Logarithmic Centering

11.4.1 The geometric mean is the natural center on logarithmic scale:

$$\log(E_0) = \frac{\log(m_e) + \log(m_\mu)}{2} \quad (69)$$

Graphically:



Alternative Notations

11.5.1 All these formulas are equivalent:

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (70)$$

$$E_0^2 = m_e \cdot m_\mu \quad (71)$$

$$\log(E_0) = \frac{1}{2}[\log(m_e) + \log(m_\mu)] \quad (72)$$

$$E_0 = \sqrt{0.511 \times 105.658 \text{ MeV}} \quad (73)$$

$$E_0 = m_e^{1/2} \cdot m_\mu^{1/2} \quad (74)$$

The Fine Structure Constant Directly

The Most Direct Formula

11.6.1 Without detour through E0:

$$\alpha = \frac{m_e \cdot m_\mu}{7500} \quad (75)$$

With fractal correction:

$$\alpha = \frac{m_e \cdot m_\mu}{7500 \times 0.986} \quad (76)$$

(Corrected on 30 Sept. 2026: previously $\alpha = \frac{m_e m_\mu}{7500} \times 0.986$ – the fractal correction multiplies α^{-1} ($138.9 \times 0.986 \approx 137$), so in α it belongs in the denominator; the old expression gives $\alpha^{-1} \approx 140.9$.)

Relation to the Other Derivations

11.7.1 The documents show various derivations of E0: - Fractally corrected, $\sqrt{m_e m_\mu / K_{\text{frac}}}$ (R72) *(Corrected on 30 Sept. 2026: previously “gravitationally-geometrically” – this derivation has been removed, see 3.2.2.)* - Through Yukawa couplings - From quantum numbers

But the simplest definition is:

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (77)$$

Meaning

11.8.1 The geometric mean is not an arbitrary choice: it is the midpoint on a logarithmic scale (Section 11.4).

.12 The Fundamental Dependence: $\alpha \sim \xi^{11/2}$

Inserting the Mass Formulas

12.1.1 From T0-theory we have the mass formulas:

$$m_e = c_e \cdot \xi^{5/2} \quad (78)$$

$$m_\mu = c_\mu \cdot \xi^2 \quad (79)$$

where c_e and c_μ are coefficients.

Calculation of E_0

12.2.1 The characteristic energy calculation:

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (80)$$

$$= \sqrt{(c_e \cdot \xi^{5/2}) \cdot (c_\mu \cdot \xi^2)} \quad (81)$$

$$= \sqrt{c_e \cdot c_\mu} \cdot \sqrt{\xi^{5/2+2}} \quad (82)$$

$$= \sqrt{c_e \cdot c_\mu} \cdot \xi^{9/4} \quad (83)$$

Calculation of α

12.3.1 The fine structure constant derivation:

$$\alpha = \xi \cdot E_0^2 \quad (84)$$

$$= \xi \cdot (\sqrt{c_e \cdot c_\mu} \cdot \xi^{9/4})^2 \quad (85)$$

$$= \xi \cdot c_e \cdot c_\mu \cdot \xi^{9/2} \quad (86)$$

$$= c_e \cdot c_\mu \cdot \xi^{1+9/2} \quad (87)$$

$$= c_e \cdot c_\mu \cdot \xi^{11/2} \quad (88)$$

Result

12.3.2 The fine structure constant depends on ξ :

$$\alpha = K \cdot \xi^{11/2} \quad (89)$$

where $K = c_e \cdot c_\mu$ is a constant.

The powers do not cancel out.

What Does This Mean?

1. Fundamental Connection

12.4.1 The fine structure constant is not independent of ξ , but rather:

$$\alpha \propto \xi^{11/2} \quad (90)$$

This means: If ξ changes, α also changes.

2. Hierarchy Problem

12.4.2 Because of the power $11/2 = 5.5$, small changes in ξ are amplified:

$$\frac{\Delta\alpha}{\alpha} = \frac{11}{2} \cdot \frac{\Delta\xi}{\xi} = 5.5 \cdot \frac{\Delta\xi}{\xi} \quad (91)$$

3. No Independence

12.4.3 One cannot choose α and ξ independently. They are firmly connected through:

$$\alpha = K \cdot \xi^{11/2} \quad (92)$$

Numerical Verification

12.5.1 With $\xi = 4/3 \times 10^{-4}$:

$$\xi^{11/2} = (1.333 \times 10^{-4})^{5.5} \quad (93)$$

$$= 4.87 \times 10^{-22} \quad (94)$$

12.5.2 For $\alpha \approx 1/137$ we would need:

$$K = \frac{\alpha}{\xi^{11/2}} \quad (95)$$

$$= \frac{7.3 \times 10^{-3}}{4.87 \times 10^{-22}} \quad (96)$$

$$= 1.5 \times 10^{19} \quad (97)$$

(Corrected on 30 Sept. 2026: previously $\xi^{11/2} = 5.19 \times 10^{-22}$, $K = 1.4 \times 10^{19} - (1/7500)^{5.5} = 4.87 \times 10^{-22}$, $7.3 \times 10^{-3} / 4.87 \times 10^{-22} = 1.5 \times 10^{19}$.)

The Units Problem

12.6.1 The large constant $K \sim 10^{19}$ points to a units problem: - The mass formulas are in natural units - Conversion to MeV requires the Planck energy - K contains these conversion factors

Alternative View: Reduction to ξ

12.7.1 If we accept that:

$$m_e \sim \xi^{5/2} \quad (98)$$

$$m_\mu \sim \xi^2 \quad (99)$$

$$\alpha \sim \xi^{11/2} \quad (100)$$

Then these three quantities are determined by the single geometric constant ξ :

$$\begin{aligned} \xi &= \frac{4}{3} \times 10^{-4} \quad (\text{Geometry}) \\ \Downarrow \\ m_e &= f_e(\xi) \\ m_\mu &= f_\mu(\xi) \\ \alpha &= f_\alpha(\xi) \end{aligned}$$

(101)

.13 Derivation of the Coefficients c_e and c_μ

Starting Point: Mass Formulas

13.1.1 The fundamental mass formulas:

$$m_e = c_e \cdot \xi^{5/2} \quad \text{and} \quad m_\mu = c_\mu \cdot \xi^2$$

Step 1: Quantum Numbers and Geometric Factors

13.2.1 The coefficients arise from T0-theory with:

$$\begin{aligned} c_e &= \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \\ c_\mu &= \frac{9}{4\pi\alpha} \end{aligned}$$

Step 2: Derivation of c_e (Electron)

13.3.1 For the electron ($n = 1, l = 0, j = 1/2$):

$$c_e = \frac{\text{Geometry factor} \times \text{Quantum number factor}}{\alpha^{1/2}}$$

$$\text{Geometry factor} = \frac{3\sqrt{3}}{2\pi}$$

$$\text{Quantum number factor} = 1 \quad (\text{for ground state})$$

$$\text{Fine structure correction} = \alpha^{-1/2}$$

$$\Rightarrow c_e = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}}$$

Step 3: Derivation of c_μ (Muon)

13.4.1 For the muon ($n = 2, l = 1, j = 1/2$):

$$c_\mu = \frac{\text{Geometry factor} \times \text{Quantum number factor}}{\alpha}$$

$$\text{Geometry factor} = \frac{9}{4\pi}$$

$$\text{Quantum number factor} = 1$$

$$\text{Fine structure correction} = \alpha^{-1}$$

$$\Rightarrow c_\mu = \frac{9}{4\pi\alpha}$$

Step 4: Physical Interpretation

13.5.1 The different α dependencies reflect:

$$c_e \sim \alpha^{-1/2} \quad (\text{weaker dependence})$$

$$c_\mu \sim \alpha^{-1} \quad (\text{stronger dependence})$$

The different α dependence reflects:

- Electron: Ground state, less sensitive to α
- Muon: Excited state, more strongly dependent on α

Step 5: Dimensional Analysis

13.6.1 Dimensional considerations:

$$[c_e] = [m_e] \cdot [\xi]^{-5/2}$$

$$[c_\mu] = [m_\mu] \cdot [\xi]^{-2}$$

Since ξ is dimensionless (in natural units), both coefficients have the dimension of mass.

Step 6: Consistency Check

13.7.1 With $\alpha \approx 1/137$:

$$c_e \approx \frac{3 \times 1.732}{2 \times 3.1416 \times 0.0854} \approx \frac{5.196}{0.537} \approx 9.67$$

$$c_\mu \approx \frac{9}{4 \times 3.1416 \times 0.0073} \approx \frac{9}{0.0917} \approx 98.1$$

These values give $c_\mu/c_e \approx 10.1$ and hence $m_\mu/m_e = (c_\mu/c_e)\xi^{-1/2} \approx 878$, not the measured 206.77. (Corrected on 30 Sept. 2026: previously "These values match the mass hierarchy $m_\mu/m_e \approx 207$ " – $98.1/9.68 = 10.14$ and $10.14 \times \xi^{-1/2} = 10.14 \times 86.60 = 878$.)

.14 Why Natural Units Are Necessary

The Problem with Conventional Units

14.1.1 In conventional units (SI, cgs) the coefficients c_e and c_μ appear as very large numbers:

$$c_e \approx 1.65 \times 10^{19}$$

$$c_\mu \approx 1.03 \times 10^{20}$$

(Note of 30 Sept. 2026: these values follow from no unit conversion of the formulas $c_e = 3\sqrt{3}/(2\pi\alpha^{1/2}) = 9.68$ and $c_\mu = 9/(4\pi\alpha) = 98.1$; their product 1.7×10^{39} also does not match the $c_e c_\mu \approx 1.5 \times 10^{19}$ required in 12.5.2.)

These large numbers are **artifactual** and arise only from the choice of units.

Natural Units Simplify Physics

14.2.1 In natural units we set:

$$\hbar = c = 1$$

Thus all quantities become dimensionless or have energy dimension.

Transformation to Natural Units

14.3.1 The transformation formulas:

$$m_e^{\text{nat}} = m_e^{\text{SI}} \cdot \sqrt{\frac{G}{\hbar c}}$$

$$m_\mu^{\text{nat}} = m_\mu^{\text{SI}} \cdot \sqrt{\frac{G}{\hbar c}}$$

$$\xi^{\text{nat}} = \xi^{\text{SI}}$$

(Corrected on 30 Sept. 2026: previously $m^{\text{nat}} = m^{\text{SI}} \cdot G/(\hbar c)$ and $\xi^{\text{nat}} = \xi^{\text{SI}} \cdot (\hbar c)^2 - G/(\hbar c)$ has dimension kg^{-2} , $m_e G/(\hbar c) = 1.9 \times 10^{-15} \text{ kg}^{-1}$; correct is $m/m_P = m\sqrt{G/(\hbar c)}$, for the electron 4.19×10^{-23} . A dimensionless ξ is not converted.)

The Coefficients in Natural Units

14.4.1 In natural units the coefficients become **order of magnitude 1**:

$$c_e^{\text{nat}} = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \approx 9.67$$

$$c_\mu^{\text{nat}} = \frac{9}{4\pi\alpha} \approx 98.1$$

Comparison of Representations

14.5.1 The difference:

	Conventional	Natural
c_e	1.65×10^{19}	9.67
c_μ	1.03×10^{20}	98.1
ξ	1.33×10^{-4}	1.33×10^{-4}

Advantages of Natural Units

14.6.1 The advantages of natural units:

1. **Elimination of artifacts**: The large numbers disappear
2. **Physical transparency**: The relationships become easier to follow
3. **Scale invariance**: Fundamental laws become scale-independent
4. **Simpler formulas**: Formulas become shorter and clearer

Example: The Mass Formula

14.7.1 In conventional units:

$$m_e = 1.65 \times 10^{19} \cdot (1.33 \times 10^{-4})^{5/2}$$

In natural units:

$$m_e = 9.67 \cdot \xi^{5/2}$$

Fundamental Interpretation

14.8.1 The coefficients $c_e \approx 9.67$ and $c_\mu \approx 98.1$ in natural units show:

- The lepton masses are **pure numbers**
- The ratio $c_\mu/c_e \approx 10.14$ is a pure number
- The fine structure constant α appears explicitly

.15 The Formula from ξ to α

Fundamental Relationship

15.1.1 The basic equation:

$$\alpha = c_e c_\mu \cdot \xi^{11/2}$$

Coefficients

15.2.1 The precise values:

$$c_e = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \quad (\text{Electron coefficient})$$

$$c_\mu = \frac{9}{4\pi\alpha} \quad (\text{Muon coefficient})$$

Product of Coefficients

15.3.1 The multiplication:

$$c_e c_\mu = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \cdot \frac{9}{4\pi\alpha} = \frac{27\sqrt{3}}{8\pi^2\alpha^{3/2}}$$

Complete Formula

15.4.1 The full expression:

$$\alpha = \frac{27\sqrt{3}}{8\pi^2\alpha^{3/2}} \cdot \xi^{11/2}$$

Solving for α

15.5.1 Rearranging:

$$\alpha^{5/2} = \frac{27\sqrt{3}}{8\pi^2} \cdot \xi^{11/2}$$

$$\alpha = \left(\frac{27\sqrt{3}}{8\pi^2} \right)^{2/5} \cdot \xi^{11/5}$$

.16 T0-Theory: Formulas and Values

In T0-Theory

16.1.1 The fundamental relations:

$$m_e \sim \xi^{5/2} \quad (\text{Electron}) \quad (102)$$

$$m_\mu \sim \xi^2 \quad (\text{Muon}) \quad (103)$$

$$\xi = \frac{4}{3} \times 10^{-4} \quad (104)$$

Correct Assignment in Natural Units

Mass Scaling Laws

16.2.1 The precise formulas:

$$m_e = c_e \cdot \xi^{5/2} \quad (105)$$

$$m_\mu = c_\mu \cdot \xi^2 \quad (106)$$

Geometric Constant

16.2.2 The fundamental parameter:

$$\xi = \frac{4}{3} \times 10^{-4} = 1.333 \times 10^{-4} \quad (107)$$

Calculation of the Characteristic Energy

16.2.3 Step-by-step derivation:

$$E_0 = \sqrt{m_e \cdot m_\mu} = \sqrt{c_e \cdot \xi^{5/2} \cdot c_\mu \cdot \xi^2} \quad (108)$$

$$= \sqrt{c_e c_\mu} \cdot \xi^{9/4} \quad (109)$$

Calculation of the Fine Structure Constant

16.2.4 Complete derivation:

$$\alpha = \xi \cdot E_0^2 = \xi \cdot \left[\sqrt{c_e c_\mu} \cdot \xi^{9/4} \right]^2 \quad (110)$$

$$= \xi \cdot c_e c_\mu \cdot \xi^{9/2} \quad (111)$$

$$= c_e c_\mu \cdot \xi^{11/2} \quad (112)$$

Numerical Values

16.2.5 With $\xi = 1.333 \times 10^{-4}$:

$$\xi^{11/2} = (1.333 \times 10^{-4})^{5.5} \approx 4.87 \times 10^{-22} \quad (113)$$

For $\alpha \approx 1/137 \approx 7.3 \times 10^{-3}$ we need:

$$c_e c_\mu = \frac{\alpha}{\xi^{11/2}} \approx \frac{7.3 \times 10^{-3}}{4.87 \times 10^{-22}} \approx 1.5 \times 10^{19} \quad (114)$$

(Corrected on 30 Sept. 2026: previously 5.19×10^{-22} and $1.4 \times 10^{19} - (1/7500)^{5.5} = 4.87 \times 10^{-22}$, $7.3 \times 10^{-3} / 4.87 \times 10^{-22} = 1.5 \times 10^{19}$.)

Interpretation

16.3.1 The large constant $c_e c_\mu \approx 10^{19}$ corresponds approximately to the ratio of Planck energy to gigaelectron volt and represents the conversion factor between natural units and GeV. (Corrected on 30 Sept. 2026: previously "Planck energy to electron volt" and "MeV" – $E_P/\text{eV} = 1.22 \times 10^{28}$, $E_P/\text{GeV} = 1.22 \times 10^{19}$; whether the closeness to $c_e c_\mu \approx 1.5 \times 10^{19}$ is meaningful remains open.)

17 Definitions

Geometric Constant

17.1.1 The fundamental constant:

$$\xi = \frac{4}{3} \times 10^{-4} = \frac{1}{7500} \quad (115)$$

Mass Formulas

17.2.1 The precise mass relationships:

$$m_e = c_e \cdot \xi^{5/2} \quad (116)$$

$$m_\mu = c_\mu \cdot \xi^2 \quad (117)$$

$$m_\tau = c_\tau \cdot \xi^{3/2} \quad (118)$$

.18 Coefficients from T0-Theory

Electron (n=1, l=0, j=1/2)

18.1.1 The electron coefficient:

$$c_e = \frac{3\sqrt{3}}{2\pi} \cdot \frac{1}{\alpha^{1/2}} \approx 1.6487 \times 10^{19} \quad (119)$$

Muon (n=2, l=1, j=1/2)

18.2.1 The muon coefficient:

$$c_\mu = \frac{9}{4\pi} \cdot \frac{1}{\alpha} \approx 1.0262 \times 10^{20} \quad (120)$$

Tauon (n=3, l=2, j=1/2)

18.3.1 The tauon coefficient:

$$c_\tau = \frac{27\sqrt{3}}{8\pi} \cdot \frac{1}{\alpha^{3/2}} \approx 6.1853 \times 10^{20} \quad (121)$$

(Note of 30 Sept. 2026: the numerical values in 18.1.1–18.3.1 do not follow from the formulas: with $\alpha = 1/137.036$ one gets $\frac{3\sqrt{3}}{2\pi}\alpha^{-1/2} = 9.68$, $\frac{9}{4\pi}\alpha^{-1} = 98.1$ and $\frac{27\sqrt{3}}{8\pi}\alpha^{-3/2} = 2985$; the values 1.6487×10^{19} , 1.0262×10^{20} and 6.1853×10^{20} cannot be derived by any unit conversion.)

.19 Mass Calculation

Electron Mass

19.1.1 Complete calculation:

$$m_e = c_e \cdot \xi^{5/2} \quad (122)$$

$$= \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \cdot \left(\frac{4}{3} \times 10^{-4}\right)^{5/2} \quad (123)$$

$$= 0.5109989461 \text{ MeV} \quad (124)$$

Muon Mass

19.2.1 Complete calculation:

$$m_\mu = c_\mu \cdot \xi^2 \quad (125)$$

$$= \frac{9}{4\pi\alpha} \cdot \left(\frac{4}{3} \times 10^{-4}\right)^2 \quad (126)$$

$$= 105.6583745 \text{ MeV} \quad (127)$$

(Note of 30 Sept. 2026: the masses given here, 0.5109989461 MeV and 105.6583745 MeV, are the measured values (anchors); the coefficient formulas themselves do not lead to them with a common anchor – their ratio gives $m_\mu/m_e \approx 878$ (section on mass ratios), they would need two different scales for m_e and m_μ .)

Tauon Mass

19.3.1 Complete calculation:

$$m_\tau = c_\tau \cdot \xi^{3/2} \quad (128)$$

$$= \frac{27\sqrt{3}}{8\pi\alpha^{3/2}} \cdot \left(\frac{4}{3} \times 10^{-4}\right)^{3/2} \quad (129)$$

$$= 1776.86 \text{ MeV} \quad (130)$$

.20 Characteristic Energy

20.1.1 The precise calculation:

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (131)$$

$$= \sqrt{c_e c_\mu} \cdot \xi^{9/4} \quad (132)$$

$$= \sqrt{\frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \cdot \frac{9}{4\pi\alpha}} \cdot \left(\frac{4}{3} \times 10^{-4}\right)^{9/4} \quad (133)$$

$$= 7.346881 \text{ MeV} \quad (134)$$

(Note of 30 Sept. 2026: With $c_e = 9.68$, $c_\mu = 98.1$ ($\alpha = 1/137$) and $\xi = 4/30000$ the second and third lines give $\sqrt{9.68 \cdot 98.1} \cdot \xi^{9/4} = 5.9 \times 10^{-8}$, not 7.346881 MeV; that value comes from $\sqrt{m_e m_\mu}$ with the T0 masses of Section 8.3.)

.21 Fine Structure Constant

21.1.1 The complete derivation:

$$\alpha = \xi \cdot E_0^2 \quad (135)$$

$$= \xi \cdot c_e c_\mu \cdot \xi^{9/2} \quad (136)$$

$$= c_e c_\mu \cdot \xi^{11/2} \quad (137)$$

$$= \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \cdot \frac{9}{4\pi\alpha} \cdot \left(\frac{4}{3} \times 10^{-4}\right)^{11/2} \quad (138)$$

.22 Numerical Values

22.1.1 Table of values:

(Corrected on 30 Sept. 2026: $\xi^{5/2}$ previously $3.098386676965933 \times 10^{-10} - (4/30000)^{5/2} = 2.0528 \times 10^{-10}$, as in Section 3 ($\frac{32}{9\sqrt{3}} \times 10^{-10}$).)

(Corrected on 30 Sept. 2026: in the table and the following sentence previously “= e (Euler’s number)” – 1.6487... = $\sqrt{e}$, not $e = 2.718...$)

The coefficients contain the constants π and α ; the numerical closeness of the mantissa of c_e to $\sqrt{e}$ is an observation whose significance is open **[S]**.

Quantity	Value	Comment
ξ	$1.33333333333333 \times 10^{-4}$	$= 4/3 \times 10^{-4}$
ξ^2	$1.77777777777778 \times 10^{-8}$	
$\xi^{5/2}$	$2.052800957118670 \times 10^{-10}$	
c_e	$1.648721270700128 \times 10^{19}$	$= \sqrt{e}$ (mantissa)
c_μ	$1.026187714072347 \times 10^{20}$	
m_e	0.5109989461 MeV	Reference value
m_μ	105.6583745 MeV	Reference value
E_0	7.346881 MeV	from T0 masses (Section 8.3)

.23 The Formula from ξ to α (Detailed)

From the Fundamental Relationship

23.1.1 Starting equation:

$$\alpha = c_e c_\mu \cdot \xi^{11/2} \quad (139)$$

Inserting the Coefficients

23.2.1 The detailed calculation:

$$c_e = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \quad (140)$$

$$c_\mu = \frac{9}{4\pi\alpha} \quad (141)$$

$$c_e c_\mu = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \cdot \frac{9}{4\pi\alpha} \quad (142)$$

$$= \frac{27\sqrt{3}}{8\pi^2\alpha^{3/2}} \quad (143)$$

Complete Formula

23.3.1 The full expression:

$$\alpha = \frac{27\sqrt{3}}{8\pi^2\alpha^{3/2}} \cdot \xi^{11/2} \quad (144)$$

Solving for α

23.4.1 Algebraic manipulation:

$$\alpha^{5/2} = \frac{27\sqrt{3}}{8\pi^2} \cdot \xi^{11/2} \quad (145)$$

$$\alpha = \left(\frac{27\sqrt{3}}{8\pi^2} \right)^{2/5} \cdot \xi^{11/5} \quad (146)$$

Numerical Values

23.5.1 Step-by-step calculation:

$$\frac{27\sqrt{3}}{8\pi^2} \approx \frac{46.765}{78.956} \approx 0.5923 \quad (147)$$

$$\left(\frac{27\sqrt{3}}{8\pi^2}\right)^{2/5} \approx (0.5923)^{0.4} \approx 0.8110 \quad (148)$$

$$\xi^{11/5} = \xi^{2.2} = \left(\frac{4}{3} \times 10^{-4}\right)^{2.2} \quad (149)$$

With $\xi = 4/3 \times 10^{-4}$

23.6.1 Final calculation:

$$\xi = 1.333333 \times 10^{-4} \quad (150)$$

$$\xi^{2.2} \approx (1.333333 \times 10^{-4})^{2.2} \quad (151)$$

$$\approx 2.984 \times 10^{-9} \quad (152)$$

$$\alpha \approx 0.8110 \times 2.984 \times 10^{-9} \quad (153)$$

$$\approx 2.42 \times 10^{-9} \quad (154)$$

$$\alpha^{-1} \approx 4.1 \times 10^8 \quad (155)$$

(Corrected on 30 Sept. 2026: previously $(0.5923)^{0.4} \approx 0.8327$, $\xi^{2.2} \approx 8.758 \times 10^{-9}$, $\alpha \approx 7.292 \times 10^{-3}$, $\alpha^{-1} \approx 137.13$ – correct are $0.5923^{0.4} = 0.8110$ and $(1/7500)^{2.2} = 2.984 \times 10^{-9}$, hence $\alpha \approx 2.42 \times 10^{-9}$; even the old product $0.8327 \times 8.758 \times 10^{-9}$ is 7.29×10^{-9} , not 7.29×10^{-3} . This chain does not yield α (cf. the note at 3.6.1).)

Symbol Explanation

23.7.1 Key symbols used:

α	Fine structure constant ($\approx 1/137.036$)
ξ	Geometric space constant ($= \frac{4}{3} \times 10^{-4}$)
c_e	Electron mass coefficient
c_μ	Muon mass coefficient
π	Pi (≈ 3.14159)
$\sqrt{3}$	Square root of 3 (≈ 1.73205)
m_e	Electron mass ($= 0.5109989461$ MeV)
m_μ	Muon mass ($= 105.6583745$ MeV)

With Fractal Correction

23.8.1 Including the fractal factor:

$$\alpha^{-1} = \frac{7500}{m_e m_\mu} \cdot \left(1 - \frac{D_f - 2}{68}\right) = 138.911 \times 0.9862 = 136.99$$

(Corrected on 30 Sept. 2026: previously $138.949 \times 0.9862 = 137.036 - 7500/(0.51099895 \times 105.6583755) = 7500/53.9905 = 138.911$ and $138.911 \times 0.9862 = 136.99$; with $K_{frac} = 1 - 100\xi = 0.98667$ one obtains 137.06.)

Final Fundamental Relationship

23.9.1 The complete formula:

$$\alpha = \left(\frac{27\sqrt{3}}{8\pi^2} \right)^{2/5} \cdot \xi^{11/5} \cdot K_{\text{frac}} \quad \text{with} \quad K_{\text{frac}} = 0.9862$$

(Note of 30 Sept. 2026: as in 3.6.1 – the formula gives $\alpha \approx 2.4 \times 10^{-9}$, and K_{frac} belongs with α^{-1} .)

.24 Observation: α Cancels Out

Equating the Formula Sets

24.1.1 Comparing two representations:

$$\begin{aligned} \text{Simple:} \quad m_e &= \frac{2}{3} \cdot \xi^{5/2} \\ \text{T0-Theory:} \quad m_e &= \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \cdot \xi^{5/2} \end{aligned}$$

After dividing by $\xi^{5/2}$:

$$\frac{2}{3} = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}}$$

Solving for α

24.2.1 Algebraic solution:

$$\alpha^{1/2} = \frac{3\sqrt{3}}{2\pi} \cdot \frac{3}{2} = \frac{9\sqrt{3}}{4\pi} \quad \Rightarrow \quad \alpha = \left(\frac{9\sqrt{3}}{4\pi} \right)^2 = \frac{243}{16\pi^2}$$

For the Muon

24.3.1 Similar analysis:

$$\begin{aligned} \text{Simple:} \quad m_\mu &= \frac{8}{5} \cdot \xi^2 \\ \text{T0-Theory:} \quad m_\mu &= \frac{9}{4\pi\alpha} \cdot \xi^2 \end{aligned}$$

After dividing by ξ^2 :

$$\frac{8}{5} = \frac{9}{4\pi\alpha} \quad \Rightarrow \quad \alpha = \frac{9}{4\pi} \cdot \frac{5}{8} = \frac{45}{32\pi}$$

The Apparent Contradiction

24.4.1 Three different values:

$$\begin{aligned} \text{From electron:} \quad \alpha &= \frac{243}{16\pi^2} \approx 1.539 \\ \text{From muon:} \quad \alpha &= \frac{45}{32\pi} \approx 0.4474 \\ \text{Experimental:} \quad \alpha &\approx 0.007297 \end{aligned}$$

Resolution in the T0 Framework

24.5.1 In the T0 framework, α is **not a free parameter** but a function of ξ :

$$\frac{2}{3} = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \quad \Rightarrow \quad \alpha = \alpha(\xi)$$

$$\frac{8}{5} = \frac{9}{4\pi\alpha}$$

The Key Elements

24.6.1 The key elements:

1. The **geometric factors** ($3\sqrt{3}/2\pi$, $9/4\pi$)
 2. The **powers of α** ($\alpha^{-1/2}$, α^{-1})
 3. The **rational coefficients** ($2/3$, $8/5$)
- are set up so that they **compensate** each other.

Meaning of the Different Representations

24.7.1 Comparative analysis:

- **Simple formulas:** $m_e = \frac{2}{3}\xi^{5/2}$, $m_\mu = \frac{8}{5}\xi^2$
 - Show the pure ξ -dependence
 - Compact and transparent
- **Extended formulas:** $m_e = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}}\xi^{5/2}$, $m_\mu = \frac{9}{4\pi\alpha}\xi^2$
 - Show the **origin** of the coefficients
 - Connect geometry (π , $\sqrt{3}$) with EM coupling (α)
 - But: α is thereby **fixed**, not freely choosable

Central Statement

24.8.1 The central insight:

In the T0 framework the lepton mass hierarchy is fixed by ξ

The different mathematical representations are equivalent descriptions of the same fundamental geometry.

Why This Insight Is Important

24.9.1 The implications:

1. **Unity:** The lepton masses follow from the one parameter ξ (plus structural choices as settings)
2. **Geometric basis:** The coefficients stem from fundamental geometry
3. **α is derived:** The fine structure constant appears as a secondary quantity
4. **Simple structure:** The formulas are compact; mathematical simplicity is a hint, not evidence

.25 Why the Extended Form Is Important

The Two Equivalent Representations

25.1.1 Comparing formulations:

$$\text{Simple form: } m_e = \frac{2}{3} \cdot \xi^{5/2}$$

$$\text{Extended form: } m_e = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}} \cdot \xi^{5/2}$$

The Apparent Contradiction

25.2.1 When equating both formulas:

$$\frac{2}{3} = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}}$$

This yields for α :

$$\alpha = \left(\frac{9\sqrt{3}}{4\pi} \right)^2 = \frac{243}{16\pi^2} \approx 1.539$$

Consequence

25.3.1 The fractions cannot simply cancel out.

The extended form shows that the apparently simple fraction $\frac{2}{3}$ is actually composed of more fundamental geometric and physical constants:

$$\frac{2}{3} = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}}$$

Mathematical Structure

25.4.1 The decomposition:

$$\frac{2}{3} = \frac{\text{Geometry factor}}{\alpha^{1/2}}$$

$$\text{with Geometry factor} = \frac{3\sqrt{3}}{2\pi} \approx 0.826$$

Physical Interpretation

25.5.1 Interpretation:

- $\frac{2}{3}$ is **not** a simple rational fraction
- In the extended form it is composed of:
 - Space geometry ($\pi, \sqrt{3}$)
 - Electromagnetic coupling (α)
 - Quantum numbers (implicit in the coefficients)
- The extended form shows this origin

Why Both Representations Are Important

25.6.1 Complementary perspectives:

Simple Form

Shows pure ξ -dependence
Compact
Practical for calculations
Hides the composition

Extended Form

Shows physical origin
Shows the physical origin
Fundamental for understanding
Shows the composition

The Actual Statement of T0-Theory

25.7.1 Core statement:

$$\frac{2}{3} \neq \text{simple fraction} \quad \text{but rather} \quad \frac{2}{3} = \frac{3\sqrt{3}}{2\pi\alpha^{1/2}}$$

The extended form is necessary to show:

1. That the fractions do **not** simply cancel
2. That the apparently simple coefficient $\frac{2}{3}$ actually has a complex structure
3. That α is part of this structure, even if it formally cancels out
4. That the geometry of space $(\pi, \sqrt{3})$ is embedded in it

Why No Fractal Correction is Needed for Mass Ratios and Characteristic Energy

1. Different Calculation Approaches

Path A: $\alpha = \frac{m_e m_\mu}{7500}$ (requires correction)

Path B: $\alpha = \frac{E_0^2}{7500}$ (requires correction)

Path C: $\frac{m_\mu}{m_e} = f(\alpha)$ (no correction needed)

Path D: $E_0 = \sqrt{m_e m_\mu}$ (no correction needed)

2. Mass Ratios Are Correction-Free

The lepton mass ratio:

$$\frac{m_\mu}{m_e} = \frac{c_\mu \xi^2}{c_e \xi^{5/2}} = \frac{c_\mu}{c_e} \xi^{-1/2}$$

Substituting the coefficients:

$$\frac{m_\mu}{m_e} = \frac{\frac{9}{4\pi\alpha}}{\frac{3\sqrt{3}}{2\pi\alpha^{1/2}}} \cdot \xi^{-1/2} = \frac{\sqrt{3}}{2\alpha^{1/2}} \cdot \xi^{-1/2}$$

(Corrected on 30 Sept. 2026: previously $\frac{3\sqrt{3}}{2\alpha^{1/2}}$ – the quotient $\frac{9}{4\pi\alpha} / \frac{3\sqrt{3}}{2\pi\alpha^{1/2}}$ is $\frac{\sqrt{3}}{2\alpha^{1/2}}$; with these coefficients $m_\mu/m_e \approx 878$.)

3. Why the Ratio is Correct

The fractal correction cancels out in the ratio.

$$\frac{m_\mu}{m_e} = \frac{K_{\text{frac}} \cdot m_\mu}{K_{\text{frac}} \cdot m_e} = \frac{m_\mu}{m_e}$$

The same correction factor affects both masses and cancels in the ratio.

4. Characteristic Energy is Correction-Free

$$E_0 = \sqrt{m_e m_\mu} = \sqrt{K_{\text{frac}} m_e \cdot K_{\text{frac}} m_\mu} = K_{\text{frac}} \cdot \sqrt{m_e m_\mu}$$

However: E_0 is itself an observable. The corrected characteristic energy is:

$$E_0^{\text{corr}} = \sqrt{m_e^{\text{corr}} m_\mu^{\text{corr}}} = K_{\text{frac}} \cdot E_0^{\text{bare}}$$

(Note of 30 Sept. 2026: $E_0^{\text{corr}} = K_{\text{frac}} \cdot E_0^{\text{bare}}$ contradicts Section 3.2.2 and R72, where $E_0 = E_0^{\text{bare}} / \sqrt{K_{\text{frac}}}$: $7.348 / \sqrt{0.98667} = 7.397 \text{ MeV}$, whereas $0.98667 \cdot 7.348 = 7.250 \text{ MeV}$. The same applies to E_0^{exp} below.)

5. Consistent Treatment

$$\begin{aligned} m_e^{\text{exp}} &= K_{\text{frac}} \cdot m_e^{\text{bare}} \\ m_\mu^{\text{exp}} &= K_{\text{frac}} \cdot m_\mu^{\text{bare}} \\ E_0^{\text{exp}} &= K_{\text{frac}} \cdot E_0^{\text{bare}} \end{aligned}$$

6. Calculating α via Mass Ratio

$$\frac{m_\mu}{m_e} = \frac{105.6583745}{0.5109989461} = 206.768282$$

Theoretical prediction (without correction):

$$\frac{m_\mu}{m_e} = \frac{8/5}{2/3} \cdot \xi^{-1/2} = \frac{12}{5} \cdot \xi^{-1/2}$$

7. Why Different Paths Require Different Treatments

No Correction Needed

Mass ratios
Characteristic energy E_0
Scale ratios
Dimensionless quantities

Correction Required

Absolute mass values
Fine structure constant α
Absolute energies
Dimensionful quantities

8. Physical Interpretation

- **Relative quantities:** Ratios are independent of absolute scale
- **Absolute quantities:** Require correction for absolute energy scale
- **Fractal dimension:** Affects absolute scaling, not ratios

9. Mathematical Reason

The fractal correction acts as a multiplicative factor:

$$m^{\text{exp}} = K_{\text{frac}} \cdot m^{\text{bare}}$$

For ratios:

$$\frac{m_1^{\text{exp}}}{m_2^{\text{exp}}} = \frac{K_{\text{frac}} \cdot m_1^{\text{bare}}}{K_{\text{frac}} \cdot m_2^{\text{bare}}} = \frac{m_1^{\text{bare}}}{m_2^{\text{bare}}}$$

10. Comparison with Experiment

$$\left(\frac{m_\mu}{m_e}\right)_{\text{exp}} = 206.768282$$

$$\left(\frac{m_\mu}{m_e}\right)_{\text{theo}} = 207.85 \quad (\text{without correction})$$

(Corrected on 30 Sept. 2026: previously theoretical value 206.768282 – that is the measured value (anchor); the prediction without correction is $\frac{12}{5}\xi^{-1/2} = 207.85$.)

Is This Indirect Proof That the Fractal Correction is Correct?

The Consistency Argument

The behaviour is consistent with the fractal correction. Since a common multiplicative factor cancels in every ratio, this is a consistency hint, but on its own not yet independent evidence.

1. The Theoretical Framework

The T0-theory proposes:

$$m_e = \frac{2}{3} \cdot \xi^{5/2} \cdot K_{\text{frac}}$$

$$m_\mu = \frac{8}{5} \cdot \xi^2 \cdot K_{\text{frac}}$$

$$\alpha = \frac{m_e m_\mu}{7500} \cdot \frac{1}{K_{\text{frac}}}$$

2. The Consistency Test

If the fractal correction is valid, then:

$$\frac{m_\mu}{m_e} = \frac{\frac{8}{5} \cdot \xi^2 \cdot K_{\text{frac}}}{\frac{2}{3} \cdot \xi^{5/2} \cdot K_{\text{frac}}} = \frac{12}{5} \cdot \xi^{-1/2}$$

3. Experimental Verification

$$\begin{aligned}\left(\frac{m_\mu}{m_e}\right)_{\text{theo}} &= \frac{12}{5} \cdot (1.333 \times 10^{-4})^{-1/2} \\ &= 2.4 \times 86.6 = 207.84 \\ \left(\frac{m_\mu}{m_e}\right)_{\text{exp}} &= 206.768\end{aligned}$$

The 0.5% difference is not broken down further in this document.

4. What Speaks for the Correction

1. **Self-consistency:** The correction cancels where the multiplicative ansatz leads one to expect it
2. **Predictive power:** Mass ratios work without correction
3. **Explanatory power:** Absolute values need correction
4. **Parameter economy:** One correction factor (K_{frac}) describes the deviations of the absolute values considered here

5. Comparison with Alternative Theories

Without fractal correction:

$$\begin{aligned}\alpha^{-1} &= 138.93 \quad (\text{calculated}) \\ \alpha^{-1} &= 137.036 \quad (\text{experimental}) \\ \text{Error} &= 1.38\%\end{aligned}$$

With fractal correction:

$$\alpha^{-1} = 138.93 \times 0.9862 = 137.01 \quad (\text{with correction})$$

(Corrected on 30 Sept. 2026: previously $= 137.036 - 138.93 \times 0.9862 = 137.01$; with $K_{\text{frac}} = 1 - 100\xi = 0.98667$ one obtains $138.91 \times 0.98667 = 137.06$.)

6. The Philosophical Argument

That the correction is needed for absolute values and cancels in ratios is compatible with its interpretation as a real physical effect. Whether it is more than a fitting factor will be decided by an independent derivation of K_{frac} .

7. Further Connections

- The correction factor $K_{\text{frac}} = 0.9862$ is motivated in the T0 framework by fractal geometry
- It connects to the effective fractal dimension $D_f^{\text{eff}} \approx 2.973$ of spacetime (Note of 30 Sept. 2026: $K_{\text{frac}} = 1 - (D_f - 2)/68 = 0.9862$ follows from the older anchor $D_f = 2.94$; with $D_f^{\text{eff}} = 2.973$ the same formula gives 0.9857.)
- For the value $C = 68$ a connection to tetrahedral symmetry is conjectured [S]

9. Remaining Open Questions

- Direct measurement of spacetime's fractal dimension
- Extension to other particle families