

T0 Model: Field-Theoretic Derivation of the β -Parameter

Contents

.1	Introduction and Motivation	1
.2	Natural Units Framework	1
.3	Fundamental Structure of the T0 Model	2
	Time-Mass Duality	2
	Fundamental Field Equation	2
.4	Geometric Derivation of the β -Parameter	2
	Spherically Symmetric Point Source	2
	Solution of the Field Equation	3
	Determination of Integration Constants	3
	The Characteristic Length Scale	3
	Definition of the β -Parameter	4
.5	Physical Interpretation of the β -Parameter	4
	Dimensional Analysis	4
	Connection to Classical Physics	4
	Limiting Cases and Application Domains	4
.6	Comparison with Established Theories	5
	Connection to General Relativity	5
	Differences from the Standard Model	5
.7	Experimental Predictions	5
	Time Dilation Effects	5
	Spectroscopic Tests	5
.8	Mathematical Consistency	5
	Conservation Laws	5
	Solution Stability	6

Correction notes

Corrections from the calculation check (30 Sept. 2026). The following places are corrected or annotated in the text; each carries a dated note with the previous value:

- Integration constant: from $4\pi\epsilon^2(-C_1/\epsilon^2) = 4\pi Gm_0/T_0$ it follows that $C_1 = -Gm_0/T_0$, not $+Gm_0/T_0$ (sign of $m(r)$ and $T(r)$ reversed); factor 2 in $\beta = 2Gm/r$ and the claim that $dt/dt_0 = 1 - 2Gm/r$ is GR to first order (GR: $1 - Gm/r$) (each annotated); table: hydrogen atom $\beta \sim 10^{-39} \rightarrow \sim 5 \times 10^{-44}$.

.1 Introduction and Motivation

The T0 model introduces a fundamentally new perspective on spacetime, where time itself becomes a dynamic field. At the center of this theory lies the dimensionless β -parameter, which characterizes the strength of the time field and establishes a direct connection between gravitational and electromagnetic interactions.

This work focuses exclusively on the mathematically rigorous derivation of the β -parameter from the fundamental field equations of the T0 model, avoiding the complexity of additional scaling parameters.

Central Result

The β -parameter is derived as:

$$\beta = \frac{2Gm}{r} \quad (1)$$

where G is the gravitational constant, m is the source mass, and r is the distance from the source.

.2 Natural Units Framework

The T0 model employs the system of natural units established in modern quantum field theory [Peskin & Schroeder(1995), Weinberg(1995)]:

- $\hbar = 1$ (reduced Planck constant)
- $c = 1$ (speed of light)

This system reduces all physical quantities to energy dimensions and follows the tradition established by Dirac [Dirac(1958)].

Dimensions in Natural Units

- Length: $[L] = [E^{-1}]$
- Time: $[T] = [E^{-1}]$
- Mass: $[M] = [E]$
- The β -parameter: $[\beta] = [1]$ (dimensionless)

.3 Fundamental Structure of the T0 Model

Time-Mass Duality

The central principle of the T0 model is the time-mass duality, which states that time and mass are inversely linked. This relationship differs fundamentally from the conventional treatment in general relativity [Einstein(1915), Misner et al.(1973)].

Theory	Time	Mass	Reference
Einstein GR	$dt' = \sqrt{g_{00}}dt$	$m_0 = \text{const}$	[Einstein(1915), Misner et al.(1973)]
Special Relativity	$t' = \gamma t$	$m_0 = \text{const}$	[Einstein(1905)]
T0 Model	$T(x) = \frac{1}{m(x)}$	$m(x) = \text{dynamic}$	This work

Table 1: Comparison of time-mass treatment in different theories

Fundamental Field Equation

The fundamental field equation of the T0 model is derived from variational principles, analogous to the approach for scalar field theories [Weinberg(1995)]:

$$\nabla^2 m(x) = 4\pi G \rho(x) \cdot m(x) \quad (2)$$

This equation shows structural similarity to the Poisson equation of gravitation $\nabla^2 \phi = 4\pi G \rho$ [Jackson(1998)], but is nonlinear due to the factor $m(x)$ on the right-hand side.

The time field follows directly from the inverse relationship:

$$T(x) = \frac{1}{m(x)} \quad (3)$$

.4 Geometric Derivation of the β -Parameter

Spherically Symmetric Point Source

For a point mass source, we use the established methodology for solving Einstein's field equations [Schwarzschild(1916), Misner et al.(1973)]. The mass density of a point source is described by the Dirac delta function:

$$\rho(\vec{x}) = m_0 \cdot \delta^3(\vec{x}) \quad (4)$$

where m_0 is the mass of the point source.

Solution of the Field Equation

Outside the source ($r > 0$), where $\rho = 0$, the field equation reduces to:

$$\nabla^2 m(r) = 0 \quad (5)$$

The spherically symmetric Laplace operator [Jackson(1998), Griffiths(1999)] yields:

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{dm}{dr} \right) = 0 \quad (6)$$

The general solution to this equation is:

$$m(r) = \frac{C_1}{r} + C_2 \quad (7)$$

Determination of Integration Constants

Asymptotic boundary condition: For large distances, the time field should assume a constant value T_0 :

$$\lim_{r \rightarrow \infty} T(r) = T_0 \quad \Rightarrow \quad \lim_{r \rightarrow \infty} m(r) = \frac{1}{T_0} \quad (8)$$

This gives us: $C_2 = \frac{1}{T_0}$

Behavior at the origin: Using Gauss's theorem [Griffiths(1999), Jackson(1998)] for a small sphere around the origin:

$$\oint_S \nabla m \cdot d\vec{S} = 4\pi G \int_V \rho(r) m(r) dV \quad (9)$$

For a small radius ϵ :

$$4\pi\epsilon^2 \left. \frac{dm}{dr} \right|_{r=\epsilon} = 4\pi G m_0 \cdot m(\epsilon) \quad (10)$$

With $\frac{dm}{dr} = -\frac{C_1}{r^2}$ and $m(\epsilon) \approx \frac{1}{T_0}$ for small ϵ :

$$4\pi\epsilon^2 \cdot \left(-\frac{C_1}{\epsilon^2}\right) = 4\pi Gm_0 \cdot \frac{1}{T_0} \quad (11)$$

This yields: $C_1 = \frac{Gm_0}{T_0}$ (Note of 30 Sept. 2026: The preceding equation gives $-C_1 = Gm_0/T_0$, hence $C_1 = -Gm_0/T_0$. The solution then reads $m(r) = \frac{1}{T_0}(1 - Gm_0/r)$ and $T(r) \approx T_0(1 + Gm_0/r)$; the signs in the following equations are reversed accordingly.)

The Characteristic Length Scale

The complete solution reads:

$$m(r) = \frac{1}{T_0} \left(1 + \frac{Gm_0}{r}\right) \quad (12)$$

The corresponding time field is:

$$T(r) = \frac{T_0}{1 + \frac{Gm_0}{r}} \quad (13)$$

For the practically important case $Gm_0 \ll r$, we obtain the approximation:

$$T(r) \approx T_0 \left(1 - \frac{Gm_0}{r}\right) \quad (14)$$

The characteristic length scale at which the time field significantly deviates from T_0 is:

$$\boxed{r_0 = Gm_0} \quad (15)$$

This scale is proportional to half the Schwarzschild radius $r_s = 2GM/c^2 = 2Gm$ in geometric units [[Misner et al.\(1973\)](#), [Carroll\(2004\)](#)].

Definition of the β -Parameter

The dimensionless β -parameter is defined as the ratio of the characteristic length scale to the actual distance:

$$\boxed{\beta = \frac{r_0}{r} = \frac{Gm_0}{r}} \quad (16)$$

This parameter measures the relative strength of the time field at a given point. For astronomical objects, we can write the more general form:

$$\boxed{\beta = \frac{2Gm}{r}} \quad (17)$$

where the factor of 2 arises from the complete relativistic treatment, analogous to the emergence of the Schwarzschild radius. (Note of 30 Sept. 2026: The derivation above yields $\beta = Gm_0/r = r_s/(2r)$, as also stated in the section Connection to classical physics. The factor 2 does not follow from it but is set here; $\beta = 2Gm/r = r_s/r$ is twice as large. The same applies to the Central Result at the beginning.)

.5 Physical Interpretation of the β -Parameter

Dimensional Analysis

The dimensionlessness of the β -parameter in natural units:

$$[\beta] = \frac{[G][m]}{[r]} = \frac{[E^{-2}][E]}{[E^{-1}]} = [1] \quad (18)$$

Connection to Classical Physics

The β -parameter shows direct connections to established physical concepts:

- **Gravitational potential:** β is proportional to the Newtonian potential $\Phi = -Gm/r$
- **Schwarzschild radius:** $\beta = r_s/(2r)$ in geometric units
- **Escape velocity:** β is related to v_{esc}^2/c^2

Limiting Cases and Application Domains

Physical System	Typical β -Value	Regime
Hydrogen atom	$\sim 5 \times 10^{-44}$	Quantum mechanics
Earth (surface)	$\sim 10^{-9}$	Weak gravitation
Sun (surface)	$\sim 10^{-6}$	Stellar physics
Neutron star	~ 0.1	Strong gravitation
Schwarzschild horizon	$\beta = 1$	Limiting case

Table 2: Typical β -values for various physical systems

(Corrected on 30 Sept. 2026: hydrogen atom previously $\sim 10^{-39} - 2Gm_p/(a_0c^2) = 4.7 \times 10^{-44}$ (without the factor 2: 2.3×10^{-44}); 10^{-39} is the ratio of gravitational to Coulomb force, not β .)

.6 Comparison with Established Theories

Connection to General Relativity

In general relativity, the parameter $rs/r = 2Gm/r$ characterizes the strength of the gravitational field. The T0 parameter $\beta = 2Gm/r$ is identical to this expression, revealing a deep connection between both theories.

Differences from the Standard Model

While the Standard Model of particle physics treats time as an external parameter, the T0 model makes time a dynamic field. The β -parameter quantifies this dynamics and represents a measurable deviation from standard physics.

.7 Experimental Predictions

Time Dilation Effects

The T0 model predicts a modified time dilation:

$$\frac{dt}{dt_0} = 1 - \beta = 1 - \frac{2Gm}{r} \quad (19)$$

This relationship is identical to the gravitational time dilation of GR in first order, but offers a fundamentally different theoretical foundation. *(Note of 30 Sept. 2026: In GR, $dt/dt_0 = \sqrt{1 - 2Gm/r} \approx 1 - Gm/r$ to first order; $1 - 2Gm/r$ is twice as large and does not agree with GR to first order. With $\beta = Gm/r$ from the derivation, $1 - \beta$ would be identical.)*

Spectroscopic Tests

The β -parameter could be tested through high-precision spectroscopy:

- Gravitational redshift in stellar spectra
- Atomic clock experiments in different gravitational potentials
- High-precision interferometry

.8 Mathematical Consistency

Conservation Laws

The derivation of the β -parameter respects fundamental conservation laws:

- **Energy conservation:** Guaranteed by the Lagrangian formulation
- **Momentum conservation:** From spatial translation invariance
- **Dimensional consistency:** Verified in all derivation steps

Solution Stability

The spherically symmetric solution is stable against small perturbations, which can be shown by linearization around the ground state solution.

Bibliography

- [Carroll(2004)] Carroll, S. M. *Spacetime and Geometry: An Introduction to General Relativity*. Addison-Wesley, San Francisco, CA (2004).
- [Dirac(1958)] Dirac, P. A. M. *The Principles of Quantum Mechanics*. Oxford University Press, Oxford, 4th edition (1958).
- [Einstein(1905)] Einstein, A. Zur Elektrodynamik bewegter Körper. *Annalen der Physik*, **17**, 891–921 (1905).
- [Einstein(1915)] Einstein, A. Die Feldgleichungen der Gravitation. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften*, 844–847 (1915).
- [Griffiths(1999)] Griffiths, D. J. *Introduction to Electrodynamics*. Prentice Hall, Upper Saddle River, NJ, 3rd edition (1999).
- [Jackson(1998)] Jackson, J. D. *Classical Electrodynamics*. John Wiley & Sons, New York, 3rd edition (1998).
- [Misner et al.(1973)] Misner, C. W., Thorne, K. S., and Wheeler, J. A. *Gravitation*. W. H. Freeman and Company, New York (1973).
- [Peskin & Schroeder(1995)] Peskin, M. E. and Schroeder, D. V. *An Introduction to Quantum Field Theory*. Addison-Wesley, Reading, MA (1995).
- [Schwarzschild(1916)] Schwarzschild, K. Über das Gravitationsfeld eines Massenpunktes nach der Einsteinschen Theorie. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften*, 189–196 (1916).
- [Weinberg(1995)] Weinberg, S. *The Quantum Theory of Fields, Volume I: Foundations*. Cambridge University Press, Cambridge (1995).