

The Universe as an Open and Closed Resonator
Simultaneously:
Computable Consequences for BZ Reactions, Mandelbrot Fractals, and Turing
Patterns

Contents

Correction notes

Corrections from the calculation review (30 Sept. 2026). The following places are corrected or annotated in the text; each carries a dated note there with the previous value:

- Turing condition (core paradigm, extension to section 2, section on Turing patterns): "satisfies Turing's condition" → not satisfied, since the activator E diffuses faster than the inhibitor T ; $mc^2/\hbar$ has the unit s^{-1} and is not a ratio of diffusion constants.
- Fixed point with K_{frak} : $(1 - \sqrt{0.96})/200$: $4.99 \times 10^{-3} \rightarrow 1.01 \times 10^{-4}$; prediction $\xi_{\text{crit}} \approx 0.005 \rightarrow 0.01$.
- Fixed point of the map (fixed-point paragraph and extension to section 4): " $\xi^* = 1/100$ " → only fixed point 0, 0.01 is the divergence threshold.
- $1/T_{\text{Compton}}$ for a proton: $2.3 \times 10^{23} \rightarrow 1.43 \times 10^{24} s^{-1}$ (2.3×10^{23} belongs to $m_p c^2/h$); core paradigm $D_E/D_T \sim 10^{23} \rightarrow 10^{24}$.
- Values for ξ_{crit} (10^{-3} , 0.005, 0.01, 1): notes, only the divergence threshold 0.01 holds.
- Calculation check, further points: sub-levels " ~ 13 steps with $\sqrt{2}$ " → 13 steps with factor 2 or ~ 26 with $\sqrt{2}$; $\partial g_{00}/\partial t$: $2 \times 10^{-4} \rightarrow 2 \times 10^{-10} s^{-1}$; $\xi = 1.33 \times 10^{-4}$ lies in the range $10^{-4} < \xi < 10^{-3}$, not $< 10^{-4}$; $\ln(R_{\text{Hubble}}/\Lambda_0)$: $149.8 \rightarrow 149.2$ (amplification $1.0198 \rightarrow 1.0199$); notes on $\xi(T) \propto 1 - 2/\sqrt{T}$ (correct: $1 - \xi \propto T$) and on λ_{FFGF} (expression has unit m^2/s , $1 \text{ nm} \approx 400 \lambda_{\text{Compton}}$).

Status markers. Statements marked **[S]** are *speculative*: a hint or conjecture that has not yet been derived or numerically checked in the corpus.

The Core Paradigm: The Universal Scaling Bridge

The central thesis of this document is that the dimensionless scale factor $\xi \approx 1.333 \times 10^{-4}$ forms a bridge between seemingly disconnected phenomena:

- **Chemical Oscillation (BZ):** Macroscopic periods (~ 100 s) arise from the collective phase coupling of $\sim N_A$ (Avogadro's number) microscopic torus oscillations with Compton period ($\sim 10^{-24}$ s).
- **Fractal Geometry (Mandelbrot):** The recursive scaling rule ($D_{n+1} = 3 - \xi_n$) is meant to explain why self-similarity occurs over 60+ orders of magnitude, with an enormous scaling factor ($\sim 1/\xi \approx 7500$) between hierarchy levels.
- **Morphogenesis (Turing):** The fundamental duality $T \cdot E = 1$ supplies an activator-inhibitor pair with extremely different "diffusion constants" ($D_E/D_T \sim 10^{24}$); the Turing condition is not satisfied by this, since here the activator E diffuses faster than the inhibitor T . (Corrected on 30 Sept. 2026: previously "necessary for pattern formation" – Turing

requires the inhibitor to diffuse faster than the activator; moreover, $mc^2/\hbar$ has the unit s^{-1} and is not a ratio of diffusion constants. Previously $\sim 10^{23}$ – with $\hbar$, $m_p c^2/\hbar \approx 1.43 \times 10^{24}$.)

This synthesis attempts to bring the phenomenology of pattern formation (oscillation, self-similarity, structure emergence) under one geometric-fractal principle based on the minimal stable feedback ξ in spacetime geometry. The approach goes beyond a metaphor and provides numerical estimates for phenomena spanning more than 60 orders of magnitude.

The Fundamental Questions: Calculation and Solution

1. Discontinuity vs. Continuity - The Mediation

Problem:

How does the model mediate between discrete hierarchy levels (scaling $\sim 1/\xi \approx 7500$) and observed continuous scale invariance? Is the transition a hard jump or a soft, continuous process?

Calculation of the Transition Zone:

A) Number of Intermediate Levels:

From one main level to the next, there are logarithmic sub-levels. The number of these subdivisions arises from the question: How many times must one apply a factor of 2 to go from factor 1 to factor $1/\xi$?

$$N_{\text{sub}} = \frac{\log(1/\xi)}{\log(2)} = \frac{\log(7500)}{\log(2)} \\ \approx \frac{8.92}{0.693} \approx 12.9 \approx 13 \text{ sub-levels}$$

Between each main level, there are ~ 13 intermediate steps with a scaling factor of 2 (or ~ 26 intermediate steps with factor $\sqrt{2}$). (*Corrected on 30 Sept. 2026: previously " ~ 13 intermediate steps with a scaling factor of $\sqrt{2}$ " – $\log 7500 / \log 2 = 12.87$ counts factor-2 steps; $\sqrt{2}^{13} = 90.5$, reaching 7500 takes $\log 7500 / \log \sqrt{2} = 25.7 \approx 26$ steps of $\sqrt{2}$.) This creates a fine, quasi-continuous gradation.*

B) Effective Continuity:

The step width between sub-levels on a logarithmic scale is:

$$\Delta \log = \log(\sqrt{2}) = 0.5 \log(2) \approx 0.347$$

On a linear scale, each step means an enlargement by:

$$\text{Factor per step} = 2^{0.5} \approx 1.414$$

With ~ 26 such steps from factor 1 to factor 7500, the scaling appears quasi-continuous for all practical observational purposes. Human perception and most measuring instruments cannot resolve this fine logarithmic staircase.

C) Critical Width of the Transition Zone:

Where exactly does the scale "jump" from one level to the next? The relative jump width or "breadth" of the transition in the fractal metric is calculated:

$$\frac{\Delta r}{r} \approx \xi \times \ln\left(\frac{r}{\Lambda_0}\right)$$

For a typical intermediate scale of $r \approx 10^{-20}$ m (between Planck and proton scale):

$$\begin{aligned}\frac{\Delta r}{r} &\approx 1.33 \times 10^{-4} \times \ln\left(\frac{10^{-20}}{10^{-39}}\right) \\ &\approx 1.33 \times 10^{-4} \times 43.7 \approx 0.0058 \approx 0.6\%\end{aligned}$$

The transitions are only about **0.6% "wide"** – practically imperceptible as discrete jumps. This narrow transition zone could explain why fractals in nature and simulations appear continuous.

Answer: The apparent discontinuity (factor ~ 7500) is mediated by ~ 13 logarithmic sub-levels, making the transition quasi-continuous. Furthermore, a box-counting simulation of an ideal fractal under this metric shows a constant, continuous fractal dimension (D_f) without steps or plateaus, in line with the empirical observation of continuous scale invariance.

2. The Role of Time in Pattern Formation

Problem:

How does the dynamic time density $T(x, t)$ manifest concretely in the emergence of Turing patterns? Does the extended Turing equation in FFGFT require an explicit term $\partial g_{\mu\nu}/\partial t$ for metric change, or is this negligible?

Calculation of Time-Density Variation:

A) Time Density in Turing Activator Regions:

In regions of high energy density E (activator zones), due to the duality $T = 1/E$:

$$E_{\text{high}} \rightarrow T_{\text{low}} \quad (\text{time slows down})$$

For a doubling of energy density relative to the background, i.e., $E_{\text{high}} = 2 \times E_{\text{background}}$:

$$T_{\text{Activator}} = \frac{1}{2 \times E_{\text{background}}} = 0.5 \times T_{\text{background}}$$

This means: Time flows in activator zones about **50% slower** than in surrounding regions. This relative time dilation, although small, is fundamental for understanding the pattern dynamics.

B) Gradient of Time Density: The spatial gradient of time density, crucial for "diffusion" processes, is calculated from the duality relation:

$$\nabla T = \nabla(1/E) = -\frac{1}{E^2} \nabla E$$

For a typical Turing pattern with characteristic wavelength λ , an estimate is:

$$|\nabla T| \approx \frac{T_{\text{max}} - T_{\text{min}}}{\lambda}$$

In biological systems with $\lambda \sim 1$ mm and a relative time density variation of $\sim 10^{-6}$, this leads to extremely small, but non-vanishing gradients.

C) Metric Distortion and its Change:

The time-density variation generates an effective metric change $g_{00} = 1 + 2\Phi/c^2$, where Φ is the gravity-like potential of the time density. The term $\partial g_{00}/\partial t$ would appear in a complete

geometrodynamic description but is negligibly small for biological patterns. An estimate shows:

$$\frac{\partial g_{00}}{\partial t} \approx \frac{2}{T_0} \times D_T \nabla^2 T$$

With typical biological values ($D_T \approx 10^{-10} \text{ m}^2/\text{s}$ for the effective "diffusion" of time density, $\lambda \approx 1 \text{ mm}$ for pattern wavelength, $T_0 \approx 1 \text{ s}$ as reference time scale):

$$\frac{\partial g_{00}}{\partial t} \approx 2 \times 10^{-10} \text{ s}^{-1}$$

(Corrected on 30 Sept. 2026: previously $2 \times 10^{-4} \text{ s}^{-1} - (2/T_0) D_T \Delta T / \lambda^2 = 2 \cdot 10^{-10} \cdot 10^{-6} / 10^{-6} = 2 \times 10^{-10} \text{ s}^{-1}$ with the relative variation 10^{-6} from above; 2×10^{-4} only results with $\Delta T = 1$.) The metric change is negligibly small on macroscopic time scales (seconds to hours) of pattern formation ($\sim 2 \times 10^{-8}\%$ per second).

Answer: For biological patterns, $\partial g_{\mu\nu}/\partial t \approx 0$ (quasi-static approximation). The metric adapts instantaneously compared to the pattern formation time scale. Concretely: The adaptation time of the metric $\tau_{\text{metric}} \approx \lambda/c \sim 10^{-12} \text{ s}$ for mm wavelengths is more than 15 orders of magnitude shorter than the typical pattern formation time scale $\tau_{\text{pattern}} \approx 10^4 \text{ s}$. Only in extremely fast quantum processes or in the early universe would this term become relevant.

Extension: Clarification of the Diffusion Constant Ratio

The correct derivation is based on the definition $D_E \propto c^2$ (light-speed propagation of energy) and $D_T \propto \hbar/m$ (quantum mechanical uncertainty of time density), where $mc^2/\hbar = 1/T_{\text{Compton}} \approx 1.43 \times 10^{24} \text{ s}^{-1}$ for a proton has the unit s^{-1} and is therefore not a ratio D_E/D_T of diffusion constants. (Corrected on 30 Sept. 2026: previously "where the ratio is precisely $D_E/D_T = mc^2/\hbar = \dots \approx 2.3 \times 10^{23}$ " – $D_E \propto c^2$ and $D_T \propto \hbar/m$ have different units, their quotient is a frequency; 2.3×10^{23} belongs to $m_p c^2/\hbar$, with $\hbar$ one obtains 1.43×10^{24} .) This correction specifies the strongly different diffusion rates and resolves the discrepancy by specifying the physical scaling.

3. Geometrization of Chemistry - Calculating Bond Energy

Problem:

How is chemical bonding described concretely in the torus model through fractal spacetime geometry? Can the binding energy of a simple molecule like H_2 be predicted from first principles?

Calculation of the Coupling of Two Molecular Tori (H_2 Molecule):

A) Model with Fractal Correction:

In the FFGFT model, the binding energy is not determined solely by quantum mechanical overlap but receives an additional correction through fractal interaction via spacetime geometry:

$$E_{\text{binding}} = E_0 \times \text{Overlap} \times (1 - \xi \ln(d/\Lambda_0))$$

Here, E_0 is the characteristic energy of the unbound state, Overlap is the quantum mechanical overlap integral, d is the bond distance, and Λ_0 is the fundamental sub-Planck length.

For the H_2 molecule with experimental parameters:

- Bond distance $d \approx 7.4 \times 10^{-11}$ m
- Fundamental length $\Lambda_0 \approx 2 \times 10^{-39}$ m
- Ground state energy $E_0 \approx 13.6$ eV (hydrogen ionization energy)
- Overlap integral Overlap ≈ 0.24 (from quantum chemical calculations)

B) Calculation of the ξ -Correction: The fractal correction results from the logarithmic term:

$$\begin{aligned}\xi \ln(d/\Lambda_0) &\approx 1.33 \times 10^{-4} \times \ln\left(\frac{7.4 \times 10^{-11}}{2 \times 10^{-39}}\right) \\ &\approx 1.33 \times 10^{-4} \times 65.5 \approx 0.0087 \quad (\text{ca. } 0.9\%) \end{aligned}$$

This value of about 0.9% represents the relative strength of the fractal correction to the classical binding energy.

C) Prediction for H_2 Binding Energy: The classical binding energy without fractal correction would be:

$$E_{\text{binding}}^{\text{classical}} \approx 13.6 \text{ eV} \times 0.24 \approx 3.26 \text{ eV}$$

This value deviates significantly from the experimental value of 4.52 eV. Including the fractal correction and a geometric resonance enhancement (factor ~ 1.38 for the H_2 resonance) yields:

$$E_{\text{binding}}^{\text{FFGFT}} \approx (3.26 \text{ eV} \times 1.38) \times (1 - 0.009) \approx 4.48 \text{ eV} \times 0.991 \approx 4.44 \text{ eV}$$

Comparison: Experimental value ≈ 4.52 eV. The deviation is 0.08 eV (ca. 1.8%). **[S]** Since the resonance factor 1.38 is not derived independently in this document, the comparison is not yet a standalone prediction; it becomes one once this factor is calculated from the resonance condition.

D) Resonance Condition:

Two molecular tori couple maximally when their winding numbers are compatible ($w_1/w_2 = \text{rational number}$). For H_2 with two electrons (spin 1/2):

$$w_1 = w_2 = 1/2 \quad \rightarrow \quad w_1/w_2 = 1 \quad (\text{resonance condition met})$$

This could explain the special stability of the H_2 bond compared to other possible dimer configurations. The additional factor 1.38 in the above calculation is attributed to the resonance condition.

Extension: Adjustment of Correction Based on Hierarchy Accumulation

An extended correction incorporating an accumulated hierarchy ($1 - 100\xi \approx 0.9867$) leads to an adjusted binding energy of about 4.41 eV, so that the deviation from the experimental value stays below 2.5%. This addition takes the fractal iteration rule into account.

4. Critical ξ for Chaos Transition

Problem:

At which critical value ξ_{crit} does the fractal spacetime fabric become unstable and potentially collapse into a chaotic regime? Is there an upper limit for ξ in a stable universe?

Calculation from the Logistic Map:

From the FFGFT iteration rule for fractal scaling $\xi_{n+1} = \xi_n(1 - 100\xi_n)$, a critical threshold for stability is derived. The change of ξ per iteration step is:

$$\left| \frac{d\xi}{dn} \right| = 100\xi^2$$

Instability occurs when this rate of change becomes greater than about 10% of ξ itself (an arbitrary but physically plausible threshold for the transition to nonlinear instability):

$$\begin{aligned} 100\xi^2 &> 0.1\xi \\ \xi &> 0.001 = 10^{-3} \end{aligned}$$

Thus, the critical value is:

$$\xi_{\text{crit}} \approx 10^{-3}$$

The physical interpretation of these different regimes:

- For $\xi > 10^{-3}$: System collapses too quickly, no stable structures can form over cosmological time scales.
- For $\xi < 10^{-4}$: System is ultra-stable, with extremely long-lived structures spanning many orders of magnitude.
- For $10^{-4} < \xi < 10^{-3}$ (our reality: 1.33×10^{-4}): Metastable phase possible, potentially with interesting transition phenomena and intermittent chaos. *(Corrected on 30 Sept. 2026: previously "our reality: 1.33×10^{-4} " was assigned to the range $\xi < 10^{-4} - 1.33 \times 10^{-4} > 10^{-4}$, so by this classification our ξ lies in the range $10^{-4} < \xi < 10^{-3}$.)*

This refines the earlier rough estimate of $\xi_{\text{crit}} \approx 0.005$ and is consistent with our universe, with $\xi = 1.333 \times 10^{-4}$, lying in the stable, but not too rigid, region. *(Note of 30 Sept. 2026: of the values given for ξ_{crit} in this document (10^{-3} , 0.005, 0.01, 1), only the divergence threshold 0.01 of the map $\xi_{n+1} = \xi_n(1 - 100\xi_n)$ holds; 10^{-3} rests on the arbitrary 10 % threshold.)*

Extension: Correction of the Critical Limit

Upon closer analysis of the logistic map $\xi_{n+1} = \xi_n(1 - 100\xi_n)$, the divergence threshold is at $\xi = 1/100 = 0.01$ (the only fixed point is $\xi^* = 0$). *(Corrected on 30 Sept. 2026: previously "the fixed point is at $\xi^* = 1/100$ " - 0.01 is not a fixed point of the map.)* The stability limit, where $|1 - 200\xi| < 1$ holds, lies at $\xi < 0.01$. This corrects the original estimate from 10^{-3} to 10^{-2} , which allows model stability over a broader range. The discrepancy arose from an approximate threshold; the exact fixed-point analysis resolves it.

5. Temperature Dependence of ξ

Problem:

Is the fundamental scale factor ξ an absolute constant or temperature-dependent? How does a possible temperature dependence influence experimental predictions, particularly for the BZ reaction at low temperatures?

Calculation of Temperature Dependence:

From the BZ period formula $T_{BZ} \propto T_{\text{Compton}} \times N_A / \sqrt{1 - \xi(T)}$ and the empirically well-established classical Arrhenius behavior ($T_{BZ} \propto 1/\sqrt{T}$ for chemical reactions), equating leads to:

$$\xi(T) \propto 1 - \frac{2}{\sqrt{T}}$$

For a reference temperature of $T_{\text{ref}} = 300$ K with $\xi(300) = \xi_0 = 1.333 \times 10^{-4}$, at low temperatures, e.g., $T = 10$ K:

$$\begin{aligned} \xi(10 \text{ K}) &= \xi_0 \times \left[1 - 2 \left(\frac{1}{\sqrt{10}} - \frac{1}{\sqrt{300}} \right) \right] \\ &\approx \xi_0 \times (1 - 0.516) \approx 0.48 \times \xi_0 \end{aligned}$$

(Note of 30 Sept. 2026: equating $T_{BZ} \propto 1/\sqrt{1 - \xi}$ and $T_{BZ} \propto 1/\sqrt{T}$ gives $1 - \xi \propto T$, not $\xi \propto 1 - 2/\sqrt{T}$; the evaluation above is moreover a linearized difference – computed as a proportionality it would give $(1 - 2/\sqrt{10})/(1 - 2/\sqrt{300}) = 0.42$ instead of 0.48.)

Prediction [S]: At low temperatures (~ 10 K), **ξ approximately halves**. In this approach, this follows from the assumed coupling between thermal excitation and fractal spacetime geometry.

Experimental Consequence for the BZ Reaction:

The BZ period should shorten upon cooling from room temperature initially according to the classical Arrhenius law (higher reaction rate at lower temperature would be unusual, so the precise form of the dependence needs checking here; alternatively: $T_{BZ} \propto \exp(E_a/kT)$ with positive E_a). However, at very low temperatures ($T < 10$ K), it should **saturate** and not shorten further, as $\xi(T)$ approaches a constant value:

$$T_{BZ}(1 \text{ K}) \approx T_{BZ}(10 \text{ K}) \quad (\text{no further significant shortening})$$

This would be a signal distinguishable from classical reaction kinetics: While classical theory would predict a steady lengthening of the period with decreasing temperature (until the reaction freezes), FFGFT predicts saturation at low temperatures. This effect is testable in a cryogenic experiment with precise temperature control and period measurement.

Extension: Alternative Form of Temperature Dependence and Divergence Avoidance

The original form $\xi(T) \propto 1 - 2/\sqrt{T}$ can become negative at low T , which is physically nonsensical. An improved form, derived from thermal vacuum excitation, is $\xi(T) = \xi_0 / \sqrt{T_{\text{ref}}/T}$. For $T=10$ K, this gives $\xi \approx 0.18\xi_0$, representing a reduction without divergence and fitting better to BZ saturation. This correction avoids the negative range; the predicted value at 10 K thereby changes from about $0.48\xi_0$ to about $0.18\xi_0$.

6. Cosmic Time-Density Variations in the CMB

[S] This section and Appendix A deal with the cosmic sector. The present corpus deliberately leaves it aside, because the Standard Model does not derive it either (P39); the cosmological statements here are therefore speculative.

Problem:

Do the cosmic microwave background (CMB) and other observations show signatures of time-density variations? Can the observed CMB dipole be modified by fractal geometry effects, and how does this relate to the alternative interpretation of the T_0 theory?

Clarification and Conflict with the T_0 Core Thesis

Within the framework of Fractal Field Geometrodynamics (FFGFT), the observed CMB dipole is interpreted primarily as a kinematic effect – a result of the solar system's motion relative to the CMB rest frame. The scale-invariant parameter ξ modifies this effect through fractal amplification over cosmological distances.

However, this interpretation contradicts the core thesis of the T_0 theory, as formulated in the accompanying document '039_Zwei-Dipole-CMB_En.pdf'. There, the CMB dipole is explicitly **not** interpreted as a Doppler shift due to motion, but as an intrinsic, static anisotropy of the fundamental ξ -field in a non-expanding universe:

> **"The CMB dipole is NOT motion, but an intrinsic anisotropy of the ξ -field. The ξ -field is the fundamental vacuum field from which the CMB emerges as equilibrium radiation."**

The "fractal amplification" of the kinematic dipole calculated here in the main document retains the paradigm of an expanding universe, where ξ is a scaling constant. The T_0 interpretation discards this paradigm in favor of a static, cyclic universe. Both approaches cannot be true simultaneously; this is a conceptual break within the theoretical framework.

Calculation of Fractal Amplification (FFGFT Approach)

Starting from the above premise, which contradicts the T_0 core thesis of a kinematic dipole, the observed dipole can be modified by a cumulative effect of fractal spacetime geometry over the Hubble distance:

$$\Delta T_{\text{obs}} = \Delta T_{\text{intrinsic}} \times \left[1 + \xi \ln \left(\frac{R_{\text{Hubble}}}{\Lambda_0} \right) \right]$$

With standard values:

- Hubble radius: $R_{\text{Hubble}} = c/H_0^{T0} \approx 1.39 \times 10^{26} \text{ m}$ (with $H_0 \approx 66.2 \text{ km/s/Mpc}$, see Document 026)
- Fundamental length: $\Lambda_0 \approx 2.15 \times 10^{-39} \text{ m}$
- Scale parameter: $\xi = 1.333 \times 10^{-4}$
the logarithmic scale factor is:

$$\ln \left(\frac{R_{\text{Hubble}}}{\Lambda_0} \right) \approx \ln(6.5 \times 10^{64}) \approx 149.2$$

and thus the total amplification:

$$\Delta T_{\text{obs}} \approx \Delta T_{\text{intrinsic}} \times (1 + 1.333 \times 10^{-4} \times 149.2) \approx \Delta T_{\text{intrinsic}} \times 1.0199$$

(Corrected on 30 Sept. 2026: previously $\ln(1.07 \times 10^{65}) \approx 149.8$ and $1.0198 - 1.39 \times 10^{26} / 2.15 \times 10^{-39} = 6.5 \times 10^{64}$, $\ln = 149.2$, amplification $1 + \xi \cdot 149.2 = 1.0199$.)

The model thus predicts an **amplification of the geometric (kinematic) dipole component by nearly 2%**. This small but measurable effect lies within the order of systematic uncertainties of high-precision CMB experiments like *Planck* and could theoretically contribute to solving anomalies.

The Empirical Problem: The Dipole Anomaly

The motivation for these considerations is a serious tension in the standard model of cosmology (Λ CDM): While the CMB dipole suggests a velocity of about 370 km/s towards the constellation Leo, dipole measurements in the distribution of quasars and radio galaxies (e.g., in the CatWISE and NVSS catalogs) show both differing directions and a significantly larger amplitude, about twice the kinematic expectation from the CMB dipole. This discrepancy is termed the "Cosmic Dipole Anomaly" and calls into question the cosmological principle of homogeneity and isotropy – a cornerstone of the Λ CDM model.

Extension: Deeper Integration of the T0 Interpretation

To resolve the conflict, the T0 theory is more fully integrated: The CMB dipole as an intrinsic ξ -anisotropy eliminates the need for kinematic amplification. Instead, a wavelength-dependent redshift emerges, explaining the dipole amplitude discrepancy (quasar dipole about twice the kinematic expectation) as a consequence of different field interactions, according to this proposal. This extends the model to a hybrid approach, where FFGFT applies on local scales and T0 on cosmological scales.

Appendix A: On the CMB Dipole Anomaly and the T₀ Solution

This appendix provides an in-depth discussion of the empirical tension mentioned in section 6 and the alternative explanation by the T₀ theory [S], as presented in the linked document.

A.1 The Empirical Tension in Detail

The CMB dipole is the dominant signal in the cosmic microwave background – about 100 times stronger than the primary anisotropies (quadrupole and higher multipoles). In the Λ CDM standard model, it is fully interpreted as a kinematic Doppler and aberration effect, indicating the motion of the solar system at about 370 km/s relative to the CMB rest frame. A fundamental postulate of the cosmological principle is that this rest frame is the same for radiation and matter.

The so-called "Ellis-Baldwin test" offers a critical check of this postulate: The same peculiar velocity causing the CMB dipole should produce a predictable, characteristic dipole in the sky distribution of very distant extragalactic sources (like quasars or radio galaxies). This matter dipole should match the CMB dipole in amplitude and direction.

Current measurements using large, statistically robust catalogs, however, find significant and growing deviations:

- **CatWISE dipole** (1.3 million quasars in the infrared): Points towards $(l, b) \approx (238^\circ, 29^\circ)$ (Secrest et al. 2021) with an amplitude about twice the kinematic expectation derived from the CMB dipole.
- **NVSS dipole** (radio galaxies): Shows a similarly large amplitude and also deviates in direction.
- **CMB dipole** (Planck satellite): Points towards **Leo** (galactic coordinates: $l \approx 264^\circ, b \approx +48^\circ$), corresponding to ~ 370 km/s.
- **Angular deviation**: The directions of the CMB dipole and the quasar dipole are about **28°** apart.

This discrepancy is now established at a significance level of **over 5σ** (see review by Sarkar et al., 2025) and constitutes a serious challenge to the cosmological principle and the Λ CDM model. More recent Bayesian analyses confirm the strong tension between datasets and largely rule out systematic errors as the sole cause.

A.2 The T_0 Approach: A Reinterpretation of the Dipole

The T_0 theory, as laid out in the document '039_Zwei-Dipole-CMB_En.pdf', proposes a reinterpretation that addresses the cause of the tension:

1. **The CMB Dipole is Not Motion:** The T_0 theory discards the kinematic interpretation. Instead, the CMB dipole is an **intrinsic, static anisotropy** of the fundamental ξ vacuum field ($\xi = \frac{4}{3} \times 10^{-4}$). The CMB temperature itself arises in this model directly from this field: $T_{\text{CMB}} = \frac{16}{9} \xi^2 \times E_\xi \approx 2.725 \text{ K}$, where E_ξ is a characteristic field energy. The dipole arises from a slight spatial variation of the ξ -field itself.
2. **Resolving the Contradiction:** If the CMB dipole is not an indicator of motion, the fundamental requirement that matter distributions must show the same dipole vanishes. The dipole measured in the quasar catalog can then either reflect a true (much larger) peculiar velocity of our Local Group or itself be a structural asymmetry in the large-scale matter distribution of the universe. The observed angular offset of about 28° between the dipoles might indicate a fundamental geometric or dynamic relationship between the ξ -field (determining radiation) and baryonic matter distribution.
3. **Consequence: A Static, Cyclic Universe:** This approach is not isolated but embedded in a larger model of a **static, cyclic universe without Big Bang expansion**. Cosmological redshift is interpreted in this model not as a Doppler effect of expansion but as a wavelength-dependent energy loss of photons during their long travel time through interaction with the ξ -field. This also offers an alternative interpretation of the "Hubble tension", the discrepancy between locally and cosmologically measured values of the Hubble constant.

A.3 Comparison of the Incompatible Explanatory Approaches

The following list summarizes the conceptual differences between the FFGFT approach taken in the main document and the T_0 interpretation. These approaches are incompatible in their basic assumptions:

- **Aspect: Nature of the CMB Dipole** - *FFGFT Approach (Main Document):* Predominantly **kinematic** (motion), fractally modified. - * T_0 Interpretation (Document 039):* **Intrinsic anisotropy** of the ξ -field, **non-kinematic**.
- **Aspect: Foundational Paradigm** - *FFGFT Approach:* Expanding universe (Big Bang, Λ CDM), ξ as a scale-invariant parameter within this framework. - * T_0 Interpretation:* **Static, cyclic universe** without expansion and without a singular beginning.
- **Aspect: Solution Strategy for the Dipole Anomaly** - *FFGFT Approach:* Small **modification** ($\approx 2\%$ amplification) of the expected kinematic signal within the standard paradigm. - * T_0 Interpretation:* **Fundamentally different interpretation**: Separation of the physical causes for radiation and matter dipoles.
- **Aspect: Predictive Statement** - *FFGFT Approach:* Slight amplification of the CMB dipole compared to the purely kinematic expectation. - * T_0 Interpretation:* **No** necessary coincidence between CMB and quasar dipoles; instead, prediction of wavelength-dependent redshifts.
- **Aspect: Consistency and Explanatory Power** - *FFGFT Approach:* Internally (mathematically) coherent, but in direct contradiction to the T_0 core thesis and does not fully explain the large anomaly amplitude. - * T_0 Interpretation:* Offers a principled proposal for solving the dipole anomaly but requires complete abandonment of the standard expansion paradigm of cosmology.

The Core Idea

The question of whether the universe is open and closed at the same time – like an open and closed resonator – touches a core point of the T_0 theory. The metaphor of the **“open and closed resonator simultaneously”** captures well how the universe functions in T_0 .

1. The Universe is Open and Closed Simultaneously

- **Open** – because the T/E-field is continuous, scale-invariant, and without a hard boundary. There is no fundamental isolation, no intrinsic discretization, and no “wall” at the Planck scale or elsewhere. The field can extend and couple fractally – ξ is scale-invariant, the duality $T \cdot E = 1$ holds over all scales.
→ Like an open pipe: Resonances can escape, propagate, excite new modes, generate diversity. No total isolation.
- **Closed** – because the minimal feedback via ξ enforces closed geometric loops. Only configurations where $\xi \cdot T \approx \text{integer/half-integer/fraction thereof}$ are stably amplified. Everything else diffuses away, becomes incoherent.
→ Like a closed pipe: Only certain wavelengths (modes) fit inside and remain stable – others interfere destructively. There are preferred, quasi-discrete states.

2. The Universe is an Open Resonator with Closed Modes

- **Open resonator** – the field as a whole is open, continuous, allows fractal propagation and coupling over all scales.
- **Closed modes** – within this open system, closed, stable resonance conditions arise through ξ -feedback (just as in a closed pipe only quarter-, half-, and full-integer wavelengths are stable).

This is exactly what happens in T_0 : The field is open (no fundamental isolation), but ξ enforces closed loops → only specific geometric ratios (resonance modes) couple coherently and become stable. Result: The universe appears quasi-discrete and quantized (preferred energy levels, spin ratios, stable scales), but leaves freedom (variations, clusters, irregularities) because ξ is minimal and continuous.

Clarification: no infinities

- The fractal dimension $D_f = 3 - \xi$ with $\xi = \frac{4}{3} \times 10^{-4}$ prevents **“true infinities”**.
- What classically appears as “infinite propagation” or “continuous spectrum” is always fractally bounded by $D_f < 3$ in FFGFT.
- The “open field” does not mean mathematically infinite, but **“no fundamental isolation”** – the field can extend fractally, but always within the fractal metric.

Computable Consequences: Connection to Belousov-Zhabotinsky, Mandelbrot, and Turing

1. Belousov-Zhabotinsky Reaction → FFGFT Torus Oscillation

BZ Reaction (classical):

Period: $T_{BZ} \approx 1 - 2$ minutes

Mechanism: Autocatalysis + Inhibition

$\text{Ce}^{3+} \leftrightarrow \text{Ce}^{4+}$ (color change)

FFGFT Equivalent:

The torus oscillation on different scales.

Computable:

A) Compton Time of the Proton as "BZ Period":

$$T_p = \frac{h}{m_p c^2} \approx 4.4 \times 10^{-24} \text{ s}$$

This is the "oscillation period" of the proton torus between two states:

- Ce^{3+} analog: low energy density (poloidal flow dominates)
- Ce^{4+} analog: high energy density (toroidal flow dominates)

B) Ratio to BZ Reaction:

$$\frac{T_{BZ}}{T_p} \approx \frac{100 \text{ s}}{4.4 \times 10^{-24} \text{ s}} \approx 2.3 \times 10^{25}$$

This is about 38 times Avogadro's number; this factor is discussed in the extension below.

Prediction [S]: Chemical oscillations (BZ) are ****collective torus resonances**** over $\sim 10^{25}$ particles. The period results from:

$$T_{BZ} = T_{\text{Compton}} \times N_A \times (\text{geometric factor})$$

Deepening on BZ Reaction and Scale Transition: The prediction $T_{BZ} \propto T_{\text{Compton}} \times N_{\text{Avogadro}}$ would, if correct, mean that the macroscopic period is a resonance phenomenon where microscopic torus oscillators synchronize via the fractality of space.

Concrete Test Suggestion: Investigate BZ-like reactions in mesoscopic systems (nano- to microdroplets) with particle numbers $N \ll N_A$. FFGFT predicts a discontinuous change in oscillation dynamics once N falls below a critical value depending on the fractal coherence length. Classical reaction kinetics would expect a continuous change.

C) Spiral Patterns in BZ → Torus Winding:

The characteristic spiral wavelength in BZ:

$$\lambda_{\text{spiral}} \approx 1 \text{ mm}$$

FFGFT prediction (with $R/r \approx 10$ for molecular tori):

$$\begin{aligned} \lambda_{\text{spiral}} &\approx R_{\text{molecular}} \times \sqrt{N_{\text{particle}}} \\ &\approx 10^{-9} \text{ m} \times \sqrt{10^{18}} \approx 10^{-3} \text{ m} \approx 1 \text{ mm} \quad (\text{agrees}) \end{aligned}$$

Experimentally testable: The spiral velocity should scale as:

$$v_{\text{spiral}} \propto \sqrt{\xi \times D_{\text{diffusion}}}$$

Extension: Resolution of the Period Discrepancy

The calculated ratio $T_{BZ}/T_p \approx 2.27 \times 10^{25}$ vs. $N_A = 6.022 \times 10^{23}$ gives a factor of ≈ 37.74 . This factor is interpreted as a geometric correction term arising from the effective volume of the BZ reaction mixture (e.g., 0.1 mol in typical volume) and torus coupling efficiency. The extended formula $T_{BZ} = T_{\text{Compton}} \times N_{\text{eff}}$ with $N_{\text{eff}} \approx 38N_A$ absorbs the discrepancy; the factor 38 has not yet been derived independently.

2. Mandelbrot Set → FFGFT Fractal Scaling

Mandelbrot Set (classical):

$$z_{n+1} = z_n^2 + c$$

Boundary between bounded/unbounded

Fractal dimension $D \approx 2$

FFGFT Equivalent:

The recursive scaling via ξ .

Computable:

A) FFGFT Iteration Rule:

Instead of $z \rightarrow z^2 + c$ we have:

$$D_{n+1} = 3 - \xi_n$$

$$\xi_{n+1} = \xi_n \times K_{\text{frak}} = \xi_n \times (1 - 100\xi_n)$$

This is a ****logistic map****.

B) Bifurcation Diagram:

The logistic equation $x_{n+1} = rx_n(1 - x_n)$ shows chaos for $r > 3.57$.

For $K_{\text{frak}} = 1 - 100\xi$:

$$\xi_{n+1} = \xi_n - 100\xi_n^2$$

With $\xi_0 = \frac{4}{3} \times 10^{-4}$:

$$\xi_1 = 1.333 \times 10^{-4} - 100 \times (1.333 \times 10^{-4})^2$$

$$\approx 1.333 \times 10^{-4} - 1.78 \times 10^{-6}$$

$$\approx 1.315 \times 10^{-4}$$

The iteration ****converges**** to a fixed point (no chaos).

Fixed Point:

$$\xi^* = \xi - 100\xi^2$$

$$100\xi^2 = 0$$

→ $\xi^* = 0$ (only fixed point; $\xi = 1/100 = 0.01$ is the divergence threshold)

(Corrected on 30 Sept. 2026: previously " $\xi^ = 0$ (trivial) or $\xi^* = 1/100$ " – $100\xi^2 = 0$ gives only $\xi^* = 0$; above 0.01, ξ_{n+1} becomes negative and the iteration diverges.)*

But: With K_{frak} -modification:

$$\xi^* = \frac{1 - \sqrt{1 - 4/100}}{200} \approx 1.01 \times 10^{-4}$$

(Corrected on 30 Sept. 2026: previously $\approx 4.99 \times 10^{-3} - (1 - \sqrt{0.96})/200 = 1.01 \times 10^{-4}$.)

Prediction: There is a ****critical scale**** at $\xi_{\text{crit}} \approx 0.01$ (divergence threshold of the map), above which the fractal structure becomes unstable. (Corrected on 30 Sept. 2026: previously $\xi_{\text{crit}} \approx 0.005$ – this value rested on the calculation error above; the divergence threshold 0.01 holds.)

Interpretation of the Mandelbrot Set: The link to the logistic map is the core point here. The FFGFT iteration rule for ξ runs towards the fixed point $\xi^* \approx 0$, which fits the observed stability of matter and scales over cosmic time.

Speculative interpretation [S]: The Mandelbrot set might be more than a model for fractality, namely the mathematical projection of the attractor dynamics of the fractal vacuum itself. The Mandelbrot boundary would then mark the transition between stably bound (bounded) and unstable, freely releasing (unbounded) energy states in $T \cdot E$ space.

C) Mandelbrot Boundary in FFGFT:

The "boundary" of the Mandelbrot set corresponds to the transition:

$$|z_n| < 2 \text{ (bounded) vs. } |z_n| \rightarrow \infty \text{ (unbounded)}$$

In FFGFT:

$$D_f > 2 \text{ (3D-like) vs. } D_f < 2 \text{ (collapsed)}$$

The critical dimension:

$$D_{\text{crit}} = 2 \rightarrow \xi_{\text{crit}} = 1$$

(Note of 30 Sept. 2026: here $\xi = 1$ is the limit $D_f = 3 - \xi = 2$, not a stability threshold of the iteration; for the latter only 0.01 holds.)

But our reality has $\xi = 1.333 \times 10^{-4} \ll 1$, thus ****far in the stable region****.

D) Calculating Self-Similarity:

The Mandelbrot set shows self-similarity with scaling factor $\sim 2 - 3$.
FFGFT scaling between levels:

$$\text{Scaling factor} = 1/\xi \approx 7500$$

Much larger. This could explain why the universe shows self-similar structures over ~ 60 orders of magnitude (Planck $\rightarrow$ Cosmos).

Clarification: no "infinite zoom" – The fractal zoom ends at the sub-Planck scale $\Lambda_0 \approx 2.15 \times 10^{-39}$ m. The Mandelbrot-like behavior is fractally bounded.

3. Turing Patterns $\rightarrow$ FFGFT Structure Formation

Turing (classical):

$$\begin{aligned} \frac{\partial a}{\partial t} &= f(a, h) + D_a \nabla^2 a \\ \frac{\partial h}{\partial t} &= g(a, h) + D_h \nabla^2 h \\ &\text{with } D_h > D_a \text{ (Inhibitor diffuses faster)} \end{aligned}$$

FFGFT Equivalent:**A) Field Equations Instead of Reaction-Diffusion:**

In FFGFT we have no separate "morphogens", but:

$$\text{Activator} = E(x, t) \quad (\text{energy density})$$

$$\text{Inhibitor} = T(x, t) \quad (\text{time density})$$

$$\text{with } T \cdot E = 1 \quad (\text{duality})$$

The "diffusion" is the fractal propagation:

$$\frac{\partial E}{\partial t} = -\nabla \cdot (c^2 \nabla T) + \xi \times (\text{nonlinear terms})$$

$$\frac{\partial T}{\partial t} = -\nabla \cdot (\nabla E / c^2) + \xi \times (\dots)$$

B) Effective Diffusion Constants:

From the time-mass duality:

$$D_E \propto c^2 \quad (\text{energy diffuses "fast"})$$

$$D_T \propto \hbar / m \quad (\text{time diffuses "slow"})$$

Ratio:

$$\frac{D_E}{D_T} \propto \frac{mc^2}{\hbar} = \frac{1}{T_{\text{Compton}}}$$

For a proton:

$$\frac{D_E}{D_T} \approx \frac{1}{7.0 \times 10^{-25} \text{ s}} \approx 1.43 \times 10^{24}$$

(Corrected on 30 Sept. 2026: previously $1/(4.4 \times 10^{-24} \text{ s}) \approx 2.3 \times 10^{23} - 4.4 \times 10^{-24} \text{ s}$ is $\hbar/(m_p c^2)$; the formula with $\hbar$ requires $T_{\text{Compton}} = \hbar/(m_p c^2) = 7.0 \times 10^{-25} \text{ s}$.)

Large difference. However, Turing's condition $D_h \gg D_a$ is not satisfied by this: here the activator E diffuses faster than the inhibitor T , and $mc^2/\hbar$ has the unit s^{-1} . (Corrected on 30 Sept. 2026: previously "This is meant to satisfy Turing's condition $D_h \gg D_a$." – the assignment activator = E , inhibitor = T with $D_E \gg D_T$ is the reverse of the Turing condition.)

C) Pattern Wavelength:

Turing wavelength:

$$\lambda_{\text{Turing}} \approx 2\pi \sqrt{D_a D_h} / \sqrt{\text{reaction rate}}$$

FFGFT equivalent:

$$\begin{aligned} \lambda_{\text{FFGF}} &\approx 2\pi \sqrt{c^2 \times \hbar / m} / \sqrt{\omega_{\text{Compton}}} \\ &\approx \lambda_{\text{Compton}} \times \text{constant factors} \end{aligned}$$

For electrons (biological systems):

$$\lambda_{\text{Compton}} \approx 2.4 \times 10^{-12} \text{ m}$$

$$\lambda_{\text{FFGF}} \approx 10^{-9} \text{ m} = 1 \text{ nm}$$

(Note of 30 Sept. 2026: with $\omega_{\text{Compton}} = mc^2/\hbar$, $2\pi \sqrt{c^2 \hbar / m} / \sqrt{\omega_{\text{Compton}}} = 2\pi \hbar / m (= 7.3 \times 10^{-4} \text{ m}^2/\text{s}$ for electrons), which is not a length; moreover 1 nm is ≈ 400 times $\lambda_{\text{Compton}} = 2.4 \times 10^{-12} \text{ m}$, contradicting " $\lambda_{\text{FFGF}} \approx \lambda_{\text{Compton}}$ ".)

That is the ****typical size of biological molecules****.

Turing Pattern Prediction Deepened: The derivation of the characteristic length $\lambda_{\text{FFGF}} \approx \lambda_{\text{Compton}}$ offers an approach to justifying the fundamental length scale of biological building blocks from first principles.

Extended Testability [S]: This leads to the conjecture that the lattice constants of molecular assemblies (cell membrane lipid bilayers, actin/tubulin spacing, chromatin fiber diameter) should all appear as integer multiples of this basic wavelength ($\lambda_{\text{FFGF}} \sim 1 \text{ nm}$), modulated by the local ξ_{eff} of the tissue.

D) Calculating Zebra Stripes:

Turing said: Stripes arise when $\lambda_{\text{Turing}} \approx$ characteristic length.

For a zebra embryo ($\sim 10 \text{ cm}$ diameter):

$$\text{Number of stripes} \approx (10 \text{ cm})/\lambda_{\text{FFGF}}$$

If λ_{FFGF} is determined by cellular scale:

$$\lambda_{\text{FFGF}} \approx 100 \text{ cells} \times 10 \mu\text{m} \approx 1 \text{ mm}$$

$$\text{Number of stripes} \approx 100 \text{ mm}/1 \text{ mm} = 100$$

The order of magnitude agrees: Zebras have $\sim 40 - 80$ stripes.

Bibliography

Bibliography

- [1] Mandelbrot, Benoit B. (1977). *The Fractal Geometry of Nature*. W.H. Freeman and Company, New York.
- [2] Falconer, Kenneth (2003). *Fractal Geometry: Mathematical Foundations and Applications* (2nd ed.). John Wiley & Sons.
- [3] Russ, John C. (1994). *Fractal Surfaces*. Plenum Press, New York.
- [4] Belousov, B. P. (1959). A periodic reaction and its mechanism. *Collection of Abstracts on Radiation Medicine*, **147**, 1.
- [5] Zhabotinsky, A. M. (1964). Periodic processes of malonic acid oxidation in a liquid phase. *Biofizika*, **9**, 306–311.
- [6] Epstein, I. R., & Pojman, J. A. (1998). *An Introduction to Nonlinear Chemical Dynamics: Oscillations, Waves, Patterns, and Chaos*. Oxford University Press.
- [7] Turing, Alan M. (1952). The Chemical Basis of Morphogenesis. *Philosophical Transactions of the Royal Society B*, **237**(641), 37–72.
- [8] Kondo, S., & Miura, T. (2010). Reaction-Diffusion Model as a Framework for Understanding Biological Pattern Formation. *Science*, **329**(5999), 1616–1620.
- [9] Meinhardt, H. (1982). *Models of Biological Pattern Formation*. Academic Press, London.
- [10] Compton, Arthur H. (1923). A Quantum Theory of the Scattering of X-Rays by Light Elements. *Physical Review*, **21**(5), 483–502.
- [11] Planck, Max (1901). On the Law of Distribution of Energy in the Normal Spectrum. *Annalen der Physik*, **4**, 553–563.
- [12] Planck Collaboration (2020). Planck 2018 results. VI. Cosmological parameters. *Astronomy & Astrophysics*, **641**, A6. <https://arxiv.org/abs/1807.06209>
- [13] Peebles, P. J. E. (1993). *Principles of Physical Cosmology*. Princeton University Press.
- [14] Nicolis, G., & Prigogine, I. (1977). *Self-Organization in Nonequilibrium Systems: From Dissipative Structures to Order through Fluctuations*. Wiley, New York.
- [15] Haken, H. (1983). *Synergetics: An Introduction* (3rd ed.). Springer-Verlag, Berlin.
- [16] Pauling, Linus (1960). *The Nature of the Chemical Bond* (3rd ed.). Cornell University Press.
- [17] Szabo, A., & Ostlund, N. S. (1996). *Modern Quantum Chemistry: Introduction to Advanced Electronic Structure Theory*. Dover Publications.

- [18] May, Robert M. (1976). Simple mathematical models with very complicated dynamics. *Nature*, **261**(5560), 459–467.
- [19] Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. (2007). *Numerical Recipes: The Art of Scientific Computing* (3rd ed.). Cambridge University Press.
- [20] Pascher, J. (2024). *Comment: CMB and Quasar Dipole Anomaly – A Dramatic Confirmation of T_0 Predictions!* (Document '039_Zwei-Dipole-CMB_En.pdf'). [PDF on GitHub]. *Contains the central thesis, diverging from the FFGFT approach, of a non-kinematic, intrinsic CMB dipole in a static T_0 universe.*
- [21] Sarkar, S., Secrest, N., et al. (2025). *Colloquium: The Cosmic Dipole Anomaly*. arXiv:2505.23526. <https://arxiv.org/abs/2505.23526>. *Current, comprehensive review outlining the empirical crisis of the cosmological principle due to the dipole anomaly at over 5σ level.*
- [22] Wikipedia contributors. (2024). *Cosmic microwave background*. In Wikipedia, The Free Encyclopedia. https://en.wikipedia.org/wiki/Cosmic_microwave_background. *Basic article on CMB, its discovery, and the standard interpretation of the dipole as a kinematic effect.*
- [23] Wen, Y. et al. (2021). *The role of T_0 in CMB anisotropy measurements*. Physical Review D, 104, 043516. <https://arxiv.org/abs/2011.09616>. *Discusses the calibrating role of the CMB monopole T_0 , which represents a central dual parameter in the T_0 theory.*
- [24] White, M., et al. (1994). *Anisotropies in the CMB*. Annual Review of Astronomy and Astrophysics, 32, 319. <https://ned.ipac.caltech.edu/level5/March02/White/White1.html>. *Shows the historical development of the interpretation of the CMB dipole and other anisotropies.*
- [25] Secrest, N. J., et al. (2021). *A Test of the Cosmological Principle with Quasars*. The Astrophysical Journal Letters, 908(2), L51. <https://iopscience.iop.org/article/10.3847/2041-8213/abdd40>. *Important original work that first robustly demonstrated the significant deviation of the quasar dipole from the CMB dipole.*
- [26] Anonymous (2024). *T_0 Framework: Fractal Field Geometry Theory*. Internal documentation.
- [27] Anonymous (2024). *Fractal Field Geometry Theory: Complete Derivation*. In: 145_FFGFT_donat-part1_En.pdf