

Correction notes

Corrections from the calculation check (30 Sept. 2026). The following places are corrected or annotated in the text; each carries a dated note with the previous value:

- Abstract (iii) and the section on the ladder prediction: $Q_{\text{FFGFT}} = 0.6677$ lies 0.16 % "below" $\rightarrow$ "above" $2/3$.

The Recurrence of $2/3$:

**Koide Projectors, Complement Step,
FFGFT Ladder and $D = 3 + \varepsilon$ Deformation**

Johann Pascher

In dialogue with P. Austin, *A Purely Algebraic Short Note on the Repeated Appearance of $2/3$* (May 2026)

May 2026

Abstract

At exactly $D = 3$, three distinct mathematical structures yield the value $2/3$: the Koide equal-projector condition $A = B$ between democratic and orthogonal projection components, the complement step $1 - 1/D$, and the geometric FFGFT lepton ladder with ratio $q = 2/3$. A $D = 3 + \varepsilon$ deformation separates the three roads. This document shows: (i) the FFGFT ladder is geometric, not arithmetic; (ii) the exponent step is tied to the integer topological $D = 3$, not to the fractal dimension D_f ; (iii) the FFGFT ladder predicts $Q \approx 0.6677$ (0.16 % above $2/3$) without Koide input (Corrected on 30 Sept. 2026: previously "below" – the ladder with $r_i = (4/3, 16/5, 25/9)$, $p_i = (3/2, 1, 2/3)$, $\xi = 4/30000$ gives $Q = 0.667742$, i.e. +0.161 % above $2/3$, consistent with $\Delta_K > 0$.) – the exact experimental value $Q = 2/3$ confirms the exponent structure from an independent direction.

Contents

Notation	3
.1 Starting Point: Three Roads to $2/3$	4
.2 The Koide Relation as a Projector Statement	4
.3 The FFGFT Ladder is Geometric, not Arithmetic	4
.4 Topological $D = 3$ versus the Fractal Dimension	5
.5 Forced or Merely Fitted? An Honest Answer	5
.6 The Explicit Map FFGFT $\rightarrow$ Koide	6
.7 Summary	7

Notation

Symbol	Meaning
$\xi = 4/30000$	Fundamental FFGFT parameter [7]
$v = 246.22 \text{ GeV}$	Higgs vacuum expectation value
$m_i = r_i \xi^{p_i} v$	FFGFT mass formula [4]
r_i	Rational prefactors of the leptons
p_i	Rational exponents of the leptons [5]
$D = 3$	Topological (integer) spatial dimension of T^4
$D_f^{\text{space}} = 3 - \xi$	Fractal spatial dimension (local metric) [7]
$D_f^{\text{eff}} \approx 2.973$	Effective fractal dimension (scale flow)
K_{frak}	Fractal correction $= (D_f^{\text{eff}}/3)^{D_f^{\text{eff}}/2}$
$x_i = \sqrt{m_i}$	Square-root lepton mass
$u = (1, 1, 1)^T$	Democratic unit vector
$P = uu^T/3$	Democratic projector
$P^\perp = I - P$	Residual projector
$A = x^T P x$	Democratic quadratic component
$B = x^T P^\perp x$	Residual quadratic component
Q	Koide quantity $= (A + B)/(3A)$ [3]
$\Delta_K = 3Q - 2$	Koide residual diagnostic
ξ_{Koide}	Value of ξ for exact $Q = 2/3$
α	Fine-structure constant

.1 Starting Point: Three Roads to $2/3$

At exactly $D = 3$, several independent principles produce the same numerical value:

$$1 - \frac{1}{D} = \frac{2}{3}, \quad \frac{2}{D} = \frac{2}{3}, \quad \frac{1}{2} \left(1 + \frac{1}{D} \right) = \frac{2}{3}. \quad (.1)$$

The useful instrument for distinguishing them is the formal deformation $D = 3 + \varepsilon$: principles that coincide at $D = 3$ separate under this continuation.

The following table summarises the three continuations:

Principle	Formula	Expansion at $D = 3 + \varepsilon$
Koide equal-projector	$\frac{2}{D}$	$\frac{2}{3} - \frac{2\varepsilon}{9} + O(\varepsilon^2)$
Complement step	$1 - \frac{1}{D}$	$\frac{2}{3} + \frac{\varepsilon}{9} + O(\varepsilon^2)$
Interval midpoint	$\frac{1}{2} \left(1 + \frac{1}{D} \right)$	$\frac{2}{3} - \frac{\varepsilon}{18} + O(\varepsilon^2)$

The roads separate at first order – the complement step rises, the Koide equilibrium road and the interval midpoint fall.

.2 The Koide Relation as a Projector Statement

For the charged leptons define $x_i = \sqrt{m_i}$ and write the Koide quantity as

$$Q = \frac{\sum_i m_i}{(\sum_i \sqrt{m_i})^2} = \frac{x^T x}{(u^T x)^2}, \quad u = (1, 1, 1)^T. \quad (.2)$$

With the democratic projector $P = uu^T/3$ and residual projector $P^\perp = I - P$:

$$A = x^T P x = \frac{(u^T x)^2}{3}, \quad B = x^T P^\perp x, \quad Q = \frac{A + B}{3A}.$$

The Koide value $Q = 2/3$ is equivalent to the **equal-projector condition** $A = B$: the square-root mass vector splits into a democratic and a residual component of equal quadratic weight.

Under formal D -dimensional extension with $Q_D = 2/D$:

$$Q_{3+\varepsilon} = \frac{2}{3+\varepsilon} = \frac{2}{3} - \frac{2\varepsilon}{9} + O(\varepsilon^2) \quad (\text{descending}). \quad (.3)$$

.3 The FFGFT Ladder is Geometric, not Arithmetic

The canonical charged-lepton parameters in FFGFT [7, 4] are:

$$(r_e, r_\mu, r_\tau) = \left(\frac{4}{3}, \frac{16}{5}, \frac{25}{9} \right), \quad (p_e, p_\mu, p_\tau) = \left(\frac{3}{2}, 1, \frac{2}{3} \right), \quad (.4)$$

with $m_i = r_i \xi^{p_i} v$, $\xi = 4/30000$, $v = 246$ GeV (Higgs vacuum value).

The exponents $(\frac{3}{2}, 1, \frac{2}{3})$ do *not* form a uniform $1/3$ ladder – the gaps are $p_e - p_\mu = 1/2$ and $p_\mu - p_\tau = 1/3$. The resolution: the ladder is **geometric** with constant ratio $q = 2/3$:

$$p_{n+1} = \frac{2}{3} p_n, \quad \frac{3}{2} \xrightarrow{\times 2/3} 1 \xrightarrow{\times 2/3} \frac{2}{3}. \quad (.5)$$

The unequal differences $1/2$ and $1/3$ are *consequences* of a constant ratio, not independent rules.

Two equivalent readings.

- **Ratio reading.** $q = 2/3$ is the inverse of the tetrahedral geometry factor $3/2$. The same number that orders the ladder is, numerically, the Koide value.
- **Complement reading.** At the $\mu \rightarrow \tau$ rung $p_\tau = 1 - 1/D = 2/3$ with $D = 3$. This places FFGFT on the complement-step road – *not* on the equal-projector road $2/D$.

The electron at $p_e = 3/2 = 1 + 1/(D-1)$ is a half-integer rung – a geometric midpoint between winding levels (the spin-winding contribution [5]), not a full winding step. That is the structural origin of the $1/2$ gap.

.4 Topological $D = 3$ versus the Fractal Dimension

FFGFT carries two derived fractal dimensions:

$$D_f^{\text{space}} = 3 - \xi \approx 2.9998667 \quad (\text{local metric; GRB time-of-flight}), \quad (.6)$$

$$D_f^{\text{eff}} \approx 2.973$$

(cumulative scale flow, Planck to electroweak).

Both are derived, not assumed.

Why no $\xi/9$ correction reaches the Koide quantity. The Koide quantity Q is a pure mass *ratio*. Every scale-dependent factor – the vacuum value v , the fractal correction $K_{\text{frak}} = (D_f^{\text{eff}}/3)^{D_f^{\text{eff}}/2}$, and any D_f -level shift – is common to all three leptons and cancels in Q . Fractal corrections attach to *absolute* scales, never to ratios.

The exponent step is therefore tied to the integer, topological $D = 3$. The shift $2/3 - \xi/9$ from the D_f road is a real effect on absolute predictions; it simply does not enter Q .

.5 Forced or Merely Fitted? An Honest Answer

What is structurally forced

The mass *ratios* are not fitted. Two points:

- m_μ/m_e follows two independent roads that agree: geometrically from the ladder ($\frac{12}{5}\xi^{-1/2} \approx 207.8$) and fractally via K_{frak} (≈ 206.8). The agreement fixes K_{frak} uniquely – a consistency test, not a fit.
- In the compact representation the coefficients are not free rationals: they are constructed so that α cancels exactly. The rationals are α -geometric invariants.

What the ladder does not force

The ladder does not force the equal-projector condition $A = B$ exactly. Inserting the FFGFT masses into the Koide formula gives [6]:

$$Q_{\text{FFGFT}} \approx 0.6677, \quad \Delta_K = 3Q_{\text{FFGFT}} - 2 \approx +0.003 \quad (.7)$$

(0.16 % above the exact value $2/3$). (Corrected on 30 Sept. 2026: previously “below” – the ladder with $r_i = (4/3, 16/5, 25/9)$, $p_i = (3/2, 1, 2/3)$, $\xi = 4/30000$ gives $Q = 0.667742$, i.e. +0.161 % above $2/3$, consistent with $\Delta_K > 0$.)

The reason is structural: the Koide denominator $(\sum \sqrt{m_i})^2$ generates cross-terms with exponents

$$\xi^{5/4}, \quad \xi^{13/12}, \quad \xi^{5/6}, \quad (.8)$$

none of which is a multiple of $1/3$. They lie *outside* the FFGFT structure space $\{p\} \subset \frac{1}{3}\mathbb{Z}$. Exact $Q = 2/3$ would require $\xi_{\text{Koide}} \approx 1.372 \times 10^{-4}$, about 2.9 % larger.

The honest position: The ladder *predicts* Q to 0.16 % from (r_i, p_i, ξ) with no Koide input. The exact experimental $Q = 2/3$ then *confirms* the exponent structure from an independent direction – neither fitted into FFGFT nor algebraically forced by it. The small residual lies precisely where the cross-term analysis predicts.

.6 The Explicit Map FFGFT $\rightarrow$ Koide

The three structures – the Koide projector form, the FFGFT ξ -ladder, and Paul’s 24-cell rank split¹ – are three distinct primitive objects. A relation between them must be exhibited by a map, not by a shared numeral.

The FFGFT-to-Koide map is:

$$(r_i, p_i, \xi) \mapsto x_i = \sqrt{r_i} \xi^{p_i/2} \sqrt{v} \mapsto (A, B) \mapsto Q_{\text{FFGFT}} = \frac{\sum_i r_i \xi^{p_i}}{(\sum_i \sqrt{r_i} \xi^{p_i/2})^2}. \quad (.9)$$

Since $\sqrt{v}$ cancels, the map is explicit and trigonometry-free.

The Δ_K discriminator:

$$\Delta_K = 3Q_{\text{FFGFT}} - 2 = \frac{B}{A} - 1 \approx +0.003 \quad (.10)$$

¹P. Phillips (IPI correspondence, May 2026 [1]) postulates a 24-dimensional mediation space with an involution J , $J^2 = I$, whose projectors $P_\pm = (I \pm J)/2$ have ranks $\text{rank}(P_-) = 16$ and $\text{rank}(P_+) = 8$. This yields the ratios $8/24 = 1/3$ and $16/24 = 2/3$ as a pure rank statement, independent of masses or exponents. The geometric motivation is the regular 24-cell polytope (24 vertices in $\mathbb{R}^4$) whose symmetry group F_4 admits a natural $8 + 16$ splitting of the weight space. A formal map to FFGFT quantities has not yet been established.

confirms that the ladder road and the equal-projector road meet near $2/3$ and separate – exactly as the ε -deformation predicts.

.7 Summary

1. At $D = 3$ three roads converge to $2/3$; the $D = 3 + \varepsilon$ deformation separates them.
2. The FFGFT ladder $(\frac{3}{2}, 1, \frac{2}{3})$ is geometric with ratio $q = 2/3$; the unequal gaps $1/2$ and $1/3$ are consequences.
3. FFGFT lies on the complement road $p_\tau = 1 - 1/D$, not on the equal-projector road $2/D$.
4. Fractal corrections (D_f) cancel in Q ; the step is tied to topological $D = 3$.
5. The ladder predicts $Q \approx 0.6677$ (0.16 %) without Koide input; the residual arises from cross-term exponents outside $\frac{1}{3}\mathbb{Z}$.
6. Exact $Q = 2/3$ would require $\xi_{\text{Koide}} \approx 1.372 \times 10^{-4}$; the agreement is a confirmation, not a fit.

Methodological classification: The algebraic derivations in Sections .3–.4 are bridges (directly from the ladder structure and torus topology). The connection to Paul's 24-cell rank split is a plausibility sketch; a formal map requires further elaboration.

Johann Pascher, Upper Austria, May 2026

Bibliography

- [1] P. Phillips, *IPI correspondence: 24-dimensional mediation space and $1/3 + 2/3$ rank split*, unpublished private communication, IPI group (May 2026).
- [2] P. Austin, *A Purely Algebraic Short Note on the Repeated Appearance of $2/3$: Koide, Projectors, Complements, FFGFT, and $n + \varepsilon$ Deformation*, IPI correspondence (May 2026).
- [3] Y. Koide, *A fermion-boson composite model of quarks and leptons*, Physics Letters B **120**(1-3), 161-165 (1983).
- [4] J. Pascher, *Dok. 006 – Fundamental Parameter Table of FFGFT*, in: FFGFT corpus, Zenodo DOI: 10.5281/zenodo.20474821 (2026).
- [5] J. Pascher, *Dok. 051 – Spin-winding contribution and electron exponent $p_e = 3/2$* , in: FFGFT corpus, Zenodo DOI: 10.5281/zenodo.20474821 (2026).
- [6] J. Pascher, *Dok. 192 – Koide analysis: $Q_{FFGFT} \approx 0.6677$ and non-closure theorem*, in: FFGFT corpus, Zenodo DOI: 10.5281/zenodo.20474821 (2026).
- [7] J. Pascher, *Fundamental Fractal-Geometric Field Theory (FFGFT) – T_0 Time-Mass Duality*, Zenodo, DOI: 10.5281/zenodo.20474821 (2026).