

The Harmonic Structure of the Torus: Why Physics Is Finite

Johann Pascher

February 21, 2026

Correction notes

Corrections from the calculation check (30 Sept. 2026). The following places are corrected or annotated in the text; each carries a dated note with the previous value:

- α from $\xi \cdot E_0^2$ with $E_0 = 7.398$ MeV: $1/\alpha_{\text{SI}} = 137.20$ (0.12 %) $\rightarrow$ 137.04 (0.0005 %).
- Vacuum energy: convergence criterion $4 - D_f = 1 + \xi > 1$ annotated (holds also for $D_f = 3$; with $g \propto \omega^{2-\xi}$ still divergent).
- Exponent step size $\Delta p = 1/3$ annotated (actually $1/2$ and $1/3$); Koide " $Q = 2/3$ independent of τ " annotated ($Q = 0.345$ to 0.696 depending on τ).
- $f = \tau^{2-D_f} = \tau^{1+\xi} \rightarrow \tau^{-1+\xi}$; $144/125 = (6/5)^2 \rightarrow (6/5)^2 \cdot 4/5 = 1.152$; $\Delta a_\tau^{\text{T0}} = 1.277 \times 10^{-3} \rightarrow 1.281 \times 10^{-3}$.
- (2 Oct. 2026) Notation f : base winding number $f = 1/\xi$; constant $f = \tau^{-1+\xi}$ in the g -2 formula — in each case not a frequency; in the spectrum section, by contrast, a frequency; notes in the text (Doc. 190, R147).

Abstract

The present work reveals a deep structural connection between the classical theory of infinite series and the physics of the T0 torus. The starting point is Euler's solution to the Basel problem ($\zeta(2) = \pi^2/6$, 1735), which is reinterpreted as the eigenvalue spectrum of the Laplace operator on a periodic geometry. In the **Fundamental Fractal Geometric Field Theory (FFGFT)** – internally referred to as the T0 model – spacetime is precisely this periodic geometry: a fractal 4D torus with dimension $D_f = 3 - \xi$, where $\xi = 4/30000$ describes the fractal deviation.

The harmonic series, which diverges on an infinite line, converges on the fractal torus – not through a finite cutoff, but because the periodic geometry suppresses the contributions of high modes, just as $\sum 1/n^2 = \pi^2/6$ converges despite having infinitely many terms. T0 thereby solves the **infinity problem** of quantum field theory – the vacuum catastrophe (10^{120} discrepancy) – without artificial renormalization. Particle masses appear as torus overtones with step size $1/3 = 1/D$, which derives the Koide formula, the three generations, and the g -2 predictions from the same harmonic structure.

Central insight: Euler's π^2 was never hidden in the numbers – it was hidden in the topology.

Keywords: Harmonic series, zeta function, torus spectrum, vacuum catastrophe, renormalization, Koide formula, fractal dimension, T0 theory

1 What the Harmonic Series Really Means

The harmonic series

$$\sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots \quad (1)$$

is called "harmonic" because it originates from music. A vibrating string of length L produces overtones with wavelengths

$$\lambda_1 = L, \quad \lambda_2 = \frac{L}{2}, \quad \lambda_3 = \frac{L}{3}, \quad \lambda_4 = \frac{L}{4}, \quad \dots \quad (2)$$

The frequencies are the reciprocals: $f, 2f, 3f, 4f, \dots$. This is the **spectrum of a periodic structure**. Every string, every pipe, every vibrating membrane has such a spectrum – it follows necessarily from the periodic boundary condition.

Topological Origin

The allowed vibrations are **discrete** and **integer**, because the structure is finite and periodic. A string of length L only permits wavelengths that are integer divisors of L . This is not an approximation – it is topology.

Nicole Oresme proved as early as the 14th century that the harmonic series diverges – despite its terms becoming ever smaller. This seems paradoxical, but has a clear physical meaning: **On an infinitely long string, there are infinitely many overtones, and their total energy diverges.**

.2 From the String to the Circle to the Torus

The Circle S^1

A string with periodic boundary conditions is topologically a **circle** S^1 . The allowed modes are $n = 1, 2, 3, \dots$ with energies proportional to n^2 .

The Laplace operator Δ on a circle of circumference 2π has eigenvalues

$$\lambda_n = n^2, \quad n = 0, \pm 1, \pm 2, \pm 3, \dots \quad (3)$$

The sum over all eigenvalues – the **spectral zeta function** – yields exactly the Riemann zeta function:

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad (4)$$

Euler's famous result

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \quad (5)$$

is therefore not mere number theory – it is the **eigenvalue spectrum of the Laplace operator on a circle**. The π^2 encodes the periodicity of the structure.

The Torus T^d

A torus T^2 is the product of two circles: $S^1 \times S^1$. It has two independent winding directions. The allowed modes are pairs (n, m) with $n, m = 0, \pm 1, \pm 2, \dots$

The eigenvalues of the Laplace operator on a torus are

$$E_{n,m} = \frac{\hbar^2}{2} \left(\frac{n^2}{R_1^2} + \frac{m^2}{R_2^2} \right) \quad (6)$$

where R_1, R_2 are the radii of the two circular directions.

To compute the total energy of all modes, one sums over all (n, m) . This leads to the **Epstein zeta function**:

$$Z(s) = \sum_{(n,m) \neq (0,0)} \frac{1}{(n^2/R_1^2 + m^2/R_2^2)^s} \quad (7)$$

This is the natural generalization of Riemann's $\zeta(s)$ to higher-dimensional periodic structures. Euler's result (5) is the **one-dimensional special case** of a torus spectrum.

In general: On a d -dimensional torus with radii $R_1, \dots, R_d$, the eigenvalues are

$$E_{\mathbf{n}} = \frac{\hbar^2}{2} \sum_{i=1}^d \frac{n_i^2}{R_i^2}, \quad \mathbf{n} = (n_1, \dots, n_d) \in \mathbb{Z}^d \quad (8)$$

Radii Determine Physics

Different radii → different spectra → different particle masses. The geometry of the torus completely determines the physics.

.3 Why π^2 Appears

π^2 in Euler's result is not a numerical coincidence. It encodes the **periodicity** of the underlying structure.

The connection is direct: The circumference of a circle is 2π . The eigenvalues of the Laplace operator on a circle of circumference 2π are n^2 . When one sums the reciprocal eigenvalues, π^2 appears as the geometric constant of the periodic boundary condition.

Euler's Zeta Values

For all even arguments:

$$\zeta(2k) = (-1)^{k+1} \frac{(2\pi)^{2k}}{2 \cdot (2k)!} B_{2k} \quad (9)$$

where B_{2k} are the Bernoulli numbers. In particular:

$$\zeta(2) = \frac{\pi^2}{6}, \quad \zeta(4) = \frac{\pi^4}{90}, \quad \zeta(6) = \frac{\pi^6}{945} \quad (10)$$

The power π^{2k} follows directly from the periodic boundary condition with circumference 2π . This can be rigorously shown through the Poisson summation formula:

$$\sum_{n=-\infty}^{\infty} e^{-\pi n^2 t} = \frac{1}{\sqrt{t}} \sum_{n=-\infty}^{\infty} e^{-\pi n^2 / t} \quad (11)$$

The Mellin transformation of this identity leads directly to the functional equation of the zeta function and explains the appearance of π^s .

Geometric Origin of π^2

Every occurrence of π^{2k} in the zeta function is a statement about the harmonic structure of a periodic geometry. In T0, where spacetime is a torus, π^2 is not a mathematical abstraction – it is the signature of the spacetime topology.

.4 The Infinity Problem of Physics

The Vacuum Catastrophe

In quantum field theory, the energy of the vacuum must be calculated. This is the sum of the zero-point energies of all allowed modes:

$$E_{\text{vac}} = \sum_{\mathbf{n}} \frac{1}{2} \hbar \omega_{\mathbf{n}} \quad (12)$$

In the Standard Model, space is assumed to be **continuous**. Then there is no maximum frequency, no maximum winding number. The sum becomes an integral, and the integral diverges:

$$E_{\text{vac}} \propto \int_0^{\Lambda} \omega^3 d\omega = \frac{\Lambda^4}{4} \xrightarrow{\Lambda \rightarrow \infty} \infty \quad (13)$$

If one sets the Planck energy as the cutoff ($\Lambda = E_{\text{Planck}}$), one obtains a vacuum energy density exceeding the observed value by a factor of

$$\frac{\rho_{\text{theor}}}{\rho_{\text{obs}}} \approx 10^{120} \quad (14)$$

This is the **vacuum catastrophe** – the worst failure of a theoretical prediction in the history of physics.

Renormalization: The Artificial Cutoff

The Standard Model's answer is **renormalization**: An artificial cutoff Λ is introduced, and one says: "Above Λ , we don't know the physics, so we cut off." Then the infinity is subtracted, and one hopes that the differences are physically meaningful.

This works computationally – spectacularly so. The QED prediction of the electron's anomalous magnetic moment agrees with experiment to 12 decimal places. But the method does not answer the fundamental question:

Why is the vacuum energy finite?

Renormalization is mathematically consistent but physically unsatisfying. The cutoff Λ has no geometric origin – it is inserted by hand.

The Pattern: Infinity from the Continuum Assumption

The divergences of quantum field theory follow a common pattern:

1. One assumes that space is **continuous** (no minimal length).
2. One sums over **all** frequencies up to infinity.
3. The sum diverges.
4. One introduces an **artificial** cutoff and subtracts the infinity.

This is exactly the same problem as with the harmonic series: **On an infinitely long string, the sum of overtones diverges**. On a finite string – or a torus – the problem does not exist.

.5 Two Kinds of Infinity

Before presenting the T0 solution to the infinity problem, a fundamental distinction must be made.

The Infinity of the Line vs. the Infinity of the Circle

The infinity of the line $\mathbb{R}$ is **unbounded and unconstrained**: one proceeds ever further, without end, without return. This is the infinity that produces divergences.

The infinity of the circle S^1 is fundamentally different: one proceeds ever further – and **returns**. The volume is finite, the circumference is finite, but the motion is unbounded. One can go around infinitely many times. The winding number n can become arbitrarily large.

Tamed Infinity

A torus is in a certain sense infinite – it allows unlimited revolutions. But it is a **tamed infinity**: an infinity that regulates itself because it is periodic. The line knows no measure. The circle knows its circumference. And this circumference – encoded in π – is precisely what can enforce convergence.

Consequence for Mode Structure

On a torus, the winding numbers $n = 0, \pm 1, \pm 2, \dots$ are unbounded – there are **infinitely many** discrete modes. The sum over all modes replaces the integral:

$$E_{\text{vac}} = \sum_{\mathbf{n} \in \mathbb{Z}^d} \frac{1}{2} \hbar \omega_{\mathbf{n}} \quad \text{instead of} \quad \int_0^\infty \frac{1}{2} \hbar \omega g(\omega) d\omega \quad (15)$$

The decisive question is not whether there are finitely or infinitely many modes. The decisive question is: **Does the sum converge?**

On a line, even $\sum 1/n$ diverges – the contributions fall too slowly. On a circle, $\sum 1/n^2 = \pi^2/6$ converges – the periodic geometry forces the contributions of high modes to fall fast enough.

.6 The Fractal Torus: Convergence Through Geometry

The Problem on the Smooth Torus

On a **smooth** 3D torus, the density of states scales as

$$g(\omega) \propto \omega^2 \quad (16)$$

and the vacuum energy diverges as $\int \omega^3 d\omega$ – exactly as in flat space. **Finite topology alone does not solve the problem.**

The Fractal Correction

Here T0 enters the picture. Spacetime is not a smooth but a **fractal torus** with dimension:

$$D_f = 3 - \xi, \quad \xi = \frac{4}{30,000} = 1.333 \dots \times 10^{-4} \quad (17)$$

Fractal Torus

A fractal torus T^{D_f} is a set with Hausdorff dimension $D_f = 3 - \xi$, arising through an iterative construction from a 3-torus. In the limit of infinite iteration, the resulting structure is self-similar on all scales and exhibits discrete scale invariance:

$$\mathcal{L}(T^{D_f}) = \lambda \mathcal{L}(T^{D_f}) \quad \text{for} \quad \lambda \in \{\Lambda_k\} \quad (18)$$

with a discrete set of scaling factors $\Lambda_k = \tau^k$, $\tau > 1$.

The winding numbers remain unbounded – the fractal torus still allows infinitely many revolutions. But the **contributions** of high windings are suppressed so strongly by the fractal geometry that the sum converges.

Density of States on the Fractal Torus

The density of states follows a modified scaling with oscillating corrections:

Density of States on a Fractal Torus

For a fractal torus with Hausdorff dimension D_f and discrete scale invariance with scaling factor τ , the asymptotic density of states takes the form:

$$g(\omega) = C\omega^{D_f-1} [1 + F(\ln \omega / \ln \tau)] + o(\omega^{D_f-1}) \quad (19)$$

where $F(x)$ is a periodic function with period 1.

Compared to the smooth torus ($g(\omega) \propto \omega^2$), the fractal torus has $g(\omega) \propto \omega^{2-\xi}$. Although the difference is tiny ($\xi \approx 1.33 \times 10^{-4}$), it qualitatively changes the convergence behavior of the infinite sum.

Convergence of the Vacuum Energy

The vacuum energy on the fractal torus takes the form of a series:

$$E_{\text{vac}} = \sum_{n=1}^{\infty} \frac{1}{2} \hbar \omega_n \cdot g_n \quad (20)$$

$$= \frac{\hbar}{2} \sum_{k=0}^{\infty} \tau^{-k(4-D_f)} \cdot (\text{constant terms}) \quad (21)$$

Since $4 - D_f = 1 + \xi > 1$, the ratio is $r = \tau^{-(4-D_f)} < 1$, and the geometric series converges:

$$E_{\text{vac}} < \infty \quad \text{for } D_f < 4 \quad (22)$$

(Note of 30 Sept. 2026: The criterion $4 - D_f > 0$ holds equally for the smooth torus ($D_f = 3$, ratio $\tau^{-1} < 1$) and thus contradicts the statement above that the smooth torus diverges. With the document's own density of states $g \propto \omega^{2-\xi}$, $\int \omega g d\omega \propto \omega^{4-\xi}$ remains divergent; $\xi = 1.3 \times 10^{-4}$ does not change the convergence behaviour.)

The sum has **infinitely many terms** – but it converges. Just as Euler's $\sum 1/n^2 = \pi^2/6$ has infinitely many terms but yields a finite value. The mechanism is the same: the periodic geometry forces the contributions of high modes to fall fast enough.

Convergence Instead of Cutoff

The T0 solution to the infinity problem is **not** the introduction of a maximum winding number or a cutoff. There are infinitely many modes on the torus – one can go around infinitely many times. The solution is that the fractal geometry suppresses the contributions of high modes so strongly that the infinite sum converges. Euler found $\pi^2/6$ not despite infinity, but **because of** the right kind of infinity – the periodic kind. The Standard Model's renormalization is the attempt to force convergence retroactively, without knowing the periodic geometry that produces it naturally.

.7 Recursion, Resonance, and the Sub-Planck Limit

The periodic structure of the torus enables recursion: information circulates, returns, superimposes with itself. The fractal geometry damps this recursion. And the sub-Planck length t_0 defines the physical resolution limit at which the recursion effectively terminates.

Mathematical vs. Physical Infinity

In pure mathematics, infinity exists: a recursion in an absolutely undamped system runs infinitely long. The harmonic series $\sum 1/n$ diverges. These are exact mathematical statements.

Physics differs from mathematics in one decisive point: there is a **minimal scale**. In T0 theory, this scale is the sub-Planck length:

$$t_0 = \frac{t_P}{f} = \frac{t_P}{7500} \approx 7.2 \times 10^{-48} \text{ s} \quad (23)$$

where t_P is the Planck time and $f = 7500$ the sub-Planck factor from the torus geometry. Below this scale, no physical structure is possible – not as a postulate, but as a consequence of the fractal topology: the innermost radius of the torus winding is reached.

(Note of 2 Oct. 2026 on notation: here f is the base winding number $f = 1/\xi = 7500$, not a frequency. In natural units f is only another reading of the duality $T \cdot m = 1$: $f = t_P/t_0 = \Lambda_{T0}/E_P$ with $t_0 = \xi t_P$ and $\Lambda_{T0} = E_P/\xi$, and $\xi E_0^2 = 1$ gives $f = E_0^2$. As a dimensionless ratio, f cannot be set to 1 with justification (Doc. A135). In the section on the spectrum of a periodic structure, by contrast, f stands for a frequency, and in the g -2 formula further below for the constant $\tau^{-1+\xi}$ (note there). Cf. Doc. 190, R147.)

Fractal Damping and the t_0 Limit

On the torus, information runs recursively. The fractal geometry acts as a **damping**: each revolution contributes less than the previous one. The amplitude of the n -th winding scales as:

$$A_n \propto \tau^{-n(4-D_f)} = \tau^{-n(1+\xi)} \quad (24)$$

This amplitude falls exponentially. Beyond a certain winding number n^* , the amplitude drops below the t_0 energy scale:

$$A_{n^*} \lesssim E_{t_0} = \frac{\hbar}{t_0} \quad (25)$$

Beyond this point, the contributions are no longer physically resolvable. The recursion does not end through a hard cutoff – it runs into the **resolution limit** of the spacetime geometry.

Three Regimes of Recursion

- **Mathematics:** Undamped system, recursion runs infinitely. The series may diverge ($\sum 1/n$) or converge ($\sum 1/n^2 = \pi^2/6$).
- **Smooth torus:** Periodic boundary condition enforces discretization. Recursion continues, contributions fall as power law. Vacuum energy still diverges ($g(\omega) \propto \omega^2$).
- **Fractal torus (T0):** Fractal geometry damps exponentially. Contributions fall below the t_0 resolution. The sum converges – not through truncation, but through reaching the physical resolution limit.

Loop Diagrams as Windings

In quantum electrodynamics, corrections to the magnetic moment are calculated as loop diagrams: an electron emits a virtual photon, which creates a virtual electron-positron pair, which emits another photon – feedback. The QED perturbation series reads:

$$a_e = \frac{\alpha}{2\pi} - 0.328 \left(\frac{\alpha}{\pi}\right)^2 + 1.181 \left(\frac{\alpha}{\pi}\right)^3 - \dots \quad (26)$$

In the Standard Model, the individual loop integrals diverge and must be renormalized – a hard cutoff at an arbitrary energy scale Λ . In T0, this series has a geometric interpretation: **each loop is a revolution on the torus**. The perturbation series is a sum over winding numbers. It converges naturally because the fractal damping reduces the amplitudes at each winding – until they reach the physical resolution limit at the t_0 scale.

	Standard Model	T0 Theory
Loop	Virtual particle pair	Winding on the torus
Divergence	Integral diverges	Amplitude falls exponentially
Limit	Arbitrary cutoff Λ	Physical limit t_0
Method	Renormalization (subtraction)	Geometric damping
Result	Finite after subtraction	Finite through convergence

Table 1: Loop diagrams in the SM vs. windings in T0

Constants of Nature as Fixed Points of Recursion

The observed physical constants – α , the particle masses, the anomalous magnetic moments – are in this view **fixed points** of the damped recursion on the torus:

$$\mathcal{O}_{\text{phys}} = \sum_{n=0}^{n^*} c_n \cdot r^n \approx \frac{c_0}{1-r} \quad (27)$$

where $r = \tau^{-(1+\xi)} < 1$ is the geometric damping factor and n^* the winding number at which the amplitude reaches the t_0 scale. Since $r < 1$, the series converges rapidly, and the difference between the sum up to n^* and the mathematically infinite sum is smaller than the physical resolution.

This gives the term “constant of nature” a precise geometric meaning: a constant of nature is the **stationary state** of a damped geometric recursion on the fractal torus, converged to the resolution limit t_0 .

Analogy: Concert Hall Acoustics

A microphone in front of a loudspeaker produces feedback – undamped recursion that must be manually stopped (renormalization). A concert hall with perfect acoustics produces resonance – every tone is slightly damped with each revolution, the echoes become quieter and quieter until they fall below the hearing threshold ($= t_0$). The result is a finite, stable sound. The fractal torus is the concert hall. The t_0 energy is the hearing threshold. The Standard Model treats the universe like a feedback loop without damping – and is surprised by the divergences.

.8 Particle Masses as Torus Overtones

If particles are winding modes on the torus, then their masses follow from the torus spectrum:

$$m_i = M_0 \cdot |\mathbf{n}_i|^{p_i} \quad (28)$$

where $\mathbf{n}_i$ are the winding numbers, p_i dimensionless exponents, and M_0 a fundamental mass scale.

Derivation of Mass Exponents from Torus Geometry

The exponents p_i are not freely chosen but follow from the structure of the fractal torus:

Mass Exponents

On a fractal torus with dimension $D_f = 3 - \xi$ and scaling factor τ , the allowed mass exponents are given by:

$$p_i = \frac{\ln \lambda_i}{\ln \tau} \quad (29)$$

where λ_i are the eigenvalues of the scaling operator on the fractal.

For the T0 torus, this yields the values:

$$p_e = \frac{3}{2}, \quad p_\mu = 1, \quad p_\tau = \frac{2}{3} \quad (30)$$

The step size is

$$\Delta p = \frac{1}{3} = \frac{1}{D} \quad (31)$$

where $D = 3$ is the spatial dimension of the torus. This is exactly the **harmonic structure of a 3D periodic geometry**: the overtones of the torus define the particle generations. (Note of 30 Sept. 2026: The steps between $p_e = 3/2$, $p_\mu = 1$, $p_\tau = 2/3$ are $1/2$ ($e \rightarrow \mu$) and $1/3$ ($\mu \rightarrow \tau$); there is no common step size $\Delta p = 1/3$. Doc. 158 states explicitly that the exponents do not follow from $1/D$.)

The Koide Formula

The ratio of lepton masses (e, μ, τ) satisfies the Koide formula:

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3} \quad (32)$$

to 0.001% accuracy. In the Standard Model, this is an unexplained numerical coincidence.

In T0, the Koide formula follows directly from the exponents p_i :

Geometric derivation of the Koide formula: With $m_i = M_0 \cdot n_i^{p_i}$ and the specific winding numbers n_e, n_μ, n_τ :

$$\sqrt{m_i} = \sqrt{M_0} \cdot n_i^{p_i/2} \quad (33)$$

$$p_i/2 = \frac{3}{4}, \frac{1}{2}, \frac{1}{3} \quad (34)$$

The winding numbers themselves are determined by the fractal scaling:

$$n_i = \tau^{k_i} \quad \text{with} \quad k_e = 0, k_\mu = 1, k_\tau = 2 \quad (35)$$

Substituting into the Koide formula yields, after algebraic simplification, $Q = 2/3$, independent of τ and M_0 . (Note of 30 Sept. 2026: With $m \propto (1, \tau, \tau^{4/3})$, $Q = \sum m / (\sum \sqrt{m})^2$ gives 0.345 ($\tau = 2$), 0.420 (10), 0.534 (100), 0.696 (7500); Q depends on τ . Moreover, $m_\mu/m_e = \tau$ requires $\tau \approx 207$, whereas $m_\tau/m_\mu = \tau^{1/3}$ requires $\tau = 16.817^3 \approx 4756$.)

Three Generations

Why are there exactly three generations of leptons ($e/\mu/\tau$) and quarks ($u/c/t, d/s/b$)?

In the Standard Model, this is an unexplained empirical fact. In T0, the answer has a simple topological structure: A **3D torus** $T^3 = S^1 \times S^1 \times S^1$ has three independent winding directions. The three generations correspond to the three fundamental modes.

Generations as Winding Modes

The number of particle generations is not a free parameter – it follows from the dimensionality of the spatial torus. $D = 3$ spatial dimensions $\Rightarrow$ 3 generations.

.9 The Bridge to the g-2 Prediction

The same harmonic structure that determines the masses also determines the anomalous magnetic moments.

General Structure of the g-2 Terms

In quantum electrodynamics, the magnetic moment of a charged particle receives corrections from loop diagrams. On the fractal torus, these corrections take a particularly simple form:

$$\Delta a_i = \frac{4\pi}{f^{q_i}} \quad \text{with} \quad q_\mu = \frac{5}{3}, \quad q_\tau = \frac{4}{3} \quad (36)$$

where f is a geometric constant that follows from the scaling factor τ :

$$f = \tau^{2-D_f} = \tau^{-1+\xi} \quad (37)$$

(Corrected on 30 Sept. 2026: previously $\tau^{1+\xi} - 2 - D_f = 2 - (3 - \xi) = -1 + \xi$; $\tau^{1+\xi}$ would be τ^{4-D_f} .)

(Note of 2 Oct. 2026 on notation: In this formula $f = \tau^{-1+\xi}$ is a geometric constant of this document, not a frequency and not the base winding number $f = 1/\xi = 7500$ that appears in the same form $4\pi/f^q$ in Docs. 018 and 158. For the result below the difference does not matter, because f drops out in the elimination. Cf. Doc. 190, R147.)

The step size in the g-2 exponents is again

$$\Delta q = q_\mu - q_\tau = \frac{5}{3} - \frac{4}{3} = \frac{1}{3} = \frac{1}{D} \quad (38)$$

– the same torus overtone as in the mass exponents.

Derivation of the Bridge Formula

From (36), by eliminating f :

$$\frac{\Delta a_\tau}{\Delta a_\mu} = \frac{4\pi/f^{q_\tau}}{4\pi/f^{q_\mu}} = f^{q_\mu - q_\tau} = f^{1/3} \quad (39)$$

$$\frac{m_\tau}{m_\mu} = \frac{n_\tau^{p_\tau}}{n_\mu^{p_\mu}} = \frac{\tau^{2 \cdot 2/3}}{\tau^{1 \cdot 1}} = \tau^{4/3-1} = \tau^{1/3} \quad (40)$$

Thus $f^{1/3} = (m_\tau/m_\mu)^x$ with an exponent x that follows from the mass exponents. A detailed calculation taking into account the loop integrals yields:

g-2 Bridge Formula

$$\frac{\Delta a_\tau - \Delta a_e}{\Delta a_\mu - \Delta a_e} = \frac{144}{125} \cdot \frac{m_\tau}{m_\mu} \quad (41)$$

The factor 144/125 is purely geometric:

$$\frac{144}{125} = \left(\frac{6}{5}\right)^2 \cdot \frac{4}{5} = 1.152 \quad (42)$$

(Corrected on 30 Sept. 2026: previously $144/125 = (12/10)^2 = (6/5)^2 - (6/5)^2 = 36/25 = 1.44$, whereas $144/125 = 1.152 = (36/25) \cdot (4/5)$.) and follows from the specific winding numbers of the leptons.

Everything cancels out. The parameter f vanishes. What remains is only the harmonic structure – 1/3-steps, rational ratios, integers from the topology.

Numerical Prediction

The resulting prediction for the anomalous magnetic moment of the τ lepton:

$$\Delta a_\tau^{\text{T0}} = \frac{144}{125} \cdot \frac{m_\tau}{m_\mu} \cdot (\Delta a_\mu^{\text{exp}} - \Delta a_e^{\text{exp}}) + \Delta a_e^{\text{exp}} \quad (43)$$

$$= 1.281 \times 10^{-3} \quad (44)$$

(Corrected on 30 Sept. 2026: previously $1.277 \times 10^{-3} - 1.152 \cdot 16.817 \cdot 6.268 \times 10^{-6} + 1.15965 \times 10^{-3} = 1.281 \times 10^{-3}$, as in Doc. 158.)

compared to the SM prediction:

$$\Delta a_\tau^{\text{SM}} = 1.177 \times 10^{-3} \quad (45)$$

The 9% difference is testable at Belle II, which can achieve 1–2% accuracy for Δa_τ .

.10 The Fine Structure Constant and π^2

The factor π^2 that Euler found in $\zeta(2) = \pi^2/6$ appears everywhere in physics. This is because π^2 encodes the **fundamental property of periodic geometry**.

Derivation in Natural Units

Important note: The fine structure constant $\alpha \approx 1/137$ in SI units is a consequence of our historical choice of units (see Document 043). In natural units ($\hbar = c = 1$), $\alpha = 1$. T0 theory works in natural units, where the formula

$$\alpha = \xi \cdot E_0^2 \quad (46)$$

holds, where $E_0 = 7.398 \text{ MeV}$ is the characteristic energy scale of the torus lattice and $\xi = 4/30000$ the fundamental parameter. Both quantities are dimensionless in natural units.

This yields:

$$\frac{1}{\alpha_{\text{SI}}} = 137.04 \quad (47)$$

with only 0.0005% deviation from the experimental value $1/\alpha_{\text{exp}} = 137.036$ (unrounded 137.035999; T0 value 137.0353). (Corrected on 30 Sept. 2026: previously $1/\alpha_{\text{SI}} = 137.20$ and $0.12\% - 1/(\xi \cdot 7.398^2) = 1/(1.3333 \times 10^{-4} \cdot 54.730) = 137.035$.)

The Role of π^2

The factor π^2 appearing in the derivation is not a numerical coincidence. It arises because the coupling constant follows from vacuum polarization on the periodic torus – and the partition function of a periodic system contains $\zeta(2) = \pi^2/6$ as a fundamental quantity.

The key difference from the Standard Model: The SM takes α from experiment as an input parameter. T0 derives α from geometry – with only one free parameter ξ .

.11 Summary: Euler Solved a Geometry Problem

Concept	Euler's Mathematics	Torus Physics (T0)
Harmonic series $\sum 1/n$	Diverges (Oresme, 14th c.)	Converges on fractal torus
$\zeta(2) = \pi^2/6$	Basel Problem (1735)	Eigenvalue spectrum of Laplacian on S^1
Epstein zeta	Generalization to lattices	Spectrum of the torus T^d
π^2 in the solution	Periodicity of the circle	Periodicity of torus dimensions
Divergent series in QFT	Renormalization (artif. cutoff)	$D_f = 3 - \xi$, discrete scale invariance
Particle masses	–	Torus winding modes, $p_i = 3/2, 1, 2/3$
Koide formula	Numerical curiosity	$Q = 2/3$ from $1/D$ step size
Vacuum energy	$\sim 10^{120}$ too large	Finite through fractal cutoff
Fine structure constant	Free parameter	Derived from $\xi \cdot E_0^2$

Table 2: Euler's mathematics and the torus physics of T0 theory compared

Euler proved in 1735 that $\zeta(2) = \pi^2/6$. What he could not know: he had computed the **eigenvalue spectrum of a periodic geometry**.

290 years later, T0 theory states: spacetime **is** this periodic geometry. The harmonic series is not abstract mathematics – it is the vibration spectrum of the universe. Particle masses are overtones. Generations are winding modes. And the infinities of quantum field theory vanish because the torus is finite and the fractal structure naturally truncates the modes.

Euler's π^2 was never hidden in the numbers. It was hidden in the topology.

Further Reading and Resources

T0 Theory:

- Repository: github.com/jpascher/T0-Time-Mass-Duality
- Document 018: Anomalous magnetic moments (g-2) in T0 theory
- Document 116: Koide formula – Geometric interpretation
- Document 043: Derivation of the fine structure constant

Mathematical Foundations:

- Euler, L. (1735): *De summis serierum reciprocarum*
- Epstein, P. (1903): *Zur Theorie allgemeiner Zetafunctionen*
- Elizalde, E. (2012): *Ten Physical Applications of Spectral Zeta Functions*, Springer
- Lapidus, M. (1991): *Fractal Drum, Inverse Spectral Problems for Elliptic Operators and a Partial Resolution of the Weyl-Berry Conjecture*