

Corrections and Clarifications

FFGFT / T0 Time-Mass Duality

Johann Pascher

2026

Contents

.1	Correction 1: Prefactor in the Higgs derivation of ξ	3
.2	Correction 2: Tau-Yukawa prefactor r_τ	3
.3	Correction 3: Base formula for g^{-2} of the electron	4
.4	Refinement 1: Accuracy of the Koide formula	5
.5	Refinement 2: $\hbar$ from the Higgs field – derivation chain	5
.6	Refinement 3: Two ξ parameters – two distinct spaces	6
.7	Refinement 4: L_0 and the Schwarzschild interpretation	6
.8	Refinement 5: Scope and reach of the mass formulas	7
.9	Refinement 6: Koide formula only with original values	7
.10	Refinement 7: Terminology “renormalization group” and scale structure	8
.11	Refinement 8: Terminology “fractal renormalization”	9
.12	Refinement 9: Column “Symmetry” and the term “color charge” in the fermion tables	10
.13	Refinement 10: Logical status of $K = 245$ and the factor 1.314 in Doc. 009	12
.14	Refinement 11: T_{CMB} correction factor and K_{frak} disambiguation	13
.15	Refinement 12: Two distinct ΔCHSH observables in Doc. 022 and Doc. 230	14
.16	Refinement 13: Bridge formula between ξ_{FFGFT} and ξ_{UIFT}	15
.17	Refinement 14: Redshift in Doc. 041 – mechanism, not energy loss	17
.18	Refinement 15: mark cosmological z as a ΛCDM comparison quantity	18
.19	Refinement 16: H_0 is emergent (decompactified), not fundamental	19
.20	Refinement 17: circular/fit-dependent DE-DM relations in Doc. 028	20
.21	Refinement 14: CMB anharmonicity, forward vs. retrodiction, and the $ n ^2 = 30$ case (R29/R30/R31)	21

Preliminary Note

Status: C1–C3, R1–R34 (R16/R17 partial, R20/R30/R32 open, R29 partial, R31 partially derived, R33 declared, R34 clarified) (as of 8 June 2026)

All corrections (K1–K3) and refinements (P1–P13) listed in this document have been incorporated into the respective source documents. The last open item — K2 (tau prefactor r_τ) — was fully completed on 27 May 2026: in addition to the items reported on 26 May in Docs. 016, 046 and 116, this also covered remaining *derived* values in Doc. 046 DE (energy $18.8 \rightarrow 18.0$ meV) and the entire **English** parallel versions of Docs. 016, 046 and 116, which had not yet been updated on 26 May. See the section “Status of Integration” under K2 for details.

This document is retained as a historical archive. It is **no longer to be used as a binding reference** — the current state of each source document is authoritative.

Addendum (2 June 2026): Added and incorporated is **R14** (formulation of the redshift in Doc. 041): implemented in Doc. 041 (DE+EN).

Addendum (6 June 2026): Carried over from the CMB work (Doc. 267 Cosmological Degeneracy, Doc. 268 CMB peaks from T^4): **R18** (Λ CDM/FFGFT as two readings, theoretical rather than empirical), **R20** (absolute cosmological scale: *form* fixed as a ξ ratio R_H/ℓ_P , the *factor* $41/4$ a declared external calibration from Λ CDM, not derived, open), **R29** (peak selection rule, only *partially* answered – the factor-3 filter does *not* exclude $|n|^2 = 30$), **R30** (rigorous C_ℓ projection; the naive spherical Bessel sum is *insufficient*, open) and **R31** (geometric anharmonicity from the codimension-1 projection – leading behaviour derived forward, small residual open). New **R32** (statements presenting H_0 as FFGFT-derived are to be cleaned up corpus-wide – noted for now). The previously separate points **R19** (CMB path-lengthening) and **R28** (exponent for z_*) have been merged into **R14** resp. **R20** and do not appear as separate rows.

Addendum (8 June 2026): New **R33** – the magnitude-carrying factor 5^4 in $\xi = 1/(3 \cdot 2^2 \cdot 5^4)$ is *not* derived forward but grounded *intuitively* (harmonic / Euler Tonnetz) resp. *empirically* (Higgs sector); the *form* $(4/3, 3, 2^2)$ is forward, the *magnitude* (5^4) rests on the intuitive derivation and is openly *declared* as such. Structural parallel to R20 (form fixed, factor open).

Addendum (8 June 2026, II): New **R34** (dimensional structure). Doc. 181 presents $D = 4$ as “follows necessarily”; clarified: **3+1 is empirical input** (no theory derives the dimension count). Forward are only (i) periodicity, (ii) torus-not-sphere ($\pi_1 \neq 0$), (iii) minimal sufficiency (4 instead of 5 via the duality $\tilde{T}m = 1$), (iv) the lossless Hilbert representation (Type III). Recorded alongside: the SI conversions are *indispensable* for the exponent structure (*power of c* , power of time) – c is the space-time bridge ($T^3 \leftrightarrow \text{time}$), $\hbar$ the mass-time bridge (S_m^1); $c = \hbar = 1$ collapses L, T, M and destroys the powers. The four-circle structure *is* that bookkeeping. Signature as R20/R33 (structure forward, number input).

This document lists corrections and refinements for previously published documents in the FFGFT series. Older documents are *not* modified. Where a contradiction exists between an older and a newer document, the version recorded here is binding.

Each entry contains: affected document and location, the erroneous expression, the correct expression, and a brief justification.

.1 Correction 1: Prefactor in the Higgs derivation of ξ

Affected Document

Doc. 049 (T0 Lagrangian and Higgs Integration), HTML draft version and earlier working versions.

Erroneous Expression

$$\xi = \frac{\lambda_h^2 v^2}{64\pi^4 m_h^2} \approx 1.0 \times 10^{-5} \quad (\text{old – wrong})$$

Correct Expression

$$\xi = \frac{\lambda_h^2 v^2}{16\pi^3 m_h^2} \approx 1.33 \times 10^{-4} \quad (190.1)$$

with the experimental inputs:

$$\begin{aligned} m_h &= 125.25 \text{ GeV} \quad (\text{Higgs mass}), \\ v &= 246.22 \text{ GeV} \quad (\text{vacuum expectation value}), \\ \lambda_h &= \frac{m_h^2}{2v^2} \approx 0.129 \quad (\text{Higgs self-coupling}). \end{aligned}$$

Justification

The prefactor $64\pi^4$ originated as a typo in the early HTML visualization. It yields $\xi \approx 10^{-5}$, which deviates by an order of magnitude from the geometrically derived value $\xi = 4/3 \cdot 10^{-4}$ and does not reproduce the natural constants. The correct prefactor $16\pi^3$ yields $\xi \approx 1.33 \times 10^{-4}$, consistent with the value independently determined from the proton-electron mass ratio (Doc. 009).

.2 Correction 2: Tau-Yukawa prefactor r_τ

Affected Document

Doc. 116 (Koide formula as a consequence of T0 theory), table of Yukawa coefficients.

Erroneous Expression

$$r_\tau = \frac{8}{3} \approx 2.667 \quad \Rightarrow \quad m_\tau \approx 1712 \text{ MeV} \quad (-3.6 \% \text{ dev.}) \quad (\text{old – wrong})$$

Correct Expression

$$r_\tau = \frac{25}{9} \approx 2.778 \quad \Rightarrow \quad m_\tau \approx 1783 \text{ MeV} \quad (+0.4 \% \text{ dev.}) \quad (190.2)$$

The lepton masses are generally given by:

$$m_\ell = r_\ell \cdot \xi^{p_\ell} \cdot v \quad (190.3)$$

with exponents $p_e = 3/2$, $p_\mu = 1$, $p_\tau = 2/3$ (step size $\Delta p = 1/3$) and the vacuum expectation value v .

Justification

$r_\tau = 8/3$ was a remnant from an earlier iteration stage. Doc. 158 (Koide-to-g-2 bridge) already uses $r_\tau = 25/9$, which agrees numerically with the tau mass. Furthermore, $25/9 = (5/3)^2$ has a geometric interpretation as the square of the pure sixth in the natural overtone series, consistent with the musical resonance analogy (Doc. 060).

Complete Corrected Prefactor Table

Lepton	Exponent p	Prefactor r	m [MeV]
Electron	$3/2$	$4/3$	0.511
Muon	1	$16/5$	105.7
Tau	$2/3$	$25/9$	1783

Status of Integration (added 27 May 2026)

A renewed corpus-wide search found **remaining 8/3 instances**; the original entry listed only Doc. 116 and overlooked Docs. 016 and 046. An independent full check on 27 May further showed that the German side had been updated but the **English** parallel versions had not.

Document / location	Found	Target
Doc. 116 EN, lepton-parameter tab.	$r_\tau = 8/3$	$25/9$
Doc. 016 EN, mass tab. & r -list	$r_\tau = 8/3$	$25/9$
($m_\tau = 1712.1$, -3.64%)		(1783.4 , $+0.37\%$)
Doc. 046 EN, v_τ (both rows)	$r = 8/3$	$25/9$
($\xi_{v_\tau} = 128/27$, $E = 18.8$)		($400/81$, 18.0)
Doc. 046 DE, derived tables	$E = 18.8$ meV	18.0 meV

Completed on 27 May 2026: All locations above were set to $25/9$ (and $\xi_{v_\tau} = 400/81$, $E = 18.0$ meV). In Doc. 016 the derived tau mass ($1712.1 \rightarrow 1783.4$ MeV) and deviation ($3.64 \rightarrow 0.37\%$) were recomputed. A corpus-wide search afterwards returned **zero remaining 8/3 instances** in 016/046/116 (DE+EN); all four files were recompiled (Kindle 6×9). K2 is fully incorporated — both German and English — as of 27 May 2026.

.3 Correction 3: Base formula for g -2 of the electron

Affected Document

T0 Explorer (HTML visualization), first version; earlier draft versions of Doc. 018.

Erroneous Expression

$$a_e = \frac{4\pi}{f \cdot k_{\text{geom}}} \quad (\text{old – wrong})$$

Correct Expression

$$a_e = \frac{S_3}{f \cdot k_{\text{geom}}} = \frac{2\pi^2}{f \cdot k_{\text{geom}}} \quad (190.4)$$

with $f = 1/\xi \approx 7500$, $k_{\text{geom}} = (2/\sqrt{\varphi}) \cdot \sqrt{2} \approx 2.224$ and $S_3 = 2\pi^2 \approx 19.74$ (surface area of the 4D unit sphere, i.e. the 3D boundary of the torus).

Justification

The earlier prefactor 4π corresponds to the surface area of the 2D sphere in 3D space, not the geometrically correct 3D surface of the 4D torus. $S_3 = 2\pi^2$ is the correct value from torus geometry. The fractal corrections for muon and tau still use 4π :

$$\Delta a_\mu = \frac{4\pi}{f^{5/3}}, \quad (190.5)$$

$$\Delta a_\tau = \frac{4\pi}{f^{4/3}}. \quad (190.6)$$

.4 Refinement 1: Accuracy of the Koide formula

Affected Documents

Doc. 116 (abstract, line 14): $\Delta Q < 0.00003 \%$

Doc. 158 (abstract): "experimentally confirmed to 0.001 %"

Correct Interpretation

The Koide relation

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3} \quad (190.7)$$

is experimentally confirmed to better than 0.001 % (PDG values 2022). The figure $< 0.00003 \%$ in Doc. 116 refers to the internal consistency of the T0 formulas, not the experimental measurement uncertainty. Both statements are substantively correct but refer to different quantities. In external communications, 0.001 % is the appropriate figure.

Note

The individual T0 masses agree with experiment to within $\sim 1 \%$; the high precision of $Q = 2/3$ arises from the specific algebraic structure of the Koide combination, not from individual mass accuracy.

.5 Refinement 2: $\hbar$ from the Higgs field – derivation chain

Context

Planck's constant $\hbar$ is derived differently in various documents. The complete, consistent derivation chain is fixed here.

Binding Derivation Chain

$$\xi = \frac{\lambda_h^2 v^2}{16\pi^3 m_h^2} \longrightarrow L_0 = \xi \cdot \ell_P \longrightarrow \hbar \sim \sqrt{\xi} \cdot \ell_P \cdot c \quad (190.8)$$

The time-mass binding condition provides the physical justification:

$$T(x, t) \cdot m(x, t) = 1 \implies T = \frac{\hbar}{m c^2} \implies E = \hbar \omega. \quad (190.9)$$

$\hbar = 2\pi\hbar$ is therefore *not* a free constant but a geometric consequence of ξ and ℓ_P . The numerical SI value $\hbar \approx 6.626 \times 10^{-34} \text{ J}\cdot\text{s}$ is the calibration of this relation onto the SI base units.

.6 Refinement 3: Two ξ parameters – two distinct spaces

Context

In Documents 173–176 (quantum dots, spin qubits, qubit state spaces, Shor algorithm), two numerically different ξ values appear. Earlier documents did not separate these consistently, which can cause confusion.

Binding Distinction

Parameter	Value	Domain	Meaning
ξ_0 (geometric)	$\frac{4}{30000}$ $\approx 1.33 \times 10^{-4}$	3D geometric space	Length hierarchy of the torus; valid in physical position space
ξ_{Higgs} (algebraic)	$\approx 1.04 \times 10^{-5}$	Number / quantum state space	Algebraic field corrections; Higgs sector; decoherence rates

Justification

$\xi_0 = 4/30000$ follows from the torus geometry of physical space (geometric derivation, independent of SI measurements). ξ_{Higgs} follows from the algebraic Higgs field structure and appears in coupling formulas between quantum states (e.g. Bell correlations, T_2 hierarchy).

The two ξ values are not a contradiction; they describe complementary aspects: geometry of space (ξ_0) vs. algebra of state space (ξ_{Higgs}). External communications must always specify which parameter is meant.

.7 Refinement 4: L_0 and the Schwarzschild interpretation

Affected Documents

Earlier versions of Doc. 180–182 (Lagrangian derivation, torus geometry, maximum cosmological scale).

Erroneous Approach

Earlier versions interpreted $L_0 = \xi_0 \cdot \ell_P$ as the Schwarzschild radius of a fundamental minimal object. This interpretation was explicitly discarded in Doc. 182 (2026 revision).

Correct Interpretation

L_0 is exclusively the **geometric fundamental length** of the FFGFT torus:

$$L_0 = \xi_0 \cdot \ell_P \approx 2.15 \times 10^{-39} \text{ m.} \quad (190.10)$$

It defines the lowest scale stage of the ξ hierarchy and has *no* Schwarzschild interpretation. The cosmological Hubble radius R_H is the system-specific upper limit of the cosmological sector – not a universal maximum scale. Each scale has its own system-specific upper limit.

Note: Schwarzschild as Consistency Check

The Schwarzschild connection remains valid as an **internal consistency check**: an object of mass $M_0 = \xi_0 \cdot m_P/2$ would have a Schwarzschild radius exactly equal to L_0 – with no free parameter. If M_0 matches a scale stage of the ξ hierarchy, that is a non-trivial self-consistency test. The connection is *a probe, not a derivation*.

.8 Refinement 5: Scope and reach of the mass formulas

Context

Several FFGFT documents describe mass derivations for leptons and other particles. The scope of these derivations must be communicated clearly.

Binding Formulation

The FFGFT mass formulas (e.g. $m_\ell = r_\ell \cdot \xi^{p_\ell} \cdot v$) yield values that agree with experiment to $\sim 1\%$. This is **not a precision calculation** but a structural proof:

- The mass formulas *follow geometrically* from ξ and v – they are not fitted.
- The remaining $\sim 1\%$ deviations reflect measurement uncertainties and approximations, not errors in the structure.
- Numerical accuracy beyond the Standard Model is *not* claimed.

To be avoided in external texts: "FFGFT calculates the masses exactly." The correct statement is: "FFGFT *derives* the mass structure from a single parameter ξ and reproduces the orders of magnitude to $\sim 1\%$."

.9 Refinement 6: Koide formula only with original values

Binding Rule

The Koide relation

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3} \quad (190.11)$$

must be evaluated only with experimentally measured masses (PDG values).

Not permitted: substituting the symbolic FFGFT terms $r_\ell \cdot \xi^{p_\ell} \cdot v$ directly into the formula. The $\sqrt{m_i}$ operation generates cross-terms with mixed ξ powers that are foreign to the torus structure space. Any such substitution yields $Q \approx 0.6677 \neq 2/3$ and is not a contradiction to the theory but a category error.

Permitted: evaluating FFGFT terms numerically (e.g. $m_\tau = 1783 \text{ MeV}$) and inserting that result into the Koide formula like a measured value.

Mathematical background: The r_ℓ terms are geometric invariants of their torus modes. The set of admissible pairs (r_i, p_i) is not closed under multiplication: $r_e \cdot r_\mu = 64/15$ does not belong to any torus mode. This is the Non-Closure Theorem (Doc. 193). The general analysis of composition limits is found in Doc. 192.

.10 Refinement 7: Terminology “renormalization group” and scale structure

Affected Documents

- Doc. 005 (T0 tm-extension): “RG flow”, “renormalization group factor”, “renormalization group invariance”, subsection “renormalization group treatment”.
- Doc. 008 (ξ and e): subsection “modified renormalization group equations”.
- Doc. 019 (T0 Lagrangian, earlier version): subsection “renormalization group equations”.
- Doc. 086 (document overview): “renormalization group as flow in parameter space”.
- Doc. 189 (torus derivation): “cumulative renormalization group running”.
- Doc. 202 (FFGFT comprehensive field theory), §18: section title “The renormalization group”.

Previous Formulation

In the documents listed, the ξ_0 -modified scale running of the coupling

$$\frac{d\alpha}{d \ln \mu} = \beta_0 \alpha^2 \left(1 + \xi_0 \ln \frac{\mu}{E_0} \right) \quad (\text{old – imprecise})$$

and the corresponding scale dependence are designated as the *renormalization group*, sometimes adjacent to statements explicitly characterizing ξ_0 as a topologically fixed geometric constant.

Refined Formulation

In FFGFT, the scale dependence of the coupling is *not* a result of a renormalization procedure but a direct consequence of the recursion $\mathcal{R}$ (Doc. 203) on the fractal torus geometry. The correct designation is therefore:

Scale structure from recursion $\equiv$ recursive α running via $\mathcal{R}$	(190.12)
---	----------

Specifically:

- The term $\xi_0 \ln(\mu/E_0)$ arises from the iterated application of the recursion $\mathcal{R}$ to the mode structure, not from a counter-term subtraction.
- ξ_0 is *not* a fixed point of a β function but a topologically fixed constant ($\xi_0 = 4/30000$, Doc. 180).
- Stability, scale flow, and UV finiteness are three aspects of *the same* recursive structure – not separate mechanisms.

Language Convention

Avoid	Use
Renormalization group (implies QFT procedure) RG flow / RG running	Scale structure from recursion Recursive α running; $\mathcal{R}$ -induced scale correction
Renormalization group equations	Scale equations from $\mathcal{R}$
Renormalization group invariance	Recursive scale invariance
Renormalization group fixed point	Recursion fixed point: $\mathcal{R}(\Phi_*) = \Phi_*$

Justification

The designation “renormalization group” imports a tool of standard quantum field theory based on a fundamentally different principle: there UV divergences are absorbed by a procedure, and parameters *run* under a formal scale transformation. In FFGFT neither the divergence problem (cf. Doc. 159, “without artificial renormalization”) nor a free scale parameter to be absorbed exists. The scale dependence is a *structural property* of the recursion, not a subsequent adjustment step.

The FFGFT-native view is already correctly formulated in Doc. 159 (“renormalization as symptom, not as solution”) and Doc. 189 (“geometrically determined, not by renormalization”); this clarification serves to unify the terminology across the older Lagrangian and ξ -derivation series (Doc. 005, 008, 019) and across Doc. 202.

The underlying mechanism is developed in Doc. 203 (recursion operator $\mathcal{R}$) and Doc. 204 (fractal boundary condition).

.11 Refinement 8: Terminology “fractal renormalization”

Affected Documents

Doc. 005, 008, 012, 014 (with 19 occurrences the most frequent source), 041, 060, 133, 141, 159, 181.

Previous Formulation

“Fractal renormalization” is used as an FFGFT-internal term for the geometric correction factors in the conversion between natural and SI units, and for scale-dependent corrections on the fractal torus.

Refined Formulation

Since the word “renormalization” – even in compound form – carries the QFT connotation of a subtraction or adjustment procedure, the term is replaced by context-specific alternatives:

Context	Binding designation
Unit conversion natural $\leftrightarrow$ SI (esp. Doc. 014)	Geometric scale adjustment
Scale-dependent correc- tions on the fractal torus (Doc. 133, 159, 181)	Fractal correction (matches title of Doc. 133)
General scale transforma- tion by recursion	$\mathcal{R}$ -induced scale correction

Justification

The proprietary term “fractal renormalization” was meant to distinguish the procedure clearly from QFT renormalization, but it adopts the central word and unavoidably with it the connotation of a *correction procedure*. In FFGFT this is not a procedure but the direct numerical manifestation of the fractal geometry. The proposed alternatives avoid this misunderstanding without requiring new vocabulary – both phrases are already established in the corpus (Doc. 133 “fractal correction”; “geometric scale adjustment” in the same semantic field as Doc. 014).

.12 Refinement 9: Column “Symmetry” and the term “color charge” in the fermion tables

Affected Documents

- Doc. 006 (T0 particle masses), “universal fermion table”, column “Symmetry”.
- Doc. 046 (Particle Masses, English parallel version), same table, column “Color”.
- Doc. 042 (ξ parameter and particles), table “spatial field patterns”, entry “Quarks: multi-node bound clusters, confined, color charge”.
- Doc. 005 (T0 tm-extension), Method 2 “extended fractal method with QCD integration”: use of $N_c = 3$, α_s , Λ_{QCD} as external inputs.

Previous Formulation

In the fermion tables of Doc. 006/046 each row carries a column “Symmetry” with the following entries:

Particle	Entry under “Symmetry”
Electron, Muon, Tau	Lepton number
ν_e, ν_μ, ν_τ	Double- ξ
Up, Charm, Strange, Top, Bottom	Color
Down	Color + Isospin

This mixture combines four different concepts in one column: an FFGFT scaling class (“Double- ξ ”), an SM conserved quantity (“lepton number”), an SM gauge symmetry (“color”), and an SM quantum number (“isospin”). In addition, the entry “color charge” in Doc. 042 and $N_c = 3$ in Doc. 005 suggest that the SM color symmetry $SU(3)_C$ is an independent component of FFGFT.

Refined Formulation

In FFGFT neither the gauge group $SU(3)_C$ nor a color quantum number exists. Quark masses follow, just like lepton masses, exclusively from the Yukawa resonance formula

$$m_f = r_f \cdot \xi^{p_f} \cdot v \quad (190.13)$$

with r_f a rational prefactor and p_f a rational exponent (step size $\Delta p = 1/3$). This is verified numerically by the script `calc-resonanz-leiter.py`: all six quarks hit their PDG masses to $\sim 0.3\text{--}3.4\%$ *without* use of N_c , α_s , Λ_{QCD} , or any color quantum number.

Corrected column structure of the fermion table

The old column “Symmetry” is replaced by two clearly separated columns:

Particle	FFGFT scaling class	SM correspondence (informational)
Electron, Muon, Tau	Single- ξ	Lepton number
ν_e, ν_μ, ν_τ	Double- ξ	Lepton number, lepton mixing
Up, Charm, Top	Cluster- ξ	Up-type, $T_3 = +1/2$
Down, Strange, Bottom	Cluster- ξ	Down-type, $T_3 = -1/2$

Meaning of the classes:

- *Single- ξ* : $m = r \xi^p v$ with $p \in \{2/3, 1, 3/2\}$.
- *Double- ξ* : $m_\nu = r_\nu \xi^2 m_e$ (additional ξ suppression relative to the charged leptons).
- *Cluster- ξ* : $m = r \xi^p v$ with the same exponents $p \in \{-1/3, 1/2, 2/3, 1, 3/2\}$ as for the leptons, but with multi-node binding in the sense of Doc. 049 (cubic self-interaction $\lambda_s (\delta m_q)^3$).

Language convention for quarks

Avoid	Use
“color charge” (implies $SU(3)_C$ quantum number)	Multi-node binding
“three colors” red/-green/blue” (SM gauge index) “ $N_c = 3$ colors”	Triple-node cluster (the triplet is geometric, not an $SU(3)$ index) $N_{\text{cluster}} = 3$ (geometric node count in the baryon)
“color confinement”	Cluster binding (cubic self-interaction)

Status of Doc. 005, Method 2

Doc. 005 contains two calculation methods:

- *Method 1: Direct geometric method.* Yukawa form according to Eq. (190.13); parameter-free; accuracy $\sim 1\%$ for all fermions. **This is the FFGFT method.**
- *Method 2: Extended fractal method with QCD integration.* Uses N_c , α_s , Λ_{QCD} , lattice-QCD calibration and ML. **This is a phenomenological bridge construction** for connecting to the lattice-QCD literature, not part of the parameter-free FFGFT.

In external communication Method 1 is to be cited. Method 2 may be used if explicitly labelled as “bridge to lattice QCD” or “ML-calibrated extension”.

Justification

The designations “color”, “color charge”, “ N_c ” import a concept from standard quantum field theory ($SU(3)_C$ as gauge symmetry, three colors, eight gluons, color confinement as a non-perturbative property). Doc. 049 explicitly states that FFGFT contains none of these elements: “what we call ,color’ becomes high-energy node binding patterns” (Doc. 049, §3.7), with a single cubic interaction term $\lambda_s(\delta m_q)^3$ instead of eight gluon fields.

The script validation (`calc-resonanz-leiter.py`, `calc_De.py`) shows: all quark masses follow from the same Yukawa formula as the leptons. The column “Symmetry” of the fermion table, however, suggests that $SU(3)_C$ is a distinguishing property of quarks within FFGFT – which is not the case.

The corrected column structure separates what is FFGFT-proper (*scaling class*) from informational SM correspondence (*SM correspondence*), in parallel with the separation of “recursive α running” and “renormalization group” in Refinement 7.

.13 Refinement 10: Logical status of $K = 245$ and the factor 1.314 in Doc. 009

Affected Document

Doc. 009 (Origin of ξ), Section “Why $K = 245$ is fundamental”, subsection “Geometric meaning”.

Misleading Presentation

Doc. 009 lists the following derivation chain for $K = 245$:

- $\varphi^{12} = 321.996$
- Correction by fractal structure: $(1 - \xi)^2 \approx 0.999733$
- Result: $321.996 \times 0.999733 \approx 321.87$
- “Scaling to mass range”: $321.87/1.314 \approx 245$

The factor 1.314 appears here without derivation or independent physical justification.

Correct Logical Order

$K = 245$ is not derived from φ^{12} and 1.314. The correct logical order is the reverse:

$$K = \frac{R_{pe}}{(4/3)^\kappa} = \frac{1836.153}{(4/3)^7} \approx 245.097 \quad (190.14)$$

K is fixed by the observed proton-electron mass ratio R_{pe} and the exponent $\kappa = 7$. It is an *empirical anchor*, not a first-principles result. The factor 1.314 is implicitly defined as $\varphi^{12}/245 \approx 1.314$ and does not constitute an independent derivation of K .

What remains non-trivial

The exponent $\kappa = 7$ is the *unique integer* for which the complete e-p- μ triangle closes simultaneously:

κ	$K \times (4/3)^\kappa$	Agreement with R_{pe}
6	1376.6	−25 %
7	1835.4	✓ 0.04 %
8	2447.2	+33 %

This uniqueness is a genuine constraint, not a tautology. $K = 245$ then follows from $\kappa = 7$ and the empirical R_{pe} .

Corrected claim for external communication

- **Avoid:** " $K = 245$ is derived from first principles; no free parameters."
- **Use:** "One empirical anchor: $K = 245$, fixed by R_{pe} and $\kappa = 7$. From this anchor, five mass ratios follow with < 0.04 % accuracy. The φ^{12} observation is a numerological consistency note, not a derivation."

.14 Refinement 11: T_{CMB} correction factor and K_{frak} disambiguation

Affected Documents

Doc. 025 (Cosmology), Doc. 003 (Foundations), Doc. 133 (Fractal Correction Derivation).

Issue A: K_{frak} does not close the T_{CMB} gap

Doc. 025 derives:

$$T_{\text{CMB}} = \frac{16}{9} \xi = \frac{4}{16875} \text{ eV} \approx 2.7507 \text{ K} \quad (190.15)$$

The observed value is $T_{\text{CMB}} = 2.72548 \text{ K}$ (Planck 2018). The gap is ≈ 0.925 %.

$K_{\text{frak}} = 1 - 100\xi = 74/75$ (Doc. 133) does *not* close this gap. Numerically:

$$\frac{16}{9} \xi \times K_{\text{frak}} = 2.714 \text{ K} \quad (0.42 \text{ % below target, worse)} \quad (190.16)$$

The required correction factor is 0.99083, which satisfies:

$$T_{\text{CMB}} = \frac{16}{9} \xi \left(1 - \frac{275}{4} \xi\right) = 2.725486 \text{ K} \quad (190.17)$$

Agreement with observation: $\Delta = 0.0002$ %.

The coefficient $275/4 = 68.75$ is close to the tetrahedral number $68 = 72 - 4$ (Doc. 003, Section 0.5) but its derivation as a second-order correction is not yet formally established. **This is an open point:** a geometric derivation of $275/4$ is required. The formula above is a numerically verified result pending full justification.

Issue B: Two inconsistent K_{frak} definitions

Doc. 003 gives:

$$K_{\text{frak}}^{(003)} = 1 - \frac{D_{\text{frak}} - 2}{68} = 1 - \frac{0.94}{68} \approx 0.98530 \quad (190.18)$$

where $D_{\text{frak}} = 2.94$ is stated without derivation from ξ .

Doc. 133 gives:

$$K_{\text{frak}}^{(133)} = 1 - 100\xi = \frac{74}{75} \approx 0.98667 \quad (190.19)$$

derived from the cumulative RG run with $D_f = 3 - \xi$.

These differ by 1.37×10^{-3} and cannot both be correct. The **primary definition** is Doc. 133: $K_{\text{frak}} = 1 - 100\xi = 74/75$. The value $D_{\text{frak}} = 2.94$ in Doc. 003 is an independent approximation not derived from ξ ; it uses a different (tetrahedral) model. Doc. 003 should be understood as a heuristic motivation, not as the defining formula.

Corrected claim for T_{CMB}

- **Avoid:** " K_{frak} closes the T_{CMB} gap."
- **Use:** " T_{CMB} requires a second-order correction $(1 - 275/4 \cdot \xi)$, giving 0.0002 % agreement. The geometric origin of $275/4$ is an open derivation."

Addendum: $\Delta\text{CHSH} \approx 10^{-5}$ in Doc. 230

Doc. 230 states $\Delta\text{CHSH} \approx 10^{-5}$ as a geometrically motivated prediction. Doc. 022 gives the formula

$$\text{CHSH}(N) = 2\sqrt{2} \cdot \exp\left(-\xi \frac{\ln N}{D_f}\right) \quad (190.20)$$

where N is the number of measurement runs. This formula gives $\Delta\text{CHSH} \approx 5 \times 10^{-4}$ at $N = 73$ (not 10^{-5}). Moreover, ΔCHSH grows monotonically with N ; at $N = 2$ (the minimum) one already obtains $\Delta\text{CHSH} \approx 8.7 \times 10^{-5}$. The value 10^{-5} from Doc. 230 is not reachable at any finite N under the Doc. 022 formula.

Key distinction (raised by Onur Teker, May 2026): $\text{CHSH}(N) \rightarrow 0$ (correlation vanishing) requires $N \rightarrow \infty$ and is a different condition from $\Delta\text{CHSH} = 10^{-5}$ (distance from the Tsirelson bound). Both conditions confirm that the Doc. 022 formula cannot reproduce the Doc. 230 claim. The $\Delta\text{CHSH} \approx 10^{-5}$ prediction requires a separate derivation or suppression mechanism not yet documented. **Open point.**

.15 Refinement 12: Two distinct ΔCHSH observables in Doc. 022 and Doc. 230

Affected Documents

Doc. 022 (QFT-ML Addendum), Doc. 230 (Hilbert Space Translation).

The apparent contradiction

Doc. 022 gives $\Delta\text{CHSH} \approx 5 \times 10^{-4}$ at $N = 73$ via the formula $\text{CHSH}(N) = 2\sqrt{2} \cdot \exp(-\xi \ln N / D_f)$. Doc. 230 states $\Delta\text{CHSH} \approx 10^{-5}$ without derivation. These differ by a factor of ~ 50 .

Proposed resolution: cumulative vs. elementary deviation

The two values refer to different observables:

- **Doc. 022** describes the *cumulative* deviation accumulated over N measurement runs. It grows monotonically with N : $\Delta\text{CHSH}(N) \sim \xi \ln(N)/D_f \times 2\sqrt{2}$. This is the experimentally relevant quantity for a Bell test with N runs.
- **Doc. 230** plausibly refers to the *elementary* deviation per single measurement — the phase cost of one θ_4 projection cycle. The natural candidate is

$$\Delta\text{CHSH}_{\text{elem}} = \frac{\xi}{2\pi} \approx 2.1 \times 10^{-5} \approx 10^{-5} \quad (190.21)$$

This is the Landauer cost per measurement expressed as a CHSH deviation: the θ_4 phase runs through a full 2π cycle per measurement, and each cycle costs ξ in geometric units.

The ratio between cumulative and elementary deviation at N runs is:

$$\frac{\Delta\text{CHSH}(N)}{\Delta\text{CHSH}_{\text{elem}}} \approx N \cdot \frac{2\pi}{D_f} \quad (190.22)$$

At $N = 73$ this gives a factor of ≈ 153 , consistent with the observed factor of ~ 50 – 100 between the two values.

Status

This resolution is a **working hypothesis**, not a proven derivation. The identification $\Delta\text{CHSH}_{\text{elem}} = \xi/(2\pi)$ requires formal justification from the T^4 projection geometry. If confirmed, there is no contradiction between Doc. 022 and Doc. 230 — they describe the same physical effect at different integration scales.

Experimental consequence: the cumulative deviation $\Delta\text{CHSH}(N) \approx 5 \times 10^{-4}$ at $N = 73$ is within the reach of next-generation loophole-free Bell tests (current resolution $\sim 10^{-2}$; required improvement: factor ~ 20). The elementary deviation $\xi/(2\pi) \approx 2 \times 10^{-5}$ would require single-shot precision beyond current technology.

.16 Refinement 13: Bridge formula between ξ_{FFGFT} and ξ_{UIFT}

Context

In the IPI discussion (May 2026), Onur Teker (UIFT) writes:

$\xi = k_B T_{\text{CMB}} \ln 2 / (m_e c^2)$, derived from three independent measurements (T_{CMB} , k_B , m_e). This is not a hyperparameter — it is a prediction.

This appears to identify ξ_{UIFT} with $\xi_{\text{FFGFT}} = 4/30000$. Numerically they are very different. This refinement documents the exact relation.

The key: FFGFT uses eV as temperature unit

In FFGFT, the natural unit system sets $\hbar = c = k_B = 1$. Temperatures are expressed in eV (not in Kelvin):

$$T_{\text{CMB}}[\text{eV}] = k_B \cdot T_{\text{CMB}}[\text{K}] = k_B \cdot 2.72548 \text{ K} = 2.3486 \times 10^{-4} \text{ eV} \quad (190.23)$$

The FFGFT formula (Doc. 025) then reads — to first order:

$$T_{\text{CMB}}[\text{eV}] = \frac{16}{9} \xi_{\text{FFGFT}} = 2.3704 \times 10^{-4} \text{ eV} \quad (\Delta = 0.93 \%) \quad (190.24)$$

and to second order (Doc. 190 R11):

$$T_{\text{CMB}}[\text{eV}] = \frac{16}{9} \xi_{\text{FFGFT}} \left(1 - \frac{275}{4} \xi_{\text{FFGFT}} \right) = 2.3486 \times 10^{-4} \text{ eV} \quad (\Delta = 0.0002 \%) \quad (190.25)$$

Bridge formula

Substituting (190.24) into Onur's expression (first order):

$$\begin{aligned} \xi_{\text{UIFT}} &= \frac{k_B T_{\text{CMB}}[\text{K}] \ln 2}{m_e c^2} = \frac{T_{\text{CMB}}[\text{eV}] \cdot \ln 2}{m_e c^2 / \text{eV}} \\ &= \frac{\frac{16}{9} \xi_{\text{FFGFT}} \cdot \ln 2}{m_e c^2 / \text{eV}} \end{aligned} \quad (190.26)$$

Rearranging:

$$\frac{\xi_{\text{FFGFT}}}{\xi_{\text{UIFT}}} = \frac{m_e c^2 / \text{eV}}{\frac{16}{9} \cdot \ln 2} = \frac{510\,999 \text{ eV}}{1.2323 \text{ eV}} \approx 414\,684 \quad (190.27)$$

SI conversion table (reconstructed from Doc. 013, archived version)

The following table was documented in older versions of Doc. 013 (SI conversion tables for the FFGFT natural unit system, subsequently condensed). It is reproduced here because it is essential for the bridge formula derivation.

Quantity	Natural units	SI / eV
Temperature unit [T]	= [E]	1 eV = $k_B^{-1} \cdot 1 \text{ eV} = 11,604 \text{ K}$
T_{CMB} [nat.]	$(16/9) \cdot \xi = 2.370 \times 10^{-4}$	$2.370 \times 10^{-4} \text{ eV}$
T_{CMB} [K] (Planck 2018)	—	2.72548 K
T_{CMB} [eV] = $k_B \cdot T[\text{K}]$	—	$2.3486 \times 10^{-4} \text{ eV}$
$m_e c^2$	= m_e (nat.)	510,999 eV = 0.511 MeV
k_B	= 1 (nat.)	$8.617 \times 10^{-5} \text{ eV/K}$
$\hbar$	= 1 (nat.)	$6.582 \times 10^{-16} \text{ eV}\cdot\text{s}$
c	= 1 (nat.)	$2.998 \times 10^8 \text{ m/s}$

Numerical verification

Quantity	Value	Unit
$\xi_{\text{FFGFT}} = 4/30000$	1.3333×10^{-4}	dimensionless
$\xi_{\text{UIFT}} = k_B T_{\text{CMB}} \ln 2 / (m_e c^2)$	3.1858×10^{-10}	dimensionless
Ratio $\xi_{\text{FFGFT}} / \xi_{\text{UIFT}}$	418 521	—
Scale factor $m_e c^2 / ((\frac{16}{9} \ln 2))$ [eV]	414 684	—
T_{CMB} [K] (Planck 2018)	2.72548	K
T_{CMB} [eV] $= k_B \cdot T_{\text{CMB}}[\text{K}]$	2.3486×10^{-4}	eV
$(16/9) \cdot \xi$ [eV] (FFGFT 1st order)	2.3704×10^{-4}	eV
$(16/9) \cdot \xi(1 - 275/4 \cdot \xi)$ [eV] (R11)	2.3486×10^{-4}	eV
$m_e c^2$	510 999	eV

The residual difference of 0.93 % between the ratio (418 521) and the scale factor (414 684) is exactly the first-order T_{CMB} error — confirming that the bridge formula is correct.

Physical interpretation

ξ_{FFGFT} and ξ_{UIFT} are **not the same quantity**. They are related by a pure SI conversion factor: the ratio of the relativistic electron energy ($m_e c^2 = 511 \text{ keV}$) to the eV scale of the FFGFT temperature formula, modulated by $(16/9) \cdot \ln 2$.

The SI conversion table above was present in earlier versions of Doc. 013 (archived March 2026, Git commit f2be9c8) and was subsequently condensed. The explicit statement $T_{\text{CMB}}^{\text{nat}} = k_B \times 2.725 \text{ K} = 2.35 \times 10^{-4} \text{ eV}$ appears verbatim in that version. The reconstruction here restores the connection between the natural-unit formula and the SI value that was implicitly assumed but no longer shown.

The Vopson-Teker experiment measures ξ_{UIFT} directly. If confirmed, it does not by itself confirm $\xi_{\text{FFGFT}} = 4/30000$ — it confirms the thermodynamic ground state at T_{CMB} . The geometric derivation of ξ_{FFGFT} from $\kappa = 7$ (Doc. 009) is a separate and independent constraint.

.17 Refinement 14: Redshift in Doc. 041 – mechanism, not energy loss

Affected document

Doc. 041 (parameter derivation), table “Cosmological Parameters”, LEVEL 3 (Redshift) – in particular the description “Photon energy loss through ξ -field”.

Previous formulation

The cosmological redshift is labelled in the comparison table as “Photon energy loss through ξ -field” – i.e. as an energy loss of the photon in the ξ -field.

Refined formulation

The physical mechanism is **fractal path-lengthening**: the geometric stretching of the propagation path in the ξ -field (torsion lattice), as derived authoritatively in Docs. 026, 149

and 182. The dependence on path length d (and on wavelength) is retained and is consistent with fractal path-lengthening – it is *not* a contradiction.

An “energy loss” of the photon is **not an independent mechanism** but an *apparent artefact*: it would arise only if the geometric path-lengthening were mistakenly not assumed. The phrase “photon energy loss” is therefore to be replaced in Doc. 041 by “fractal path-lengthening (geometric path stretching)”; the “energy loss” may at most be named explicitly as the artefact that would result if path-lengthening were omitted.

Reasoning

Static FFGFT cosmology explains the redshift purely geometrically (Doc. 182: “not through intrinsic energy loss of the photon, but through geometric stretching of the propagation path”; tired light does not occur). The comparison table in Doc. 041 compresses this mechanism to “energy loss”, evoking precisely the tired-light connotation FFGFT does not intend. The refinement aligns Doc. 041 with the authoritative derivation (Docs. 026/149/182) without altering the path-length dependence.

Status of integration

Incorporated (as of 2 June 2026). Implemented in Doc. 041 (DE+EN).

.18 Refinement 15: mark cosmological z as a Λ CDM comparison quantity

Affected documents

Doc. 061 (temperature units / CMB), section “Modified Recombination History” and abstract/summary; Doc. 041 (Λ CDM vs. T_0 comparison tables).

Previous formulation

A z -dependent recombination calculation ($z_{\text{rec}}^{T_0} \approx 1089.5$, $T(z) = T_0(1+z)[\dots]$) was stated as “In T_0 theory, recombination occurs at ...” – i.e. as an FFGFT-intrinsic quantity. Likewise “CMB at $z \approx 1100$ ” in the abstract/summary.

Refined formulation

The static ξ universe has *no* cosmological z in the expansion sense; FFGFT’s redshift is fractal path-lengthening, not a scale factor. Hence z in these calculations is a **Λ CDM pipeline quantity** and must be marked explicitly as a *comparison calculation*, not as an FFGFT result. In FFGFT the CMB arises from stationary thermalization in an infinitely old universe, not from a decoupling at any z . Doc. 061 now frames the section accordingly; the abstract/summary lines are changed to “without a decoupling at any z ”. In Doc. 041 the $z = 1100$ values are marked “(Λ CDM)” (they already sit in the Λ CDM comparison column).

Reasoning

Unmarked comparison calculations create the impression of FFGFT-intrinsic results and violate the standing caution that cosmological pipeline quantities (H_0 , z , Λ_{obs}) must not be

carried as FFGFT statements without labelling. The marking restores the distinction without removing the comparison numbers.

Status of integration

Incorporated (as of 2 June 2026). Docs. 061 and 041 (DE+EN) marked. Further unmarked pipeline comparisons across the corpus are to be checked separately (ongoing).

.19 Refinement 16: H_0 is emergent (decompactified), not fundamental

Affected documents

Doc. 026 (Geometric Cosmology), l. 116/117 DE+EN; with residues in Doc. 064 (l. 104) and Doc. 181 (l. 158). *Already implemented correctly in Doc. 263* (H_0 as an emergent, decompactified quantity); the remaining places are only *flagged* here, not yet changed.

Previous formulation (outdated)

Doc. 026 calls H_0 a “fundamental constant describing the interaction of light with the geometry of the T0 vacuum.” This wording stems from an earlier stage that still understood redshift as an *energy loss* of light passing through the vacuum (a “tired-light”-like picture). Doc. 064 (“measures energy loss in the ξ -field”) and Doc. 181 (“geometric energy loss”) carry the same residue.

Refined view

Two independent reasons argue against treating H_0 as a fundamental constant:

- **Energy loss is outdated (cf. P14/R14):** redshift is fractal path-lengthening, not energy loss. The “vacuum as a traversed medium” is itself already a decompactified description and not part of the compactified T^4 view.
- **H_0 is emergent, not fundamental:** Doc. 016 records that age, critical density and Hubble length are *Friedmann concepts without a direct T0 counterpart*; Doc. 262, that $\hbar, \varepsilon_0$ become *real only upon decompactification*. Since $H_0 = E_H/\hbar$ contains the (only-then-real) $\hbar$, H_0 does not exist in the compactified T^4 form. H_0 is an emergent quantity of the decompactified description; the non-integer exponent $41/4$ expresses this scale transition, not a fundamental geometric constant. (The integer ansatz $n = 10$ in Doc. 165 should therefore be treated with caution – it forces an emergent transition quantity into a compact geometry.)

Status of integration

Partially incorporated (as of 2 June 2026). Implemented in Doc. 263 (H_0 emergent/decompactified). **Open:** Doc. 026 (DE+EN, l. 116/117) is an *outdated* document (energy-loss/vacuum picture) and needs a more fundamental revision or a historical marking; likewise the residues in Docs. 064/181. Flagged as a larger task.

.20 Refinement 17: circular/fit-dependent DE-DM relations in Doc. 028

Affected document

Doc. 028 ("7 Questions"), sections "Riddle 4: MOND" and "Riddle 5: Dark Energy and Dark Matter". Flagged only, not yet changed.

Finding (recomputed)

- **DE/DM ratio: circular as a *prediction*, admissible as a *conversion*.** Doc. 028 states $\rho_{\text{DE}}/\rho_{\text{DM}} = \xi^\alpha$ with $\alpha = \ln(2.5)/\ln(\xi)$. Substituting this α into ξ^α yields *exactly* 2.5 for any ξ (checked for $\xi = 10^{-3}, 10^{-5}, 0.5$). As a *prediction* from the ξ -geometry the relation is therefore circular (violates Doc. 101). However: the target value 2.5 ($\Omega_\Lambda/\Omega_{\text{DM}}$) is itself not a model-neutral measurement but a Λ CDM-pipeline output (standing caution). Read *not* as a prediction but as a *conversion factor* between the FFGFT geometry and the Λ CDM Ω bookkeeping – analogous to c (metre/second) and to z (cf. R15) – the circular calibration via 2.5 is *admissible*. **Restriction:** unlike c (both sides share the same real counterpart), the Λ CDM side carries with ρ_{DE} a model artefact (Λ does not exist in FFGFT). The factor thus translates partly a real quantity (the DM bookkeeping $\leftrightarrow \xi$ effect) and partly an artefact (ρ_{DE}). Hence: admissible, but only *partially* meaningful – as far as the Λ CDM quantity is artefact-free.
- **The MOND scale is fit-dependent.** $a_0/(cH_0) = \xi^{1/4} \cdot K_M$ matches the measured 0.176 only with $K_M = 1.637$. The $\xi^{1/4} = 0.1075$ is genuine; K_M is a *free fit factor* without derivation. Partially derived, not parameter-free.
- **Two incompatible a_0 formulas; rotation curve open.** Besides Doc. 028, Doc. 201 (§4.3) gives a *different* form: $a_0 = c^2\xi/(4\lambda) \approx 1.2 \times 10^{-10} \text{ m/s}^2$. It is *not computable*: for the correct unit (m/s^2) λ would have to be a *length* ($\sim 2.5 \times 10^{22} \text{ m}$), whereas elsewhere in Doc. 201 λ is, via $m_T = \lambda/\xi$, an *energy/mass* – the same symbol in two incompatible meanings. Without an independent λ , a_0 is either undetermined or (back-solved from the measured value) circular; the length $2.5 \times 10^{22} \text{ m}$ matches no known FFGFT scale. *Consequence*: both a_0 formulas (028: fit-dependent; 201: dimensionally inconsistent) fail to carry. The *flat rotation curve* follows qualitatively as a MOND consequence of the ξ geometry (static, no DM substance; Doc. 201/145/156), but the actual profile $v(r)$ and the SPARC comparison are *not computed* – remains open.

Delimitation (what remains valid)

The CMB relations are unaffected and remain robust: $T_{\text{CMB}} = \frac{16}{9}\xi^2 E_\xi$ and $|\rho_{\text{Casimir}}|/\rho_{\text{CMB}} = \pi^2/(240\xi) \approx 308$ contain ξ alone, with no fitted exponent or factor.

Status of integration

Flagged (as of 2 June 2026), not yet changed. Recommendation: the DE/DM ratio (Riddle 5) should be marked *not* as a prediction but as a restricted conversion factor to the Λ CDM Ω bookkeeping (admissible like c/z , but bounded by the ρ_{DE} artefact); the MOND scale (Riddle 4) as partially derived with a free K_M . The *interpretation* of the dark sector is settled – DM is a ξ -geometric effect (artefact of the Λ CDM view, like H_0 and z), DE is eliminated (no Λ). *Quantitatively* open: no model-neutral derived Ω_{DM} , no computed rotation curve.

.21 Refinement 14: CMB anharmonicity, forward vs. retrodiction, and the $|n|^2 = 30$ case (R29/R30/R31)

An explicit forward investigation (Doc. 268, Step 17; scripts Dok267_268_Skripte/) has clarified the status of the CMB peak sector and split it into three points.

(1) The direction of inference (R29). The earlier presentation suggested that $\{1, 6, 14, 26\}$ and hence $\{3, 18, 42, 78\}$ followed forward from the T^4 geometry. In fact $\{1, 6, 14, 26\}$ is obtained by squaring the *measured* CMB ratios (Doc. 268, Step 4). It is a **retrodiction** (consistency check), not a parameter-free prediction. The genuine forward maxima of the full T^4 spectrum are dense ($|n|^2 = 3, 6, 10, 14, 18, 22, 26, 30, \dots$).

(2) The $|n|^2 = 30$ case (R29, sharpened by O. Teker). $30 = 3 \cdot 10$ satisfies the factor-3 filter, since $|n|^2 = 10$ is a local maximum. Thermally one even has

$$I(30) = 13.2 > I(42) = 7.6 > I(78) = 1.7,$$

so 30 ($\ell \approx 696$) would be *stronger* than two visible peaks, yet shows no CMB peak (it sits in the trough between peaks 2 and 3). The earlier “strongest local maximum in range” argument fails here. The selection mechanism is thus *not* supplied by the factor 3 alone. *Open.*

(3) The naive C_ℓ Bessel projection is insufficient (R30). The computation $C_\ell = \sum_n N_{BE} |j_\ell(k_{obs} D_A)|^2$ does *not* reproduce $\{3, 18, 42, 78\}$; it washes out to a quasi-continuum dominated by the lowest modes. The earlier R30 wording (“a feasible computation, no fundamental obstacle”) is withdrawn: a physical source/window function is missing. *Open.*

(4) The forward content – geometric anharmonicity (new, R31). Forward, the symmetric T^4 resonance produces the *harmonic* backbone $1 : 2 : 3 : 4 : 5$ ($k_{obs}^2 = 3n^2$). The CMB is *anharmonic* ($1 : 2.44 : 3.68 : 5.09$). This anharmonicity follows forward and parameter-free from the codimension-1 projection:

$$D_{eff} = D_f - 1 = 2 - \xi \approx 2 \Rightarrow \nu = \frac{D_{eff}}{2} - 1 = -\frac{\xi}{2} \approx 0 \Rightarrow J_0 \text{ membrane} \Rightarrow 1 : 2.30 : 3.60 : 4.90,$$

matching to $\sim 5\%$, *without* mode selection. The radial Bessel index is $\nu = D/2 - 1$; anharmonicity is maximal at $D = 2$ and vanishes at $D = 1$ and $D = 3$. The CMB sits at the dimension of maximal anharmonicity. The match is to the *flat* membrane (J_0), not the curved 2-sphere ($Y_{\ell m}$, which gives $1 : 1.73 : \dots$, wrong). This is the conceptually clean, forward statement; it removes the circularity objection because no selection from data is needed.

(5) The residual – not from ξ . The best fit lies at $D \approx 1.86$ (just below 2). A spectral dimension $d_s < D_f$ would be the obvious source but fails: the FFGFT fractal is too weakly fractal ($d_w = 2 + O(\xi) \Rightarrow d_s \approx 2$); for $d_s = 1.86$ one would need $d_w \approx 2.15$ ($\sim 1000 \times \xi$). The deviation is not derivable from ξ and is of the order of the measurement uncertainty (J_0 is consistent with peaks 3,4 within $\sim 2\sigma$; only peak 2 at $\sim 4\sigma$). Pointing to an exact value would be numerology. *Open, likely partly measurement noise.*

Consequence. The defensible core reads: harmonic T^4 backbone (forward) + geometric anharmonicity from the codimension-1 projection (forward, $\sim 5\%$). The ratio match $\{3, 18, 42, 78\}$ is retrodiction; the rigorous individual-peak selector and the exclusion of $|n|^2 = 30$ remain open. The particle sector (α , lepton masses, Koide) is *unaffected* – it rests on direct ξ powers, not on mode counting.

Updated Summary

No.	Doc.	Erroneous / unclear	Correct / refined
C1	049 (HTML)	$\xi = \lambda_h^2 v^2 / (64\pi^4 m_h^2)$	$\xi = \lambda_h^2 v^2 / (16\pi^3 m_h^2)$
C2	116 (table)	$r_\tau = 8/3$	$r_\tau = 25/9$
C3	018 (Explorer)	$a_e = 4\pi / (f \cdot k)$	$a_e = 2\pi^2 / (f \cdot k)$
R1	116 / 158	$\Delta Q < 0.00003\%$ vs. 0.001%	Both correct; 0.001% for external use
R2	several	$\hbar$ as a postulate	$\hbar$ follows from ξ via $L_0 = \xi \cdot \ell_P$
R3	173–176	A single ξ value	Two parameters: ξ_0 (geom.) and ξ_{Higgs} (alg.)
R4	180–182	L_0 as Schwarzschild radius	$L_0 = \text{geom. fundamental length}$
R5	several	"FFGFT calculates masses exactly"	Mass structure from $\xi_i \sim 1\%$ agreement
R6	116 / 158	Koide with FFGFT symbolic terms	Koide only with numerical values (non-closure, Doc. 193)
R7	005, 008, 019, 086, 189, 202	"Renormalization group" as FFGFT tool	Scale structure from recursion $\mathcal{R}$
R8	005, 008, 012, 014, 041, 060, 133, 141, 159, 181	"Fractal renormalization" as proprietary term	Geometric scale adjustment / fractal correction
R9	005, 006, 042, 046	Column "Symmetry" with "Color"; $N_c = 3$ as external SM input	Columns "Scaling class" + "SM correspondence"; quarks via $m = r \xi^P v$
R10	009	$K = 245$ derived from $\varphi^{12}/1.314$; "no free parameters"	$K = R_{pe}/(4/3)^7$: one empirical anchor; factor 1.314 implicitly defined, not derived
R11	025, 003, 133	K_{frak} closes T_{CMB} gap; one definition of K_{frak}	Correct: $(1 - 275/4 \cdot \xi)$, $\Delta = 0.0002\%$; Doc. 003 and Doc. 133 use different models
R12	022, 230	$\Delta\text{CHSH} \approx 10^{-5}$ (Doc. 230) vs. 5×10^{-4} at $N = 73$ (Doc. 022)	Two distinct observables: cumulative (Doc. 022) vs. elementary $\xi/(2\pi)$ (Doc. 230); 10^{-5} not reachable at any finite N from Doc. 022
R13	013, 025, 061	$\xi_{\text{FFGFT}} = \xi_{\text{UIFT}}$; SI conversion table missing	$\xi_{\text{FFGFT}}/\xi_{\text{UIFT}} = m_e c^2 / (\frac{16}{9} \ln 2 \cdot \text{eV}) \approx 414\,684$; table in Doc. 013 (archived) reconstructed
R14	041	Redshift as "photon energy loss through ξ -field"	Fractal path-lengthening (geometric path stretching); energy loss only an artefact if path-lengthening omitted; path-length dependence retained (Docs. 026/149/182); the same path-lengthening underlies the CMB treatment in Doc. 268 (previously listed separately as R19)
R15	061 / 041	Cosmological z ($z \approx 1100$, $z_{\text{rec}} \approx 1089.5$) unmarked as an FFGFT quantity	Marked as a ΛCDM comparison quantity; static universe has no cosmological z ; CMB from stationary thermalization
R16	026 / 064 / 181	H_0 as a "fundamental constant" (energy-loss/vacuum picture)	H_0 is emergent (decompactified), not fundamental; implemented in Doc. 263, Docs. 026/064/181 flagged as outdated
R17	028	DE/DM ratio $\xi^{\ln(2.5)/\ln \xi}$ presented as a prediction; MOND with fit factor K_M	DE/DM relation is circular (yields 2.5 for any ξ); MOND partially derived ($\xi^{1/4}$ genuine, K_M free); CMB relations unaffected; flagged
R18	267	Relation ΛCDM expansion vs. FFGFT time-unfolding treated as an empirical question	Both pictures are <i>not</i> distinguishable through the observed ratios; the decision is theoretical, not empirical (conceptual placement)

continued on next page

(continued)

No.	Doc.	Erroneous / unclear	Correct / refined
R20	267 / 268	Absolute cosmological scale (R_H , D_A , z_*) presented as an FFGFT result	The <i>form</i> is fixed: a dimensionless ξ ratio (R_H/ℓ_p). The <i>factor</i> (exponent $41/4$, $E_H = E_0 \xi^{41/4}$, Doc. 182) is <i>not</i> derived from T^4 geometry and must not be fitted (fitting = retrodiction). FFGFT is not the traditional yardstick; the factor is therefore adopted as a <i>declared external calibration</i> from Λ CDM – a choice of units, not an adoption of its physics – to be inserted with the converged value once the H_0 tension (~ 67 vs. ~ 73) is resolved; parametrised until then. The imported value does <i>not</i> count as FFGFT confirmation. Affects only absolute positions ($\ell_1 \approx 220$, $z_* \approx 875$), not the ratios; the internal forward derivation of the factor remains the goal. <i>Open</i>
R29	268	Selection rule: why does the CMB show only the subset $\{3, 18, 42, 78\}$ of the T^4 peaks?	<i>Only partially answered.</i> The factor 3 from the 3D projection ($3 \cdot \{1, 6, 14, 26\}$) is plausible, but the sequence $\{1, 6, 14, 26\}$ is <i>read off the data</i> (Doc. 268, Step 4), not produced forward. In particular $ n ^2 = 30 = 3 \cdot 10$ is <i>not</i> excluded: $ n ^2 = 10$ is a local maximum, and thermally $I(30) = 13.2 > I(42) = 7.6 > I(78) = 1.7$ – so 30 ($\ell \approx 696$) should be a peak, yet sits in a CMB trough. The “strongest in range” argument fails. <i>Open</i> , sharpened by O. Teker
R30	268	Rigorous justification of the selection mechanism (full C_ℓ projection) missing	The <i>naive</i> spherical Bessel sum $C_\ell = \sum_n N_{BE} j_\ell(k_{\text{obs}} D_A) ^2$ does <i>not</i> reproduce $\{3, 18, 42, 78\}$ – it washes out to a quasi-continuum dominated by low modes. The earlier wording (“a feasible computation, no fundamental obstacle”) was too optimistic: a physical source/window function selecting the peaks is missing. <i>Open</i>
R31	268	Anharmonicity of the CMB peaks: whence the stretch relative to harmonic $1 : 2 : 3 : 4 : 5$?	<i>Leading behaviour derived forward.</i> Forward, T^4 gives the harmonic backbone $1 : 2 : 3 : 4 : 5$ (symmetric resonance $k^2 = 3n^2$). The observed anharmonicity $1 : 2.44 : 3.68 : 5.09$ follows from the codimension-1 projection: last-scattering surface $\Rightarrow D_{\text{eff}} = D_f - 1 = 2 - \xi \approx 2 \Rightarrow$ Bessel index $\nu = -\xi/2 \approx 0 \Rightarrow J_0$ membrane $\Rightarrow 1 : 2.30 : 3.60 : 4.90$, matching to $\sim 5\%$, parameter-free, without mode selection. The small residual ($D \approx 1.86$) is <i>not</i> derivable from ξ (the spectral dimension of the weakly fractal space stays ≈ 2) and is of the order of the measurement uncertainty. <i>Partially derived</i> ; remainder open
R32	corpus-wide	Statements presenting H_0 or the absolute scale as <i>derived by FFGFT</i>	Follows from R20: H_0 is an <i>imported</i> external calibration, not derived. All passages presenting H_0 as an FFGFT result are to be cleaned up (“imported/declared” instead of “derived”). <i>Noted for now</i> ; corpus-wide cleanup deferred. <i>Open</i>

continued on next page

(continued)

No.	Doc.	Erroneous / unclear	Correct / refined
R33	009 / 042 / 181	Magnitude factor 5^4 in $\xi = 1/(3 \cdot 2^2 \cdot 5^4)$ presented as <i>derived</i>	The <i>form</i> of ξ is forward: the harmonic $4/3$ (perfect fourth, Docs. 042/159) gives the leading digit, 3 (space dimensions) and $2^2=4$ (spacetime) are read off the geometry. The <i>magnitude-carrying</i> factor $5^4=625$ ($\rightarrow 10^{-4}$), however, is <i>not</i> derived forward: it is merely <i>asserted</i> ("emerges from the fractal structure", Doc. 009; "encodes the symmetry structure", Doc. 181 – there circular via $30000/4$) or the value is <i>empirical</i> (Doc. 042: measured from the Higgs sector 1.3194×10^{-4} , $\sim 1\%$ off the ideal $4/30000$). It remains the <i>intuitive</i> (harmonic / Euler Tonnetz) resp. empirical grounding of the magnitude; this is openly <i>declared</i> as such, not presented as a forward derivation. Structural parallel to R20 (form fixed, factor open), one level deeper. <i>Declared/noted</i>
R34	181 / 013	$D = 4$ presented as "follows necessarily / not a postulate"; role of the SI conversions for the c -time-power	$3+1$ is <i>empirical input</i> – no theory derives the dimension count; the earlier "follows necessarily" claim (Doc. 181) is overstated. Forward are (i) periodicity (divergence-freedom), (ii) torus-not-sphere ($\pi_1 \neq 0$, stable windings), (iii) <i>minimal sufficiency</i> – 4 instead of a naive 5, because the duality $\tilde{T} \cdot m = 1$ lets one circle carry mass <i>and</i> time, (iv) the lossless (bijective) Hilbert representation (Type III, Docs. 270/206/230). The <i>bare number</i> 4 is <i>not</i> forced forward (the "4 from ξ " argument in Doc. 181 is circular resp. base-dependent). <i>Connection with the units</i> : the four-circle structure is at the same time the exponent book-keeping of the SI conversions – c = space-time bridge (the three T^3 circles against the time direction), $\hbar$ = mass-time bridge ($S_m^1, \tilde{T}m = 1$). Hence the SI factors are indispensable for the <i>power of c</i> and the <i>power of time</i> : $c = \hbar = 1$ collapses L, T, M to one axis and destroys exactly the exponent structure one wants to read off. T^4 (mass reading) and T^3 +time (time reading) are the same object. Signature as R20/R33: structure/form forward, number input. <i>Clarified</i>