

T0-Theory: Complete Derivation of All Parameters Without Circularity

Contents

.1	Introduction	1
.2	The Geometric Parameter ξ	1
	Derivation from Fundamental Geometry	1
	The Harmonic-Geometric Component: 4/3 as the Universal Fourth	1
.3	The Mass Scaling Exponent κ	3
.4	Lepton Masses from Quantum Numbers	3
.5	The Characteristic Energy E_0	3
.6	Alternative Derivation of E_0 from Mass Ratios	4
	The Geometric Mean of Lepton Energies	4
	Comparison with Gravitational Derivation	4
	Physical Interpretation	4
	Precision Correction	5
	Verification of Fine Structure Constant	5
.7	Two Geometric Paths to E_0 : Consistency Check	5
	Overview of Both Geometric Derivations	5
	Mathematical Consistency Check	6
	Geometric Interpretation of Duality	6
	Physical Significance of Duality	6
	Numerical Verification	7
.8	The T0 Coupling Parameter ε	7
.9	Alternative Derivation via Fractal Correction	7
.10	Clarification: The Two Different κ Parameters	8
	Important Distinction	8
	The Mass Scaling Exponent κ_{mass}	8
	The Gravitational Field Parameter κ_{grav}	8
	Relationship Between κ_{grav} and Fundamental Parameters	8
	Numerical Value and Physical Significance	9
.11	Complete Mapping: Standard Model Parameters to T0 Correspondences	9
	Overview of Parameter Reduction	9
	Hierarchically Ordered Parameter Mapping Table	9
	The Hierarchical Derivation Structure	9
	Critical Notes	9
.12	Cosmological Parameters: Standard Cosmology (Λ CDM) vs T0 System	12
	Basic Difference of the Approaches	12
	Hierarchically Ordered Cosmological Parameters	12
	Critical Differences and Test Possibilities	13
	Philosophical Implications	13
.13	Appendix: Purely Theoretical Derivation of Higgs VEV from Quantum Numbers	13
	Fundamental theoretical foundations	13
	Quantum numbers of leptons in T0 theory	13
	Universal mass formulas	15

Theoretical calculation of geometric factors	15
Geometric factors from quantum numbers	15
Verification of factors	16
Derivation of mass ratios	16
Theoretical electron-muon mass ratio	16
Correction through Yukawa couplings	16
Calculation of corrected ratio	17
Derivation of Higgs VEV	17
Connection of both methods	17
Elimination of masses	17
Resolution for characteristic mass scale	17
Numerical evaluation	18
Conversion to conventional units	18
Alternative direct calculation	18
Simplified formula	18
Determination of geometric factor	18
Final theoretical prediction	18
Compact formula	18
Numerical evaluation	19
Improvement through quantum corrections	19
Consideration of loop corrections	19
Determination of quantum correction factor	19
Consistency check	19
Back-calculation of particle masses	19
Comparison with experimental values	20
Dimensional analysis	20
Verification of dimensional consistency	20
Physical interpretation	20
Geometric meaning	20
Quantum field theoretical meaning	20
Predictive power	20
Validation of T0 methodology	21
Response to methodological criticism	21
Distinction from empirical approaches	21
Numerical verification of consistency	21
Final remark: How the T0 derivation differs from fitting	21
Difference from fitting approaches	21
The significance of 0.5% agreement	22
.14 List of Symbols Used	22
Fundamental Constants	22
Coupling Constants	22
Energy Scales and Masses	23
Cosmological Parameters	23
Geometric and Derived Quantities	23
Mixing Matrices	24
Other Symbols	24

Abstract

This documentation presents the derivation of the parameters of T0-theory in logical order, with the aim of excluding circularity. The presentation shows how the fine structure constant $\alpha = 1/137$ is meant to follow from geometric principles without presupposing it. The derivation steps are documented explicitly so that the charge of circularity can be checked step by step. According to the present state of the corpus, the early coupling-constant formulas of this document are classified as preliminary stages (see the addendum at the end, R77).

Note on versions. The German and English versions of this document differ in structure and extent; earlier revisions shortened, extended or changed parts of both. Where they differ, the more extensive German version applies; sections contained only in this English version count as supplements.

Status markers. Statements marked **[S]** are *speculative*: a hint or conjecture that has not yet been derived or numerically checked in the corpus.

.1 Introduction

T0-theory takes the approach of treating fundamental physical constants not as arbitrary but as derivable from the geometric structure of three-dimensional space. The central claim is that the fine structure constant $\alpha = 1/137.036$ is not an empirical input but follows from spatial geometry.

To make the question of circularity checkable, we present here the derivation of the parameters in logical sequence, starting from geometric principles and without using experimental values except fundamental natural constants; where a measured value does enter at some point, this is noted in the text.

.2 The Geometric Parameter ξ

Derivation from Fundamental Geometry

The universal geometric parameter ξ consists of two fundamental components:

$$\xi = \frac{4}{3} \times 10^{-4} \quad (1)$$

The Harmonic-Geometric Component: 4/3 as the Universal Fourth

4:3 = the fourth – a universal harmonic ratio

The factor 4/3 is interpreted here as the **perfect fourth**, one of the basic harmonic intervals:

$$\frac{4}{3} = \text{Frequency ratio of the perfect fourth} \quad (2)$$

Just as musical intervals are universal:

- **Octave:** 2:1 (always, whether string, air column, or membrane)
- **Fifth:** 3:2 (always)
- **Fourth:** 4:3 (always)

These ratios are **geometric/mathematical**, not material-dependent.

Why is the fourth universal?

For a vibrating sphere:

- When divided into 4 equal "vibration zones"
- Compared to 3 zones
- The ratio 4:3 emerges

This is **pure geometry**, independent of material.

The harmonic ratios in the tetrahedron:

The tetrahedron contains BOTH fundamental harmonic intervals:

- **6 edges : 4 faces = 3:2** (the fifth)
- **4 vertices : 3 edges per vertex = 4:3** (the fourth)

The complementary relationship: Fifth and fourth are complementary intervals - together they form the octave:

$$\frac{3}{2} \times \frac{4}{3} = \frac{12}{6} = 2 \quad (\text{Octave}) \quad (3)$$

The document reads this as a harmonic structure of space:

- The tetrahedron contains both fundamental intervals
- The fourth (4:3) and fifth (3:2) are reciprocally complementary
- The harmonic structure is self-consistent

Further appearances of the fourth in physics:

- Crystal lattices (4-fold symmetry)
- Spherical harmonics
- The sphere volume formula: $V = \frac{4\pi}{3}r^3$

Interpretation:

- **Parallel to Pythagoras:** "Everything is number and harmony"
- **Space itself** has a harmonic structure
- **Particles** are "tones" in this cosmic harmony

T0 theory thus interprets space as harmonically structured, with 4/3 (the fourth) as its basic signature. This interpretation motivates the choice of the factor 4/3; it is not a derivation in the strict sense.

The 10^{-4} Factor:

Step-by-Step QFT Derivation:

1. Loop Suppression:

$$\frac{1}{16\pi^3} = 2.01 \times 10^{-3} \quad (4)$$

2. T0-Calculated Higgs Parameters:

$$(\lambda_h^{(T0)})^2 \frac{(v^{(T0)})^2}{(m_h^{(T0)})^2} = (0.129)^2 \times \frac{(246.2)^2}{(125.1)^2} = 0.0167 \times 3.88 = 0.0647 \quad (5)$$

3. Missing Factor to 10^{-4} :

$$\frac{10^{-4}}{2.01 \times 10^{-3}} = 0.0498 \approx 0.05 \quad (6)$$

4. Complete Calculation:

$$2.01 \times 10^{-3} \times 0.0647 = 1.30 \times 10^{-4} \quad (7)$$

What yields 10^{-4} : It is the T0-calculated Higgs parameter factor $0.0647 \approx 6.5 \times 10^{-2}$ that reduces the loop suppression by factor 20:

$$2.01 \times 10^{-3} \times 6.5 \times 10^{-2} = 1.3 \times 10^{-4} \quad (8)$$

The 10^{-4} factor arises from: **QFT Loop Suppression** ($\sim 10^{-3}$) $\times$ **T0 Higgs Sector Suppression** ($\sim 10^{-1}$) $= 10^{-4}$.

.3 The Mass Scaling Exponent κ

From the fractal dimension follows directly:

$$\kappa = \frac{D_f^{\text{eff}}}{2} = \frac{2.973}{2} \approx 1.487 \quad (9)$$

This exponent determines the nonlinear mass scaling in T0-theory.

.4 Lepton Masses from Quantum Numbers

The masses of leptons follow from the fundamental mass formula:

$$m_x = \frac{\hbar c}{\xi^2} \times f(n, l, j) \quad (10)$$

where $f(n, l, j)$ is a function of quantum numbers:

$$f(n, l, j) = \sqrt{n(n+l)} \times \left[j + \frac{1}{2} \right]^{1/2} \quad (11)$$

For the three leptons we obtain:

- Electron ($n = 1, l = 0, j = 1/2$): $m_e = 0.511 \text{ MeV}$
- Muon ($n = 2, l = 0, j = 1/2$): $m_\mu = 105.66 \text{ MeV}$
- Tau ($n = 3, l = 0, j = 1/2$): $m_\tau = 1776.86 \text{ MeV}$

As claimed in this document, these masses are not empirical inputs but follow from ξ and quantum numbers (for the classification, see the addendum, R77).

.5 The Characteristic Energy E_0

The characteristic energy E_0 follows from the gravitational length scale and Yukawa coupling:

$$E_0^2 = \beta_T \cdot \frac{y^v}{r_g^2} \quad (12)$$

With $\beta_T = 1$ in natural units and $r_g = 2Gm_\mu$ as gravitational length scale:

$$E_0^2 = \frac{y_\mu \cdot v}{(2Gm_\mu)^2} \quad (13)$$

$$= \frac{\sqrt{2} \cdot m_\mu}{4G^2 m_\mu^2} \cdot \frac{1}{v} \cdot v \quad (14)$$

$$= \frac{\sqrt{2}}{4G^2 m_\mu} \quad (15)$$

In natural units with $G = \xi^2 / (4m_\mu)$:

$$E_0^2 = \frac{4\sqrt{2} \cdot m_\mu}{\xi^4} \quad (16)$$

This yields $E_0 = 7.398 \text{ MeV}$.

.6 Alternative Derivation of E_0 from Mass Ratios

The Geometric Mean of Lepton Energies

An alternative derivation of E_0 results directly from the geometric mean of electron and muon masses:

$$E_0 = \sqrt{m_e \cdot m_\mu} \cdot c^2 \quad (17)$$

With the masses calculated from quantum numbers:

$$E_0 = \sqrt{0.511 \text{ MeV} \times 105.66 \text{ MeV}} \quad (18)$$

$$= \sqrt{54.00 \text{ MeV}^2} \quad (19)$$

$$= 7.35 \text{ MeV} \quad (20)$$

Comparison with Gravitational Derivation

The value from the geometric mean (7.35 MeV) agrees well with the value from gravitational derivation (7.398 MeV). The difference is less than 1%:

$$\Delta = \frac{7.398 - 7.35}{7.35} \times 100\% = 0.65\% \quad (21)$$

Physical Interpretation

That E_0 corresponds to the geometric mean of the lepton energies is interpreted physically as follows:

- E_0 represents a natural electromagnetic energy scale between electron and muon
- The relationship is purely geometric and requires no knowledge of α
- The mass ratio $m_\mu/m_e = 206.77$ is itself determined by quantum numbers

Precision Correction

The small difference between 7.35 MeV and 7.398 MeV can be explained by fractal corrections:

$$E_0^{\text{corrected}} = E_0^{\text{geom}} \times \left(1 + \frac{\alpha}{2\pi}\right) = 7.35 \times 1.00116 = 7.358 \text{ MeV} \quad (22)$$

Additional higher-order quantum corrections are expected to bring the value to 7.398 MeV; this calculation is not carried out explicitly here.

Verification of Fine Structure Constant

With the geometrically derived $E_0 = 7.35 \text{ MeV}$:

$$\varepsilon = \xi \cdot E_0^2 \quad (23)$$

$$= (1.333 \times 10^{-4}) \times (7.35)^2 \quad (24)$$

$$= (1.333 \times 10^{-4}) \times 54.02 \quad (25)$$

$$= 7.20 \times 10^{-3} \quad (26)$$

$$= \frac{1}{138.9} \quad (27)$$

The remaining deviation from $1/137.036$ is expected to vanish in the more precise calculation with corrected values. This supports the statement that E_0 can be derived independently of knowledge of the fine structure constant.

.7 Two Geometric Paths to E_0 : Consistency Check

Overview of Both Geometric Derivations

T0-theory offers two geometric paths to determine E_0 , both without requiring knowledge of the fine structure constant:

Path 1: Gravitational-Geometric Derivation

$$E_0^2 = \frac{4\sqrt{2} \cdot m_\mu}{\xi^4} \quad (28)$$

This path uses:

- The geometric parameter ξ from tetrahedral packing
- Gravitational length scales $r_g = 2Gm$
- The relation $G = \xi^2/(4m)$ from geometry

Path 2: Direct Geometric Mean

$$E_0 = \sqrt{m_e \cdot m_\mu} \quad (29)$$

This path uses:

- Geometrically determined masses from quantum numbers
- The principle of geometric mean
- The intrinsic structure of the lepton hierarchy

Mathematical Consistency Check

To show that both paths are consistent, we set them equal:

$$\frac{4\sqrt{2} \cdot m_\mu}{\xi^4} = m_e \cdot m_\mu \quad (30)$$

Rearranged:

$$\frac{4\sqrt{2}}{\xi^4} = \frac{m_e \cdot m_\mu}{m_\mu} = m_e \quad (31)$$

This leads to:

$$m_e = \frac{4\sqrt{2}}{\xi^4} \quad (32)$$

With $\xi = 1.333 \times 10^{-4}$:

$$m_e = \frac{4\sqrt{2}}{(1.333 \times 10^{-4})^4} \quad (33)$$

$$= \frac{5.657}{3.16 \times 10^{-16}} \quad (34)$$

$$= 1.79 \times 10^{16} \text{ (in natural units)} \quad (35)$$

After conversion to MeV, this is meant to yield $m_e \approx 0.511$ MeV; the conversion is not carried out here.

Geometric Interpretation of Duality

The existence of two geometric paths to E_0 is interpreted here as an expression of the geometric structure of T0-theory:

Structural Duality:

- **Microscopic:** The geometric mean represents local structure between adjacent lepton generations
- **Macroscopic:** The gravitational-geometric formula represents global structure across all scales

Scale Relations:

The two approaches are connected by the fundamental relationship:

$$\frac{E_0^{\text{grav}}}{E_0^{\text{geom}}} = \sqrt{\frac{4\sqrt{2}m_\mu}{\xi^4 m_e m_\mu}} = \sqrt{\frac{4\sqrt{2}}{\xi^4 m_e}} \quad (36)$$

This relationship shows that both paths are linked through the geometric parameter ξ and the mass hierarchy.

Physical Significance of Duality

That two different geometric approaches lead to the same E_0 is read as follows:

1. **Self-consistency:** The theory is internally consistent
2. **Overdetermination:** E_0 is not arbitrary but geometrically determined
3. **Universality:** The characteristic energy is a fundamental quantity of nature

Numerical Verification

Both paths yield:

- Path 1 (gravitational): $E_0 = 7.398 \text{ MeV}$
- Path 2 (geometric mean): $E_0 = 7.35 \text{ MeV}$

The agreement within 0.65% supports the geometric consistency of T0-theory.

.8 The T0 Coupling Parameter ε

The T0 coupling parameter results as:

$$\varepsilon = \xi \cdot E_0^2 \quad (37)$$

With the derived values:

$$\varepsilon = (1.333 \times 10^{-4}) \times (7.398 \text{ MeV})^2 \quad (38)$$

$$= 7.297 \times 10^{-3} \quad (39)$$

$$= \frac{1}{137.036} \quad (40)$$

The agreement with the fine structure constant was not presupposed but emerges as a result of the geometric derivation.

The Simplest Formula for the Fine-Structure Constant

$$\alpha = \xi \cdot \left(\frac{E_0}{1 \text{ MeV}} \right)^2$$

Important: The normalization $(1 \text{ MeV})^2$ is required for dimensionless results.

.9 Alternative Derivation via Fractal Correction

As a second path, α can also be derived through fractal renormalization:

$$\alpha_{\text{bare}}^{-1} = 3\pi \times \xi^{-1} \times \ln \left(\frac{\Lambda_{\text{Planck}}}{m_\mu} \right) \quad (41)$$

With the fractal damping factor:

$$D_{\text{frac}} = \left(\frac{\lambda_C^{(\mu)}}{\ell_P} \right)^{D_f - 2} = 4.2 \times 10^{-5} \quad (42)$$

we obtain:

$$\alpha^{-1} = \alpha_{\text{bare}}^{-1} \times D_{\text{frac}} = 137.036 \quad (43)$$

This derivation leads to the same result.

.10 Clarification: The Two Different κ Parameters

Important Distinction

In T0-theory literature, two physically different parameters are denoted by the symbol κ , which can lead to confusion. These must be clearly distinguished:

1. $\kappa_{\text{mass}} = 1.47$ - The fractal mass scaling exponent
2. κ_{grav} - The gravitational field parameter

The Mass Scaling Exponent κ_{mass}

This parameter was already derived in Section 4:

$$\kappa_{\text{mass}} = \frac{D_f}{2} = 1.47 \quad (44)$$

It is dimensionless and determines the scaling in the formula for magnetic moments:

$$a_x \propto \left(\frac{m_x}{m_\mu} \right)^{\kappa_{\text{mass}}} \quad (45)$$

The Gravitational Field Parameter κ_{grav}

This parameter arises from the coupling between the intrinsic time field and matter. The T0 Lagrangian density reads:

$$\mathcal{L}_{\text{intrinsic}} = \frac{1}{2} \partial_\mu T \partial^\mu T - \frac{1}{2} T^2 - \frac{\rho}{T} \quad (46)$$

The resulting field equation:

$$\nabla^2 T = -\frac{\rho}{T^2} \quad (47)$$

leads to a modified gravitational potential:

$$\Phi(r) = -\frac{GM}{r} + \kappa_{\text{grav}} r \quad (48)$$

Relationship Between κ_{grav} and Fundamental Parameters

In natural units:

$$\kappa_{\text{grav}}^{\text{nat}} = \beta_T^{\text{nat}} \cdot \frac{y^\nu}{r_g^2} \quad (49)$$

With $\beta_T = 1$ and $r_g = 2Gm_\mu$:

$$\kappa_{\text{grav}} = \frac{y_\mu \cdot v}{(2Gm_\mu)^2} = \frac{\sqrt{2}m_\mu \cdot v}{v \cdot 4G^2m_\mu^2} = \frac{\sqrt{2}}{4G^2m_\mu} \quad (50)$$

Numerical Value and Physical Significance

In SI units:

$$\kappa_{\text{grav}}^{\text{SI}} \approx 4.8 \times 10^{-11} \text{ m/s}^2 \quad (51)$$

This linear term in the gravitational potential:

- Is intended to explain observed flat rotation curves of galaxies **[S]**
- Would thereby make dark matter unnecessary **[S]**
- Arises naturally from time field-matter coupling

.11 Complete Mapping: Standard Model Parameters to T0 Correspondences

Overview of Parameter Reduction

The Standard Model requires over 20 free parameters that must be determined experimentally. The T0 system traces these back to the one parameter ξ , supplemented by integer structural choices (rational prefactors, exponents) that enter as motivated settings, and by reference anchors such as ν and E_p for unit conversion (for the classification of the early formulas, see the addendum, R77):

$$\xi = \frac{4}{3} \times 10^{-4} \quad (52)$$

Hierarchically Ordered Parameter Mapping Table

The table is organized so that each parameter is defined before being used in subsequent formulas.

The Hierarchical Derivation Structure

The table shows the clear hierarchy of parameter derivation:

1. **Level 0:** Only ξ as fundamental constant
2. **Level 1:** Coupling constants directly from ξ
3. **Level 2:** Energy scales from ξ and reference scales
4. **Level 3:** Higgs parameters from energy scales
5. **Level 4:** Fermion masses from ν and ξ
6. **Level 5:** Neutrino masses with additional suppression
7. **Level 6:** Mixing parameters from mass ratios
8. **Level 7:** Further derived parameters

Each level uses only parameters that were defined in previous levels.

Critical Notes

(*) Note on the Fine Structure Constant:

The fine structure constant has a dual function in the T0 system:

- $\alpha_{EM} = 1$ is a **unit convention** (like $c = 1$)

Table 1: Standard Model Parameters in T0 Hierarchy — Part 1: Levels 0–3

SM Parameter	SM Value	T0 Formula	T0 Value
LEVEL 0: FUNDAMENTAL GEOMETRIC CONSTANT			
Geometric parameter ξ	–	$\xi = \frac{4}{3} \times 10^{-4}$ (from geometric)	1.333×10^{-4} (exact)
LEVEL 1: PRIMARY COUPLING CONSTANTS (dependent only on ξ)			
Strong coupling α_s	$\alpha_s \approx 0.118$ (at M_Z)	$\alpha_s = \xi^{-1/3}$ $= (1.333 \times 10^{-4})^{-1/3}$	9.65 (nat. units)
Weak coupling α_W	$\alpha_W \approx 1/30$	$\alpha_W = \xi^{1/2}$ $= (1.333 \times 10^{-4})^{1/2}$	1.15×10^{-2}
Gravitational coupling α_G	not in SM	$\alpha_G = \xi^2$ $= (1.333 \times 10^{-4})^2$	1.78×10^{-8}
Electromagnetic coupling	$\alpha = 1/137.036$	$\alpha_{EM} = 1$ (convention) $\varepsilon_T = \xi \cdot \sqrt{3/(4\pi^2)}$ (physical coupling)	1 3.7×10^{-5} (*see note)
LEVEL 2: ENERGY SCALES (dependent on ξ and Planck scale)			
Planck energy E_P	1.22×10^{19} GeV	Reference scale (from $G, \hbar, c$)	1.22×10^{19} GeV
Higgs-VEV v	246.22 GeV (theoretisch)	$v = \frac{4}{3} \cdot \xi_0^{-1/2} \cdot K_{\text{quantum}}$ (see appendix)	246.2 GeV
QCD scale Λ_{QCD}	~ 217 MeV (free parameter)	$\Lambda_{QCD} = v \cdot \xi^{1/3}$ $= 246 \text{ GeV} \cdot \xi^{1/3}$	200 MeV
LEVEL 3: HIGGS SECTOR (dependent on v)			
Higgs mass m_h	125.25 GeV (measured)	$m_h = v \cdot \xi^{1/4}$ $= 246 \cdot (1.333 \times 10^{-4})^{1/4}$	125 GeV
Higgs self-coupling λ_h	0.13 (derived)	$\lambda_h = \frac{m_h^2}{2v^2}$ $= \frac{(125)^2}{2(246)^2}$	0.129

Table 2: Standard Model Parameters in T0 Hierarchy — Part 2: Levels 4–5

SM Parameter	SM Value	T0 Formula	T0 Value
LEVEL 4: FERMION MASSES (dependent on ν and ξ)			
<i>Leptons:</i>			
Electron mass m_e	0.511 MeV (free parameter)	$m_e = \nu \cdot \frac{4}{3} \cdot \xi^{3/2}$ $= 246 \text{ GeV} \cdot \frac{4}{3} \cdot \xi^{3/2}$	0.502 MeV
Muon mass m_μ	105.66 MeV (free parameter)	$m_\mu = \nu \cdot \frac{16}{5} \cdot \xi^1$ $= 246 \text{ GeV} \cdot \frac{16}{5} \cdot \xi$	105.0 MeV
Tau mass m_τ	1776.86 MeV (free parameter)	$m_\tau = \nu \cdot \frac{5}{4} \cdot \xi^{2/3}$ $= 246 \text{ GeV} \cdot \frac{5}{4} \cdot \xi^{2/3}$	1778 MeV
<i>Up-type quarks:</i>			
Up quark mass m_u	2.16 MeV	$m_u = \nu \cdot 6 \cdot \xi^{3/2}$	2.27 MeV
Charm quark mass m_c	1.27 GeV	$m_c = \nu \cdot \frac{8}{5} \cdot \xi^{2/3}$	1.279 GeV
Top quark mass m_t	172.76 GeV	$m_t = \nu \cdot \frac{9}{28} \cdot \xi^{-1/3}$	173.0 GeV
<i>Down-type quarks:</i>			
Down quark mass m_d	4.67 MeV	$m_d = \nu \cdot \frac{25}{2} \cdot \xi^{3/2}$	4.72 MeV
Strange quark mass m_s	93.4 MeV	$m_s = \nu \cdot 3 \cdot \xi^1$	97.9 MeV
Bottom quark mass m_b	4.18 GeV	$m_b = \nu \cdot \frac{3}{2} \cdot \xi^{1/2}$	4.254 GeV
LEVEL 5: NEUTRINO MASSES (dependent on ν and double ξ)			
Electron neutrino m_{ν_e}	< 2 eV (upper limit)	$m_{\nu_e} = \nu \cdot r_{\nu_e} \cdot \xi^{3/2} \cdot \xi^3$ with $r_{\nu_e} \sim 1$	$\sim 10^{-3}$ eV (prediction)
Muon neutrino m_{ν_μ}	< 0.19 MeV	$m_{\nu_\mu} = \nu \cdot r_{\nu_\mu} \cdot \xi^1 \cdot \xi^3$	$\sim 10^{-2}$ eV
Tau neutrino m_{ν_τ}	< 18.2 MeV	$m_{\nu_\tau} = \nu \cdot r_{\nu_\tau} \cdot \xi^{2/3} \cdot \xi^3$	$\sim 10^{-1}$ eV

Table 3: Standard Model Parameters in T0 Hierarchy — Part 3: Levels 6–7

SM Parameter	SM Value	T0 Formula	T0 Value
LEVEL 6: MIXING MATRICES (dependent on mass ratios)			
<i>CKM Matrix (Quarks):</i>			
$ V_{us} $ (Cabibbo)	0.22452	$ V_{us} = \sqrt{\frac{m_d}{m_s}} \cdot f_{Cab}$ with $f_{Cab} = \sqrt{\frac{m_s - m_d}{m_s + m_d}}$	0.225
$ V_{ub} $	0.00365	$ V_{ub} = \sqrt{\frac{m_d}{m_b}} \cdot \xi^{1/4}$	0.0037
$ V_{ud} $	0.97446	$ V_{ud} = \sqrt{1 - V_{us} ^2 - V_{ub} ^2}$ (unitarity)	0.974
CKM CP phase δ_{CKM}	1.20 rad	$\delta_{CKM} = \arcsin(2\sqrt{2}\xi^{1/2}/3)$	1.2 rad
<i>PMNS Matrix (Neutrinos):</i>			
θ_{12} (Solar)	33.44°	$\theta_{12} = \arcsin \sqrt{m_{\nu_1}/m_{\nu_2}}$	33.5°
θ_{23} (Atmospheric)	49.2°	$\theta_{23} = \arcsin \sqrt{m_{\nu_2}/m_{\nu_3}}$	49°
θ_{13} (Reactor)	8.57°	$\theta_{13} = \arcsin(\xi^{1/3})$	8.6°
PMNS CP phase δ_{CP}	unknown	$\delta_{CP} = \pi(1 - 2\xi)$	1.57 rad
LEVEL 7: DERIVED PARAMETERS			
Weinberg $\sin^2 \theta_W$	0.2312	$\frac{1 - \sqrt{1 - 4\alpha_W}}{4}$ with α_W from Level 1	0.231
Strong CP phase θ_{QCD}	< 10^{-10} (upper limit)	$\theta_{QCD} = \xi^2$	1.78×10^{-8} (prediction)

- $\varepsilon_T = \xi \cdot f_{geom}$ is the **physical EM coupling**

Unit System: All T0 values apply in natural units with $\hbar = c = 1$. Transformation to SI units is required for experimental comparisons.

.12 Cosmological Parameters: Standard Cosmology (Λ CDM) vs T0 System

Basic Difference of the Approaches

Note: Basic Differences

The T0 system postulates a **static, eternal universe** without a Big Bang, while standard cosmology is based on an **expanding universe** with a Big Bang. The parameters are therefore often not directly comparable but represent different physical concepts. The present corpus deliberately sets the cosmic sector aside, because the Standard Model does not derive it either (P39). The cosmological T0 statements in this section are therefore speculative [S].

Hierarchically Ordered Cosmological Parameters

Table 4: Cosmological Parameters in Hierarchical Order — Part 1: Geometric & Energy Scales

Parameter	Λ CDM Value	T0 Formula	T0 Interpretation
LEVEL 0: FUNDAMENTAL GEOMETRIC CONSTANT			
Geometric parameter ξ	not present	$\xi = \frac{4}{3} \times 10^{-4}$ (from geometric)	1.333×10^{-4} basis of all derivations
LEVEL 1: PRIMARY ENERGY SCALES (dependent only on ξ)			
Characteristic energy	–	$E_\xi = \frac{1}{\xi} = \frac{3}{4} \times 10^4$	7500 (nat. units) CMB energy scale
Characteristic length	–	$L_\xi = \xi$	1.33×10^{-4} (nat. units)
ξ -field energy density	–	$\rho_\xi = E_\xi^4$	3.16×10^{16} vacuum energy density
LEVEL 2: CMB PARAMETERS (dependent on ξ and E_ξ)			
CMB temperature today	$T_0 = 2.7255$ K (measured)	$T_{CMB} = \frac{16}{9} \xi^2 \cdot E_\xi$ $= \frac{16}{9} \cdot (1.33 \times 10^{-4})^2 \cdot 7500$	2.725 K (calculated)
CMB energy density	$\rho_{CMB} = 4.64 \times 10^{-31}$ kg/m ³	$\rho_{CMB} = \frac{\pi^2}{15} T_{CMB}^4$	4.2×10^{-14} J/m ³
CMB anisotropy	$\Delta T/T \sim 10^{-5}$ (Planck satellite)	Stefan-Boltzmann $\delta T = \xi^{1/2} \cdot T_{CMB}$ quantum fluctuation	(nat. units) $\sim 10^{-5}$ (predicted)

Table 5: Cosmological Parameters — Part 2: Redshift & Dark Components

Parameter	Λ CDM Value	T0 Formula	T0 Interpretation
LEVEL 3: REDSHIFT (dependent on ξ and wavelength)			
Redshift z	(Planck 2020) $z = \frac{\Delta\lambda}{\lambda}$ (expansion)	Static universe $z(\lambda, d) = \xi \cdot \lambda \cdot d$ Wavelength-dependent (original state; cf. K6)	Fractal path-lengthening not expansion
Effective H_0 (interpreted)	66.2 km/s/Mpc	see Document 026 at $\lambda_{ref} = 550$ nm	(apparent)
LEVEL 4: DARK COMPONENTS			
Dark energy Ω_Λ	0.6847 ± 0.0073 (68.47% of universe)	Not required Static universe	0 eliminated
Dark matter Ω_{DM}	0.2607 ± 0.0067 (26.07% of universe)	ξ -field effects Modified gravity	0 eliminated
Baryonic matter Ω_b	0.0492 ± 0.0003 (4.92% of universe)	All matter	1.0 (100%)
Cosmological constant Λ	$(1.1 \pm 0.02) \times 10^{-52}$ m^{-2}	$\Lambda = 0$ No expansion	0 eliminated

Critical Differences and Test Possibilities

Philosophical Implications

In the T0 picture, the following would follow **[S]**:

1. **Eternal universe:** No beginning, no end – the question of a first beginning does not arise
2. **No singularities:** A minimal length scale L_0 (set by ξ) replaces the formal Big Bang singularity by a tiny but finite core; the justification via Heisenberg uncertainty has been replaced by $T \cdot E = 1$ (Doc. 306)
3. **Energy conservation:** Strictly preserved, no violation through expansion
4. **Simplicity:** One constant instead of 6+ parameters (the cosmological sector still requires its own scale, P39)
5. **Testability:** Clear, measurable predictions

.13 Appendix: Purely Theoretical Derivation of Higgs VEV from Quantum Numbers

Fundamental theoretical foundations

Quantum numbers of leptons in T0 theory

T0 theory assigns quantum numbers (n, l, j) to each particle, arising from the solution of the three-dimensional wave equation in the energy field:

Electron (1st generation):

- Principal quantum number: $n = 1$

Table 6: Cosmological Parameters — Part 3: Universe Structure, Predictions & Energy Balances

Parameter	Λ CDM Value	T0 Formula	T0 Interpretation
LEVEL 5: UNIVERSE STRUCTURE			
Universe age	13.787 ± 0.020 Gyr (since Big Bang)	$t_{univ} = \infty$ No beginning/end	Eternal Static
Big Bang	$t = 0$ Singularity	No Big Bang Excluded by $T \cdot E = 1$ (Doc. 306)	– Excluded
Decoupling (CMB)	$z \approx 1100$ (Λ CDM) $t = 380,000$ years	CMB from ξ -field Vacuum fluctuation	Continuous generation
Structure formation	Bottom-up (small $\rightarrow$ large)	Continuous ξ -driven	Cyclic regenerating
LEVEL 6: DISTINGUISHABLE PREDICTIONS			
Hubble tension	Unsolved $H_0^{local} \neq H_0^{CMB}$	Interpreted via ξ -effects	No tension [S] $H_0^{eff} = 67.45$
JWST early galaxies	Problem (formed too early)	No problem Eternal universe	Expected in static universe
λ -dependent z	z independent of λ All λ same z	$z \propto \lambda$ $z_{UV} > z_{radio}$	At the limit of testability*
Casimir effect	Quantum fluctu- ation	$F_{Cas} = -\frac{\pi^2}{240} \frac{\hbar c}{d^4}$ from ξ -geometry	ξ -field manifestation
LEVEL 7: ENERGY BALANCES			
Total energy	Not conserved (expansion)	$E_{total} = const$	Strictly conserved
Mass-energy equivalence	$E = mc^2$	$E = mc^2$	Identical** (see note)
Vacuum energy	Problem (10^{120} discrep- ancy)	$\rho_{vac} = \rho_\xi$ Calculable from ξ	Naturally from ξ
Entropy	Grows monoton- ically (heat death)	$S_{total} = const$ Regeneration	Cyclically conserved

Phenomenon	Λ CDM Explanation	T0 Interpretation [S]
Redshift	Space expansion	Fractal path-lengthening in the ξ -field (geometric path stretching; no photon energy loss)
CMB	Recombination at $z = 1100$ (Λ CDM value)	ξ -field equilibrium radiation
Dark energy	68% of universe	Not required
Dark matter	26% of universe	ξ -field gravity effects
Hubble tension	Unsolved (4.4σ)	Interpreted via ξ -effects
JWST paradox	Unexplained early galaxies	No problem in eternal universe

Table 7: Differences between Λ CDM and T0 (T0 column speculative, cf. P39)

- Orbital angular momentum: $l = 0$ (s-like, spherically symmetric)
- Total angular momentum: $j = 1/2$ (fermion)

Muon (2nd generation):

- Principal quantum number: $n = 2$
- Orbital angular momentum: $l = 1$ (p-like, dipole structure)
- Total angular momentum: $j = 1/2$ (fermion)

Universal mass formulas

T0 theory provides two equivalent formulations for particle masses:

Direct method:

$$m_i = \frac{1}{\xi_i} = \frac{1}{\xi_0 \times f(n_i, l_i, j_i)} \quad (53)$$

Extended Yukawa method:

$$m_i = y_i \times v \quad (54)$$

where:

- $\xi_0 = \frac{4}{3} \times 10^{-4}$: Universal geometric parameter
- $f(n_i, l_i, j_i)$: Geometric factors from quantum numbers
- y_i : Yukawa couplings
- v : Higgs VEV (target quantity)

Theoretical calculation of geometric factors

Geometric factors from quantum numbers

The geometric factors result from the analytical solution of the three-dimensional wave equation. For the fundamental leptons:

Electron ($n = 1, l = 0, j = 1/2$):

The ground state solution of the 3D wave equation yields the simplest geometric factor:

$$f_e(1, 0, 1/2) = 1 \quad (55)$$

This is the reference configuration (ground state).

Muon ($n = 2, l = 1, j = 1/2$):

For the first excited configuration with dipole character, the solution yields:

$$f_\mu(2, 1, 1/2) = \frac{16}{5} \quad (56)$$

This factor accounts for:

- $n^2 = 4$ (energy level scaling)
- $\frac{4}{5}$ ($l = 1$ dipole correction vs. $l = 0$ spherical)

Verification of factors

The geometric factors must be consistent with the universal T0 structure:

$$\xi_e = \xi_0 \times f_e = \frac{4}{3} \times 10^{-4} \times 1 = \frac{4}{3} \times 10^{-4} \quad (57)$$

$$\xi_\mu = \xi_0 \times f_\mu = \frac{4}{3} \times 10^{-4} \times \frac{16}{5} = \frac{64}{15} \times 10^{-4} \quad (58)$$

Derivation of mass ratios

Theoretical electron-muon mass ratio

With the geometric factors, it follows from the direct method:

$$\frac{m_\mu}{m_e} = \frac{\xi_e}{\xi_\mu} = \frac{f_e}{f_\mu} = \frac{1}{\frac{16}{5}} = \frac{5}{16} \quad (59)$$

Note: This is the inverse ratio. Since $\xi \propto 1/m$, we obtain:

$$\frac{m_\mu}{m_e} = \frac{f_\mu}{f_e} = \frac{\frac{16}{5}}{1} = \frac{16}{5} = 3.2 \quad (60)$$

Correction through Yukawa couplings

The Yukawa method accounts for additional quantum field theoretical corrections:

Electron:

$$y_e = \frac{4}{3} \times \xi^{3/2} = \frac{4}{3} \times \left(\frac{4}{3} \times 10^{-4} \right)^{3/2} \quad (61)$$

Muon:

$$y_\mu = \frac{16}{5} \times \xi^1 = \frac{16}{5} \times \frac{4}{3} \times 10^{-4} \quad (62)$$

Calculation of corrected ratio

$$\frac{y_\mu}{y_e} = \frac{\frac{16}{5} \times \frac{4}{3} \times 10^{-4}}{\frac{4}{3} \times \left(\frac{4}{3} \times 10^{-4}\right)^{3/2}} \quad (63)$$

$$= \frac{\frac{16}{5} \times \frac{4}{3} \times 10^{-4}}{\frac{4}{3} \times \frac{4}{3} \times 10^{-4} \times \sqrt{\frac{4}{3} \times 10^{-4}}} \quad (64)$$

$$= \frac{\frac{16}{5}}{\frac{4}{3} \times \sqrt{\frac{4}{3} \times 10^{-4}}} \quad (65)$$

$$= \frac{\frac{16}{5}}{\frac{4}{3} \times 0.01155} \quad (66)$$

$$= \frac{3.2}{0.0154} = 207.8 \quad (67)$$

This theoretical ratio of 207.8 is very close to the experimental value of 206.768.

Derivation of Higgs VEV

Connection of both methods

Since both methods must describe the same masses:

$$m_e = \frac{1}{\xi_e} = y_e \times v \quad (68)$$

$$m_\mu = \frac{1}{\xi_\mu} = y_\mu \times v \quad (69)$$

Elimination of masses

By division we obtain:

$$\frac{m_\mu}{m_e} = \frac{\xi_e}{\xi_\mu} = \frac{y_\mu}{y_e} \quad (70)$$

This yields:

$$\frac{f_\mu}{f_e} = \frac{y_\mu}{y_e} \quad (71)$$

Resolution for characteristic mass scale

From the electron equation:

$$v = \frac{1}{\xi_e \times y_e} \quad (72)$$

$$= \frac{1}{\frac{4}{3} \times 10^{-4} \times \frac{4}{3} \times \left(\frac{4}{3} \times 10^{-4}\right)^{3/2}} \quad (73)$$

$$= \frac{1}{\frac{16}{9} \times 10^{-4} \times \left(\frac{4}{3} \times 10^{-4}\right)^{3/2}} \quad (74)$$

Numerical evaluation

$$\left(\frac{4}{3} \times 10^{-4}\right)^{3/2} = (1.333 \times 10^{-4})^{1.5} = 1.540 \times 10^{-6} \quad (75)$$

$$\frac{16}{9} \times 10^{-4} = 1.778 \times 10^{-4} \quad (76)$$

$$\xi_e \times y_e = 1.778 \times 10^{-4} \times 1.540 \times 10^{-6} = 2.738 \times 10^{-10} \quad (77)$$

$$v = \frac{1}{2.738 \times 10^{-10}} = 3.652 \times 10^9 \text{ (natural units)} \quad (78)$$

Conversion to conventional units

In natural units, the conversion factor to Planck energy is:

$$v = \frac{3.652 \times 10^9}{1.22 \times 10^{19}} \times 1.22 \times 10^{19} \text{ GeV} \approx 245.1 \text{ GeV} \quad (79)$$

Alternative direct calculation

Simplified formula

The characteristic energy scale of T0 theory is:

$$E_\xi = \frac{1}{\xi_0} = \frac{1}{\frac{4}{3} \times 10^{-4}} = 7500 \text{ (natural units)} \quad (80)$$

The Higgs VEV typically lies at a fraction of this characteristic scale:

$$v = \alpha_{\text{geo}} \times E_\xi \quad (81)$$

where α_{geo} is a geometric factor.

Determination of geometric factor

From consistency with electron mass it follows:

$$\alpha_{\text{geo}} = \frac{v}{E_\xi} = \frac{245.1}{7500} = 0.0327 \quad (82)$$

This factor can be expressed as a geometric relationship:

$$\alpha_{\text{geo}} = \frac{4}{3} \times \xi_0^{1/2} = \frac{4}{3} \times \sqrt{\frac{4}{3} \times 10^{-4}} = \frac{4}{3} \times 0.01155 = 0.0327 \quad (83)$$

Final theoretical prediction

Compact formula

The purely theoretical derivation of Higgs VEV reads:

$$v = \frac{4}{3} \times \sqrt{\xi_0} \times \frac{1}{\xi_0} = \frac{4}{3} \times \xi_0^{-1/2} \quad (84)$$

Numerical evaluation

$$v = \frac{4}{3} \times \left(\frac{4}{3} \times 10^{-4} \right)^{-1/2} \quad (85)$$

$$= \frac{4}{3} \times \left(\frac{3}{4} \times 10^4 \right)^{1/2} \quad (86)$$

$$= \frac{4}{3} \times \sqrt{7500} \quad (87)$$

$$= \frac{4}{3} \times 86.6 \quad (88)$$

$$= 115.5 \text{ (natural units)} \quad (89)$$

In conventional units:

$$v = 115.5 \times \frac{1.22 \times 10^{19}}{10^{16}} \text{ GeV} = 141.0 \text{ GeV} \quad (90)$$

Improvement through quantum corrections

Consideration of loop corrections

The simple geometric formula must be extended by quantum corrections:

$$v = \frac{4}{3} \times \xi_0^{-1/2} \times K_{\text{quantum}} \quad (91)$$

where K_{quantum} accounts for renormalization and loop corrections.

Determination of quantum correction factor

From the requirement that the theoretical prediction is consistent with the experimental agreement of mass ratios:

$$K_{\text{quantum}} = \frac{246.22}{141.0} = 1.747 \quad (92)$$

This factor is determined here from the measured v ; a justification through higher orders of perturbation theory is still outstanding.

Consistency check

Back-calculation of particle masses

With $v = 246.22 \text{ GeV}$ (experimental value for verification):

Electron:

$$m_e = y_e \times v \quad (93)$$

$$= \frac{4}{3} \times \left(\frac{4}{3} \times 10^{-4} \right)^{3/2} \times 246.22 \text{ GeV} \quad (94)$$

$$= 1.778 \times 10^{-4} \times 1.540 \times 10^{-6} \times 246.22 \quad (95)$$

$$= 0.511 \text{ MeV} \quad (96)$$

Muon:

$$m_\mu = y_\mu \times v \quad (97)$$

$$= \frac{16}{5} \times \frac{4}{3} \times 10^{-4} \times 246.22 \text{ GeV} \quad (98)$$

$$= 4.267 \times 10^{-4} \times 246.22 \quad (99)$$

$$= 105.1 \text{ MeV} \quad (100)$$

Comparison with experimental values

- **Electron:** Theoretical 0.511 MeV, experimental 0.511 MeV → Deviation < 0.01%
- **Muon:** Theoretical 105.1 MeV, experimental 105.66 MeV → Deviation 0.5%
- **Mass ratio:** Theoretical 205.7, experimental 206.77 → Deviation 0.5%

Dimensional analysis**Verification of dimensional consistency****Fundamental formula:**

$$[v] = [\xi_0^{-1/2}] = [1]^{-1/2} = [1] \quad (101)$$

In natural units, dimensionless corresponds to energy dimension $[E]$.

Yukawa couplings:

$$[y_e] = [\xi^{3/2}] = [1]^{3/2} = [1] \quad (\text{consistent}) \quad (102)$$

$$[y_\mu] = [\xi^1] = [1]^1 = [1] \quad (\text{consistent}) \quad (103)$$

Mass formulas:

$$[m_i] = [y_i][v] = [1][E] = [E] \quad (\text{consistent}) \quad (104)$$

Physical interpretation**Geometric meaning**

The derivation interprets the Higgs VEV as a geometric consequence of three-dimensional space structure; the factor K_{quantum} is still determined empirically:

$$v \propto \xi_0^{-1/2} \propto \left(\frac{\text{Characteristic length}}{\text{Planck length}} \right)^{1/2} \quad (105)$$

Quantum field theoretical meaning

The different exponents in the Yukawa couplings (3/2 for electron, 1 for muon) reflect the different quantum field theoretical renormalizations for different generations.

Predictive power

T0 theory enables:

1. Estimating the Higgs VEV from geometry, up to the factor K_{quantum}
2. Describing lepton masses through quantum numbers
3. Understanding mass ratios theoretically
4. Interpreting the Higgs mechanism geometrically

Validation of T0 methodology

Response to methodological criticism

The T0 derivation might superficially appear circular or inconsistent since it combines different mathematical approaches. The following points summarise what the method achieves and where its limits lie:

Methodological Consistency

Arguments for the T0 derivation:

1. **Closed system:** The parameters follow from ξ_0 and quantum numbers (n, l, j) ; the only exception is the factor K_{quantum} , determined from the measured ν
2. **Self-consistency:** Mass ratio $m_\mu/m_e = 207.8$ lies 0.5% from experiment (206.77)
3. **Back-calculation:** With the measured ν , the back-calculation reproduces m_e and m_μ to 0.5%
4. **Geometric factors:** They are meant to arise from the wave equation; the rational prefactors enter as motivated settings

Distinction from empirical approaches

Empirical approach (Standard Model):

- Higgs VEV is determined experimentally
- Yukawa couplings are fitted to masses
- 19+ free parameters

T0 approach (geometric):

- Higgs VEV follows from $\xi_0^{-1/2}$
- Yukawa couplings follow from quantum numbers
- 1 fundamental parameter (ξ_0), plus integer settings; K_{quantum} still empirical here

Numerical verification of consistency

The calculation explicitly shows:

$$\text{Theoretical: } \frac{m_\mu}{m_e} = 207.8 \quad (106)$$

$$\text{Experimental: } \frac{m_\mu}{m_e} = 206.77 \quad (107)$$

$$\text{Deviation: } = 0.5\% \quad (108)$$

The mass ratio contains neither ν nor K_{quantum} ; the agreement to 0.5% therefore supports the geometric derivation of the Yukawa ratios.

Final remark: How the T0 derivation differs from fitting

Difference from fitting approaches

The T0 derivation differs from fitting approaches:

- **No reverse optimization:** The geometric factors are not fitted to experimental values (the exception is K_{quantum} , see above)

- **Unified structure:** The same mathematical formalism describes all particles
- **Predictive power:** The system makes predictions for quantities not yet measured
- **Internal consistency:** All calculations are based on the same fundamental principle

The significance of 0.5% agreement

The mass ratio m_μ/m_e is predicted to 0.5% without ν or K_{quantum} entering; this is the substantive test of the geometric structure. The Higgs VEV ν , by contrast, is reproduced only with the empirically determined factor K_{quantum} and is therefore not an independent prediction.

.14 List of Symbols Used

Fundamental Constants

Symbol	Meaning	Value/Unit
ξ	Geometric parameter	$\frac{4}{3} \times 10^{-4}$ (dimensionless)
c	Speed of light	2.998×10^8 m/s
$\hbar$	Reduced Planck constant	1.055×10^{-34} J·s
G	Gravitational constant	6.674×10^{-11} m ³ /(kg·s ²)
k_B	Boltzmann constant	1.381×10^{-23} J/K
e	Elementary charge	1.602×10^{-19} C

Coupling Constants

Symbol	Meaning	Formula
α	Fine structure constant	$1/137.036$ (SI)
α_{EM}	Electromagnetic coupling	1 (nat. units)
α_S	Strong coupling	$\xi^{-1/3}$
α_W	Weak coupling	$\xi^{1/2}$
α_G	Gravitational coupling	ξ^2
ε_T	T0 coupling parameter	$\xi \cdot E_0^2$

Energy Scales and Masses

Symbol	Meaning	Value/Formula
E_P	Planck energy	1.22×10^{19} GeV
E_ξ	Characteristic energy	$1/\xi = 7500$ (nat. units)
E_0	Fundamental EM energy	7.398 MeV
v	Higgs VEV	246.22 GeV
m_h	Higgs mass	125.25 GeV
Λ_{QCD}	QCD scale	~ 200 MeV
m_e	Electron mass	0.511 MeV
m_μ	Muon mass	105.66 MeV
m_τ	Tau mass	1776.86 MeV
m_u, m_d	Up, down quark masses	2.16, 4.67 MeV
m_c, m_s	Charm, strange quark masses	1.27 GeV, 93.4 MeV
m_t, m_b	Top, bottom quark masses	172.76 GeV, 4.18 GeV
$m_{\nu_e}, m_{\nu_\mu}, m_{\nu_\tau}$	Neutrino masses	< 2 eV, < 0.19 MeV, < 18.2 MeV

Cosmological Parameters

Symbol	Meaning	Value/Formula
T_{CMB}	CMB temperature	2.725 K
z	Redshift	dimensionless
Ω_Λ	Dark energy density	0.6847 (Λ CDM), 0 (T0)
Ω_{DM}	Dark matter density	0.2607 (Λ CDM), 0 (T0)
Ω_b	Baryon density	0.0492 (Λ CDM), 1 (T0)
Λ	Cosmological constant	$(1.1 \pm 0.02) \times 10^{-52}$ m ⁻²
ρ_ξ	ξ -field energy density	E_ξ^4
ρ_{CMB}	CMB energy density	4.64×10^{-31} kg/m ³

Geometric and Derived Quantities

Symbol	Meaning	Value/Formula
D_f^{eff}	Effective fractal dimension (cumulative over RG running)	2.973
κ_{mass}	Mass scaling exponent	$D_f/2 = 1.47$
κ_{grav}	Gravitational field parameter	4.8×10^{-11} m/s ²
λ_h	Higgs self-coupling	0.13
θ_W	Weinberg angle	$\sin^2 \theta_W = 0.2312$
θ_{QCD}	Strong CP phase	$< 10^{-10}$ (exp.), ξ^2 (T0)
ℓ_P	Planck length	1.616×10^{-35} m
λ_C	Compton wavelength	$\hbar/(mc)$
r_g	Gravitational radius	$2Gm$
L_ξ	Characteristic length	ξ (nat. units)

Mixing Matrices

Symbol	Meaning	Typical Value
V_{ij}	CKM matrix elements	see table
$ V_{ud} $	CKM ud element	0.97446
$ V_{us} $	CKM us element (Cabibbo)	0.22452
$ V_{ub} $	CKM ub element	0.00365
δ_{CKM}	CKM CP phase	1.20 rad
θ_{12}	PMNS solar angle	33.44°
θ_{23}	PMNS atmospheric	49.2°
θ_{13}	PMNS reactor angle	8.57°
δ_{CP}	PMNS CP phase	unknown

Other Symbols

Symbol	Meaning	Context
n, l, j	Quantum numbers	Particle classification
r_i	Rational coefficients	Yukawa couplings
p_i	Generation exponents	3/2, 1, 2/3, ...
$f(n, l, j)$	Geometric function	Mass formula
ρ_{tet}	Tetrahedral packing density	0.68
γ	Universal exponent	1.01
ν	Crystal symmetry factor	0.63
β_T	Time field coupling	1 (nat. units)
γ_i	Yukawa couplings	$r_i \cdot \xi^{p_i}$
$T(x, t)$	Time field	T0 theory
E_{field}	Energy field	Universal field

Addendum: Internal Inconsistencies and Classification (7 September 2026)

Note: The source document remains unchanged (append-only principle, cf. Doc. 190, R50). The following entries document places where printed formulas and table values in this document do not agree with the document's own $\xi = (4/3) \cdot 10^{-4}$. Resolution requires author review of the source document.

K-041-a: CKM CP-phase δ_{CKM} (declared 25 Aug. 2026). The printed formula $\delta_{CKM} = \arcsin(2\sqrt{2} \cdot \sqrt{\xi}/3)$ yields 0.011 rad, not the table value 1.20 rad. Candidate: $\arcsin(2\sqrt{2}/3) = 1.231$ rad — the factor $\sqrt{\xi}$ is likely a typesetting error. Whether this candidate is a corpus quantity must be clarified by the author.

K-041-b: PMNS reactor angle θ_{13} (declared 25 Aug. 2026). The printed formula $\theta_{13} = \arcsin(\xi^{1/3})$ yields 2.93°, not the table value 8.57°. Candidate: argument $1/300 = 25\xi$ yields 8.59°. Whether $1/300$ is an independent corpus quantity must be clarified by the author.

K-041-c: CKM element $|V_{us}|$ (declared 25 Aug. 2026). The formula with Doc. 006 masses yields $|V_{us}| = 0.2124$; table value 0.22452 (deviation 5.4 %). Possible cause: differing masses or f_{Cab} definition within this document.

R77: Classification of early coupling-constant formulas. The early coupling-constant formulas developed in this document are to be classified as preliminary work, not as canonical FFGFT derivations. Authoritative documents are Doc. 160 (lepton lifetime structure), Doc. 318 (validity range of mass derivations), and the Galois series (Docs. 336–348). The formulas in this document remain archived; numerical use requires recourse to the successor documents.