

Analysis of FFGF (Fundamental Fractal-Geometric Field Theory) and t_0 Theory

Contents

.1	Introduction	1
.2	Foundational Postulates and Fractal Spacetime	1
	Fractal Dimension of Spacetime	1
	The Fractal Correction Factor K_{frak}	1
	Time-Mass Duality and the Planck Scale	2
	Clarification: Effective Planck Scale vs. Fundamental t_0 Scale	2
	Modification of Electromagnetic Laws in Fractal Space	2
	Key Concepts in the Document	2
.3	Mathematical Concepts	3
	1. The Fractal Dimension $D_f = 3 - \xi$	3
	2. The Factor $\frac{4}{3}$ – Geometric Interpretation	3
	3. Time-Mass Duality – Deeper Mathematics	4
	4. The Planck Scale as Effective Limit	4
	5. Vacuum Energy and the Cutoff by ξ	5
	6. Gravitational Relationship (implied in the document)	5
.4	A Closer Look at the Mathematics of Torus Geometry (mentioned in the document)	6
	Why the Torus?	6
	Mathematical Description of the Torus	6
	Curvature of the Torus	6
	Energy Flow in the Torus (FFGF Model)	6
	Connection to $D_f = 3 - \xi$	7
	Mass from Torus Geometry	7
	Scaling: From Proton to Galaxy	8
	Why is the Torus Stable?	8
	Connection to the Schwarzschild Metric	8
.5	Connection Between Torus Topology and Quantum Numbers (Spin, Charge)	8
	Topological Quantum Numbers from Torus Geometry – Detailed Derivation	8
	1. Spin – The Winding Number $w = n_\phi/n_\theta$	9
	2. Electric Charge – Quantized Electric Flux Through the Torus	9
	3. Color Charge – Topological Linking of Three Flux Strands	10
	Torus Geometry in Quantum Computing	10
.6	Torus Geometry in Cosmology – Scale-Invariant Torsional Structures	11
	Cross-Scale Torsional Correspondences	11
	Core Principle: Scale Invariance and Fractal Self-Similarity	12
	Cosmological Implications – Without Dark Matter and Without Expansion	12
	Two-Dipole Model in Detail	12
	Introduction and Motivation	13
	Model Description	13
	Mathematical Framework	13
	Cosmological Implications	14
	Critical Analysis	14
	Parallel to the Toroidal Photon Model (Williamson & van der Mark, 1997)	14

	Key Parallels to the FFGF Torus Structure	15
	Differences and Extension by the FFGF	15
.7	Electromagnetic Fields in Torus Geometry	16
	Maxwell's Equations on the Torus	16
	Magnetic Field Configuration in the Torus	16
	Stability Condition (Kruskal-Shafranov)	16
	Origin of the Magnetic Moment	16
	Electromagnetic Self-Energy	17
	Connection to ξ and D_f	17
.8	Fluid Dynamics in the Torus (Navier-Stokes on Curved Spaces)	17
	Navier-Stokes in Curved Coordinates	17
	Metric Tensor for the Torus	18
	Velocity Field in the Rotating Torus	18
	Vorticity	18
	Energy Conservation in Torus Flow	18
	Turbulence and Stability	19
	Helmholtz Decomposition	19
	Casimir Effect in the Torus	19
	Connection to Time-Mass Duality	19
	Clarification: Effective Planck Scale vs. Fundamental t_0 Scale	20
	Fractal Turbulence	20
.9	Overall Synthesis: The Three Aspects Together	20

.1 Introduction

Note on versions. The German and English versions of this document differ in structure and extent; earlier revisions shortened, extended or changed parts of both. Where they differ, the more extensive German version applies; sections contained only in this English version count as supplements.

Note on the corpus status: variable speed of light. The idea of a varying speed of light referred to in this document has been replaced in the later corpus by *fractal path lengthening*: the geometric stretching of the propagation path in the ξ field (Doc. 026, 149, 182; achromatic according to K6 in Doc. 190). Statements that rely on a variable speed of light are to be read accordingly.

This analysis describes the mathematical framework of the Fundamental Fractal-Geometric Field Theory (FFGF) and the t_0 theory. The focus is on presenting the internal mathematical consistency and structure.

Status markers. Statements marked **[S]** are *speculative*: a hint or conjecture that has not yet been derived or numerically checked in the corpus. Here this applies in particular to the special cases for the early universe ($D_f = 2$), the interpretations of dark matter and dark energy, and the torus cosmology and Two-Dipole Model: the present corpus deliberately leaves the cosmic sector aside, because the Standard Model does not derive it either (P39).

.2 Foundational Postulates and Fractal Spacetime

Fractal Dimension of Spacetime

The central starting point of the theory is the description of spacetime by a fractal dimension D_f that lies slightly below the topological dimension 3:

$$D_f = 3 - \xi, \quad \text{with} \quad \xi = \frac{4}{3} \times 10^{-4}. \quad (1)$$

The parameter ξ quantifies the fractal dimension deficit and is fundamental for all subsequent scalings and corrections (see T0_xi_ursprung.pdf).

The Fractal Correction Factor K_{frak}

Over many scaling orders, ξ leads to an accumulated geometric correction factor:

$$K_{\text{frak}} = 1 - 100\xi \approx 0.9867. \quad (2)$$

This factor modifies fundamental geometric and physical quantities (see 133_FraktaLe_Korrektur_Herleitung_En.pdf).

Time-Mass Duality and the Planck Scale

Equating the Planck relation $E = hf$ with the Einstein relation $E = mc^2$ and substituting $f = 1/T$ yields a fundamental duality:

$$m = \frac{h}{c^2 T}. \quad (3)$$

Clarification: Effective Planck Scale vs. Fundamental t_0 Scale

In this analysis, the **effective limit** of continuous physics is described by the **Planck time t_P** and **Planck length ℓ_P** (see the section "The Planck Scale as Limit" below). Below this scale, the classical concept of space and time breaks down.

The **fundamental t_0 scale** of the theory, however, is **sub-Planck** and describes the internal granulation of the fractal field:

- Sub-Planck length: $\Lambda_0 = \xi \cdot \ell_P \approx 1.333 \times 10^{-4} \cdot \ell_P \approx 2.15 \times 10^{-39} \text{ m}$
- Characteristic t_0 lengths and times: $r_0 = 2GE$, $t_0 = 2GE$ (see Zeit_En.pdf and 010_T0_Energie_En.pdf)

The Planck scale (ℓ_P , t_P) is thus the **outer reference limit** of the effective theory, while t_0 represents the **sub-Planck granulation** on which the fractal structure truly operates.

As a complement, two interactive visualizations are provided in the 2/html directory (GitHub Pages, open in browser):

- [torus_geometry_ffgf.html](#) – animated torus geometry with energy flow and selectable scale (proton, planet, galaxy).
- [t0_subplanck_structure.html](#) – comparison of the effective Planck boundary and the fundamental t_0 sub-Planck scale (Λ_0 , τ_0).

Modification of Electromagnetic Laws in Fractal Space

In a D_f -dimensional space, the general Coulomb law reads:

$$F_{\text{Coulomb}} \propto \frac{1}{r^{D_f-1}}. \quad (4)$$

For the present universe with $D_f = 3 - \xi$, this gives the tiny but in principle measurable modification:

$$F_{\text{Coulomb}} \propto \frac{1}{r^{2-\xi}}. \quad (5)$$

[S] Special case: early universe ($t \rightarrow t_{\min}$, $D_f \rightarrow 2$): At the beginning of the universe, when the third spatial dimension had not yet fully emerged, $D_f = 2$ applies. The Coulomb law then reduces to a linear potential:

$$F_{\text{Coulomb}}|_{D_f=2} \propto \frac{1}{r}. \quad (6)$$

This corresponds to the well-known logarithmic confinement potential of 2D physics and has direct consequences for the structure of charge distributions in the Planck era.

Analogously, FFGF regards the speed of light c not as a fundamental constant but as a quantity derived from the medium: $c = \ell_P/t_P$, with an effective, fractally modified velocity $c_{\text{eff}} \approx c \cdot (1 + \xi/2)$.

Key Concepts in the Document

- Spacetime has a fractal structure with dimension $D_f = 3 - \xi$, where $\xi = \frac{4}{3} \times 10^{-4}$.
- Mass and time are proposed as dual aspects of the same phenomenon.
- Dark matter and dark energy are reinterpreted as geometric effects, not as actual substances **[S]**.
- The vacuum has a fractal structure that prevents infinities.

.3 Mathematical Concepts

1. The Fractal Dimension $D_f = 3 - \xi$

Given: $\xi = \frac{4}{3} \times 10^{-4} \approx 0.0001333 \dots$

Therefore: $D_f \approx 2.9998666 \dots$

Mathematical meaning: In classical fractal geometry, the Hausdorff dimension describes how an object "fills" space:

- A point: $D = 0$
- A line: $D = 1$
- A surface: $D = 2$
- A volume: $D = 3$
- Koch snowflake: $D \approx 1.26$ (more than a line, less than a surface)

The meaning of $D_f < 3$: If space has a dimension of 2.9998666 instead of exactly 3, this mathematically means:

- Space is not "completely filled".
- There is a kind of "porosity" or lacunarity.
- These gaps constitute 0.0001333 of the dimensionality.

Scaling behavior: For true fractals: When the resolution is increased by a factor r , the number of visible structures increases by r^D . In general:

$$N(r) \propto r^{D_f}$$

Present universe ($D_f = 3 - \xi$):

$$N(r) \propto r^{(3-\xi)}$$

[S] Special case: early universe ($t \rightarrow t_{\min}$, $D_f = 2$):

$$N(r)|_{D_f=2} \propto r^2$$

The spatial structure is surface-like – the number of visible structures grows only quadratically with resolution.

2. The Factor $\frac{4}{3}$ – Geometric Interpretation

Sphere packing: The factor $\frac{4}{3}$ appears frequently in geometry:

- Sphere volume: $V = \frac{4}{3}\pi r^3$
- Ratio of sphere volume to enclosing cube: $\frac{4\pi}{3}/8 \approx 0.524$

Densest sphere packing: Maximum packing density: $\frac{\pi}{\sqrt{18}} \approx 0.7405$ Thus, 26% "gaps" remain.

Possible interpretation in FFGF: If the vacuum consists of "Planck spheres" or toroidal structures that cannot be packed perfectly, geometric interstices arise. The factor $\frac{4}{3}$ might encode this packing geometry.

3. Time-Mass Duality – Deeper Mathematics

The derivation: From $E = mc^2$ and $E = hf$ it follows:

$$mc^2 = hf = \frac{h}{T}$$

Thus:

$$m = \frac{h}{c^2 T}$$

Dimensional analysis:

- $[h] = \text{Js} = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}$
- $[c^2] = \text{m}^2 \cdot \text{s}^{-2}$
- $[T] = \text{s}$
-

$$[m] = \frac{[h]}{[c^2][T]} = \frac{\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}}{(\text{m}^2 \cdot \text{s}^{-2})(\text{s})} \quad (7)$$

$$= \frac{\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}}{\text{m}^2 \cdot \text{s}^{-1}} = \text{kg} \quad (\text{satisfied}) \quad (8)$$

Frequency interpretation: If we substitute $f = \frac{1}{T}$:

$$m = \frac{hf}{c^2}$$

This is the Compton relation in inverse form. The Compton wavelength of a particle is:

$$\lambda_C = \frac{h}{mc}$$

Inserting the above relation $m = \frac{hf}{c^2}$, we get:

$$\lambda_C = \frac{h}{\left(\frac{hf}{c^2}\right)c} = \frac{c}{f}$$

This shows that the Compton wavelength corresponds to the wavelength of the oscillation that generates the mass.

What is new in the FFGF interpretation? Standard QFT says: Particles have a Compton wavelength based on their mass.

FFGF reverses it: The high-frequency oscillation in the fractal field generates the mass.

4. The Planck Scale as Effective Limit

Planck units (from $\hbar, G, c$):

$$\ell_P = \sqrt{\frac{\hbar G}{c^3}} \approx 1.616 \times 10^{-35} \text{ m} \quad (9)$$

$$t_P = \sqrt{\frac{\hbar G}{c^5}} \approx 5.391 \times 10^{-44} \text{ s} \quad (10)$$

$$m_P = \sqrt{\frac{\hbar c}{G}} \approx 2.176 \times 10^{-8} \text{ kg} \quad (11)$$

The speed of light from these:

$$c = \frac{\ell_P}{t_P} \approx 2.998 \times 10^8 \text{ m/s} \quad (\text{satisfied})$$

FFGF interpretation: In FFGF, these values are traced back to the geometry of the fractal lattice. The Planck length is the "lattice spacing" of the effective theory, the Planck time is the "tick" of the continuous description. Below this scale, the fundamental t_0 granulation operates (see above).

5. Vacuum Energy and the Cutoff by ξ

The catastrophe problem: The zero-point energy of a harmonic oscillator:

$$E_0 = \frac{1}{2} \hbar \omega$$

Summed over all modes up to the Planck frequency:

$$\rho_{\text{vac}} \sim \int_0^{\omega_P} \omega^{D_f} d\omega \sim \omega_P^{D_f+1}$$

For the present universe ($D_f = 3 - \xi \approx 3$):

$$\rho_{\text{vac}}|_{D_f \approx 3} \sim \omega_P^4 \sim \left(\frac{c}{\ell_P}\right)^4$$

[S] Special case: early universe ($t \rightarrow t_{\min}$, $D_f = 2$):

$$\rho_{\text{vac}}|_{D_f=2} \sim \omega_P^3 \sim \left(\frac{c}{\ell_P}\right)^3$$

This yields: $\rho_{\text{vac}} \sim 10^{113} \text{ J/m}^3$

Observed: $\rho_{\text{dark energy}} \sim 10^{-9} \text{ J/m}^3$

Discrepancy: Factor 10^{122} (The largest mismatch in physics)

FFGF approach with ξ **[S]**: In a fractal space with $D_f = 3 - \xi$, not all modes fit:

$$\rho_{\text{eff}} = \rho_{\text{Planck}} \times (\xi)^n$$

Where n is a scaling exponent. With $\xi \sim 10^{-4}$, one could indeed achieve a drastic suppression factor after multiple scaling (over 30 orders of magnitude from Planck to cosmological scale).

Mathematically:

$$(10^{-4})^{30} \sim 10^{-120}$$

This would be close to the observed order of magnitude; the exponent n is not derived here.

6. Gravitational Relationship (implied in the document)

Although not explicitly stated, FFGF suggests that gravity follows from geometry:

$$\text{Einstein: } R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \frac{8\pi G}{c^4}T_{\mu\nu}$$

FFGF would propose: Curvature arises from the local variation of D_f :

$$D_f(r) = 3 - \xi(r)$$

Where $\xi(r)$ depends on energy density. High mass density $\rightarrow$ larger $\xi \rightarrow$ stronger deviation from $D = 3 \rightarrow$ stronger "curvature".

.4 A Closer Look at the Mathematics of Torus Geometry (mentioned in the document)

Why the Torus?

In FFGF, the torus is chosen as the geometrically natural form for a self-sustaining energy flow in a fractal field.

Topological properties:

- Closed: No boundaries, energy can circulate endlessly
- Two independent circles: Poloidal (small) and toroidal (large) circulation
- Non-trivial topology: Genus value $g = 1$ (one "hole")

Mathematical Description of the Torus

Parametric equations:

$$x(\theta, \phi) = (R + r \cos \theta) \cos \phi \quad (12)$$

$$y(\theta, \phi) = (R + r \cos \theta) \sin \phi \quad (13)$$

$$z(\theta, \phi) = r \sin \theta \quad (14)$$

Where:

- R = Major radius (distance from center to tube center)
- r = Tube radius (thickness of the "tube")
- $\theta \in [0, 2\pi]$ = Poloidal angle (around the tube)
- $\phi \in [0, 2\pi]$ = Toroidal angle (around the main axis)

Geometric quantities:

- Surface area: $A = 4\pi^2 Rr$
- Volume: $V = 2\pi^2 Rr^2$
- Ratio: $\frac{V}{A} = \frac{r}{2}$

Important: the ratio depends only on the tube radius.

Curvature of the Torus

Gaussian curvature:

$$K(\theta) = \frac{\cos \theta}{r(R + r \cos \theta)}$$

Critical observation:

- On the inner side ($\theta = 0$): $K > 0$ (positive curvature, like a sphere)
- On the outer side ($\theta = \pi$): $K < 0$ (negative curvature, like a saddle)
- Top/bottom ($\theta = \pm\pi/2$): $K = 0$

The torus thus has regions with different curvature; this property is central to FFGF.

Energy Flow in the Torus (FFGF Model)

The document describes a poloidal and toroidal flow:

- Poloidal flow (θ -direction):
 - Energy flows through the "tube"
 - At the center: Contraction (inflow)
 - At the edge: Expansion (outflow)
- Toroidal flow (ϕ -direction):
 - Rotation around terleitherrleitung- main axis
 - Generates angular momentum
 - Stabilizes the structure

Vector field for energy flow:

$$\vec{v}(\theta, \phi) = v_\theta \vec{e}_\theta + v_\phi \vec{e}_\phi$$

Wheirleithe the velocities depend on local curvature.

Connection to $D_f = 3 - \xi$

The fractal dimension influences the torus structure:

In a perfect 3D space ($D = 3$), a torus could shrink to $r \rightarrow 0$ (singularity).

With $D_f = 3 - \xi$ there is a minimal tube radius:

$$r_{\min} \propto \frac{\ell_{\text{Planck}}}{\xi^{1/3}}$$

With $\xi = \frac{4}{3} \times 10^{-4}$:

$$r_{\min} \sim \frac{\ell_{\text{Planck}}}{(10^{-4})^{1/3}} \sim \ell_{\text{Planck}} \times 10^{4/3} \sim 21 \times \ell_{\text{Planck}}$$

Interpretation: The fractal structure prevents the torus from collapsing to a point. There is a natural lower limit.

[S] Special case: early universe ($t \rightarrow t_{\min}$, $D_f = 2$): For $D_f = 2$, the third spatial dimension is absent as a stabilizing reservoir. The minimal torus radius collapses to the Planck scale itself:

$$r_{\min}|_{D_f=2} \sim \ell_{\text{Planck}}$$

This is the absolute lower bound – the torus cannot be smaller than one Planck length in radius. The earliest universe thus had a maximally compact torus topology at $r \sim \ell_p$.

Mass from Torus Geometry

The FFGF thesis: A particle (e.g., a proton) is a high-frequency rotating torus on the Planck scale.

Angular momentum in the torus: For a rotating mass in the torus:

$$L = 2\pi^2 R r^2 \rho \omega$$

Where:

- ρ = Energy density
- ω = Rotation frequency

Mass from rotation: If we equate $E = mc^2$ with the rotational energy:

$$E_{\text{rot}} = \frac{1}{2} I \omega^2$$

For the torus, the moment of inertia is:

$$I = \pi^2 R r^2 \left(R^2 + \frac{3r^2}{4} \right) \rho$$

The relationship to time: With $\omega = \frac{2\pi}{T}$ and the previously derived relationship $m = \frac{h}{c^2 T}$:

$$T = \frac{h}{mc^2}$$

Inserting this for a proton ($m_p \approx 1.67 \times 10^{-27}$ kg):

$$T_p \approx \frac{6.6 \times 10^{-34}}{1.67 \times 10^{-27} \times 9 \times 10^{16}} \approx 4.4 \times 10^{-24} \text{ s}$$

This is the Compton time of the proton. In FFGF, the torus rotates with this frequency.

Scaling: From Proton to Galaxy

The fractal self-similarity means:

Scale	R (Major radius)	r (Tube)	Mass/System
Proton	$\sim 10^{-15} \text{ m}$	$\sim 10^{-16} \text{ m}$	$1.67 \times 10^{-27} \text{ kg}$
Atom	$\sim 10^{-10} \text{ m}$	$\sim 10^{-11} \text{ m}$	Electrons in orbitals
Planet	$\sim 10^6 \text{ m}$	$\sim 10^5 \text{ m}$	Magnetic field torus
Star	$\sim 10^9 \text{ m}$	$\sim 10^8 \text{ m}$	Convection currents
Galaxy	$\sim 10^{20} \text{ m}$	$\sim 10^{19} \text{ m}$	Spiral arms

The ratio R/r often remains constant (typically $R/r \approx 3 - 10$), which FFGF reads as an expression of self-similarity.

Why is the Torus Stable?

Energy minimum: The torus minimizes energy for a given volume and topology:

$$E_{\text{total}} = E_{\text{Surface}} + E_{\text{Curvature}} + E_{\text{Rotation}}$$

According to the FFGF assumption, the torus is the most stable form for certain boundary conditions (constant flux, angular momentum); the corresponding variational calculation is not carried out in this document [S].

In the fractal field: The dimension $D_f = 3 - \xi$ means energy experiences "resistance" when flowing. The torus is the path of least resistance for circulating energy.

Connection to the Schwarzschild Metric

Considering the Kerr metric (rotating black hole), one also finds a torus structure:

Ergosphere: The region around a rotating black hole where nothing can stand still has a toroidal form.

FFGF interprets this as a hint that the black hole is a torus on a larger scale [S].

.5 Connection Between Torus Topology and Quantum Numbers (Spin, Charge)

Topological Quantum Numbers from Torus Geometry – Detailed Derivation

FFGF and t_0 theory trace the fundamental quantum numbers of elementary particles (spin, electric charge, and color charge) back to the topological structure of the torus. The torus is considered a stable, natural geometric form for closed, self-consistent energy flows. All quantum numbers arise from the properties of closed flux lines that must wind on the torus surface or through the torus and close exactly to form stable configurations.

The central idea is that particles are not understood as point particles but as topologically stable vortex and flow structures in the fractally modified torus field. In this picture, quantization arises from the closure conditions of these flux lines – similar to quantized magnetic fluxes or the Aharonov-Bohm effect, but on a fundamental geometric level.

1. Spin – The Winding Number $w = n_\phi / n_\theta$

The spin of a particle corresponds to the **winding number** of the closed flux lines on the torus. This is defined as the ratio of revolutions in the two non-trivial directions of the torus:

$$w = \frac{n_\phi}{n_\theta} \quad (15)$$

where

- n_ϕ is the number of revolutions in the **toroidal direction** (around the major radius R),
- n_θ is the number of revolutions in the **poloidal direction** (around the tube radius r).

A flux line is only stable if it closes exactly after an integer number of windings. The simplest non-trivial closed orbits occur for rational values of w .

The physical assignment is:

- $w = 1$ (full revolution before closure) $\rightarrow$ **Boson spin** (integer: 0, 1, 2, ...)
- $w = 1/2$ (half revolution before closure) $\rightarrow$ **Fermion spin** (half-integer: 1/2, 3/2, ...)

This topological definition offers an intuitive interpretation of the spin-statistics theorem: Fermions require two half revolutions (720°) to return to the original state, while bosons are identical after 360°. The minimal winding number is limited by the stability condition $r_{\min} \approx 21 \ell_{\text{Planck}}$; smaller values lead to unstable configurations.

2. Electric Charge – Quantized Electric Flux Through the Torus

The electric charge directly correlates with the number of closed electric flux lines that **traverse** the torus (i.e., run from the inner to the outer region or vice versa).

The quantization condition is:

$$\Phi = n \cdot \frac{h}{e} \quad (16)$$

where

- Φ is the magnetic flux through a suitable cross-section of the torus,
- h is Planck's constant,
- e is the elementary charge,
- $n \in \mathbb{Z}$ is the integer number of traversing flux lines (positive or negative depending on direction).

Physical interpretation:

- $n = +1$ $\rightarrow$ Charge $+e$ (e.g., proton, positron)
- $n = -1$ $\rightarrow$ Charge $-e$ (e.g., electron)
- $n = 0$ $\rightarrow$ Electrically neutral (e.g., neutron, neutrino, photon)
- $n = +2, -2, \dots$ $\rightarrow$ Higher charges (possible in theory but energetically unfavorable or unstable on low scales)

The quantization is topologically protected because the torus has two non-contractible loops (toroidal and poloidal). The flux through these loops is invariant under continuous deformations – therefore the charge cannot vary continuously.

3. Color Charge – Topological Linking of Three Flux Strands

The color charge (quantum number of the strong interaction) arises from the **topological linking** of exactly **three flux strands** that wind around each other and around the torus. These three strands represent the three colors of QCD: red, green, blue.

The linking configuration determines the color properties:

- Three different colors (red–green–blue) in non-trivial linking $\rightarrow$ **Quark** (Color charge 1 in each color)
- Three identical colors (e.g., red–red–red) $\rightarrow$ **Antiquark** (Color charge -1 in each color)
- One color + its anticolor (e.g., red + antired) $\rightarrow$ **Gluon** (Color neutral but color-anticolor combination)
- All three colors simultaneously balanced (red + green + blue) $\rightarrow$ **Baryon** (Color overall white/neutral)

According to the theory, exactly **eight** non-trivial linking states of the three strands are possible (plus the trivial white state). These eight states are assigned to the **eight generators of SU(3) color symmetry**; a topological justification of the gauge group $SU(3)_C$ of the strong interaction is thereby indicated, but not carried out in this document [S].

Torus Geometry in Quantum Computing

The toroidal structure used in FFGF theory can be carried over to quantum information processing. In quantum computing applications (Quantum Computing in T0 Framework, 2025), the torus manifests through:

1. **Qubit State Space:** Qubits reside on the torus surface, with state described by position (z, r, θ) in local cylindrical coordinates.
2. **Local Approximation:** For single-qubit operations, the large toroidal radius R allows a cylindrical approximation:

$$R \gg r \Rightarrow \text{Torus} \approx \text{Cylinder locally}$$

3. **Global Topology:** Multi-qubit entanglement preserves the toroidal topology (Genus-1), enabling:
 - Charge quantization via flux through torus hole
 - Spin quantization via winding numbers
 - Topologically protected quantum information
4. **Bell Correlations:** The ξ -damping assumed for Bell tests in the T0 framework is traced back to the fractal modification of torus geometry.

Quantitative Example:

For a proton modeled as a torus:

$$R_{\text{proton}} \sim 10^{-15} \text{ m} \quad (\text{major radius}) \quad (17)$$

$$r_{\text{proton}} \sim 21\ell_P \approx 10^{-34} \text{ m} \quad (\text{tube radius}) \quad (18)$$

$$R/r \sim 10^{19} \quad (\text{aspect ratio}) \quad (19)$$

A qubit encoded in this structure experiences:

$$\text{Curvature correction} \sim \frac{r}{R} \sim 10^{-19} \ll \xi \sim 10^{-4}$$

Thus, the cylindrical approximation is valid for quantum gates, while the toroidal topology remains crucial for fundamental properties (charge, spin, entanglement structure).

.6 Torus Geometry in Cosmology – Scale-Invariant Torsional Structures

[S] This section belongs to the cosmic sector, which the present corpus deliberately leaves aside (P39); its statements are speculative.

A further aspect of the Fundamental Fractal-Geometric Field Theory (FFGF) and the t_0 theory is the assumption that torus geometry, beyond the Planck scale and the scale of elementary particles, can be continued ****self-similarly and scale-invariantly**** up to the largest observable cosmic structures.

The theory postulates that on every physical scale – from protons to stars and black holes to galaxies and the large-scale cosmic web – the dominant energy and momentum dynamics can be described by ****torsion-like, vortex-shaped flow structures**** that topologically correspond to a torus. These structures are characterized by the major radius R (toroidal great circle radius) and the tube radius r and are modified by the fractal dimension deficit ξ .

Cross-Scale Torsional Correspondences

The following overview summarizes the most important cosmological correspondences as described in the documents:

- **Elementary Particle Scale (Planck to Hadron scale)**
 $R \sim 10^{-15}$ m (proton radius), $r \sim 10^{-16}$ m to $21 \ell_p$
 Stabilized energy vortex ("mass torus") with Compton frequency.
 Poloidal and toroidal flows generate rest mass, spin, and internal quantum numbers.
 Primary source: 006_T0_Teilchenmassen_En.pdf
- **Star and Black Hole Scale**
 $R \approx$ Schwarzschild radius $r_S = 2GM/c^2$
 Rotating spacetime vortex corresponding to the Kerr metric.
 The accretion disk and the ergosphere together form a macroscopic torus in which kinetic energy, angular momentum, and gravitational binding energy circulate.
 The torus is meant to stabilize the extreme rotational and gravitational fields and to explain the existence of stable rotating black holes without additional exotic matter.
 Primary source: T0_Kosmologie.pdf
- **Galactic Scale**
 $R \sim 10^{20}$ m (typical radius of the bulge / central region)
 $r \sim 10^{19}$ m (effective thickness of the galactic disk)
 Large-scale filamentary vortices in the cosmic web.
 The spiral arms are interpreted as standing density waves within a torsional base structure.
 The total galactic angular momentum ensures long-term stabilization of the torus configuration.
 The flat rotation curve and observed distribution of star velocities are meant to arise geometrically from the fractal modification of torus volume and curvature distribution – without additional dark matter.
 Primary sources: T0_Kosmologie.pdf, 145_FFGFT_donat-teil1_En.pdf
- **Cosmological Large Structure Scale (cosmic web, filaments, void structures)**
 $R \sim 10^{23}$ – 10^{24} m (order of magnitude of the largest observed filaments and superclusters)
 $r \sim 10^{22}$ – 10^{23} m (thickness of filaments)
 The cosmic web is interpreted as a hierarchical system of nested torsion-like vortices.
 The large-scale structures (filaments, walls, voids) correspond to the stable nodes and empty spaces of a huge, fractally modulated torus network.
 The observed anisotropy (e.g., CMB dipole, Hubble tension, large-scale flows) is interpreted as a consequence of asymmetric torsional flow dynamics – without cosmic expansion or Λ CDM parameters.
 Primary sources: 039_Zwei-Dipole-CMB_En.pdf, T0_Kosmologie.pdf

Core Principle: Scale Invariance and Fractal Self-Similarity

The torus geometry is ****scale-invariant**** in FFGF/ t_0 theory:

$$\frac{R}{r} \approx \text{constant} \quad \text{over many orders of magnitude}$$

(typical values range between 5 and 50, depending on the scale considered).

The fractal dimension deficit $\xi = 4/3 \times 10^{-4}$ ensures that the effective geometric quantities (surface area A_{frak} , volume V_{frak} , curvature K_{frak}) are consistently modified on every scale – with which the theory aims at a unified description from micro- to macrocosm.

Cosmological Implications – Without Dark Matter and Without Expansion

The theory puts forward the following theses **[S]**:

- Galaxy rotation curves arise purely from fractal-torsional geometry (no additional invisible mass needed).
- The Hubble tension (discrepancy between local and CMB-based H_0) is treated in Document 026.
- The CMB dipole and large-scale flows are manifestations of a global torsional flow ("Two-Dipole Model").
- The universe is static on the largest scale – expansion is not necessary.

These theses are treated in more detail in:

- T0_Kosmologie.pdf
- 145_FFGFT_donat-teil1_En.pdf
- 039_Zwei-Dipole-CMB_En.pdf

Torus cosmology thus represents an attempt to derive the entire hierarchy of cosmic structures from a single geometric basic form (the fractally modified torus) – an approach that consciously distinguishes itself from the metric-dynamic description of General Relativity.

Two-Dipole Model in Detail

The Two-Dipole Model is a central element of the Fundamental Fractal-Geometric Field Theory (FFGF) and the t_0 theory, developed to describe anomalies in the Cosmic Microwave Background radiation (CMB). It is presented in the repository documents as a geometric approach that interprets the observed CMB dipole without cosmic expansion or dark energy **[S]**. Instead, the dipole is interpreted as a manifestation of two superimposed torsional flows arising from the fractal torus structure of spacetime. The detailed derivations are found primarily in 039_Zwei-Dipole-CMB_En.pdf, supplemented by cosmological sections in T0_Kosmologie.pdf and 145_FFGFT_donat-teil1_En.pdf.

Introduction and Motivation

The standard Λ CDM model interprets the CMB dipole (a temperature anisotropy of $\Delta T/T \approx 10^{-3}$) primarily as a kinematic effect due to the peculiar motion of the Milky Way relative to the CMB rest frame (with $v \approx 370$ km/s). However, there are persistent discrepancies: The dipole appears stronger and more asymmetric than expected and does not perfectly correspond to large-scale flows (e.g., Shapley Attractor, Laniakea Supercluster). Additionally, the dipole contributes to the Hubble tension (H_0 discrepancy between local and CMB-based measurements of about 5σ).

The Two-Dipole Model addresses these problems by modeling the dipole as the superposition of **two geometric components**:

- **Kinematic dipole:** Local motion effects (similar to the standard model).
- **Intrinsic geometric dipole:** Fractal-torsional asymmetry of spacetime itself arising from the ξ -modified torus structure.

This approach leads to a static universe where apparent expansion effects are geometric – without a Big Bang or dark energy.

Model Description

The model is based on the assumption that spacetime on a cosmic scale possesses a **global torsional structure** that is self-similar to torus geometry on smaller scales (elementary particles, black holes, galaxies). The CMB dipole arises from two superimposed poles:

1. **Local dipole**: Generated by the motion of the Local Group (Milky Way) in a torsional flow field. This corresponds to the standard dipole but modified by fractal corrections.

2. **Global dipole**: An intrinsic effect of fractal spacetime resulting from the asymmetry of the cosmic torus network. The global flow is scale-invariant and connects the Planck scale (ℓ_p) with the Hubble scale (c/H_0).

The superposition of the two dipoles is meant to explain the observed asymmetries: The local dipole dominates on small scales, while the global one becomes visible on large scales (e.g., in CMB multipoles).

Mathematical Framework

The dipole moment is modeled as a vector sum:

$$\vec{D}_{\text{total}} = \vec{D}_{\text{kin}} + \vec{D}_{\text{geo}} \quad (20)$$

- **Kinematic dipole** $\vec{D}_{\text{kin}}$:

$$\Delta T(\hat{n}) = T_0 \frac{\vec{v} \cdot \hat{n}}{c} \Rightarrow D_{\text{kin}} \approx 3.35 \text{ mK}$$

(with $T_0 \approx 2.725 \text{ K}$, $v \approx 370 \text{ km/s}$, $\hat{n}$ line of sight).

- **Geometric dipole** $\vec{D}_{\text{geo}}$: It arises from the fractal modification of the spacetime metric:

$$D_{\text{geo}} \sim \xi \cdot \ln\left(\frac{L_{\text{Hubble}}}{\ell_p}\right) \cdot T_0 \approx 0.1 \text{ mK}$$

where $\xi = 4/3 \times 10^{-4}$ is the dimension deficit, and the logarithm accounts for the scale hierarchy over ~ 60 orders of magnitude.

The direction of the global dipole aligns with the axis of the cosmic torus flow, deviating from the galactic dipole by $\sim 48^\circ$ – with which the model interprets the observed misalignment.

$$H_0 \approx 66.2 \text{ km/s/Mpc} \quad (\text{see Document 026})$$

This value is calibrated to the measured H_0 (-1.9% deviation from Planck). The exponent $41/4$ requires justification from ξ -field theory.

Cosmological Implications

- **Interpretation of the Hubble tension** **[S]**: Local measurements ($H_0 \approx 73 \text{ km/s/Mpc}$) see the kinematic dipole, CMB measurements ($H_0 \approx 67 \text{ km/s/Mpc}$) see the geometric one – the discrepancy arises from the superposition.

- **Static universe** **[S]**: No expansion needed; redshift z results from fractal energy loss:

$$z \approx \xi \cdot \ln(d/\ell_p)$$

(with d distance).

- **CMB anomalies** **[S]**: The model interprets the dipole, quadrupole weakness, and hemispherical asymmetry as torsional effects.

- **Quantitative predictions** **[S]**: Dipole amplitude $\Delta T \approx 3.36 \text{ mK}$ (consistent with Planck data), misalignment angle 48° (consistent with observations).

Critical Analysis

The model describes several anomalies geometrically, with ξ as the only parameter (plus the settings named). However, a formal derivation from field equations is lacking (compared to standard cosmology). Experimental validation is pending; it departs from the Λ CDM model. Further details are in the sources.

Parallel to the Toroidal Photon Model (Williamson & van der Mark, 1997)

Since 1997, an independent semi-classical approach has existed in the literature describing the electron as a circulating, topologically closed photonic entity with toroidal character. The original paper is titled:

Is the electron a photon with toroidal topology?

J. G. Williamson and M. B. van der Mark

Annales de la Fondation Louis de Broglie, Vol. 22, No. 2, 1997, pp. 133–167

The full text is freely available online at:

https://fondationlouisdebroglie.org/IMG/pdf/22_2_133.pdf

A clear and didactically prepared popular-science explanation of this model can be found in the following video:

Is the Electron a Photon with Toroidal Topology?

YouTube video by *Physics Explained* (2021)

<https://www.youtube.com/watch?v=hYyrgDEJL0A>

Although this model was developed independently of the FFGF/ t_0 theory, it exhibits structural parallels to the toroidal geometry presented here — especially in the derivation of charge, spin, and magnetic moment from a closed, double-loop field configuration.

Key Parallels to the FFGF Torus Structure

• Torus Topology and Double Loop

In the referenced model, a circularly polarized electromagnetic field of exactly one Compton wavelength λ_C is folded into a closed double loop (double helix / double loop). This corresponds to the toroidal + poloidal circulation postulated in the FFGF: energy flows both toroidally (ϕ -direction, large circle) and poloidally (θ -direction, around the tube). The double circulation (4π instead of 2π) leads — in both approaches — to half-integer spin ($w = 1/2$ in the FFGF winding-number definition).

• Electric Field and Charge as Topological Property

In the toroidal model, the electric field vector consistently points inward on the outside (electron) or outward (positron) because field rotation is commensurate with the geometry. This is structurally similar to the FFGF derivation: electric charge arises from the quantized number of closed electric flux lines threading the torus ($\Phi = n \cdot h/e$). The direction (inward/outward) is topologically fixed and reflects the orientation of the poloidal/toroidal flux components.

• Magnetic Moment from Toroidal Magnetic Field Configuration

Both approaches derive the magnetic dipole moment from closed magnetic field lines running parallel to the torus surface (toroidal B_ϕ field in FFGF). The net moment along the torus axis arises from the asymmetry of the internal rotation — analogously, in the FFGF the intrinsic magnetic moment of the electron ($\mu_e = e\hbar/(2m_e)$) follows from rotational energy in the torus.

- **Compton Scale as Intrinsic Size**

In the external model, the Compton wavelength $\lambda_C = h/(m_e c)$ determines the length of the closed path and thus the effective size of the object ($\sim \lambda_C/(4\pi)$ for the core radius). This agrees with the FFGF, where the Compton time $T = h/(mc^2)$ sets the fundamental rotation period of the torus and the minimal stable tube radius $r_{\min} \sim 21 \ell_P$ is limited by the fractal correction ξ . Both approaches thereby avoid the infinite self-energy of a point particle.

- **Two Chiral Spin States**

The toroidal model distinguishes two non-superimposable chiral variants (handedness) that only return to themselves after 720° rotation — analogously, in the FFGF spin-1/2 arises from the winding number $w = n_\phi/n_\theta = 1/2$ and fermions require two full rotations to return to the original state.

Differences and Extension by the FFGF

While the 1997 model remains semi-classical and leaves the self-confinement mechanisms (nonlinear effects, topological stability) largely open, the FFGF/ t_0 theory offers a further-reaching attempt at a justification:

- The fractal dimension modification $D_f = 3 - \xi$ is meant to prevent collapse below $r_{\min} \approx 21 \ell_P$ and to explain stability without additional nonlinear vacuum effects.
- Energy flow is explicitly poloidal + toroidal and fractally modulated ($\vec{v}(\theta, \phi)$ depending on local curvature $K(\theta)$).
- Quantum numbers (including color charge) are assigned topologically to linking numbers and winding numbers — a generalization that extends beyond the pure electron model.
- Mass is traced back to the inertia of the inner T_0 -scale flow ($m = h/(c^2 T)$ with T as Compton time), not only to confined field energy.

The parallels are a classification made by this document, not a statement by Williamson and van der Mark.

.7 Electromagnetic Fields in Torus Geometry

Maxwell's Equations on the Torus

In curved coordinates, Maxwell's equations must be adapted:

In torus coordinates (θ, ϕ, ψ) :

$$\nabla \times \vec{E} = -\frac{\partial \vec{B}}{\partial t} \quad (21)$$

$$\nabla \times \vec{B} = \mu_0 \vec{j} + \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t} \quad (22)$$

$$\nabla \cdot \vec{E} = \frac{\rho}{\epsilon_0} \quad (23)$$

$$\nabla \cdot \vec{B} = 0 \quad (24)$$

The nabla operator in torus coordinates is more complex:

$$\nabla = \frac{1}{h_\theta} \frac{\partial}{\partial \theta} \vec{e}_\theta + \frac{1}{h_\phi} \frac{\partial}{\partial \phi} \vec{e}_\phi + \frac{1}{h_\psi} \frac{\partial}{\partial \psi} \vec{e}_\psi$$

Where h_θ, h_ϕ, h_ψ are the metric factors.

Magnetic Field Configuration in the Torus

- Poloidal magnetic field B_θ : Runs around the tube. Arises from toroidal currents.
- Toroidal magnetic field B_ϕ : Runs around the main axis. Arises from poloidal currents.

The total field configuration:

$$\vec{B} = B_\theta(r, \theta)\vec{e}_\theta + B_\phi(r, \theta)\vec{e}_\phi$$

Stability Condition (Kruskal-Shafranov)

For a stable torus plasma (as in fusion reactors) it must hold:

$$q = \frac{rB_\phi}{RB_\theta} > 1$$

This is the safety factor q .

In FFGF it is assumed that elementary particles are stable because their torus configuration satisfies $q > 1$.

Origin of the Magnetic Moment

A rotating torus with charge generates a magnetic dipole moment:

$$\mu = I \times A = \left(\frac{Q}{T}\right) \times \pi r^2$$

Where:

- Q = Charge
- T = Rotation period
- r = Tube radius

For an electron:

$$\mu_e = \frac{e\hbar}{2m_e} = \text{Bohr magneton}$$

This is the intrinsic magnetic moment of the electron.

Electromagnetic Self-Energy

The energy stored in the electromagnetic field of a torus:

$$E_{\text{em}} = \frac{\epsilon_0}{2} \int E^2 dV + \frac{1}{2\mu_0} \int B^2 dV$$

For a torus with radius R and r :

$$E_{\text{em}} \propto \frac{e^2}{r} \times f\left(\frac{R}{r}\right)$$

Where $f(R/r)$ is a geometric factor.

This energy contributes to mass:

$$m_{\text{em}} = \frac{E_{\text{em}}}{c^2}$$

A portion of the electron mass ($\sim 0.1\%$) stems from this electromagnetic self-energy.

Connection to ξ and D_f

In a D_f -dimensional space, the general Coulomb force reads:

$$F \propto \frac{1}{r^{D_f-1}}$$

Standard physics ($D = 3$):

$$F \propto \frac{1}{r^2}$$

Present universe, fractal space ($D_f = 3 - \xi$):

$$F \propto \frac{1}{r^{2-\xi}}$$

For $\xi = \frac{4}{3} \times 10^{-4}$:

$$F \propto \frac{1}{r^{1.9998667...}}$$

On large scales, this leads to a tiny modification that could explain "dark energy" effects [S].

[S] Special case: early universe ($t \rightarrow t_{\min}$, $D_f = 2$):

$$F|_{D_f=2} \propto \frac{1}{r}$$

At the beginning of the universe there was no $1/r^2$ law, but a linear confinement potential. This is physically equivalent to the 2D Coulomb potential and had fundamental consequences for structure formation in the Planck era.

.8 Fluid Dynamics in the Torus (Navier-Stokes on Curved Spaces)

Navier-Stokes in Curved Coordinates

The Navier-Stokes equations describe the flow of fluids (or in FFGF: the dynamics of the vacuum "fluid").

Standard form:

$$\rho \left(\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \nabla) \vec{v} \right) = -\nabla p + \eta \nabla^2 \vec{v} + \vec{f}$$

In torus coordinates: we must use the covariant derivative:

$$\rho \left(\frac{\partial v^i}{\partial t} + v^j \nabla_j v^i \right) = -\nabla^i p + \eta g^{ij} \nabla_j \nabla_k v^k + f^i$$

Where:

- g^{ij} = Metric tensor
- ∇_j = Covariant derivative
- η = Viscosity of the vacuum medium

Metric Tensor for the Torus

For a torus in standard position:

$$ds^2 = d\theta^2 + (R + r \cos \theta)^2 d\phi^2$$

Metric tensor:

$$g = \begin{bmatrix} 1 & 0 \\ 0 & (R + r \cos \theta)^2 \end{bmatrix}$$

Determinant:

$$\sqrt{g} = R + r \cos \theta$$

Velocity Field in the Rotating Torus

Assumption: Steady rotation with constant angular velocity ω .

Poloidal component:

$$v_\theta(r, \theta) = v_0 \sin(n\theta)$$

Where n is the number of vortices.

Toroidal component:

$$v_\phi(r, \theta) = \omega(R + r \cos \theta)$$

Vorticity

The vorticity is:

$$\vec{\omega} = \nabla \times \vec{v}$$

In torus coordinates:

$$\omega_r = \frac{1}{h_\theta h_\phi} \left[\frac{\partial(h_\phi v_\phi)}{\partial \theta} - \frac{\partial(h_\theta v_\theta)}{\partial \phi} \right]$$

For a stable torus vortex: The vorticity must remain positive everywhere (no backflows).

Energy Conservation in Torus Flow

The kinetic energy of the flow:

$$E_{\text{kin}} = \frac{\rho}{2} \int v^2 dV$$

For a torus:

$$E_{\text{kin}} = \frac{\rho}{2} \times 2\pi^2 R r \times \langle v^2 \rangle$$

Dissipation due to viscosity:

$$\frac{dE}{dt} = -\eta \int (\nabla \times \vec{v})^2 dV$$

Equilibrium: If energy input (through vacuum fluctuations on the Planck scale) balances dissipation, the torus is stable.

Turbulence and Stability

The Reynolds number for a torus:

$$Re = \frac{\rho v R}{\eta}$$

Critical value: $Re_{\text{crit}} \approx 2300$

For $Re < Re_{\text{crit}}$: Laminar flow (stable)

For $Re > Re_{\text{crit}}$: Turbulent flow (unstable)

In FFGF: The "viscosity" η of the vacuum is determined by ξ :

$$\eta \propto \frac{\hbar}{\ell_{\text{Planck}}^3 \times \xi}$$

With $\xi = \frac{4}{3} \times 10^{-4}$ results in a very low viscosity $\rightarrow$ the vacuum behaves like a superfluid.

Helmholtz Decomposition

Any vector field can be decomposed into:

$$\vec{v} = \nabla\varphi + \nabla \times \vec{A}$$

- Potential part ($\nabla\varphi$): Compressible flow
- Vortex part ($\nabla \times \vec{A}$): Incompressible rotation

In the torus, the vortex part dominates; FFGF sees this as the reason for stability.

Casimir Effect in the Torus

In a D_f -dimensional space, the Casimir pressure scales generally as:

$$P_{\text{Casimir}} \propto -\frac{\hbar c}{d^{D_f+1}}$$

Present universe ($D_f \approx 3$):

$$P_{\text{Casimir}} = -\frac{\pi^2 \hbar c}{240 d^4}$$

Where d is the distance (here: tube radius $2r$).

In FFGF, this pressure is meant to stabilize the torus against collapse.

[S] Special case: early universe ($t \rightarrow t_{\text{min}}$, $D_f = 2$):

$$P_{\text{Casimir}}|_{D_f=2} \propto -\frac{\hbar c}{d^3}$$

The weaker distance exponent (d^{-3} instead of d^{-4}) means that Casimir pressure in the Planck era acted more strongly at large separations and more weakly at small separations – with direct consequences for torus stability at the beginning of the universe.

Connection to Time-Mass Duality

The effective flow velocity in the torus on the Planck scale is:

$$v \sim \frac{\ell_{\text{Planck}}}{t_p} = c$$

This corresponds to the speed of light and is compatible with the interpretation that c emerges as an effective velocity from the Planck scale.

On the fundamental t_0 scale (sub-Planck), however:

$$v_0 \sim \frac{\Lambda_0}{t_0} = \frac{\xi \cdot \ell_{\text{Planck}}}{t_0}$$

where t_0 is the sub-Planck time (2GE). Mass arises from the inertia of this internal flow at the t_0 granulation level.

Clarification: Effective Planck Scale vs. Fundamental t_0 Scale

To avoid confusion: In this analysis, the **effective limit** of continuous physics is described by the **Planck length ℓ_P** and **Planck time t_P** . The minimal stable torus tube is at $r_{\min} \approx 21\ell_P$, i.e., significantly above ℓ_P .

The **fundamental t_0 scale**, however, is **sub-Planck** and describes the internal granulation of the fractal field:

- Sub-Planck length: $\Lambda_0 = \xi \cdot \ell_P \approx 1.333 \times 10^{-4} \cdot \ell_P \approx 2.15 \times 10^{-39} \text{ m}$
- Characteristic t_0 lengths and times: $r_0 = 2GE$, $t_0 = 2GE$ (see Zeit_En.pdf and 010_T0_Energie_En.pdf)

The Planck scale is thus the **outer reference limit** of the effective theory, while t_0 represents the **sub-Planck granulation** on which the fractal structure truly operates.

Fractal Turbulence

In a general D_f -dimensional space, the turbulence energy spectrum follows:

$$E(k) \propto k^{-(D_f+2)/3}$$

Kolmogorov spectrum ($D = 3$):

$$E(k) \propto k^{-5/3}$$

Present universe, fractal spectrum ($D_f = 3 - \xi$):

$$E(k) \propto k^{-(5/3-\xi/3)}$$

This could be measurable in cosmic plasma structures.

[S] Special case: early universe ($t \rightarrow t_{\min}$, $D_f = 2$):

$$E(k)|_{D_f=2} \propto k^{-4/3}$$

The shallower exponent $-4/3$ means that in the Planck era energy cascaded less efficiently to small scales – a clear difference from present-day cosmic turbulence.

.9 Overall Synthesis: The Three Aspects Together

- Fluid dynamics generates stable vortices (torus form)
- Electromagnetic fields arise from the rotation of charged vortices
- Quantum numbers are topological properties of linking

Everything is connected through:

- The fractal dimension $D_f = 3 - \xi$
- The Planck time t_0 as fundamental rhythm
- The torus geometry as the most stable form (as assumed in FFGF)