

The Zeta Function of the T^4/Z_3 Torus:

Orbifold Projection, Dirichlet L-Functions and
Sector Pairing of the $GF(27)$ Classes

Johann Pascher
ORCID 0009-0000-6518-4064

2 September 2026

Contents

The Zeta Function of the T^4/Z_3 Torus	2
.1 Status markers and starting point	1
.2 Theorem A: Orbifold projection and Dirichlet L-function	1
Decomposition of the zeta by Z_3 classes	1
Principal character $\chi_0 \bmod 3$	1
Non-trivial character $\chi_1 \bmod 3$	1
.3 Theorem B: Hierarchical Euler product by $k = \text{ord}_p(3)$	2
Fractal suppression of higher Euler factors	2
.4 Theorem C: New zeros from the orbifold projection	2
Zeros of $(1 - 3^{-s})$	2
Complete zero structure of the orbifold zeta	3
.5 Theorem D: The four classes under sector pairing	3
The Galois character does not distinguish the classes	3
Sector pairing reduces four to two	3
Negation couples order 26 to order 13	3
Open question	4
.6 Theorem E: Prime patterns in the $GF(3^k)$ hierarchy	4
The harmonic primes are uniquely forced	4
Necessary congruence condition	4
.7 Theorem F: Limits of distinguishability and computational cost	4
Exact threshold: Euler factor vs. fractal correction	4
Galois products vs. random products: hit rate	5
Structural computational limit: Weyl obstruction and quantum computers	5
.8 In plain terms: what the limit means	6
.9 In plain terms II: synthesis, analysis, factorisation — and coupling	6
Synthesis is unlimited, analysis is limited	6
Factorisation is analysis	7
Coupling: the same threefold division, not the same law	7
FFGFT is dynamically deeply subcritical	8
.10 Summary	8

The Zeta Function of the T^4/Z_3 Torus

Abstract

The Galois factorisation classes of Doc. 342 and the harmonic series on the T^4 torus of Doc. 159 together yield a precise statement about the zeta function of the FFGFT spacetime.

Four results: **(A)** The Z_3 orbifold projection of T^4/Z_3 transforms the full Riemann zeta into a Dirichlet L-function: $\zeta_{T^4/Z_3}(s) = L(s, \chi_0) = (1 - 3^{-s})\zeta(s)$ **[B]**. **(B)** The Euler product of this L-function is hierarchically ordered by the entry depth $k = \text{ord}_p(3)$; the physically visible primes $\{2, 3, 5, 7, 11, 13\}$ are exactly those with $k \leq 6$ **[B]**. **(C)** The orbifold projection generates additional zeros at $s = 2\pi i k / \ln 3$ ($k \in \mathbb{Z} \setminus \{0\}$) on the imaginary axis $\text{Re}(s) = 0$; these are the algebraic signature of the Z_3 sector separation **[B]**. **(D)** The four Frobenius conjugacy classes of primitive elements in $\text{GF}(27)^*$ (Doc. 342, Theorem A) are paired into two classes by the FFGFT sector pairing $k \leftrightarrow -k$ (Doc. 336; on $\text{GF}(27)^*$: $x \mapsto x^{-1}$), $f_1 \leftrightarrow f_4$ and $f_2 \leftrightarrow f_3$; negation $x \mapsto -x = x^{13}x$ couples each primitive class to exactly one order-13 class **[B]**. The Galois character of $\text{Gal}(\text{GF}(27)/\text{GF}(3)) \cong \mathbb{Z}_3$ does not distinguish the classes; the distinction lies in $(\mathbb{Z}/26)^*/\langle 3 \rangle \cong \mathbb{Z}_4$. Whether exactly one second $\text{GF}(27)^*$ layer besides the lepton layer is physically realised remains open **[S]**.

Two further results: **(E)** The harmonic primes $\{5, 7, 11, 13\}$ are uniquely forced by the prime factorisations of $3^k - 1$ for $k = 3, 4, 5, 6$; $p = 13$ is the worldwide unique prime with $\text{ord}_p(3) = 3$ **[B]**. **(F)** The Euler factor threshold at $p^* \approx 9.76$ ($s = 2$) marks the 7-limit exactly; from $p_{\max} \approx 30$ the Galois product density lies at chance level; the Weyl obstruction (Doc. 316) shows that the limit lies in the object — no quantum computer of any size circumvents it **[B]**.

Check scripts: `pruef_343_zeta_galois.py`, `pruef_343b_klassen_symmetrie.py`, `pruef_344_primzahlmuster.py`.

.1 Status markers and starting point

The following status markers are used throughout:

- [K]** *Confirmed* — fully proved from ξ or a corpus-established derivation; all check-script assertions passed.
- [B]** *Established* — numerically secured; analytical proof present or trivial.
- [S]** *Speculative* — indication or conjecture, not yet fully proved.

Doc. 159 shows that the spectral zeta function of the Laplace operator on a torus T^d is the Epstein zeta function, whose one-dimensional special case is the Riemann zeta $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$. Doc. 342 shows that primes appear in the $\text{GF}(3^k)$ hierarchy of the T^4/Z_3 orbifold ordered by entry depth $k = \text{ord}_p(3)$. This document combines both: what does the $\text{GF}(3^k)$ hierarchy imply for the zeta function itself?

.2 Theorem A: Orbifold projection and Dirichlet L-function

Decomposition of the zeta by Z_3 classes

On T^4/Z_3 the allowed modes $n \in \mathbb{Z}^4$ are structured by Z_3 invariance. In the one-dimensional analogue $\zeta(s)$ splits into three partial sums by $n \bmod 3$ **[B]**:

$$\zeta(s) = \underbrace{\sum_{n=1}^{\infty} \frac{1}{n^s}}_{=3^{-s}\zeta(s)} + \underbrace{\sum_{\substack{n=1 \\ 3 \nmid n}}^{\infty} \frac{1}{n^s}}_{L_1(s)} + \underbrace{\sum_{\substack{n=1 \\ n \equiv 2(3)}}^{\infty} \frac{1}{n^s}}_{L_2(s)} \quad (1)$$

The fixed-point sector ($n \equiv 0$) contributes exactly $3^{-s}\zeta(s)$; the twist sectors ($n \equiv \pm 1$) together $(1 - 3^{-s})\zeta(s)$.

Principal character $\chi_0 \bmod 3$

The Dirichlet character $\chi_0 \bmod 3$ with $\chi_0(n) = 0$ if $3 \mid n$, otherwise $\chi_0(n) = 1$, gives **[B]**:

$$L(s, \chi_0) = \sum_{\gcd(n,3)=1} \frac{1}{n^s} = (1 - 3^{-s})\zeta(s) = \prod_{p \neq 3} \frac{1}{1 - p^{-s}} \quad (2)$$

This is the **spectral zeta function of the T^4/Z_3 torus**: the characteristic prime $p = 3$ does not appear as an Euler factor — it is the geometric foundation of the orbifold and stands outside the prime hierarchy.

Non-trivial character $\chi_1 \bmod 3$

The character $\chi_1 \bmod 3$ with $\chi_1(1) = +1$, $\chi_1(2) = -1$, $\chi_1(0) = 0$ measures the *asymmetry* of the twist sectors **[B]**:

$$L(s, \chi_1) = \sum_{\substack{n=1 \\ n \equiv 1(3)}}^{\infty} \frac{1}{n^s} - \sum_{\substack{n=1 \\ n \equiv 2(3)}}^{\infty} \frac{1}{n^s} \quad (3)$$

with the exact value (Dirichlet 1837): $L(1, \chi_1) = \pi/(3\sqrt{3})$.

Physical meaning **[S]**: $L(s, \chi_1)$ measures the difference between winding numbers $w = +1$ and $w = -1$ on the torus, i.e. the Z_3 sector asymmetry. Whether $L(s, \chi_1)$ has a direct physical observable in FFGFT (particle–antiparticle asymmetry in the twisted sector?) remains open **[S]**.

.3 Theorem B: Hierarchical Euler product by $k = \text{ord}_p(3)$

The primes in $L(s, \chi_0) = \prod_{p \neq 3} (1 - p^{-s})^{-1}$ are not equivalent. By the entry-depth rule $k = \text{ord}_p(3)$ from Doc. 342, Theorem B, the Euler product is hierarchically ordered **[B]**:

$$L(s, \chi_0) = \prod_{k=1}^{\infty} \prod_{\substack{p \neq 3 \\ \text{ord}_p(3)=k}} \frac{1}{1 - p^{-s}} \quad (4)$$

Table 1: Hierarchy of Euler factors by $k = \text{ord}_p(3)$

k	new prime(s)	Euler factor at $s = 2$	harmonics	FFGFT role
1	2	$4/3 = 1.333$	octave	Z_2 fixed point, massive sector
3	13	≈ 1.006	thirteenth	$GF(27)^*$, leptons
4	5	≈ 1.042	major third	factor in $(m_\mu/m_e)^2$
5	11	≈ 1.008	eleventh	neutrinos (Doc. 340)
6	7	≈ 1.021	seventh	top quark (Doc. 289)
≥ 7	$\geq 41, 1093, \dots$	$\approx 1 + p^{-2}$	no small harmonics	suppressed

Fractal suppression of higher Euler factors

On the fractal T^4 torus (Doc. 159) high-mode contributions are suppressed by the damping factor $\tau^{-(1+\xi)k}$. This transfers to the Euler factors; the *weighted* Euler product reads

$$\zeta_{T^4/Z_3}^{\text{frac}}(s) \cong \prod_{k=1}^{\infty} \prod_{\substack{p \neq 3 \\ \text{ord}_p(3)=k}} \left(\frac{1}{1 - p^{-s}} \right)^{\tau^{-k(1+\xi)}} \quad (5)$$

Since $\tau^{-k(1+\xi)} \rightarrow 0$ as $k \rightarrow \infty$, each Euler factor is pushed toward 1 for large k : primes with $\text{ord}_p(3) \geq 7$ make no physical contribution **[B]**. This is Theorem D of Doc. 342 in the language of the zeta function.

.4 Theorem C: New zeros from the orbifold projection

Zeros of $(1 - 3^{-s})$

The factor $(1 - 3^{-s})$ in $L(s, \chi_0) = (1 - 3^{-s})\zeta(s)$ has zeros at **[B]**:

$$3^{-s} = 1 \quad \Longleftrightarrow \quad s = \frac{2\pi i k}{\ln 3}, \quad k \in \mathbb{Z} \setminus \{0\} \quad (6)$$

These lie on $\text{Re}(s) = 0$ (imaginary axis), *not* on the critical line $\text{Re}(s) = 1/2$ of the Riemann hypothesis. They are absent from the Riemann zeta spectrum; they arise solely from the Z_3 orbifold projection.

The zero spacing $2\pi/\ln 3 \approx 5.72$ encodes the periodicity of the Z_3 lattice structure. A resonance test of the first eight known Riemann zeros γ_n against this unit shows no significant coincidence (deviations 0.15–0.47) **[B]**: the full Riemann zeta encodes the distribution of *all* primes, not only the Z_3 -compatible ones.

Complete zero structure of the orbifold zeta

$L(s, \chi_0)$ has three types of zeros:

- **Trivial zeros of $\zeta(s)$** : at $s = -2, -4, -6, \dots$
- **Non-trivial Riemann zeros**: on $\text{Re}(s) = 1/2$ (assumed), unchanged
- **Orbifold zeros**: at $s = 2\pi ik/\ln 3, k \neq 0$ — new, generated by the Z_3 projection **[B]**

.5 Theorem D: The four classes under sector pairing

The Galois character does not distinguish the classes

The Galois group $\text{Gal}(\text{GF}(27)/\text{GF}(3)) = \langle F \rangle \cong \mathbb{Z}_3$ with $F: x \mapsto x^3$ has three characters $\chi_j(F) = \omega^j$, $j = 0, 1, 2$. All four primitive classes $f_1, \dots, f_4$ of Doc. 342 are *regular* F -orbits of length 3: F acts on each as a 3-cycle, and the associated character is the same for all four. The Galois character therefore does not separate the classes **[B]**.

The distinction lies in the discrete logarithm: with a generator x of $\text{GF}(27)^*$, each class is an orbit $\{x^k, x^{3k}, x^{9k}\}$ with $\gcd(k, 26) = 1$, i.e. a coset of $\langle 3 \rangle = \{1, 3, 9\}$ in $(\mathbb{Z}/26)^* \cong \mathbb{Z}_{12}$. There are $12/3 = 4$ cosets **[B]**:

Table 2: The four primitive classes as cosets of $\langle 3 \rangle$ in $(\mathbb{Z}/26)^*$ (basis $x^3 + 2x + 1 = 0$)

Class	Minimal polynomial	Exponents k	Role
f_2	$x^3 + 2x + 1$	$\{1, 3, 9\}$	Doc. 341, lepton layer
f_3	$x^3 + 2x^2 + 1$	$\{17, 23, 25\}$	$= f_2^{-1}$
f_1	$x^3 + 2x^2 + x + 1$	$\{5, 15, 19\}$	
f_4	$x^3 + x^2 + 2x + 1$	$\{7, 11, 21\}$	$= f_1^{-1}$

The quotient $(\mathbb{Z}/26)^*/\langle 3 \rangle \cong \mathbb{Z}_4$ is cyclic; inversion $k \mapsto -k$ is its element of order 2.

Sector pairing reduces four to two

Doc. 336 identifies the FFGFT sector pairing $k \leftrightarrow -k$ with the Frobenius automorphism; at the level of elements of $\text{GF}(27)^*$ it is inversion $x \mapsto x^{-1} = x^{25}$. It permutes the classes as **[B]**

$$f_1 \leftrightarrow f_4, \quad f_2 \leftrightarrow f_3. \quad (7)$$

If sector pairing is a symmetry of the mass layer — as it is in the corpus — there are not four but **two** physically distinguishable $\text{GF}(27)^*$ layers: $\{f_2, f_3\}$ (leptons, Doc. 341) and $\{f_1, f_4\}$.

Negation couples order 26 to order 13

Since $-1 = x^{13}$ in $\text{GF}(27)$, the map $x \mapsto -x$ sends every element of order 26 to one of order 13 and vice versa. The four order-13 classes $g_1, \dots, g_4$ of Doc. 342 are therefore not an independent structure but the sign partners of the primitive classes **[B]**:

$$f_1 \leftrightarrow g_1, \quad f_2 \leftrightarrow g_3, \quad f_3 \leftrightarrow g_2, \quad f_4 \leftrightarrow g_4. \quad (8)$$

The candidate “sub-lepton structure” from the order-13 polynomials, left open in Doc. 342, is thereby eliminated.

Open question

The reduction $4 \rightarrow 2$ sharpens the question of Doc. 342: not “do f_2, f_3, f_4 carry further layers”, but “is there exactly one second $\text{GF}(27)^*$ layer $\{f_1, f_4\}$ besides the lepton layer $\{f_2, f_3\}$, and which particles carry it?” This remains open **[S]**.

.6 Theorem E: Prime patterns in the $\text{GF}(3^k)$ hierarchy

The harmonic primes are uniquely forced

A prime p satisfies $\text{ord}_p(3) = k$ if and only if p divides $3^k - 1$ and $p \nmid 3^j - 1$ for all $j < k$. The prime factorisations of the first relevant values give **[B]**:

Table 3: Prime factors of $3^k - 1$ and the forced physical primes

k	$3^k - 1$	prime factors	phys. prime	note
1	2	$\{2\}$	2	octave
3	26	$\{2, 13\}$	13	unique prime with $\text{ord}_p(3) = 3$ worldwide
4	80	$\{2, 5\}$	5	only non-2 factor
5	242	$\{2, 11\}$	11	only non-2 factor
6	728	$\{2, 7, 13\}$	7	(13 has $\text{ord}_{13}(3) = 3$, not 6)

The strongest result concerns $p = 13$ **[B]**: since $3^3 - 1 = 26 = 2 \cdot 13$ and $\text{ord}_2(3) = 1$, $p = 13$ is the *unique* prime worldwide with $\text{ord}_p(3) = 3$. There is no other, and there can be none. The physical primes $\{5, 7, 11, 13\}$ are not selected by harmonic considerations, but are completely and uniquely forced by the prime-factor structure of $3^k - 1$ for $k = 3, 4, 5, 6$.

Necessary congruence condition

From $\text{ord}_p(3) = k$ with $3 \mid k$ it follows that $3 \mid (p - 1)$, i.e. $p \equiv 1 \pmod{3}$ **[B]**. In particular $p = 13$ (at $k = 3$) and $p = 7$ (at $k = 6$) are both $\equiv 1 \pmod{3}$ — exactly the $\mathbb{Z}_3$ -compatible primes of the corpus (Doc. 342, Theorem B).

.7 Theorem F: Limits of distinguishability and computational cost

Exact threshold: Euler factor vs. fractal correction

The Euler factor $1/(1 - p^{-s})$ exceeds the fractal correction 1.05 % (Doc. 338, bare to measured) exactly for $p < p^*$ with $p^* = (0.0105)^{-1/s}$ **[B]**:

Table 4: Exact threshold p^* by value of s

s	p^*	primes $\leq p^*$	harmonic limit	Euler factor $\geq 1.05\%$
1	95.2	$\{2, 3, 5, \dots, 89\}$	—	all small primes
2	9.76	$\{2, 3, 5, 7\}$	7-limit	octave, fifth, third, seventh
3	4.57	$\{2, 3\}$	3-limit	octave and fifth only

For $s = 2$ (reference value of the spectral zeta) the threshold is $p^* \approx 9.76$: primes $\{2, 5, 7\}$ have Euler factors exceeding the fractal correction; $p = 11$ and all larger primes fall below it. This is exactly the harmonic 7-limit. $p = 11$ appears in the corpus (neutrinos, Doc. 340) via the structural formula $\Delta m^2 = (11^2 - 1)m_\nu^2$, not as a Galois product hit.

Galois products vs. random products: hit rate

Forming all products of up to three Galois building blocks with exponents $\{-1, 1, 2\}$ for $p \leq p_{\max}$, the product set grows rapidly and the hit rate for arbitrary targets reaches chance level **[B]**:

Table 5: Hit rate for physical vs. random targets at $\pm 0.5\%$

$p_{\max}$	values	43200	m_μ/m_e	random avg
13	1334	2	0	0.92
30	5802	3	2	3.60
50	8946	3	2	5.33
100	13398	4	3	7.66
300	24992	8	12	14.15

From $p_{\max} \approx 30$ the hit rate lies at chance level ($\pm 0.5\%$); at the strict threshold $\pm 0.05\%$ already from $p_{\max} \approx 13$. The identity $43200 = |\text{GF}(9)^*|^2 \cdot 5^2 \cdot |\text{GF}(27)|$ (Doc. 338) uses only building blocks with $p \leq 5$ and is derived *geometrically* from ξ , not found by product search.

Structural computational limit: Weyl obstruction and quantum computers

The distinguishability threshold at $p_{\max} \approx 13$ is a *physical* limit (fractal correction). Independently of this, the corpus contains a sharper *mathematical* limit that applies to every technology **[B]**:

Weyl obstruction, Doc. 316

There is no compact Riemannian manifold — of any dimension, curvature, topology, or orbifold structure — whose Laplace spectrum consists of the Riemann zeros. Reason: $T \ln T$ is not a power $T^{d/2}$.

From Doc. 147, an n -qubit register is a compact state space with eigenvalue density $N(\lambda) \sim \lambda^{d/2}$; arithmetic counting densities grow as $T \ln T$. A finite state space cannot fully carry such a spectrum, *independently of the register size* **[B]**.

Doc. 075 names the consequence for Shor's algorithm:

Method	Limit	Type
Trial division	$O(\sqrt{n})$	computational
Pollard ρ	$O(n^{1/4})$	computational
Shor's QFT	resource limit	Weyl obstruction

The central statement of Doc. 316 holds technology-independently:

The limit lies not in the tool but in the object. Transformation (Fourier, QFT, optics) projects and generates no information. Relaxation finds what is already in the potential. Both presuppose that the sought structure is already present.

Physical limit ($p^* \approx 9.76$, Theorem F.1) and mathematical limit (Weyl obstruction, Theorem F.3) thus appear in the same picture: beyond a certain point the prime structure is hidden in the noise, and no tool — including an arbitrarily large quantum computer — retrieves it.

.8 In plain terms: what the limit means

Question. So arbitrarily dense harmonics can be computed — but beyond a certain depth they can no longer be distinguished?

Answer. Yes, exactly that.

Computation can go arbitrarily far. $\text{GF}(3^k)$ for $k = 7, 8, 12, 100$ — the algebra does not stop. Every k yields new primes, new polynomial classes, new chords. At $k = 12$ all four harmonic primes 5, 7, 11, 13 sit in one field simultaneously; at $k = 7$ the Wieferich prime 1093 enters. These structures are exactly computable and mathematically all present.

But beyond a certain depth they are physically meaningless, for two reasons that coincide.

First, the grid becomes too fine. For $k \leq 3$ the reachable values lie about 3 % apart — if 43200 hits a mass value to 1 %, that is a result. For $k \geq 5$ they lie less than 0.5 % apart. Then every number hits something, and a hit means nothing. The control test in Theorem F shows it: two hundred random numbers hit as often as the actual mass ratios.

Second — and this is the physical reason — the fractal correction of about 1 % (Doc. 338, bare to measured) is itself already a statement about higher winding modes. Once the Galois grid is finer than this correction, nature itself no longer distinguishes neighbouring Galois values. The structure is not gone; it has dissolved into the noise of the higher modes.

The analogy with harmonics is exact: everyone hears the overtones 2:1 and 3:2. 17:16 is a semitone, 33:32 a quarter-tone. From roughly the 250th overtone the spacing falls below 5 cents — mathematically exactly present, acoustically no longer distinguishable. The series becomes a continuum.

The consequence for FFGFT. The exact Galois layer reaches to $\text{GF}(27)$, i.e. $k = 3$: the lepton layer, 43200, the four classes, $p = 13$ as the unique prime with $\text{ord}_p(3) = 3$. Above that ($k = 4$, $\text{GF}(81)$, $p = 5$) is already the transition. From $k = 5$ on, anything found by search is indistinguishable from chance — and must receive no status unless it comes directly from ξ .

This is not a deficiency of the theory but its consistency: Doc. 159 says mode convergence, Doc. 316 says Weyl obstruction, Doc. 342 and this document say the same thing algebraically. There is a limit, it lies at $k \approx 3-4$, and it comes from the subject itself, not from our computational capacity.

.9 In plain terms II: synthesis, analysis, factorisation — and coupling

Question. What does the limit mean for synthesis and analysis — and for factorisation? And how does coupling enter: Doc. 328 has three coupling regimes; do pure numbers show the same picture as physical processes?

Synthesis is unlimited, analysis is limited

Synthesis — adding overtones, multiplying primes, forming $\text{GF}(3^k)$ chords from partitions — goes arbitrarily far and arbitrarily dense. Nothing prevents superposing the 250th overtone

with the 251st or multiplying two 300-digit primes. Synthesis does not know the limit of Theorem F.

Analysis — which overtones are in the sound, which primes in the number, which Galois class carries the mass — hits exactly this limit. It is not a matter of computational cost but of resolution: to separate overtone n from $n+1$ one needs a time window of about n periods; to find the period r of a number, a frequency resolution $\Delta f \approx f_0/n^2$ (Doc. 075, precision barrier). Beyond a certain density the spacing lies below the noise, and the decomposition is no longer unique — not because it does not exist, but because many decompositions fit equally well. That is the chance-level finding of Theorem F in other words.

Factorisation is analysis

Factorising $N = p \cdot q$ means reading the two fundamentals out of a beat. For small p, q — the 7-limit range, $k \leq 3$ — this works; those are exactly the primes visible in the FFGFT spectrum at all. RSA primes with 300 digits sit at $k = \text{ord}_p(3)$ of the order of p itself, i.e. at a depth undercutting the grid spacing of Theorem F by hundreds of orders of magnitude. Doc. 316 puts it thus: Fourier, QFT and optics project, they generate no information; the period must already be in the prepared states, and this preparation (modular exponentiation, 95 % of the gate cost, Doc. 147) is exactly the problem one wanted to solve.

Two things follow. First: FFGFT does not break RSA — and that is not a deficiency but a consistent result. The asymmetry synthesis-easy / analysis-hard on which RSA lives is the same asymmetry that ends the Galois layer at $k \approx 3-4$. Second, for instrument building: an instrument can *produce* overtones arbitrarily densely; ear and analysis *hear* them sharply only up to about the 7-limit. What lies above acts as timbre, not as pitch — as the higher winding modes in FFGFT act as a 1 % correction, not as a separate mass layer.

Caveat [S]. Transferring Theorem F to $\mathbb{Z}/N\mathbb{Z}$ is an analogy. Doc. 075 records that ξ has no function in $\mathbb{Z}/N\mathbb{Z}$ and the period is fixed purely arithmetically by $\text{lcm}(p-1, q-1)$. The analogy holds; the fractal correction as the *mechanism* of the factorisation limit does not.

Coupling: the same threefold division, not the same law

Doc. 328 classifies couplings relative to loss rates: subcritical ($\kappa < 1$, one maximum), critical ($\kappa = 1$, maximally flat), supercritical ($\kappa > 1$, splitting). Locking onto a frequency ratio p/q occurs in Arnold tongues of width $\Delta\Omega(p/q) \propto K^q$.

For pure numbers the tolerance ε takes the role of the coupling strength: a real number “locks” onto p/q if $|x - p/q| < \varepsilon$. Fractions with denominator $\leq Q$ cover the interval with measure $\approx 2\varepsilon \cdot 3Q^2/\pi^2$; coverage 1 gives the critical denominator height **[B]**

$$Q_c = \frac{\pi}{\sqrt{6}\varepsilon}. \quad (9)$$

For $\varepsilon = 1.05\%$ (fractal correction), $Q_c \approx 12.5$ — the same order as $p_{\max} \approx 13$ in Theorem F. The Galois version confirms it numerically (pruef_343c): at $\varepsilon = 1.05\%$ the Galois grid with $p_{\max} = 7$ is subcritical (coverage 0.52), with $p_{\max} = 13$ critical (0.92), from $p_{\max} \approx 30$ supercritical (1.00).

The limit $k \approx 3-4$ of Theorem F is the critical point $\kappa = 1$ in the sense of Doc. 328 [B].

So the threefold division is the same in arithmetic and physics. *The law behind it is not [B]:* Arnold tongues fall exponentially in q ($\Delta\Omega \propto K^q$), Farey spacings only polynomially ($1/q^2$). At $K = 0.3$ and $q = 5$ the tongue is already 10^{-5} narrower than the arithmetic capture interval; at $q = 13$ by 10^{-15} . Physical locking is exponentially selective, arithmetic distinguishability polynomial. Physics “hears” fewer harmonics than arithmetic permits — the edge is the same, but physically sharper.

FFGFT is dynamically deeply subcritical

With $K_{\text{eff}} = 2\pi\xi \approx 8.4 \times 10^{-4}$ (Doc. 328) the widest non-trivial Arnold tongue is $\Delta\Omega(1/2) \approx 5.6 \times 10^{-8}$, the total measure of all tongues $\approx 8 \times 10^{-4}$. Both lie five to two orders of magnitude below the fractal correction **[B]**. No mass ratio is pulled onto a p/q by resonant locking.

This yields a clarification that was so far only implicit in the corpus: the exact rationals — 43200, $8^2 \cdot 5^2 \cdot 27$, the winding numbers (r_i, p_i) — are *topological* (integer windings on T^4), not *dynamical* (locking). Discreteness and coupling are two separate mechanisms in FFGFT. This is consistent with Doc. 328, where φ lies outside all tongues: the coupling is too weak to hold ratios; what is held, topology holds.

.10 Summary

Table 6: Results and status

Theorem	Content	Status
A	$\zeta_{T^4/\mathbb{Z}_3}(s) = (1 - 3^{-s})\zeta(s) = L(s, \chi_0)$	[B]
A'	$L(s, \chi_1)$ measures twist-sector asymmetry	[S]
B	Euler product hierarchical by $k = \text{ord}_p(3)$	[B]
B'	Fractal damping suppresses Euler factors $k \geq 7$	[B]
C	Orbifold zeros at $s = 2\pi i k / \ln 3$ on $\text{Re}(s) = 0$	[B]
C'	No resonance of Riemann zeros with $\mathbb{Z}_3$ unit	[B]
D	Galois character does not separate; classes $= (\mathbb{Z}/26)^* / \langle 3 \rangle \cong \mathbb{Z}_4$	[B]
D'	Sector pairing $k \leftrightarrow -k$: $f_1 \leftrightarrow f_4, f_2 \leftrightarrow f_3$ — 4 classes $\rightarrow 2$	[B]
D''	Negation couples order 26 to 13; no sub-lepton layer	[B]
D'''	Second layer $\{f_1, f_4\}$ physically realised?	[S]
E	$p = 13$ unique prime with $\text{ord}_p(3) = 3$ worldwide; primes forced by $3^k - 1$	[B]
E'	$\text{ord}_p(3) = k, 3 k \Rightarrow p \equiv 1 \pmod{3}$ ($\mathbb{Z}_3$ -compatible)	[B]
F	Euler factor threshold $p^* \approx 9.76$ ($s = 2$) = 7-limit boundary	[B]
F'	Product density: chance level from $p_{\text{max}} \approx 30$ ($\pm 0.5\%$)	[B]
F''	Weyl obstruction: limit lies in the object, not the tool (Doc. 316)	[B]
G	Coupling: $Q_c = \pi/\sqrt{6\varepsilon} \approx 12.5$ at 1.05 %; limit $k \approx 3-4$ = critical point $\kappa = 1$	[B]
G'	Same threefold division, different law: Arnold K^q vs. Farey $1/q^2$	[B]
G''	FFGFT dynamically deeply subcritical ($K_{\text{eff}} = 2\pi\xi$); discreteness topological, not dynamical	[B]
G'''	Theorem F on $\mathbb{Z}/N\mathbb{Z}$ (RSA) only as analogy	[S]

The central statement: the full Riemann zeta $\zeta(s)$ belongs to the flat $\mathbb{R}^4$ spectrum. The zeta of the $T^4/\mathbb{Z}_3$ orbifold is $L(s, \chi_0) = (1 - 3^{-s})\zeta(s)$ — a filtered version in which the 3-Euler factor is separated out and the primes are weighted by the Galois hierarchy $k = \text{ord}_p(3)$. This is the zeta-function form of the entry rule of Doc. 342.

Check scripts

Directory 2/python/Dok343_Skripte/, wrapper `run_all_343.sh`:

- `pruef_343_zeta_galois.py` — Theorems A–D; partial zeta decomposition, L-functions, zeros, resonance test, weighted Euler product.
- `pruef_343b_klassen_symmetrie.py` — Theorem D; explicit Frobenius classes in $\text{GF}(27)^*$, action of inversion and negation, cosets of $\langle 3 \rangle$ in $(\mathbb{Z}/26)^*$, exact values of $L(s, \chi_1)$.
- `pruef_344_primzahlmuster.py` — Theorems E–F; patterns in $\text{ord}_p(3)$, exact threshold p^* , Galois vs. random hit rate, Euler factors.

- `pruef_343c_kopplung_zahlen.py` — Plain-terms II section; Farey coverage and $Q_c(\varepsilon)$, law comparison Arnold K^q vs. Farey $1/q^2$, regime of $K_{\text{eff}} = 2\pi\xi$, Galois coverage by p_{max} . All assertions passed.

Note on the use of AI assistance

This document was prepared with the assistance of Claude (Anthropic). All numerical computations are verified by Python check scripts. The physical interpretations and conclusions are those of Johann Pascher (ORCID 0009-0000-6518-4064).

Bibliography

- [1] J. Pascher, *Doc. 057: Relational number system*, FFGFT corpus.
- [2] J. Pascher, *Doc. 159: Harmonic structure of the torus*, FFGFT corpus.
- [3] J. Pascher, *Doc. 328: Coupling regimes, resonance and synchronisation*, FFGFT corpus, August 2026.
- [4] J. Pascher, *Doc. 330: Three operations from T^4 to the observation space*, FFGFT corpus, August 2026.
- [5] J. Pascher, *Doc. 336: The algebraic bridge: FFGFT and $\mathbb{Z}_3\mathbb{C}$* , FFGFT corpus, August 2026.
- [6] J. Pascher, *Doc. 075: RSA and period-finding*, FFGFT corpus, August 2026.
- [7] J. Pascher, *Doc. 147: Quantum computing and $T0$ time-mass duality*, FFGFT corpus, August 2026.
- [8] J. Pascher, *Doc. 316: FFGFT and the Riemann zeta function*, FFGFT corpus, August 2026.
- [9] J. Pascher, *Doc. 342: Factorisation classes in $G(6)$ and the harmonics of the primes*, FFGFT corpus, 2 September 2026.
- [10] P. G. L. Dirichlet, *Über die Bestimmung der Dichte einer unendlichen Anzahl von Punkten*, 1837.