

Anomalous Magnetic Moments in FFGFT Theory

Geometric Derivation from Time-Mass Duality

Purely Geometric Formulas and Precise Ratio Predictions

Revision 10

Johann Pascher

2026 February 21

Contents

.1	Introduction: Geometric vs. Semi-Empirical Approaches	2
	The Philosophy of T0 Theory	2
	Consistency with Mass Predictions	2
.2	Physical Fundamentals	3
	What is the Anomalous Magnetic Moment?	3
	T0 Interpretation: Windings in the Torsion Lattice	3
.3	Geometric Formulas	3
	Fundamental Parameters	3
	The Real Sub-Planck Factor: $f = 7500$	3
	The Symmetry Breaking: The Role of the Golden Ratio	4
	Electron: Base Winding	4
	Muon: Fractal Additional Winding	5
	Tau: More Complex Fractal Structure	6
.4	Two Classes of Predictions: Absolute Values vs. Ratios	6
	Why 2% Deviation for Absolute Values?	6
	Ratios are Mathematically Exact	6
	Analogy to the Koide Formula	7
.5	Precise Ratio Predictions	7
	Analogy to the Koide Formula	7
	The Ratio of Differences	7
	Numerical Verification	8
	Testable Prediction for Tau	8
.6	Why 2% Deviation?	9
	Higher-Order Quantum Effects	9
	Discrete Lattice Structure	9
	Pentagonal Symmetry Breaking	9
.7	Experimental Tests	9
	Belle II (2027–2028)	9
	Fermilab/J-PARC	10
.8	Comparison with Other Approaches	10
.9	Reconstruction of the Correction Factor from Experimental Data	10
	The Central Observation	10
	Reconstruction of k_{geom}	10
	Using the Reconstructed Correction Factor	11
	Alternative: Directly from Ratio Relation	11
	Two Complementary Tau Predictions	11
	What does this mean for Belle II?	11
.10	Important Note: No α in the T0 g-2 Formulas	12

Correction notes

Corrections from the calculation review (30 Sept. 2026). The following places are corrected or annotated in the text; each carries a dated note there with the previous value:

- a_τ from the ratio: $1.280 \rightarrow 1.277 \times 10^{-3}$ (summation error; at all five places). The abstract value 1.282 follows with the exact measured values (Rev. 12, Doc. 158).
- Mass deviation: $0.87\text{--}2.16\% \rightarrow 0.4\text{--}1.2\%$ (the document's own table).
- Pentagonal symmetry breaking: $f = f_{\text{ideal}} - 5\varphi$ (0.1 %) $\rightarrow f = f_{\text{ideal}} = 7500$, breaking in k_{geom} (about 2 %); likewise in the list of causes " $\Delta = 5\varphi$ leads to 0.1 %".
- Literature examples for the exponents: $p_\mu = 5/3$ now the Flory value of the self-avoiding walk in 3D (previously 2D Brownian motion, random walk, Koch curve); $p_\tau = 4/3$ now the frontier of planar Brownian motion and the planar self-avoiding walk (previously Koch curve, Mandelbrot set). The exponents themselves are settings.

Abstract

In the present work, the fundamental architecture of spacetime is reinterpreted within the framework of **Fundamental Fractal Geometric Field Theory (FFGFT)** – internally referred to as the T0 model (B18). The central idea is the transition from a point-like to a purely geometric description of the vacuum as a four-dimensional **Gyral Torus**.

Note on versions. The German version of this document is more extensive than this English one. Earlier revisions shortened or changed parts of the English version. Where the two differ, the more extensive German version applies.

Geometric Structure: The theory is based on the fractal-geometric foundation with the parameter $\xi \approx (4/3) \times 10^{-4}$ and the densest local sphere packing by regular **Tetrahedra**. This tetrahedral basis forms the stable foundation for the low generations (electron, muon, proton/neutron) as well as the local 3D crystal structure of the torus. Building upon this, the ideal sub-Planck factor

$$f = 7500,$$

emerges through fractal branching and pentagonal symmetry breaking, representing an exactly 7500-fold reduction compared to the conventional Planck scale (t_0) and following directly from the geometric winding density $30000/4$.

g-2 Anomaly: A core element of the work is the transparent geometric derivation of the anomalous magnetic moments of leptons. While the Standard Model relies on numerous perturbative terms, in FFGFT the electron anomaly follows directly from the base winding (tetrahedral projection). The muon and tau anomalies arise from fractal branchings with Hausdorff dimensions $p \approx 5/3$ and $4/3$, respectively. With the ideal value $f = 7500$, the purely geometric predictions achieve an accuracy of about 2 %. By reconstructing the projection factor k_{geom} , the deviation for the muon drops below 0.2 %. The most precise, k_{geom} -independent prediction for the tau anomaly is

$$a_\tau \approx 1.282 \times 10^{-3},$$

which follows exclusively from the exact ratio $f^{1/3} - 1$.

Geometric Proportionality: All physical base quantities (constants, masses, couplings) stand in fixed geometric ratios, considerably reducing the number of free parameters compared to the Standard Model. The T0 theory thus offers a transparent geometric

description and provides concrete, experimentally testable predictions – particularly for the tau anomaly as an important test at Belle II.

Status markers. Statements marked **[S]** are *speculative*: a hint or conjecture that has not yet been derived or numerically checked in the corpus.

Note on Older Documents

Previous versions of the g-2 analysis (018_T0_Anomale-g2-10_En.pdf) used semi-empirical factors. The present formulation uses **exclusively geometric factors** and states the 2% deviation openly, which is consistent with the precision of all T0 predictions. Python scripts available at: github.com/jpascher/T0-Time-Mass-Duality

Keywords: Anomalous magnetic moment, g-2, T0 theory, Time-Mass Duality, Torsion lattice, Ratio predictions, Koide formula

.1 Introduction: Geometric vs. Semi-Empirical Approaches

The Philosophy of T0 Theory

The T0 theory is based on the principle that **all** physical constants should follow from the geometric structure of a 4-dimensional torsion lattice. For anomalous magnetic moments this means:

- **No** hidden fit parameters
- **Only** geometric factors: φ, ξ, f
- Open statement of precision limits
- Consistency with other predictions

Consistency with Mass Predictions

The T0 theory predicts lepton masses with 0.4–1.2% deviation:

Lepton	T0 [MeV]	Exp [MeV]	Deviation
Electron	0.505	0.511	1.18%
Muon	105.0	105.7	0.66%
Tau	1783	1777	0.37%

Table 1: Lepton masses in T0

Expectation: g-2 should have similar precision (1%).

Since the masses already deviate by 1%, no closer agreement is to be expected for g-2.

.2 Physical Fundamentals

What is the Anomalous Magnetic Moment?

The magnetic moment of a charged spin-1/2 particle is:

$$\mu = g \cdot \frac{e}{2m} \cdot \frac{\hbar}{2} \quad (1)$$

where g is the gyromagnetic factor (g-factor).

Dirac Prediction: For a point-like particle: $g = 2$

Quantum Effects: Vacuum polarization, vertex corrections $\Rightarrow g \neq 2$

Anomaly: $a = (g - 2)/2$

QED Expectation: $a \approx \alpha/(2\pi) + \mathcal{O}(\alpha^2) \approx 0.00116$

T0 Interpretation: Windings in the Torsion Lattice

In T0 theory, leptons are **winding structures** in the 4D torsion lattice:

- **Electron:** Simple winding (1st generation)
- **Muon:** Winding with fractal branching (2nd generation)
- **Tau:** More complex fractal structure (3rd generation)

The anomalous moment arises from:

1. The **rotation** of the winding (spin)
2. The **charge distribution** on the winding
3. The **projection** 4D $\rightarrow$ 3D

$\Rightarrow$ **No** point-like charge $\Rightarrow a \neq 0$

.3 Geometric Formulas

Fundamental Parameters

The T0 theory uses exclusively three geometric fundamental constants:

$$\varphi = \frac{1 + \sqrt{5}}{2} = 1.618 \dots \quad (\text{Golden Ratio}) \quad (2)$$

$$\xi = \frac{4}{3} \times 10^{-4} = 1.333 \times 10^{-4} \quad (\text{Torsion constant}) \quad (3)$$

$$f = 7500 \quad (\text{Sub-Planck factor}) \quad (4)$$

The Real Sub-Planck Factor: $f = 7500$

Now we put everything together: The ideal crystal remains intact, the symmetry breaking only affects the projection factors:

$$\boxed{f = 7500} \quad (5)$$

This is a **central number of the T0 theory**. It appears in almost all formulas and describes:

- The number of Sub-Planck cells per Planck length
- The density of the torsion lattice
- The fundamental frequency of all geometric resonances

The Symmetry Breaking: The Role of the Golden Ratio

A perfect, ideal crystal would be completely symmetric. Yet our world shows symmetry breaking on all levels:

- Matter dominates over antimatter
- The weak interaction violates parity symmetry

- The neutron is heavier than the proton
- The three lepton generations have different masses

In the T0 theory, all these symmetry breakings have a single, geometric origin: the pentagonal symmetry of the crystal, embodied by the **golden ratio** φ . The golden ratio $\varphi = (1 + \sqrt{5})/2 = 1.618033989 \dots$ is the irrational number describing pentagonal symmetry. In a perfect pentagon, φ appears everywhere: The ratio of diagonal to side is exactly φ . Why pentagonal symmetry specifically? For deep mathematical reasons, pentagonal symmetry is the first one that **cannot tile the plane periodically**. This leads to *quasicrystals* – structures that are ordered but not periodic. Exactly such a quasicrystalline structure is postulated by T0 theory for the Sub-Planck scale. The symmetry breaking is quantified in the theory not by a direct subtraction of 5φ from the ideal anchor number 7500. Instead, it is hidden in the **ca. 2% deviations** that appear in the calculations of the anomalous magnetic moments (g-2 anomalies). This deviation arises from the pentagonal projection in the geometric factor k_{geom} :

$$k_{\text{geom}} = \frac{2}{\sqrt{\varphi}} \times \sqrt{2} \approx 2.22357, \quad (6)$$

which projects the 4D torsion onto the 3D world. The version reconstructed from experimental data deviates by about 2% ($k_{\text{geom}}^{\text{rec}} \approx 2.26955$), reflecting the actual symmetry breaking – a slight distortion by the pentagonal geometry that breaks perfect symmetry without changing the ideal value $f = 7500$.

From the ideal 7500 remained the ideal 7500. In the T0 model this number becomes the fundamental constant. It determined how densely the lattice was packed, how quickly torsion could propagate, which resonances were possible. In the T0 picture the observed particle masses and force strengths are meant to go back to this single geometric story, and likewise – **[S]**, the cosmic sector is deliberately set aside in the current corpus (P39) – the cosmological constants: From perfect crystal to pentagonally broken reality, with the breaking hidden in the 2%.

Electron: Base Winding

Formula:

$$a_e = \frac{S_3/f}{k_{\text{geom}}} \quad (7)$$

where:

- $S_3 = 2\pi^2 = 19.739$: 3D surface of the 4D winding
- $f = 7500$: Sub-Planck scaling
- k_{geom} : Geometric projection factor

Geometric Projection Factor:

$$k_{\text{geom}} = \frac{2}{\sqrt{\varphi}} \times \sqrt{2} \quad (8)$$

Explanation of Factors:

- $2/\sqrt{\varphi} = 1.572$: Pentagonal projection (from ξ -structure)
- $\sqrt{2} = 1.414$: Diagonal projection 4D $\rightarrow$ 3D
- $k_{\text{geom}} = 2.224$: determined purely geometrically

Numerical Calculation:

$$k_{\text{geom}} = \frac{2}{\sqrt{1.618}} \times \sqrt{2} = 2.224 \quad (9)$$

$$a_e = \frac{19.739/7500}{2.224} \quad (10)$$

$$a_e = 1.184 \times 10^{-3} \quad (11)$$

Comparison:

- T0: $a_e = 1.184 \times 10^{-3}$
- Experiment: $a_e = 1.160 \times 10^{-3}$
- Deviation: **2.03%**

Muon: Fractal Additional Winding**Formula:**

$$a_\mu = a_e + \Delta a_{\text{fractal}} \quad (12)$$

with

$$\Delta a_{\text{fractal}} = \frac{4\pi}{f p_\mu} \quad (13)$$

where:

- $p_\mu = 5/3$: Fractal Hausdorff dimension
- 4π : Complete torsion revolution

Meaning of $p_\mu = 5/3$:

This is the well-known Hausdorff dimension of:

- the self-avoiding random walk in 3D (Flory value $1/\nu = 5/3$) (*Corrected on 30 Sept. 2026: previously "Brownian motion in 2D, self-avoiding random walk, Koch curve" – misattributed: planar Brownian motion has dimension 2 (its frontier $4/3$), the planar self-avoiding walk $4/3$, the Koch curve $\log 4 / \log 3 = 1.262$; $5/3$ is the Flory value of the self-avoiding walk in 3D. The exponent itself is a setting and is not affected.*)

⇒ Physically plausible for a "partially branched winding".

Numerical Calculation:

$$\Delta a_{\text{fractal}} = \frac{4\pi}{7500^{5/3}} = 4.373 \times 10^{-6} \quad (14)$$

$$a_\mu = 1.184 \times 10^{-3} + 4.373 \times 10^{-6} \quad (15)$$

$$a_\mu = 1.188 \times 10^{-3} \quad (16)$$

Comparison:

- T0: $a_\mu = 1.188 \times 10^{-3}$
- Experiment: $a_\mu = 1.166 \times 10^{-3}$
- Deviation: **1.89%**

Tau: More Complex Fractal Structure**Formula:**

$$a_\tau = a_e + \frac{4\pi}{f p_\tau} \quad (17)$$

where:

- $p_\tau = 4/3$: Stronger fractal branching

Meaning of $p_\tau = 4/3$:

This is the Hausdorff dimension of the frontier of planar Brownian motion and of the planar self-avoiding walk. (Corrected on 30 Sept. 2026: previously "box-counting dimension of many fractals (e.g., Koch curve, Mandelbrot set)" – the Koch curve has $\log 4 / \log 3 = 1.262$, the boundary of the Mandelbrot set 2.)

Numerical Calculation:

$$\Delta a_{\text{fractal}} = \frac{4\pi}{7500^{4/3}} = 8.560 \times 10^{-5} \quad (18)$$

$$a_\tau = 1.184 \times 10^{-3} + 8.560 \times 10^{-5} \quad (19)$$

$$a_\tau = 1.269 \times 10^{-3} \quad (20)$$

Status: This is a **prediction** – tau-g-2 has not been measured yet.

.4 Two Classes of Predictions: Absolute Values vs. Ratios

Why 2% Deviation for Absolute Values?

The T0 theory uses exclusively geometric factors without adjustment parameters. The 2% deviation for absolute g-2 values is:

- **Consistent** with all T0 predictions (masses: 0.4–1.2%) (Corrected on 30 Sept. 2026: previously "0.87–2.16%" – contradicted the document's own table of lepton masses with 0.34–1.17 %.)
- **Expected** for a purely geometric description
- **Comparable** to α^2 effects in QED (1–2%)
- **A property** of a description at leading geometric order

Causes of the 2% Deviation:

1. **Higher-order quantum effects:** T0 captures the leading geometric structure, but not all loop corrections
2. **Discrete lattice structure:** The torsion lattice is discrete, not continuous
3. **Pentagonal symmetry breaking:** acts in the projection factor k_{geom} (about 2 %), $f = 7500$ stays unchanged (Corrected on 30 Sept. 2026: previously " $\Delta = 5\varphi$ leads to 0.1% corrections" – aligned with $f = f_{\text{ideal}} = 7500$, see section "Pentagonal Symmetry Breaking".)

Ratios are Mathematically Exact

In contrast to absolute values, **ratios of differences** are structurally exact:

$$\frac{\Delta a(\tau - \mu)}{\Delta a(\mu - e)} = \frac{4\pi/f^{4/3} - 4\pi/f^{5/3}}{4\pi/f^{5/3}} = f^{1/3} - 1 \quad (21)$$

Why is this exact?

- The common factor 4π cancels out
- The projection factor k_{geom} cancels out
- Only the fractal exponents (5/3 and 4/3) determine the ratio
- The result depends **only** on f : $f^{1/3} - 1 = 18.57$

Important**Fundamental Distinction Absolute values:**

- Depend on k_{geom} , f_i and SI conversion
- 2% deviation due to higher-order quantum effects
- Consistent with all T0 predictions

Ratios:

- Depend **only** on f
- k_{geom} and SI factors cancel out
- Mathematically exact from fractal exponents
- Difference $< 10^{-13}$ (numerical precision)

⇒ Within the model, the ratio prediction is **not an approximation**, but an **exact geometric relation**.

Analogy to the Koide Formula

This behavior is analogous to the Koide formula for lepton masses:

- **Individual masses:** 0.4–1.2% deviation
- **Koide ratio:** $\pm 0.0004\%$ precision

The ratio is **more robust** than absolute values because systematic factors cancel out.

For g-2 in T0:

- **Absolute values:** 2% deviation
- **Ratio** $\Delta a(\tau - \mu)/\Delta a(\mu - e)$: Exactly $= f^{1/3} - 1$

The difference between the two classes reflects the **geometric structure** of the theory.

.5 Precise Ratio Predictions**Analogy to the Koide Formula**

The Koide formula for lepton masses:

$$\frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3} \pm 0.0004\% \quad (22)$$

shows: **Ratios** are more precise than absolute values.

Question: Does this also hold for g-2?

The Ratio of Differences

Define the differences:

$$\Delta a(\mu - e) = a_\mu - a_e = \frac{4\pi}{f^{5/3}} \quad (23)$$

$$\Delta a(\tau - \mu) = a_\tau - a_\mu = \frac{4\pi}{f^{4/3}} - \frac{4\pi}{f^{5/3}} \quad (24)$$

Ratio:

$$\frac{\Delta a(\tau - \mu)}{\Delta a(\mu - e)} = \frac{4\pi/f^{4/3} - 4\pi/f^{5/3}}{4\pi/f^{5/3}} \quad (25)$$

$$= \frac{f^{5/3}}{f^{4/3}} - 1 \quad (26)$$

$$= f^{5/3-4/3} - 1 \quad (27)$$

$$= f^{1/3} - 1 \quad (28)$$

Important

Core Prediction

$$\boxed{\frac{\Delta a(\tau - \mu)}{\Delta a(\mu - e)} = f^{1/3} - 1 = 18.57} \quad (29)$$

This relation is:

- **Depends only on** f (no further parameter)
- **Independent** of k_{geom}
- **Exact** (difference $< 10^{-13}$)
- **Testable** at Belle II

Numerical Verification

With $f = 7500$:

$$f^{1/3} = 7500^{1/3} = 19.57 \quad (30)$$

$$f^{1/3} - 1 = 18.57 \quad (31)$$

From T0 values:

$$\Delta a(\mu - e) = 4.373 \times 10^{-6} \quad (32)$$

$$\Delta a(\tau - \mu) = 8.123 \times 10^{-5} \quad (33)$$

$$\text{Ratio} = \frac{8.123 \times 10^{-5}}{4.373 \times 10^{-6}} = 18.57 \quad (34)$$

Agreement: identical, as expected algebraically.

Testable Prediction for Tau

With experimental values for e and μ :

$$a_e^{\text{exp}} = 1.160 \times 10^{-3} \quad (35)$$

$$a_\mu^{\text{exp}} = 1.166 \times 10^{-3} \quad (36)$$

$$\Delta a(\mu - e)^{\text{exp}} = 6.000 \times 10^{-6} \quad (37)$$

Prediction:

$$\Delta a(\tau - \mu) = \Delta a(\mu - e)^{\text{exp}} \times (f^{1/3} - 1) \quad (38)$$

$$= 6.000 \times 10^{-6} \times 18.57 \quad (39)$$

$$= 1.114 \times 10^{-4} \quad (40)$$

$$a_\tau^{\text{predicted}} = 1.166 \times 10^{-3} + 1.114 \times 10^{-4} \quad (41)$$

$$= 1.277 \times 10^{-3} \quad (42)$$

(Corrected on 30 Sept. 2026: previously 1.280×10^{-3} – the sum $1.166 \times 10^{-3} + 1.114 \times 10^{-4}$ gives 1.277×10^{-3} .)

.6 Why 2% Deviation?

Higher-Order Quantum Effects

QED calculates g-2 as a perturbation series:

$$a = \frac{\alpha}{2\pi} + \mathcal{O}(\alpha^2) + \mathcal{O}(\alpha^3) + \dots \quad (43)$$

T0 captures the **geometric basic structure**, but not all higher-order quantum corrections.
 $\Rightarrow$ [S] The 2% are attributed here to higher-order effects; a quantitative estimate is still missing.

Discrete Lattice Structure

The torsion lattice is **discrete**, not continuous.

This leads to small corrections compared to continuous QFT.

Pentagonal Symmetry Breaking

$$f = f_{ideal} = 7500 \quad (44)$$

(Corrected on 30 Sept. 2026: previously $f = f_{ideal} - 5\varphi$ and "(0.1%)" – contradicted the section on the role of the golden ratio, according to which the breaking does not arise by subtracting 5φ but resides in the projection factor k_{geom} (about 2 %) and leaves $f = 7500$ unchanged.)

[S] This symmetry breaking (in k_{geom} , about 2 %) is meant to explain:

- Matter-antimatter asymmetry
- Generation structure
- Small corrections to idealized values

.7 Experimental Tests

Belle II (2027–2028)

Belle II expects sensitivity of $\sim 10^{-7}$ for a_τ .

Test 1: Absolute value

- T0 prediction: $a_\tau = 1.269 \times 10^{-3}$
- From ratio: $a_\tau = 1.277 \times 10^{-3}$ (Corrected on 30 Sept. 2026: previously 1.280×10^{-3} – summation error, see section "Testable Prediction for Tau".)
- Difference: 1%

Test 2: Ratio

- T0 prediction: $\Delta a(\tau - \mu) / \Delta a(\mu - e) = 18.57$
- This is the **more precise** prediction
- Independent of absolute calibration

Possible outcomes:

1. **Confirmation:** Ratio ≈ 18.6
 $\Rightarrow$ Supports the fractal structure hypothesis
2. **Deviation:** Ratio $\neq 18.6$
 $\Rightarrow$ Different fractal dimensions or additional physics
3. **Null result:** $a_\tau < 10^{-8}$
 $\Rightarrow$ T0 contributions suppressed or theory needs revision

Fermilab/J-PARC

Further precision improvements for a_μ :

- Reduction of experimental uncertainties
- Clearer determination of SM discrepancy
- Refinement of $\Delta a(\mu - e)$ measurement

.8 Comparison with Other Approaches

Approach	Precision	Parameters	Explainable
QED (SM)	Very high	Many	Yes
T0 (semi-empirical)	0.1%	1 adjusted	Partially
T0 (geometric)	2%	0 adjusted	Geometric

Table 2: Comparison of different approaches

Guideline: Here **explainability** takes priority over precision.

.9 Reconstruction of the Correction Factor from Experimental Data

The Central Observation

The ratio $\Delta a(\tau - \mu)/\Delta a(\mu - e) = f^{1/3} - 1$ is **mathematically exact** because the correction factor k_{geom} cancels out completely.

Since experimental measurements of a_e and a_μ are more precise (10^{-10}) than our geometric derivation of k_{geom} (2%), we can determine this factor **backwards from experiments**.

Reconstruction of k_{geom}

From the experimental electron value:

$$k_{\text{geom}}^{(\text{reconstructed})} = \frac{S_3/f}{a_e^{(\text{exp})}} = \frac{2\pi^2/7500}{1.160 \times 10^{-3}} = 2.269 \quad (45)$$

Comparison:

- Geometrically derived: $k_{\text{geom}} = (2/\sqrt{\varphi}) \times \sqrt{2} = 2.224$
- Reconstructed from experiment: $k_{\text{geom}}^{(\text{rec})} = 2.269$
- Difference: 2.0% (within the expected uncertainty range)

Using the Reconstructed Correction Factor

When we use the reconstructed value $k_{\text{geom}}^{(\text{rec})} = 2.269$:

Lepton	With $k = 2.224$	With $k = 2.269$	Experiment	Dev.
Electron	1.184×10^{-3}	1.160×10^{-3}	1.160×10^{-3}	0%
Muon	1.188×10^{-3}	1.164×10^{-3}	1.166×10^{-3}	0.2%
Tau	1.269×10^{-3}	1.246×10^{-3}	(not measured)	Prediction

Table 3: Absolute values with geometric vs. reconstructed k_{geom}

Important

Crucial Point With the reconstructed correction factor $k_{\text{geom}}^{(\text{rec})} = 2.269$, the deviations drop considerably:

- Electron: 0% deviation (by definition, since reconstructed from a_e)
- Muon: 0.2% deviation (reduced from 2% to 0.2%)
- Tau: New prediction $a_\tau = 1.246 \times 10^{-3}$

This suggests that the 2% deviation stems **mainly** from the uncertainty in deriving k_{geom} and not from the fractal base structure.

Alternative: Directly from Ratio Relation

Independent of k_{geom} is the calculation directly from the exact ratio:

$$\Delta a(\mu - e)^{(\text{exp})} = a_\mu^{(\text{exp})} - a_e^{(\text{exp})} = 6.000 \times 10^{-6} \quad (46)$$

$$\Delta a(\tau - \mu) = \Delta a(\mu - e)^{(\text{exp})} \times (f^{1/3} - 1) \quad (47)$$

$$= 6.000 \times 10^{-6} \times 18.57 = 1.114 \times 10^{-4} \quad (48)$$

$$a_\tau^{(\text{Ratio})} = a_\mu^{(\text{exp})} + \Delta a(\tau - \mu) \quad (49)$$

$$= 1.166 \times 10^{-3} + 1.114 \times 10^{-4} \quad (50)$$

$$= \boxed{1.277 \times 10^{-3}} \quad (51)$$

(Corrected on 30 Sept. 2026: previously 1.280×10^{-3} – summation error, see section “Testable Prediction for Tau”.)

Note: This prediction is **independent** of k_{geom} and uses only the exact geometric ratio structure.

Two Complementary Tau Predictions

(Corrected on 30 Sept. 2026: row “From ratio” previously 1.280×10^{-3} – summation error, see section “Testable Prediction for Tau”.)

What does this mean for Belle II?

If Belle II measures:

1. $a_\tau \approx 1.28 \times 10^{-3}$:

Method	a_τ Prediction	Dependent on
Purely geometric	1.269×10^{-3}	$k_{\text{geom}} = 2.224$ (geometric)
With rec. k_{geom}	1.246×10^{-3}	$k_{\text{geom}} = 2.269$ (from a_e)
From ratio	1.277×10^{-3}	Only f (exact)
Range	$1.25\text{--}1.28 \times 10^{-3}$	$\pm 1.5\%$

Table 4: Three T0 predictions for a_τ

- Supports the exact ratio relation $f^{1/3} - 1$
 - Indicates that experimental a_μ and ratio structure fit together
 - → **Clearest test in favour of T0 geometry**
2. $a_\tau \approx 1.25 \times 10^{-3}$:
- Supports the reconstructed value $k_{\text{geom}} = 2.269$
 - Hint that a_e, a_μ are both slightly shifted
 - → Consistent with T0, but different ratio interpretation
3. $a_\tau \approx 1.27 \times 10^{-3}$:
- Supports the purely geometric value $k_{\text{geom}} = 2.224$
 - ? Ratio deviates → fractal exponent $p_\tau \neq 4/3$?
4. a_τ **outside** 1.25–1.28:
- Not satisfied: T0 structure needs revision

Key Statement

According to this analysis, the 2% deviation of the purely geometric T0 predictions stems **mainly** from the uncertainty in deriving k_{geom} .

When we reconstruct k_{geom} from experimental data, the deviations drop considerably:

- Electron: 0% (by definition)
- Muon: 0.2% (instead of 2%)

This suggests that the **fractal T0 structure holds** and only the derivation of the projection factor $k_{\text{geom}} = (2/\sqrt{\varphi}) \times \sqrt{2}$ has a 2% uncertainty.

The k_{geom} -independent T0 prediction for tau uses the exact ratio relation:

$$a_\tau = 1.277 \times 10^{-3} \quad (52)$$

(Corrected on 30 Sept. 2026: previously 1.280×10^{-3} – summation error, see section "Testable Prediction for Tau".)

.10 Important Note: No α in the T0 g-2 Formulas

Note: The T0 formulas for g-2 contain **no** α .

In natural units ($\hbar = c = \alpha = 1$):

$$a_\ell = f(\varphi, \xi, f, \text{generation quantum numbers})$$

The anomalous moment is a **purely geometric quantity**, following from the winding structure in the torsion lattice.

- Ratios like $\Delta a(\tau - \mu)/\Delta a(\mu - e) = f^{1/3} - 1$ are **independent** of: • α (fine-structure constant)
- SI conversion factors • k_{geom} (projection factor)
- They depend only on the fractal structure.

Further Reading and Resources

T0 Theory and Python Scripts:

- Repository: github.com/jpascher/T0-Time-Mass-Duality
- Python scripts: github.com/jpascher/T0-Time-Mass-Duality/blob/main/2/python/
- Time-Mass Duality documentation
- Fundamental Fractal Geometric Field Theory (FFGFT)

Experimental Results:

- Fermilab Muon g-2 (2025): muon-g-2.fnal.gov
- Theory Initiative White Paper
- Belle II: www.belle2.org

Related T0 Documents:

- Lepton masses: Systematic derivation from quantum numbers
- Koide formula in T0: Geometric interpretation
- Fractal spacetime: $D_f = 3 - \xi$