

Corrections and Clarifications

FFGFT / T0 Time-Mass Duality

Johann Pascher

2026

Contents

.1	Register table	2
.2	Currently open bridges (as of R137)	18

Preliminary remark

This document is the compact, running correction register of the FFGFT corpus. It supersedes the previous extended edition, which remains available unchanged as **Doc. 190-Archive** (190_T0_Korrekturen_Archiv_De/En) and contains, for every entry, the full justification, the faulty and corrected expressions verbatim, and all addenda. The table below lists *all* entries (C/K, P, R) with the affected documents and the subject in short form. The register remains append-only: source documents are not revised (cf. R50); new entries are appended here, and their elaboration will henceforth be written directly in this document.

.1 Register table

C/K = correction, P = clarification, R = revision/register entry. Full texts: Doc. 190-Archive.

No.	Affects	Subject (short form; details: Archive)
C1	049 (HTML)	$\xi = \lambda_h^2 v^2 / (16\pi^3 m_h^2)$ [Addendum 28 July 2026, cf. R66: reason for the correction — $16\pi^3 = (4\pi) \cdot (4\pi^2) = S^2 \cdot 2S^3$ is the sphere count of the compact T^4 boundary geometry (Doc. 310); the earlier version counted one 2-sphere too many]
C2	116 (tab.)	$r_\tau = 25/9$
C3	018 (Explorer)	$a_e = 2\pi^2 / (f \cdot k)$
C4	267 (DE+EN), 268 (DE)	Factor 3 = three observable space dimensions: the observer measures $k_{\text{obs}}^2 = (n_1^2 + n_2^2 + n_3^2) / L^2$ (the fourth torus direction is time unfolding).
C5	275 (script v3)	All three fixed in v4 (11 June 2026): T^4 as kNN in genuine 4D coordinates with dimension-matched 4D nulls (result: $1,329 \pm 0,025$ inside the band $[0,690, 2,406]$); HLV ensemble with $\sigma = 0,05$ ($0,710 \pm 0,008$, in the band); the shell detect ...
P1	116 / 158	both correct; 0,001 % for external use
P2	several	$\hbar$ follows from ξ via $L_0 = \xi \cdot \ell_P$
P3	173–176	two parameters: ξ_0 (geom.) and ξ_{Higgs} (alg.)
P4	180–182	L_0 = geom. fundamental length
P5	several	mass structure from $\xi_i \sim 1\%$ agreement
P6	116 / 158	Koide only with numerical values (non-closure, Doc. 193)
P7	005, 008, 019, 086, 189, 202	scale structure from the recursion $\mathcal{R}$
P8	005, 008, 012, 014, 041, 060, 133, 141, 159, 181	geometric scale adjustment / fractal correction
P9	005, 006, 042, 046	columns "scaling class" + "SM correspondence"; quarks via $m = r \xi^p v$
P10	009	$K = R_{pe} / (4/3)^7$: an empirical anchor; factor 1,314 defined implicitly, not derived
P11	025, 003, 133	correct: $(1 - 275/4 \cdot \xi)$, $\Delta = 0,0002\%$; Doc. 003 and Doc. 133 use different models
P12	022, 230	two different observables: cumulative (Doc. 022) vs. elementary $\xi/(2\pi)$ (Doc. 230); 10^{-5} unreachable at any finite N from Doc. 022
P13	013, 025, 061	$\xi_{\text{FFGFT}} / \xi_{\text{UIFT}} = m_e c^2 / (\frac{16}{9} \ln 2 \cdot \text{eV}) \approx 414\,684$; table in Doc. 013 (archived) reconstructed

No.	Affects	Subject (short form; details: Archive)
P14a	041	fractal path lengthening (geometric path stretching); energy loss only an artifact when the path lengthening is omitted; path-length dependence remains (Doc. 026/149/182); the same path lengthening carries the CMB treatment in Doc. 268 (formerly tracked separately as P19)
P15	061 / 041	marked as a Λ CDM comparison quantity; the static universe has no cosmological z ; CMB by stationary thermalization
P16	026 / 064 / 181	H_0 is emergent (decompactified), not fundamental; implemented in Doc. 263, Docs. 026/064/181 noted as outdated
P17	028	the DE/DM relation is circular (gives 2.5 for any ξ); MOND partially derived ($\xi^{1/4}$ genuine, K_M free); CMB relations unaffected; noted
P18	267	the two pictures are <i>not</i> distinguishable through the observed ratios; the decision is theoretical, not empirical (conceptual placement)
P20	267 / 268	the <i>form</i> is fixed: a dimensionless ξ ratio (R_H/ℓ_p).
P29	268	<i>Only partially answered.</i> The factor 3 from the 3D projection ($3 \cdot \{1, 6, 14, 26\}$) is plausible, but the sequence $\{1, 6, 14, 26\}$ is <i>read off the data</i> (Doc. 268, step 4), not produced forward.
P30	268	the <i>naive</i> spherical Bessel sum $C_t = \sum_n N_{BE} j_t(k_{\text{obs}} D_A) ^2$ does <i>not</i> reproduce $\{3, 18, 42, 78\}$ – it washes out to a quasi-continuum dominated by low modes.
P31	268	<i>Leading behaviour derived forward.</i> Forward, T^4 gives the harmonic backbone $1 : 2 : 3 : 4 : 5$ (symmetric resonance $k^2 = 3n^2$).
P32	corpus-wide	follows from P20: H_0 is an <i>imported</i> external calibration, not derived.
P33	009 / 042 / 181	the <i>form</i> of ξ is forward: the harmonic $4/3$ (perfect fourth, Doc. 042/159) gives the leading digit, 3 (space dimensions) and $2^2=4$ (spacetime) are read off the geometry.
P34	181 / 013	$3+1$ is <i>empirical input</i> – no theory derives the dimension count; the earlier “follows necessarily” claim (Doc. 181) is overstated.
P35	corpus-wide (esp. 182, 250s, scale tables)	ξ^N triviality : “ $X = \xi^N$ ” is by itself <i>vacuous</i> , since any magnitude can be written as a ξ power – an exponent alone is not a prediction (cf. P10, P20, P33: form forward, number input).
P36	182 / cosmol. sector	R_H interpretation refined : $R_H = c/H_0$ is the <i>Hubble length</i> – the local expansion rate expressed as a distance – <i>not</i> the size of the universe and <i>not</i> the observable horizon (the latter $\approx 3 R_H$ in Λ CDM).
P37	009, 188, 271, 272, 275	Three levels clearly separated:
P38	275	(1) Microscopic – input: $\xi = 4/30000$ (fractal dimension).
P39	182, 267, 277, corpus-wide (H_0/R_H)	Status clarification: (a) $1/\varphi$ is a <i>pass-through point</i> , not a fixed point – the only fixed point of $r \mapsto r(1 - \xi)$ is 0. H_0/R_H status refined (shared reference point, not an FFGFT gap): the relation Planck length $\leftrightarrow R_H$ (exponent $41/4$, $R_H/L_0 = 6,49 \times 10^{64}$) is derived from first principles by <i>no</i> framework – why the universe is $\sim 10^{60}$ Plan...
C6	004, 025, 030, 039, 041, 053, 061, 063, 068, 081	Redshift is achromatic. The fractal path-lengthening is purely geometric and stretches <i>all</i> wavelengths by the same factor: $1 + z = e^{\xi x}$, independent of λ .
P40	005, 006, 011, 012, 133, 275	Only ratios are exact. Absolute values carry the fractal/SI correction, absorbed into the bridge constants v (masses $m = r \xi^p v$), $E_0 = 7,398 \text{ MeV}$ (α) and $3,521 \times 10^{-2}/2,843 \times 10^{-5}$ (G; calc_De.py); in ratios they cancel ($m_\mu/m_e = 206,768 \dots$

No.	Affects	Subject (short form; details: Archive)
P41	267, 280, 221 (cos-mological sector)	Checked and not booked (degeneracy risk at the shared scale). Λ CDM signal $\Delta v = c \dot{z} \Delta t / (1+z)$ over 25 years: +6 cm/s ($z=0.5$), sign change at $z_0 = 1.92$, -14 cm/s ($z=4$), -20 cm/s ($z=5$); ELT/ANDES order $\sigma \sim 2$ cm/s.
P42	282, 292 (lepton sector)	One declared reference point: the ratio m_μ/m_e. The framework fundamentally keeps the purely theoretical derivation from ξ ; the lepton sector receives – analogous to P39 (H_0/R_H as shared empirical reference) – exactly <i>one</i> declared reference point...
P43	006, 011, 292 (calc_De.py)	No rework; recorded as a note. Verified: calc_De.py does <i>not</i> use $K_{\text{frak}} = 1 - 100\xi$ explicitly.
R41	284 (cf. 270, 249)	The loss is not the SI conversion. Per Doc. 270 the Type I duality decompactification $S_m^1 \rightarrow \mathbb{R}_t$ ($\tilde{T} \cdot m = 1$) is a <i>mode-preserving embedding</i> (lossless); compactness survives as the discrete spectrum and as physical observables (T_{rev} ...
R42	270 (cf. 248)	Not an open geometric question – it is the measurement. In the compact view the four circles are symmetric (no space/-time distinction); time is not selected by the geometry but is <i>relational</i> – the symmetry-breaking <i>is</i> the measurement (the Type I decompact ...
R43	285, 283	Corrected to containment. $C_3 < A_5$ (3-fold and 5-fold both permute the same icosahedron, recompute_reconcile.py) and $T^7 = T^4 \times T^3$ ($7 = 4 + 3$), so HLV $\supset$ FFGFT at the group/dimension level – the models are reconcilable, not inco ...
R44	272, 190 (P35), 285	Sharpened (no contradiction). φ acquires meaning <i>not</i> inside FFGFT but as the derived eigenvalue of HLV's 5-fold / perp sector (golden inflation $\varphi^2 = \varphi + 1$) – a marker of the boundary HLV \ FFGFT.
R45	285, 282–284	Sharpened. A quasicrystal has two spectra: the <i>structure / diffraction</i> spectrum is pure point (sharp Bragg peaks, $\approx 21\times$ above a random null, quasicrystal_information.py) – that is the information and exactly HLV's substrate-distinguishabilit ...
R46	296 (cf. 025, 026, 063, 156)	Open check-point (not yet processed). This intermediate-scale formulation potentially runs <i>counter</i> to the already established corpus line, which treats dark matter/energy not as an open signature but as a <i>derived consequence</i> : Doc. 025 reads dark matter as a ...
R47	285, 297, 298 (cf. 286, 284, 295)	In the time sector: containment, not enlargement. In unified notation – both D_6 +time, Marcel as D_6 +spiral-time, FFGFT in Hilbert space as D_6 +recursion-time – the spiral-time window is not additive but <i>internal</i> : a bounded section (window) of the unrolled re ...
R48	304 (cf. 295, 301, 303)	Note withdrawn, document made precise (not repaired). The overstatement lay solely in the <i>mail</i> and was withdrawn without reservation; the document was written throughout from the FFGFT reading and already booked the bridge as restricted.
R49	304, 305 (cf. 295, 303, 272)	Vocabulary cited, marked as analysis, abstract de-symbolised — notation not renamed. A consolidated first-use footnote attributes U1/U2/U3 and M_r/M_B to Krüger (2026) with the primary source (doi:10.5281/zenodo.21302278) and documented lineage (Me ...
R50	025, 063, 061, 064, 078 (superseded by 306; cf. 003, 010, 295, 298, 302)	Justification replaced, conclusion unchanged. The conclusion (no singular beginning; a finite core with a minimal scale L_0 set by ξ) stands.

No.	Affects	Subject (short form; details: Archive)
R51	049 (cf. 095, 306, 267, 302; notation)	A notational relic, not a substantive contradiction — and nonetheless no longer authoritative. (a) <i>The resolution:</i> in the original T0 view, T_0 never denoted the instant zero but the <i>smallest possible time interval</i> .
R52	101, 165 (numerical values; cf. 180, 182, 013, 250, 290)	Values corrected; the main relation is unaffected. (i) <i>Wrong</i> L_0 : correct is $L_0 = \xi \ell_P = \frac{4}{30000} \cdot 1.616 \times 10^{-35} \text{ m} = \boxed{2.155 \times 10^{-39} \text{ m}}$ — as in Doc. 180 (the dedicated derivation), Docs. 013, 181, 182, 187 ...
R53	267 §518 (table row <i>source of the scale</i> ; cf. 182, 165, P20/P35/P36)	Corrected reading — and the symmetry becomes sharper, not weaker. (a) On the FFGFT side, the table row <i>source of the scale</i> correctly reads not “from ξ ” but “ set / calibrated to H_0 ”.
R54	292, 306 (cf. 307)	(a) N_0 is not a base unit. $N_0 \approx 38.6$ is a <i>ladder-internal</i> , generation-counting factor ($N_g = g N_0$, Dok. 292); its value is <i>open</i> (P35: $100/e$, $100/\varphi^2$, 39 do not hit it reliably).
R55	292 §Cross-ladder coupling (cf. R54, P35, P40/P41)	The law stands; its evidential breadth is one entry, not three. (i) <i>The third entry is forced arithmetically.</i> From $m_\tau/m_e = (m_\tau/m_\mu)(m_\mu/m_e)$ a uniformly multiplicative correction satisfies $K^{p_3} = K^{p_1} K^{p_2}$, hence $p_3 = p_1 + p_2$ <i>identically</i> ...
R56	188 §Special role of $n = 3$ (cf. 189 §Derivation hierarchy, 248, 230/231/232, R55)	The three elements are hereby booked uniformly as foundational assumptions. (i) <i>The declared framework.</i> FFGFT posits three structural elements: the fundamental relation $\tilde{T} \cdot m = 1$ (Doc. 189, level 1), the topology T^4 (Doc. 248), and the order Z_3 ...
R57	A015 (cf. 020, 030, A-Series)	Demand dissolved — ξ is a fundamental parameter. A proof is neither possible nor necessary; the demand was wrongly posed.
R58	A142, A140 (cf. 127, 163)	Gravitational quantisation not required. $G = \xi^2/(4m_{\text{char}})$ is intrinsic to the torus geometry; a separate quantum gravity does not arise, because the geometry T^4/Z_3 is pre-ordered and not dynamical in the same sense as quantum-field-theoretic fields.
R59	A130 (cf. 011, 043, 101; P40)	Structurally explained [S]. Two causes: (i) SI values are scheme-dependent (1-loop, $\overline{\text{MS}}$), i.e. the EFT precision values are not theory-independent raw data; (ii) fractal corrections in v are not cleanly separable from the loop structure of the SM ...
R60	A160, A165 (cf. 020, 022, 023, 230; P35)	Both factors now geometrically derived [B]. (i) 2π is the circumference of the unit circle — the natural conversion between the torus winding phase and the measurement angle; $\xi/(2\pi)$ is thus the phase correction per measurement.
R61	174, 175 (cf. A142, 147, 162; R57, P35)	Data-driven value selection — not a second parameter; withdrawn. (i) <i>The genesis:</i> ξ_{Higgs} was selected <i>because</i> it lands in the measured window; the two-spaces doctrine is rationalisation after the selection, not independent justification.
R62	005 §Baryon worked example (proton) (cf. R50, P35)	The anchor of the formula line is $\pi^2/2$, not m_μ. (i) <i>Diagnosis.</i> The missing factor identifies the error as a single symbol: $(\pi^2/2)/m_\mu = 46.705$ hits the missing factor to 0.07 %.

No.	Affects	Subject (short form; details: Archive)
R63	253 § ξ reso- nance heuris- tic (code: rsa/ factorization_ benchmark_ library.py; cf. 024, 075, 076, 176; R57, P35)	ξ is inert in the “optimal” range; the optimisation switched the filter off. (i) <i>Inertness</i> . The code threshold is 1/1000.
P44	253, 190 (R63); cf. A135, P35	Pre-registered, tested, and refuted for this design; the window width is a property of the measurement setup. (i) <i>Design</i> . Shor phase estimation, 12 counting qubits (phase resolution $2^{-12} = 2.44 \times 10^{-4}$), 8 shots, 10 % fully random outcomes, 300 ...
R64	2/python/t0_no_fall	The procedure is right, the separating cut too coarse, and the label wrong. (i) <i>What holds</i> . The counters t0_success_count and fallback_success_count are incremented at separate places (l. 376 resp. 280/300), get_statistics() report...
R65	024, 075, 076, 147, 176, 253 as well as the	No procedure distinguishable from Shor exists; the method claim is withdrawn. (i) <i>What is withdrawn</i> . (a) The claim of an independent T0 resp. ξ factorisation procedure.
R66	097 (DE/EN and the standalone versions under	The $16\pi^3$ count is right — but not for the reason given in Doc. 097.
C7	the fourteen HTML pages under rsa/	Rebuilt, not annotated: the pages now say what they do. No correction box was inserted; what was corrected is the computation, the labelling and the method text.
R67	A040 (fractal correction), A270 (bulk exponent 36),	(i) With $D_f^{\text{eff}} = 2.973$ (three decimals) the rounding interval leaves $n \in [98.3, 102.0]$ open; even $D = 2.973$ exactly gives $n = 100.12$.
R68	279 (H_0 chain), 308 (Λ reading,	(i) <i>Identification</i> : the Λ function is carried by the derived quantity $\Lambda^* := 1/R_H^2 = (\pi/2)^2 \xi^{20} / \bar{\lambda}_e^2 = 5.218 \times 10^{-53} \text{ m}^{-2}$ (“closure scale”).
R69	308 § a_0 derivation; cf. P20, R68;	(i) From $\partial T^4 = \emptyset$ (Doc. 306) an isotropic torus yields a compact time cycle $\tau_c = 2\pi/H_0 = 91.9 \text{ Gyr}$ [K P20]; the energy quantum is exactly $\hbar H_0$, identical to the momentum quantum of the spatial cycles.
R70	061 (tempera- ture/CMB), A085 (natural units),	T_0 scale gap
R71	309 (scale anchor, inheritance prob- lem);	sharpening: side-taking in the Hubble tension
R72	A010 (glossary), A080 (units), A265	E_0 : bare and corrected
R73	026 (geometric cosmology), 279	H_0 notations, exponent 41/4
R74	025 (Cosmology), 063 (Cosmic), DE+EN each,	lithium problem: mechanism, not solution
R75	024, 034, 075, 076, 147, 176: ξ name collision and rewrite of the Shor documents	In all affected documents the search parameter is now denoted σ and placed as a sampling grid without physical meaning.
R76	318	Scope of mass derivations
R77	041, 160, 318	Classification of early coupling constant formulas

No.	Affects	Subject (short form; details: Archive)
R78	160, 005	α_s and running coupling
R79	321, 319, 049	Algebraic derivation of the $SU(3)_c$ gauge group
R81	323, 321, 160, 320	Weinberg angle at M_Z from ξ
R82	322, 320	Hilbert-space embedding and spectral theory
R83	319, 321, 323, 322, 320	Status update: closed bridges
R84	324, 321, 009	Casimir derivation of ξ ; ξ not a G(6) spectral value
R85	313, 325, 257, 321, 322	Hawking mechanism closed microscopically
R86	Docs. 180, 322, 314, 321, 325	Self-adjointness of $\hat{F}$ fully closed; $\xi = \lambda_{\min}(\hat{F}_{D_4})$ [B]
R87	Docs. 320, 322, 340	Δm_{32}^2 : numerical values corrected, dev. +16.0 %; candidate analysis added [K] . Closed by Doc. 340 (R103): Galois derivation $\Delta m_{atm}^2 = 120 m_\nu^2$ from $GF(27)^*$ orbits, dev. 1.0 % [K] .
R88	Docs. 326, 329	n_{thresh} scale-dependent: $n(L) = L/(\sqrt{2}\ell_p)$; Matzke's scale = $2\pi\ell_p/\ln 2$ [B] , universal number [X]
R89	Docs. 180, 019, 201, 032	$m_T = M_{\text{base}}/\xi$; one structure, sector-dependent base mass; range = $\xi \cdot \lambda_C(M)$ [B]
R90	Docs. 180, 019, 201	Exponent in $L_0 = \xi^1 \ell_p$ derived from $m_T \sim 1/\xi$, not by analogy [B]
R91	Docs. 019, 201	λ in $m_T = \lambda/\xi$ is a base mass, not the Higgs quartic — label misleading [K]
R92	Docs. 250, 190	Doc. 250: ξ/ℓ is the coupling g_{T_i} , not m_{T_i} ; Schwarzschild check for L_0 is an identity [B]
R93	Docs. 325, 329	$S_{\text{BH}}(M_{\text{coll}}) = 2\pi \text{nat} = n_{\text{thresh}}$ (Matzke) in bits; residual entropy of the endpoint [B]
R94	Doc. 325	$I_{\text{sector}}/I_{\text{thermal}} = \log_2 3$ for all M ; structural constant of the $\mathbb{Z}_3$ geometry [B]
R95	Docs. 330, 270, 314, 327	Two wording refinements (flatness: $K=0$ pointwise via metric descent; s.a.: bounded <i>plus</i> symmetric); Type-I theorem [sketch]→ [B] (function pullback p^*); D_4 specificity test open [S]
R96	Docs. 330, 332, 270	Type-I follow-up refinement after second Krüger feedback: $\text{Per}_{R_m}(\mathbb{R}_t)$ with per-period norm is isometrically isomorphic to $L^2(S_m^1)$, not to $L^2(\mathbb{R}_t)$ – pullback remains a change of representation, not a new sector [B] ; Nash–Kuiper attribution in Doc. 330/R95(i) corrected (existence theorem for C^1 , no C^2 non-embeddability claim) [K]
R97	Dok. 324	Vacuum operator correction following Doug Matzke's objection (IPI-Mail 26.Aug.2026): Matzkes $V = N_1^+ N_2^+ N_3^+ N_1^- N_2^- N_3^- = -64 \cdot P_{N_1} P_{N_2} P_{N_3}$ is an involution in $\mathbb{Z}_3\mathbb{C}$ ($V^2 \equiv -V \pmod{3}$) [B] ; Dok. 324 incorrectly identified the Pp-product $P_{P_1} P_{P_2} P_{P_3}$ (complementary sector, different ray: $ \langle \text{vac}_V \text{vac}_{P_P} \rangle = 0$) with Matzke's V ; numerically verified by <code>pruef_324_vakuum_involution.py</code> (22 Assertions); Trine-Theorem $T_k^3 = \mathbf{1}$ und ξ -Nicht-Spektralwert remain correct [B]

No.	Affects	Subject (short form; details: Archive)
R98	Doc. 333, 332, 315, 295, 285	Two meanings of “unrolling” in the corpus distinguished and the recursion bound to the $\tilde{T} \cdot m$ duality (Doc. 333, 27 Aug. 2026): (i) Type-I (measurement act, lossy, prior corpus usage Doc. 285): $S_m^1 \rightarrow \mathbb{R}_t$, winding number discarded; (ii) Type-III (canonical function pullback, lossless, Doc. 332/R96): $L^2(S_m^1) \cong \text{Per}_{R_m}(\mathbb{R}_t)$, exact change of representation, m and t unroll together [B] . K_{frak} plays no role in the Type-III pullback; it sits in the choice of R_m via the mass formula and becomes visible only at the SI transition [B] . The recursion $\xi_{n+1} = \xi_n(1 - 100 \xi_n)$ (Doc. 295/146) follows necessarily from $\tilde{T} \cdot m = 1$: every mass change forces a time change, which feeds back recursively. Case B (stable particle, $K_{\text{frak}} = 74/75$) is the projection of Case C (running recursion, log-spiral) onto the static approximation [B] . Doc. 332 reduced to a short cross-reference.
R99	Doc. 334, 307, 306, 285, 174	Superposition without time (Doc. 334, 27 Aug. 2026): (i) classical physics and QM inherit tacit instantaneity — time runs externally, calculations have no duration [K] ; (ii) hidden inconsistency: state $\psi(t)$ is instantaneous, but probabilistic evaluation $ \psi(t) ^2$ requires real time for verification — this tension is absent from the formalism [K] ; (iii) superposition is doubly timeless: no intrinsic “when”, no verification without time expenditure; (iv) collapse has no duration in the QM formalism; (v) Pauli theorem does not apply to compact $\tilde{T}$ (angle variable); (vi) $\tilde{T} \cdot m = 1$ gives every mode an intrinsic time scale — superposition within one mode has scale $\tilde{T}_i$, cross-mode gives incompatible $\tilde{T}_i \neq \tilde{T}_j$ [B]
R100	Doc. 336, 324, 329	Notation updated to symmetric $\mathbb{Z}_3\mathbb{C}$ (after Matzke, IPI mail 28 Aug 2026): elements $a + bi$ with $a, b \in \{-1, 0, +1\}$ instead of $\{0, 1, 2\}$; $+1 + 1 = -1$ (ternary inline carry) [E] . Frobenius fixed points $= \{0, +1, -1\} = \text{FFGFT sectors } \Delta_w$ [B] ; norm classes $\{+1, -1\}$ instead of $\{1, 2\}$; tripotent eigenvalues $\{0, +1, -1\}$ [B] . Verification scripts 324–327 updated to symmetric $\mathbb{Z}_3\mathbb{C}$ (all assertions passed).
R101	Doc. 317, 336, 338	$\xi = 4/30000$ derived as Galois number [K] : core identity $(r_\mu/r_e)^2/\xi = \text{GF}(9)^* ^2 \cdot 5^2 \cdot \text{GF}(27) = 43200$ (pruef_331). All six winding numbers $(n_{\theta,g}, n_{\phi,g})$ from $\text{GF}(3^n)$ orders; recursion $n_{\theta,g} = n_{\theta,g-1} + n_{\phi,g-1}$ [K] .
R102	Doc. 338	$1/\alpha = 3700/27 = 137.037$ (dev. 7.6 ppm) from Galois group orders [K] (pruef_332): new observation $m_e \cdot m_\mu = 54 = \text{GF}(3)^* \cdot \text{GF}(27) \text{ MeV}^2$ (0.016 %; Doc. 011 gives geometric derivation). $ \text{GF}(27) $ cancels; no ξ , no v , no m_e as explicit input.
R103	Doc. 339, 340	Frobenius decomposition $\text{GF}(27)^* \cong \mathbb{Z}_2 \times \mathbb{Z}_{13}$ (Doc. 339): fixed points $\{+1, -1\} = \text{massive sector}$; 8 three-orbits = gluons; $\text{Orbit}_4 = 2^{-1} \cdot \text{Orbit}_1$ [B] . Neutrino mass hierarchy (Doc. 340): $\Delta m_{\text{atm}}^2 = 120 m_\nu^2$ (1.0 %) [K] ; $\Delta m_{\text{sol}}^2 = (11/3) m_\nu^2$ (0.46 %) [B] ; $p_{\text{vent}} = 5/24$ via branching rule [B] ; $\theta_{13}: (25\xi)^{1/3}$ (1.4 %) [K] and $2/3 \cdot R_v$ (7.4 %) [K] ; $\text{gcd}(13, 24) = 1$, spiral lcm = 312 [B] . Check scripts: pruef_340 (13/13), pruef_340b (6/6).
R104	Doc. 006, 338, 319, 340	$v = m_e/((4/3)\xi^{3/2}) = 248.3 \text{ GeV}$ (0.85 %) [K] ; $G_F = 1/(\sqrt{2} v^2)$ (1.7 %) [K] ; $\tau_n \propto G_F^{-2} = 909 \text{ s}$ (3.4 %) [K] ; $\Delta m = \frac{39}{8} m_e - (27/3700) \cdot \Lambda_{\text{QCD}} \cdot \frac{5}{6} = 1.2912 \text{ MeV}$ (0.17 %) [K] : QCD part from r_d, r_u, r_e (Doc. 006); EM part from $\alpha = 27/3700$ (R102), $\Lambda_{\text{QCD}} = \hbar c/\text{fm}$ (Doc. 319), $5 = A_5 : A_4 $ (Doc. 340), $6 = r_u$ (Doc. 006). Check script: pruef_319 (7/7).

No.	Affects	Subject (short form; details: Archive)
R105	Doc. 006, 011, 070, 182, 231, 257, 285, 306, 307	Retrospective marker certification — documents created before the marker system (1 Sept 2026). These documents predate the introduction of the marker system (A010) and carry no markers of their own. Their derivations are internally consistent and are necessary chain links for the Galois programme (Docs. 336, 338, 339) and further documents. Retrospective status: Doc. 006 (particle masses, winding numbers r_i, p_i) [K] ; Doc. 011 (fine structure α, E_0 , fundamental relations) [K] ; Doc. 231 (Hilbert space extension, L^2 structure) [B] ; Doc. 257 (information unit, bit-energy scale) [K] ; Doc. 285 (FFGFT-HLV dimension bridge) [K] ; Doc. 306 (native time-energy reciprocity from $\tilde{T} \cdot m = 1$) [K] ; Doc. 307 (time in state space) [K] ; Doc. 182 (maximum universe scale from ξ , Schwarzschild radius and Hubble scale) [K] ; Doc. 070 (mathematical structure: K_{frak} cancels exactly in ratios, absolute values carry the correction) [B] . Dependent documents may treat these results as a certified chain link by citing R105, without modifying the original documents.
R106	Docs. 342, 343	Primes forced by $\text{GF}(3^k)$ [B] : the physical primes $\{5, 7, 11, 13\}$ are completely forced by the prime factorisations of $3^k - 1$ ($k = 3, 4, 5, 6$); $p = 13$ is the worldwide unique prime with $\text{ord}_p(3) = 3$. Euler factor threshold $p^* \approx 9.76$ ($s = 2$) = 7-limit; Galois product density at chance level from $p_{\text{max}} \approx 30$ (pruef_342, pruef_344).
R107	Docs. 343, 338, 162, 186, 328	ξ as resolution floor [B/S] : the limit $k \approx 3-4$ (Theorem F) holds for $\varepsilon = 1.05\%$, one fractal level ($\approx 100\xi$, Doc. 162). At $\varepsilon = \xi$ (torus-mode linewidth, Doc. 328) the Galois grid stays resolved to $k \geq 8$, $p^* \approx 87$. Coupling: Farey $Q_c = \pi/\sqrt{6\varepsilon} \approx 12.5$; Arnold K^q vs. Farey $1/q^2$ — same tripartition, different law [B] ; FFGFT with $K_{\text{eff}} = 2\pi\xi$ dynamically deeply subcritical, discreteness topological [B] . TFLN $Q > 10^6$ resolves ξ , not ξ^2 . Doc. 186 preliminary note: locking [B] , Fibonacci [B] , B meson [X] corrected; renamed to _De/_En. Resonance factoriser capped at $N_{\text{max}} \approx 1/\varepsilon$ (13–26 bit); quantum computers/photonics as analogue computers with QM precision, curve at $N = 21$ since 2012 [X] (pruef_343c-e).
R108	Docs. 340, 342, 343	Dirac/Majorana partition of the $\text{GF}(27)^*$ classes [B] : sector pairing $k \mapsto -k$ and Majorana criterion $k \mapsto k^{-1} \pmod{13}$ partition the four primitive classes identically — $\{f_1, f_2\}$ (orbits O_1, O_3 , quadratic residues mod 13) Majorana, $\{f_3, f_4\}$ (O_2, O_4) Dirac. Canonical: Majorana layer = kernel of the quadratic character mod 13. Numbering unified per the polynomial table of Doc. 342 (sector pairing $f_1 \leftrightarrow f_2, f_3 \leftrightarrow f_4$; negation $f_i \leftrightarrow g_i$); exponent list in Doc. 342 corrected (f_3, f_4 were the same orbit). Consequence for Doc. 340: $v_1 \in O_1, v_2 \in O_3$ Majorana; $v_3 \in O_4$ Dirac neutrino [B] — no Majorana phase, m_{ee} from v_1, v_2 only (≤ 14.4 meV), inverted hierarchy more strongly excluded; sterile neutrinos drop out as a separate prediction. Doc. 340 revised directly (pruef_343f). Precursor: Doc. 007 derives $m_{\nu_1} = 4.54$ meV from $\xi^2/2 \times m_e$ [K] and motivates the double ξ suppression physically (neutrino = almost photon + geometric decoupling); flavour hierarchy first in Doc. 340. Addendum inserted in Doc. 007.

No.	Affects	Subject (short form; details: Archive)
R110	Docs. 320, 022	Addenda on ν_3 Dirac neutrino (Doc. 343 Theorem D''', R108): Doc. 320 (spectral theory) — fixed-point derivation F_2, F_5, F_7 provides no Majorana/Dirac character; this follows from the Galois algebra of Doc. 340/343. Doc. 022 (QFT-ML addendum) — ξ^3 sterile suppression in DUNE row not to be interpreted as a sterile signal; Majorana layer fully occupied by ν_1, ν_2 . Authoritative: Doc. 340/343.
R111	Doc. 344, 345, 346, 347, 348	Charge quantisation, Gell-Mann–Nishijima, and generation structure from GF(27)* (4 Sept. 2026). Doc. 344: $\text{Su}[0]=\text{Vss}[0]$ exactly — neutrino = vacuum of Clifford-Fock space [B] ; sector pairing $k \mapsto -k$ has no fixed point among primitive classes $\Rightarrow$ all neutrinos Dirac [B/S] ; nEXO (5–20 meV) decides. Doc. 345: literature comparison Furey/Dixon/Froggatt-Nielsen; FN: ε^{q_i} free vs. ξ^{p_i} forced [K] ; after Docs. 346/347 equal to Furey/Dixon at charge content. Doc. 346: two Galois properties classify all SM fermions [B] : (i) quadratic residue mod 13 $\Leftrightarrow$ electrically neutral; (ii) $N \bmod 3 \Leftrightarrow$ colour charge; forbidden combination QR+triplet algebraically forced; Legendre symbol $(k/13)$ gives $Q \in \{0, \pm\frac{1}{3}, \pm\frac{2}{3}, \pm 1\}$ [B] . Doc. 347: $Q = I_3 + Y/2$ algebraically derived [B] : $Y = -1 + \frac{4}{3}\text{NQR}$ from Legendre symbol (Doc. 346), $I_3 = \pm\frac{1}{2}$ from jE7 projector (Doc. 344); all four fermion charges exact. Theorem E [S] of Doc. 346 closed. Doc. 348: root orbits f_1-f_4 are genuine Frobenius orbits; 3 elements = 3 generations [B] . Trace bilinear form: $f_3 \times f_4$ (charged leptons $\times$ quarks) = permutation matrix (Gen.2 $\leftrightarrow$ Gen.3 forced) [B] ; $f_1 \times f_3$: entries $1/\sqrt{2}$ [B] , consistent with large PMNS angles. CKM complexity: $\delta_{CP} \approx 68^\circ$ not derivable from GF(27)* ($\mathbb{Z}_3$ phases give 120°); usable as phase anchor (same status as $\nu = 246$ GeV). Check scripts: pruef_346, pruef_347, pruef_348, all assertions passed.
R109	Doc. 006 (restriction of R105)	Neutrino part of Doc. 006 not tenable [X]. The “direct method” $E_i = 1/\xi_i$ is not a computation: $1/(\xi_{r_i})$ gives $e:\mu = 1:0.42$ (inverse hierarchy, dimensionless); the MeV values shown alongside are measured values. The neutrino masses $9.1/1.9/18.0$ meV follow from three incompatible prescriptions ($\nu_e = m_e \xi^2$; $\nu_\mu = r_\mu \xi^3 \nu_\tau$ from none); the exponent $p_{\nu_\tau} = 25/9$ is r_τ copied into the exponent column. The Yukawa method $m_i = r_i \xi^{p_i} \nu$ remains valid for charged leptons and quarks ($< 1.2\%$). The [K] certification in R105 applies to Doc. 006 only for r_i, p_i and the Yukawa method, not for neutrinos and not for the claimed equivalence of both methods. Authoritative for neutrinos: Doc. 340 (R103, R108). Doc. 343 D''' records the discrepancy [S] .
R112	Doc. 006, 319, R104; Doc. 351	$\nu = 246$ GeV is not a measurement but an SM definition [Q] (5 Sept. 2026). $\nu \equiv (\sqrt{2} G_F)^{-1/2}$ follows from $M_W = g\nu/2$ and $G_F/\sqrt{2} = g^2/(8M_W^2)$; only $\tau_\mu = 2.196\,980\,3(22)$ μs (MuLan) is measured, G_F follows via $\tau_\mu^{-1} = G_F^2 m_\mu^5 / (192\pi^3) (1 + \Delta q)$ with $\Delta q = -0.44\%$ (QED $O(\alpha^3)$, hadr. VP with 4.7σ CMD-3 tension; Eberhart et al. 2026) and Sirlin absorption of EW corrections. The irrational factors $192\pi^3$ and $\sqrt{2}$ come from the SM formulation, not from measurement. Corpus-wide wording correction: “ $\nu = 246$ GeV (measured)” $\rightarrow$ “ $\nu = 246$ GeV (SM definition from G_F)”; the FFGFT value $\nu = 245.6$ GeV (R104) is a comparison of structure against convention. Convention-free test only at the τ_μ level — open [S] .

No.	Affects	Subject (short form; details: Archive)
R113	Doc. 338, 317, 011, R100, R101; Doc. 351	Anchor m_μ/m_e is QED-dependent; comparison values are not theory-free [Q] (5 Sept. 2026). $m_\mu/m_e = 206.768\,277(24)$ derives solely from muonium HFS (LAMPF 1999, Liu et al.); extraction via $\nu_F = \frac{16}{3}\alpha^2 c R_\infty (m_e/m_\mu)(m_r/m_e)^3$ — the anchor implicitly contains α^2; Eides (2026): CODATA underestimates the theory uncertainty by a factor of 5–10 (271–515 Hz instead of 51 Hz). α: Rb (Morel 2020) and Cs (Parker 2018) disagree at $> 5\sigma$; the CODATA average covers neither. m_r: two methods (threshold scan BESIII 1776.91, pseudomass Belle II 1777.09 with $m_{\nu_e} = 0$) differ by 0.18 MeV; PDG average 1776.93(9), FFGFT 1776.97 lies 0.4σ — test undecided. Exact computation: the residual $43200 - 42753.12 = 446.9$ in $(m_\mu/m_e)^2$ is $240\,000\sigma$ of the CODATA uncertainty; no measurement error explains it; it is an empirical discrepancy [K] (Doc. 352, §9–10); physical attribution (R113 bridge) open [S]. Wording correction: “model-independently measured” → “least model-dependent available value”. Bookkeeping rule: every prediction states anchor, number of steps, corrections applied, extraction method of the comparison value. All 21 sources verified (Doc. 351) Three Galois identities compared in rational arithmetic (verification script <code>pruef_351_rationale_vergleiche.py</code>): (1) $(m_\mu/m_e)^2 = 43200$: residual -1.034% ($240\,000\sigma$), physical [B]. (2) m_μ/m_e vs. $\sqrt{43200} = 120\sqrt{3}$: residual -0.52% is <i>not an independent finding</i> — $\sqrt{43200}$ is irrational; the residual is exactly half the quadratic residual (artefact of taking the square root). (3) $m_e \cdot m_\mu = 54\text{ MeV}^2$ ($= GF(3)^* \cdot GF(27) = 2 \cdot 27$): residual -0.016% (7386σ), physical, 65 times smaller than the quadratic residual; no computational artefact (rounding uncertainty 2.2% of measurement uncertainty), not covered by measurement tolerance [K]. Bookkeeping rule: Galois identities should be compared only in their natural algebraic form — where no irrational numbers appear.

No.	Affects	Subject (short form; details: Archive)
R114	Doc. 351 (update), Doc. 352 (new); R113	Lepton residuals: lepton ladder, Galois and Koide compared; R113 attribution clarified [Q] (5 September 2026). Doc. 352 compares three FFGFT structures (lepton ladder Docs. 006/317, Galois identities Doc. 351, Koide relation) with PDG and expresses all residuals in units of ξ. Main findings: (1) Koide matches PDG to $0.046 \cdot \xi$ — most precise prediction [K]. (2) Galois difference residual $-77.6 \cdot \xi = -2 \times (+39.1 \cdot \xi)$ lepton ladder algebraically forced [B]. (3) Generation-dependent corrections ε_i (order $73-192 \cdot \xi$, decreasing with generation) explain transition $Q_{\text{bare}}(+8 \cdot \xi) \rightarrow Q_{\text{PDG}}(-0.05 \cdot \xi)$ to 101% [K]. (4) Z_3 circulant: angle $\theta = 2/9$ from Doc. A110 [K]; amplitude $d/c = \sqrt{2}$ (Koide condition) and θ are orthogonal parameters [B] — $Q = \frac{1}{3} + \frac{d^2}{6c^2}$ does not contain θ. (5) Status separation of the ε_i (clarified 6 September 2026): $\varepsilon_i = m_i^{\text{PDG}}/m_i^{\text{bare}} - 1$ is a <i>computed residual</i> [K] from PDG [Q] against FFGFT bare [K] — not a number taken from QED literature; its ξ scaling is observed [K]; the assignment to QED extraction + SI projection is an <i>attribution</i> [S], not a citation; the reason for the falling structure $\varepsilon_e > \varepsilon_\mu > \varepsilon_\tau$ is open [S]. The QED self-energy formula $\delta m/m \approx \frac{3\alpha}{4\pi} \ln(m/\mu)$ in Doc. 352 §10 is [Q] (standard formula) but yields only the order of magnitude, not the ε_i values themselves. Column “Origin” in Doc. 352 §9 → “Presumed origin [S]”. Doc. 351 updated: three occurrences of “physical (R113)” → “empirical discrepancy [K]” (Doc. 352, §9–10); physical attribution (R113 bridge) open [S]. Verification script <code>pruef_352_leptonreste.py</code> (32/32) [K].
R120	Docs. 180, 201, 321, 307	Status of Maxwell dynamics: imported input, no derivation (14 September 2026). An external reading attributed to the corpus a “sketched derivation of the full dynamical Maxwell system” with status [S], citing Docs. 201 and 307. Check against the chapter files: neither document treats Maxwell. Doc. 201 takes the field-strength term $-\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$ over as a component of the extended Lagrangian; Doc. 180 explicitly sets the Standard Model Lagrangian including this term as its starting point; Doc. 307 contains only the Kaluza–Klein reading of charge quantisation. Retroactive status assignment: the kinetic term and the field equations that follow from it are [E] in Docs. 180 and 201 (established physics, imported) — neither [S] nor an open bridge. What the corpus does derive is the gauge group $U(1)_Y$ from the fourth torus direction [B] (Doc. 321, R80), charge quantisation and Gell-Mann–Nishijima [B] (Docs. 346/347, R111), and α from ξ [K]. A derivation of Maxwell dynamics from the T^4/Z_3 geometry is neither claimed in the corpus nor listed as open.

No.	Affects	Subject (short form; details: Archive)
R121	Doc. 352 (§ 10), Doc. 369 (new)	Origin of the ε_i: second reading without QED bridge (20 September 2026). Doc. 352 § 10 takes the T0 ladder as fundamental and attributes the generation-dependent residuals ε_i to QED extraction and SI projection (attribution [S], R113 bridge open). Doc. 367 permits a second reading: the Koide formula with $\theta = 2/9$ is fundamental and parameter-free from the icosahedron; the T0 ladder $r_k \xi^{p_k}$ is then a rational approximation to the transcendental Koide amplitudes, and the ε_i are the distance between cosine and power law — without PDG, without QED [K]. In this reading the ε_i differ because the three cosine values differ; the decreasing trend needs no external physics. Both readings reproduce the same numbers. Status: the open point “why do ε_i decrease” (R113/R114) is answered in Reading B; what remains open instead is why the rational approximation parameters are exactly the $GF(3^n)$ winding numbers [S]. The R113 bridge thereby moves from “needed” to “needed only in Reading A”. Doc. 352 § 10 remains valid as Reading A; Doc. 369 § 4 places both side by side.
R122	Doc. 041, 005 (Higgs mass, Λ_{QCD}); Doc. 372 (§ 3.6)	Higgs mass not derived (21 September 2026) [X]. The parameter table of Doc. 041 gives the Higgs mass as a power of ξ times v and lists the measured value beside it; recomputed, the formula yields barely a fifth of that. The same table gives Λ_{QCD} as another power of ξ times v ; the formula is off by almost two orders of magnitude. Doc. 005 gives the Higgs mass as top mass times golden ratio with a fractal correction factor; that factor would have to exceed the scale ξ by several thousand, hence it is a fitted parameter, not a derivation. Both documents remain unchanged; the status of the Higgs mass in the corpus is [X]: no viable derivation. The A_5 matrix-element spectrum (Doc. 372) contains no matching value either. The remaining rows of the Doc. 041 table are unaffected; v is governed by Doc. 006 (R104). Status (Doc. 373, 21 September 2026): For the Higgs mass there are now two compatible candidates [K]: the Galois chain linking Z , Higgs and top, and the trace rule, by which the Higgs particle is the residual stiffness of the lattice left over by the gauge bosons. Trace conservation, counting and normalisation factor are justified; together with the Weinberg angle of Doc. 372 the electroweak boson spectrum follows from v alone at tree level. The formulas in Doc. 041/005 remain wrong. The trace rule applies to the lattice modes only; for the top the Galois chain applies, the reading of an equal fermion budget is discarded. Still open: Λ_{QCD} from ξ and the choice of v (R104/R112).
R123	Doc. 160; Doc. 133, Clarification 11	Naming conflict K_{frak} in Doc. 160 (21 September 2026). Doc. 160 uses the label K_{frak} for a hadronic factor that does not agree with the primary K_{frak} of Doc. 133 (fixed in Clarification 11), lies on the other side of one, and derives from a Koide-type relation. It is a different quantity; in the sense of Doc. 160 it is to be read as a hadronic factor K_{had} , while the primary K_{frak} remains that of Doc. 133. The factor was selected in Doc. 160 by a scan over combinations; the result for the lifetime ratio τ_μ/τ_e is therefore [K] with a selection caveat, not parameter-free. Doc. 160 remains unchanged. Consequence: calculations using K_{frak} (e.g. v in R104 or a future derivation of τ_μ) use exclusively the value of Doc. 133. Open bridge: geometric derivation of the hadronic factor [S].

No.	Affects	Subject (short form; details: Archive)
R124	Doc. 374	Withdrawal of the #740 statement (22 September 2026) [X]. The result section of Doc. 374 concludes that the bridge to Observer Patch Holography is closed under the reading $E_* = E_p$. That statement goes too far: the coefficient has meanwhile been derived on the other side, while the length identity, the clock and a matching factor are still carried there as open, as are most of the dependencies of the item itself. What remains valid are the bridge equations B1–B4 and the absolute scale through the FFGFT chain; the reading is no longer posited on the other side but derived. Doc. 374 has been corrected directly; the theorem and the status statements are in Doc. 376.
R125	Docs. 085, 176, 179, 186	Bibliographic data and one standard-physics statement in the photonics documents (24 September 2026). Checking the cited photonics literature against the publisher records revealed four points that do not affect any derivation. (1) Doc. 085: "Photonics for sustainable AI" appeared in <i>Communications Physics</i> , not in <i>Nature</i> . (2) Doc. 176: the article number given for Wang et al. (2022) differs from the DOI record; the DOI is authoritative. The paper is a gedanken experiment with a simulation for a small example number; it supports the Fourier step with classical light conceptually, not the statements on speed and energy efficiency of the optical Fourier transform, which rest on the Optica source alone. (3) Doc. 186: the placeholder "K. Name et al." stands for Shimura et al. (TDK), as already given correctly in Doc. 187. (4) Doc. 179, SPDC section [STANDARD] : the perpendicular polarization of the two photons in type-II conversion follows from phase matching in the birefringent crystal, not from angular-momentum conservation. The documents remain unchanged; FFGFT statements are not affected.
R126	Doc. 230 (sections on states and measurement)	Cylinder representation of the qubit: placement in the literature and justification of the Born statistics (24 September 2026). Doc. 230 introduced the cylindrical coordinates (z, r, θ) without classification and attributed the Born statistics of measurement to the unknown phase θ . Refined: (z, r, θ) is the Bloch sphere in cylindrical coordinates; measurement drives the point deterministically to one of the poles, which pole is reached being decided by the hidden quantity λ along the axis; $(1+z)/2$ is not due to θ but is a geometric quotient (axis fraction, Archimedes' area preservation, only for complex amplitudes), not at root a probability, consistent with Docs. 365, 367 and 370. The closest relative of the measurement reading is the hidden-measurement model. For multi-qubit systems the deterministic reading is retained but, by Bell and Kochen–Specker, is necessarily non-local or contextual; the T^4 reading satisfies this. Sources [Q] added to Doc. 230; measurement section, state comparison and numerical equivalence reworded accordingly (the earlier identification of r^2 with $ \alpha ^2$ was incorrect).

No.	Affects	Subject (short form; details: Archive)
R127	Doc. 043, 044; Doc. 261 (units table, ϵ_0 paragraph)	$\alpha = 1$ is not a choice of units (25 September 2026) [X]. Doc. 044 (section “Why can $\alpha = 1$ be set?”) and Doc. 043 justify $\alpha = 1$ as the result of redefining the unit of charge; Doc. 261 adopts this as a convention for the written value. This does not hold: α is dimensionless and stays unchanged under any redefinition of the unit of charge, because e and ϵ_0 transform together. The step $e := \sqrt{4\pi\epsilon_0\hbar c}$ at fixed ϵ_0 is not a choice of units; it sets e equal to the Planck charge. Of e , $\hbar$, c and the Coulomb constant, at most three can be normalised to one simultaneously. For FFGFT this is a prerequisite, not a contradiction: α is derived as a dimensionless quantity from the dimensionless ξ , and it is precisely the unit independence of α that gives this derivation its meaning [B]. The physical invariant α obtained from ξ is authoritative (Doc. 261). The documents remain unchanged.
R128	Doc. 221 (Signature 3, falsification criterion 3)	No deviation of the supernova time dilation from $(1+z)$ at high z (28 Sept. 2026) [X]. Doc. 221 predicts for $z > 2$ a light-curve stretching $e^{\xi_0 x} = 1 + z + z^2/2 + \dots$ instead of $(1+z)$ and presents this as a discriminating test against standard cosmology. This does not hold: the series arises only from inserting the small-distance approximation $\xi_0 x \approx z$ into the exact law; moreover, the document uses two incompatible definitions of the redshift side by side. From $E \propto e^{-\xi_0 x}$ it follows exactly that $1+z = e^{\xi_0 x}$, and because of $\tilde{T} \cdot m = 1$ wavelength and clock rate are booked together, so the time dilation equals $(1+z)$ for all z [B], in agreement with Doc. 312 and A265. Light-curve stretching is therefore not a discriminator between FFGFT and expansion, only against tired light without time dilation. By Doc. 267 this holds for all frequency and clock-rate signatures of Doc. 221: they are degenerate and not falsification criteria; a static picture with $\tilde{T} \cdot m = 1$ is therefore not excluded. Doc. 221 remains unchanged.
R129	Doc. 006, 046 (universal quantum number table)	Unbooked early changes to the quantum number table (28 Sept. 2026, recorded retroactively). The table with (n, l, j, r_i, p_i) for all particles was changed during revisions in 2025 without a register entry: the prefactors r_i for tau, charm and strange and p for ν_τ were replaced, and the boson rows (Higgs, Z , W , photon, gluon) were removed. The new r_i are corrections; with the earlier values tau and charm lay clearly off the measured values. The p value for ν_τ is not tenable by R109; the boson masses have since been treated in Doc. 372 and 373. Clarification on reading the table: (n, l, j) coincide per generation for leptons and quarks and fix the generation and the exponent p_i , but not the prefactor r_i . Whether the quark r_i are geometrically forced remains open (Doc. 189, 373) [S]. Doc. 006 and 046 remain unchanged.

No.	Affects	Subject (short form; details: Archive)
R130	Doc. 343 (Theorem A)	Theorem A: scope of the statement (29 Sept. 2026). Theorem A derives $(1 - 3^{-s})\zeta(s) = L(s, \chi_0)$ in the one-dimensional analogue (decomposition by $n \bmod 3$) and calls the result the spectral zeta function of the $T^4/\mathbb{Z}_3$ torus. For the orbifold of the corpus (fixed-point-free $\mathbb{Z}_3$ on D_4 , Doc. 314/330) the spectral zeta function is one third of the four-dimensional Epstein zeta of D_4 , with the Weyl pole at $s = 2$ and without the factor $(1 - 3^{-s})$ [B] . Theorem A remains valid as a statement about the 1D analogue or $L(s, \chi_0)$ [B] ; the identification with the spectral zeta of the orbifold is withdrawn [K] . The same holds for Theorem B (Euler product by $\text{ord}_p(3)$) and Theorem C (zeros from the factor $(1 - 3^{-s})$): valid as statements about $L(s, \chi_0)$ and arithmetic modulo 3, not about the spectrum of the orbifold. The question whether the $\text{GF}(3^k)$ reading has a counterpart on the four-dimensional object is decided negatively: the Euler factors of the orbifold have the same form for every odd prime, including $p = 3$ [X] .
R131	Doc. 091 (appendix zeta regularisation)	Zeta regularisation of the mode sum closed for the corpus lattice (29 Sept. 2026). Doc. 091 carries the zeta regularisation of the zero-point energy only as a sketch, with a general $\zeta(-1)$ and no specified momentum lattice. For the momentum lattice D_4 of the corpus, $\zeta(-1)$ is expressed in closed form through values of the Riemann zeta function and is negative, so the Casimir term is attractive [B] . For the orbifold $T^4/\mathbb{Z}_3$ the value is one third of this, because the $\mathbb{Z}_3$ acts freely on all momenta $\neq 0$; the nine fixed points in position space are unaffected [B] . At equal volume of the fundamental cell, D_4 has the smallest magnitude among the comparison lattices, in the same order as the lattice energy in Doc. 380 [K] . The point carried as a sketch is thereby closed; the coefficient is lattice-dependent. Doc. 091 remains unchanged.
R132	Doc. 077, 078, 134, 145	Variable speed of light replaced by fractal path lengthening (29 Sept. 2026). The listed documents work with a varying speed of light: $c = L/T$ as a ratio that varies with position, time or mass scale (Doc. 077, 078, 134), or an effective speed depending on the medium (Doc. 145). In the later corpus this idea is replaced by fractal path lengthening: the effects in question are described as a geometric stretching of the propagation path in the ξ field (Doc. 026, 149, 182), which is achromatic according to K6. The four documents carry a corresponding note after the abstract; statements and predictions that rely on a variable speed of light are to be read accordingly. Doc. 030 mentions variable-speed-of-light theories only as a comparison and is not affected.

No.	Affects	Subject (short form; details: Archive)
R133	Doc. 019, 030, 033, 049, 053, 059, 070, 081, 095; A138	Muon $g - 2$ anomaly no longer significant (29 Sept. 2026) [X]. The listed documents use the deviation between the measured a_μ and the Standard Model prediction as of 2021 as evidence, as the target of a T0 contribution, or as a benchmark. After the Fermilab final result and the 2025 White Paper of the Muon $g - 2$ Theory Initiative this deviation is no longer significant; the statement that the ξ term explains the muon anomaly is outdated. The load-bearing ratio line (Doc. 018 Rev. 12, 158, A138) is not affected: its ratios are exact, and its absolute values follow by anchoring to measured values, which have not changed. The listed documents carry a note after the abstract; the current status and assessment are given in Doc. 382. It remains open whether the FFGFT yields a contribution within today's allowed range [S].
R134	Doc. 008, 018 (Rev. 10–12), 033, 046, 049, 059, 070, 081; R109	Lepton masses and individual $g - 2$ formulas: calculation errors in older documents (29 Sept. 2026). Recalculation with conversion and the respective anchor. The logarithmic symmetry of the muon mass holds only with the additional factor of the lepton ladder (Doc. 352); without it the calculation in 008 and 033 is wrong, as a bare value the result is correct. Individual formulas for a_μ , a_τ and an equal contribution for all leptons (049, 059, 081) do not give the stated numbers or are excluded by the measurement of a_e ; authoritative is the ratio line (Doc. 158, 382). In 018 sums and roundings of the a_τ values are inconsistent (Rev. 11: precision, not an error); in 070 measured values appear as calculation results and several intermediate steps are wrong. R109 applies equally to the neutrino values in 046; the 4th generation extrapolated there is experimentally excluded. The documents carry a note before the first section [X].
R135	Doc. 012, 020, 041, 056, 059, 068, 070, 077, 081, 105, 127, 149, 167, 180	Constants, couplings and G: calculation and factual errors; SI conversion of G (29 Sept. 2026). The short formula for G in Doc. 012 reproduces the measured value, the step chains in 012 and 127 do not. The conversion factor C_{conv} is not a pure unit conversion but contains the Planck energy and thus G ; the statement in 180 that ℓ_p follows entirely from ξ is therefore circular [X]. Open remains a derivation of G whose conversion uses only $\hbar$, c and the energy unit [S]. Further points: $G = \xi^2 c^3 / \hbar$ (081) dimensionally wrong; θ_{QCD} or θ_{CP} from ξ (041, 059) and $m_h/M_p = \sqrt{\xi}$ (068, 167) excluded by measurements; numerical errors in UV cutoff, couplings and α (020, 070, 081, 105); c and μ_0 are SI definitions or measured quantities respectively, not derivations (056, 077). The documents carry a note before the first section.

No.	Affects	Subject (short form; details: Archive)
R136	Doc. 033, 061, 063, 091, 145; R70, R131	CMB and Casimir calculations (29 Sept. 2026) [X]. According to the literature, the formula with denominator 240 is the Casimir pressure, not the energy density; 061, 063 and 091 compare it with the energy density of the CMB, which changes the ratio by a factor of three. The comparison value called experimental is the same formula, not a measured value. In 145 the Casimir pressure is attractive and not stabilising; the exact torus evaluation (R131) confirms the sign. Further calculation errors: CMB temperature in 033 (exponent lost), in 063 measured values instead of formula values, missing c for the frequency, non-reproducible ρ_{CMB} number, significance in 061. The cosmic sector remains bracketed (P39). The documents carry a note before the first section.
R137	Doc. 018, 146, 149, 150, 155, 183, 210	Numerical and structural statements; $f = 1/\xi$ instead of $f = 1/(4\xi)$ (29 Sept. 2026). The form $f = 1/(4\xi)$ given in 210 as the binding reading is miscalculated; correct is $f = 1/\xi$, and the justification of the factor four from four-dimensionality falls away. 149 and 150 are affected as well. In 149 the number of divisors, the deviation of G and the quark formulas are also miscalculated. Further points: literature examples for fractal dimensions misattributed (018), activator and inhibitor of the Turing condition swapped and inconsistent critical values (146), number of levels and total factor contradictory (155), ξ as an approximation of a natural value instead of an exact setting (183). The documents carry a note before the first section [X].

.2 Currently open bridges (as of R137)

- τ_μ directly from FFGFT, without detour via G_F and ν — the only convention-free test of the weak sector (R112, Doc. 351) [S]
- Why do ε_i decrease with generation? In Reading A (T0 fundamental) attributed to QED + SI projection, R113 bridge open [S]; in Reading B (Koide fundamental, Doc. 367) answered: distance cosine–power law, no QED needed [K]. What remains open in Reading B is why the rational approximation carries exactly the $\text{GF}(3^n)$ winding numbers (Doc. 369 § 4, R121) [S]
- Koide amplitude $d/c = \sqrt{2}$ from FFGFT geometry — orthogonal to $\theta = 2/9$ [B], not yet anchored (Doc. 352, § 8) [S]
- Charge quantisation $Q \in \{0, \pm\frac{1}{3}, \pm\frac{2}{3}, \pm 1\}$: closed by Doc. 346, R111 [B] —
- Gell-Mann–Nishijima $Q = I_3 + Y/2$: closed by Doc. 347, R111 [B] —
- Δm_{32}^2 atmospheric (+16.0 %): closed by Doc. 340, R103 [K] —
- Geometric derivation of the hadronic factor in Doc. 160 (K_{had} , R123) [S]
- Λ_{QCD} from ξ — not derivable from the ξ ladder (R122, Doc. 372) [X]; Higgs mass: candidates via Galois chain and trace rule (Doc. 373) [K]; residual of the trace rule presumably from the definition of ν (R112) [S]
- 2-loop corrections to $\alpha_s(M_Z)$ [S]
- m_{Pl} and $\alpha_{\text{em}}(M_Z)$ from ξ (external inputs in R81) [S]
- Quark/hadron sector — Doc. 318, R76

- Greybody factors; back-reaction on orbifold fixed points — R85 [S]
- Forward derivation of the cosmic exponent $41/4$ (P20); the 10^{123} question, $R_H \rightarrow \ell_1$ and $\kappa = \Lambda_{\text{obs}} R_H^2$ hang on it
- Light elements without a hot early phase: abundances of ^4He and D as dimensionless number ratios, not reinterpretable via frequency degeneracy (Doc. 267, 222; R128) [S]
- CMB peak selection: generate $\{1, 6, 14, 26\}$ forward; the $|n|^2=30$ case (P29/P31)
- C_ℓ projection: physical source/window function (P30)
- Ω_{DM} quantitatively, model-neutral (P17)
- Spectral dimension $d_s = 1.86$ vs. FFGFT ≈ 2
- Corpus-wide H_0 language cleanup (P34); fundamental rework of Doc. 026 (P16)
- Justify the base mass $M_{\text{base}} = m_P$ in the fundamental sector — Doc. 180, R90 [S]
- ξ as line width (P44, pre-registered); ι embedding $\text{HLV} \subset \text{FFGFT}$ in the time sector (Doc. 297)
- Adversarial D_4 operator-specificity test against matched comparison carriers (GAE logic) — R95 [S]

R86 – Self-adjointness of $\hat{F}$ (August 2026): Doc. 327 closes the gap declared as [S] in R82 completely. The finiteness of the scale ladder is guaranteed by $L_0 = \xi \cdot \ell_P$ (Doc. 180 [K]) – no infinities in FFGFT, hence $\hat{F}$ is bounded and $\mathcal{D}(\hat{F}) = \mathcal{H}$. The symmetry $\hat{F} = \hat{F}^\dagger$ follows from the $\mathbb{Z}_3$ pairing $(k, -k)$ (the same as in Doc. 325 [B]). Deficiency indices $(0, 0)$: exactly one self-adjoint realisation [B]. Fractal measure (finite 100-step recursion), $\mathbb{Z}_3$ restriction and all three χ -twist classes preserve self-adjointness [B]. $\xi = \lambda_{\min}(\hat{F}_{D_4})$ well-defined and truncation-stable (min-max) [B]. No remaining cases. Verification script: 327_selbstadjungiertheit_F_verify.py (21 assertions).

Declared 23 August 2026

R87 – Δm_{32}^2 : corrected numerical values and candidate analysis (August 2026): In Doc. 320 the evaluation of $\xi^{9/5}$ was faulty (8.717×10^{-8} instead of 1.059×10^{-7}); the mass formula $m_{\nu_i} = m_e \xi^{p_i}$ and the exponents $p_i \in \{9/4, 2, 9/5\}$ were always correct. Docs. 320 and 322 have been brought to the correct numerical values: $m_{\nu_3} = 54.11$ meV [K], $\Delta m_{32}^2 = 2.846 \times 10^{-3}$ eV² [K] (deviation +16.0 % against NuFIT 5.3), $\sum m_\nu = 64.17$ meV [K] (still within the Planck limit). Unaffected: m_{ν_1} , m_{ν_2} , Δm_{21}^2 (+8.3 %) and the fixed-point assignment. Doc. 320 was at the same time extended by a candidate analysis for the remaining deviation: K_{frak} at fixed point F_7 (+12.9 %, insufficient), mixing term $F_5 - F_7$ (right sign, complete mass matrix outstanding – the most promising candidate), alternative exponents (unlikely: $p_3^{\text{PDG}} = 1.8081$ lies only 0.45 % above $9/5$). Δm_{32}^2 remains [S]. Verification script: 320_dm32_neutrino_verify.py (10 assertions).

Declared 23 August 2026

R88 – n_{thresh} is scale-dependent; Matzke's scale reconstructed (August 2026): Doc. 329 settles numerically the question booked as [S] in Doc. 326. The collapse condition Compton = Schwarzschild yields $M_{\text{coll}} = m_P/\sqrt{2}$ purely geometrically and free of any bit value [B]. With the system-dependent bit energy $E_{\text{bit}}(L) = \hbar c/L$ (Doc. 257) one obtains exactly $n_{\text{thresh}}(L) = L/(\sqrt{2} \ell_P)$ – linear in the scale [B]. Matzke's value 6.4097 corresponds exactly to $L = 2\pi \ell_P / \ln 2 = 9.0647 \ell_P$ [B]; the "universal" bit value is the FFGFT bit energy at that scale. Across the scales of the corpus n_{thresh} varies by a factor of 4.6×10^{29} ; a 10 % temperature deviation flips the stability claim for $\text{Cl}(6)$. Hence $n_{\text{thresh}} = 6.41$ as a universal number is negatively closed [X]. Untouched: Compton = Schwarzschild, r_s without field equations, $\text{Cl}(6)$ structure [B]. Verification script: 326_nthresh_skalenabhaengigkeit.py (12 assertions).

Declared 24 August 2026

R89 – Time-field mediator mass: one structure, sector-dependent base mass (August 2026): The corpus uses m_T with widely differing numerical values (Docs. 019/180/201: $m_T = \lambda/\xi$; Doc. 032: effective torsion mass ≈ 5.220 GeV). This is no contradiction: both follow the rule $m_T = M_{\text{base}}/\xi$, whence the range of the massive time field is $\hbar/(m_T c) = \xi \cdot \lambda_C(M_{\text{base}})$ **[B]** — universal form, sector-dependent value, structurally identical with $E_{\text{bit}} = \hbar c/L$ (Docs. 257, 329). Lepton sector ($M = m_e$): $m_e/\xi = 3.83$ GeV, with the $O(1)$ factors of Doc. 032 ($\sin \pi \xi$, π^2 , $\sqrt{\alpha/K_{\text{frak}}}$, R_f) 5.220 GeV — range 3.78×10^{-17} m. Fundamental sector ($M = m_p$): $m_T = m_p/\xi = 7500 m_p$ — range 2.155×10^{-39} m = L_0 exactly. Doc. 032 itself calls its quantity an *effective* torsion mass; sameness of name is not sameness of quantity. Verification script: 180_mT_struktur_L0_verify.py (17 assertions).
Declared 24 August 2026

Placement after Doc. 287: The Lagrangian description ($m_T = M_{\text{base}}/\xi$) and the bit-energy description ($E_{\text{bit}} = \hbar c/L$) are *different representations of the same floor structure* – in the terms of Doc. 287, presumably a **bijection**: lossless and mutually translatable, but neither more complete than the other. The generative core that loses nothing is the T^4 torus (Doc. 287, Doc. 243); all other descriptions stand to it in one of three relations: identity, bijection or projection.

R90 – The exponent in $L_0 = \xi \cdot \ell_P$ is derived (August 2026): Doc. 180 justifies $L_0 = \xi \cdot \ell_P$ by the ξ scaling of the coupling term $g_T^\ell = \xi m_\ell$ — an argument by analogy. The exponent however follows *necessarily* from the Lagrangian mass term: $m_T = M_{\text{base}}/\xi^1$ together with $\lambda_C \sim 1/m$ gives range $\sim \xi^1$ **[B]** (R89). Counter-check: $m_T \sim \xi^{-n}$ would give $L_0 = \xi^n \ell_P$; the exponent of the range is the negative exponent of m_T . L_0 is thus the Compton range of the time-field mediator in the fundamental sector — below it the time field can no longer mediate, which is why “distance” loses its meaning. This also explains *why* a floor exists at all: because the mediator is massive. What remains open is only the choice $M_{\text{base}} = m_p$ for the fundamental sector; it introduces no new parameter ($\lambda = 1$ in Planck units) but is not independently justified [S]. Without consequences for Doc. 327 (R86): boundedness requires only a finite floor **[B]**.
Declared 24 August 2026

R91 – λ in $m_T = \lambda/\xi$ is a base mass (August 2026): Docs. 019 and 201 describe λ as the “Higgs coupling parameter”. This is misleading: (i) Dimensionally λ must be a *mass*, since m_T is a mass and ξ is dimensionless **[B]**; the Higgs quartic $\lambda_h \approx 0.129$ is dimensionless. (ii) λ_h runs under the RGE and is a low-energy parameter of the electroweak scale; a value measured there cannot be inserted at the Planck scale. (iii) Interpreted as a dimensionless factor in Planck units, λ_h would give the range $7.75 \cdot L_0$ — missing L_0 . Correct is $\lambda = M_{\text{base}}$ of the respective sector, in the fundamental case m_p (R89/R90). A Higgs connection can at most be an effective statement in the electroweak sector, never the definition. The label in Docs. 019/201 is to be corrected [S].
Declared 24 August 2026

R92 – Two misattributions concerning the time field (August 2026): (i) Doc. 250: The sentence “The term $-\frac{1}{2}m_T^2(\partial m)^2$ with $m_T = \xi/\ell$ (Doc. 201)” contains two errors. In natural units $m = 1/\ell$, so $\xi/\ell = \xi \cdot m$ — this is the *coupling* $g_T^\ell = \xi m_\ell$, not the mediator mass $m_T = M_{\text{base}}/\xi$ **[B]**. Furthermore the mass term reads $-\frac{1}{2}m_T^2(\Delta m)^2$, not $(\partial m)^2$; the latter would contribute to the kinetic term. The physical point (the mass term provides the stiffness against divergence of Δm) remains correct; only the formula is misquoted [S]. (ii) Doc. 190, *Clarification 4*: The statement recorded as an internal consistency check, that an object with $M_0 = \xi m_p/2$ would have Schwarzschild radius exactly L_0 , is an *identity*: from $r_s = 2GM/c^2$ and $\ell_P = Gm_p/c^2$ it follows that $r_s(xm_p/2) = x\ell_P$ for any x and any exponent **[B]**. It tests neither ξ nor the exponent in L_0 ; classifying it as a “check” overstates it. Non-trivial would be only the condition

appended there, whether M_0 coincides with a rung of the ξ hierarchy — open [S].

Declared 24 August 2026

R93 – $S_{\text{BH}}(M_{\text{coll}}) = 2\pi \text{ nat} = n_{\text{thresh}}$ (Matzke) in bits (August 2026): Doc. 325 §5.3 (new) and Doc. 329 [B] jointly yield: $r_s(M_{\text{coll}}) = \sqrt{2} \ell_P$ (geometric, Compton = Schwarzschild); $S_{\text{BH}}(M_{\text{coll}}) = 4\pi \cdot 2\ell_P^2/(4\ell_P^2) = 2\pi \text{ nat}$ exactly. In bits: $2\pi/\ln 2 = 9.0647 \text{ bit} = n_{\text{thresh}}$ (Matzke) [B]. Interpretation: Matzke's stability threshold is the residual entropy of the evaporation endpoint — the same geometry in two languages. Matzke's universality claim remains closed negative [X] (R88); this identity explains *why* the value 9.065 appears in both contexts without making it universal. Verification script: 325_informationsbilanz.py (9 assertions, assertions 2–3).

Declared 25 August 2026

R94 – $I_{\text{sector}}/I_{\text{thermal}} = \log_2 3$ for all M (August 2026): From Doc. 325 §5.1 [B] ($\log_2 3$ bits of sector information per quantum, orthogonal $\mathbb{Z}_3$ pair states) and §Building Block 2 [B] (1 nat thermal area bookkeeping per quantum via Clausius):

$$\frac{I_{\text{sector}}}{I_{\text{thermal}}} = \log_2 3 \approx 1.585 \quad \text{for every mass } M > M_{\text{coll}}.$$

The ratio is mass-independent — a structural constant of the $T^4/\mathbb{Z}_3$ geometry. Total budget: $N_{\text{emit}} = S_{\text{BH}}(M_{\text{init}}) - 2\pi \text{ nat}$, $I_{\text{sector}} = N_{\text{emit}} \cdot \log_2 3$ bits. Verification script: 325_informationsbilanz.py (assertions 6–7).

Declared 25 August 2026

K-041-a/b/c – Internal inconsistencies in Doc. 041 (August 2026): Doc. 041 remains unchanged. The following notes document places where printed formulas and table values in the same document disagree under the document's own $\xi = (4/3) \cdot 10^{-4}$ (Doc. 328 §7.4b). Resolution requires author review of the source document.

K-041-a (δ_{CKM}): Formula $\delta_{\text{CKM}} = \arcsin(2\sqrt{2} \cdot \sqrt{\xi}/3)$ yields 0.0109 rad; table value 1.20 rad. Candidate: $\arcsin(2\sqrt{2}/3) = 1.231 \text{ rad}$ — factor $\sqrt{\xi}$ likely a typesetting error.

K-041-b (θ_{13}): Formula $\theta_{13} = \arcsin(\xi^{1/3})$ yields 2.93° ; table value 8.57° . Candidate: argument $1/300 = 25\xi$ yields 8.59° — whether $1/300$ is a corpus quantity is to be clarified at the source document.

K-041-c ($|V_{us}|$): Formula with Doc. 006 masses yields 0.2124; table value 0.22452 (dev. 5.4 %). Possible cause: differing masses or f_{Cab} definition in Doc. 041.

Declared 25 August 2026

R95 – Doc. 330: two wording refinements and Type-I clarification after external review (August 2026): External review (M. Krüger, IPI, 25 August 2026) of Doc. 330 identified two overly superficial formulations; the corpus itself (Docs. 270/314/327, R41, R86) was correct and more precise in each case. Doc. 330 remains unchanged; the refinements apply at the register level.

(i) *Intrinsic flatness of T^4* (Doc. 330 §2.2). **Too superficial:** “positive and negative curvature contributions cancel in the sum” — this conflates the *embedded* torus in $\mathbb{R}^3$ ($K > 0$ outside, $K < 0$ inside, only globally $\int K dA = 0$ via Gauss–Bonnet, $\chi = 0$) with the *flat* torus $\mathbb{R}^4/\mathbb{Z}^4$. **Precise:** The flat metric δ_{ij} is $\mathbb{Z}^4$ -invariant and descends undistorted to the quotient; $\Gamma_{ij}^k = 0$, Riemann tensor pointwise zero, $K = 0$ everywhere — there is nothing to cancel. Triply verified constructively ($\Gamma = 0$, angle sum π , trivial holonomy). The conclusion of Doc. 330 (intrinsically flat, Fourier basis without curvature corrections, Type III bijective) remains unchanged [B]. Nash–Kuiper side note: the flat torus is not C^2 -isometrically embeddable in $\mathbb{R}^3$ — the embedding question does not arise for the fundamental level anyway.

(ii) *Self-adjointness of $\hat{F}$* (Doc. 330 §Question 3). **Too superficial:** “bounded and therefore self-adjoint” — boundedness alone does not imply self-adjointness (counterexample: shift

operator). **Precise (= R86 [B])**: bounded *plus* symmetric via the $\mathbb{Z}_3$ pairing $(k, -k)$ on a complete Fourier basis $\Rightarrow$ self-adjoint; deficiency indices $(0, 0)$; boundedness ($\|\hat{F}\| \leq \sum_n \xi^n = \xi/(1-\xi)$, convergent even for $N \rightarrow \infty$) secures $\mathcal{D}(\hat{F}) = \mathcal{H}$, while the symmetry argument carries the burden independently. Status **[B]** unchanged.

(iii) *Type-I direction clarification (no error; confirmation of R41)*. The review objection that the unrolling $S_m^1 \rightarrow \mathbb{R}_t$ has a lifting problem (covering $p : \mathbb{R} \rightarrow S^1$ without canonical inverse) concerns the *lifting of points* — Type I, however, lifts no points but is the *function pullback* $\Phi = p^* : L^2(S_m^1) \rightarrow \text{Per}_{R_m}(\mathbb{R}_t)$, $(\Phi f)(t) = f(t \bmod R_m)$ — canonical, no additional datum. Precisely the reverse direction $\mathbb{R} \rightarrow S^1$ is the lossy one (R41, declared 19 June 2026). The external six-point checklist is answered completely: injective; surjective onto Per_{R_m} (not onto $L^2(\mathbb{R}_t)$ — that is the embedding statement of Doc. 270); isometry per period (Haar $\leftrightarrow$ Lebesgue/period); inverse = restriction to an arbitrary period (period choice = deck-transformation gauge freedom, not a missing datum); spectrum discrete $\leftrightarrow$ discrete frequencies n/R_m ; $U(1)$ translation $\leftrightarrow$ time translation commute with Φ . The proof sketch of Doc. 270 (theorem “Type I is information-preserving in mode content”) is thereby completed to a full theorem: status change [sketch] $\rightarrow$ **[B]**.

(iv) *Accepted as open*: An adversarial operator-specificity test for D_4 against matched comparison carriers (GAE logic) does not exist; $5 \nmid |\text{Aut}(D_4)| = 1152$ is a lattice-specific algebraic fact (excluding e.g. E_8 with $5 \mid |\text{Aut}(E_8)|$), not a specificity proof. Carried as an open bridge **[S]**.

Verification scripts: `pruef_krueger_einwaende.py` (16/16), `pruef_k1_tief.py` (8/8), `pruef_k3_tief.py` (6/6), `pruef_k4_tief.py` (5/5).

Declared 25 August 2026

R96 – Type-I follow-up refinement: pullback is a change of representation, not a new sector (August 2026): Second feedback (M. Krüger, IPI, 26 August 2026) on the R95(iii) clarification accepts the function pullback $\Phi = p^*$ as the correct mathematical object and the refutation of the original lifting objection, but refines a remaining point: Φf is R_m -periodic, so $\int_{\mathbb{R}} |\Phi f|^2 dt = \infty$ for any $f \neq 0$ — Φf does not lie in ordinary $L^2(\mathbb{R}_t)$ with Lebesgue measure over the whole line. The correct target space $\text{Per}_{R_m}(\mathbb{R}_t)$ (periodic L^2_{loc} , norm over one period) is isometrically isomorphic to $L^2(S_m^1)$ itself — the same countable Fourier basis $\{e_n\}_{n \in \mathbb{Z}}$, the same Parseval identity. Formally:

$$L^2(S_m^1) \cong \text{Per}_{R_m}(\mathbb{R}_t) \quad (\text{per-period norm}), \quad \text{not} \quad L^2(S_m^1) \cong L^2(\mathbb{R}_t).$$

Algebraically confirmed (`pruef_332_krueger_periodizitaet.py`): divergence of the whole-line integral for any nontrivial periodic function (translation-invariant Lebesgue measure, each period contributes the same positive amount); isomorphism of the Fourier bases of both spaces via $\langle e_{n_1}, e_{n_2} \rangle = R_m \delta_{n_1 n_2}$ over one period. Status **[B]**.

Assessment: This is not a correction to Doc. 330/R95(iii), but a precisification of what the status change [sketch] $\rightarrow$ [B] there mathematically carries: an information-preserving representation of the compact sector on the universal cover, not a genuinely noncompact state space. Doc. 332 (Section 2, “canonical function pullback”) already formulated this restraint correctly (“change of representation of the same function algebra”, no extension), but without explicitly separating the norm/Hilbert-space level from the algebra level; Doc. 332 was accordingly extended with a clarifying paragraph (Section 2.1, “continuum of the coordinate representation” vs. “continuum of the state space”) and recompiled.

Side correction (Nash–Kuiper): The Nash–Kuiper side note in Doc. 330/R95(i) is refined: Nash–Kuiper is an existence theorem for C^1 -isometric embeddings, not a non-existence theorem for C^2 embeddings of the flat 2-torus in $\mathbb{R}^3$ — these two statements (existence of C^1 isometries and the C^2 obstruction question) must be kept separate. For the fundamental level T^4 the embedding question into $\mathbb{R}^3$ does not arise dimensionally anyway; the point was

not load-bearing for the main R95(i) conclusion (intrinsic flatness) and does not change its status **[B]**.

Verification script: `pruef_332_krueger_periodizitaet.py` (4/4 assertions).

Declared 26 August 2026

Declared 25 August 2026