

Furey Fermions in GALG and FFGFT: N-Operator, Tripotents, and Neutrino-Vacuum Identity

Abstract

Doug Matzke's IPI mail of 3 September 2026 presents concrete GALG results on Furey fermions in $Cl(6)$: complete Up/Down fermion sets Su/Sd , idempotents N_1, N_2, N_3 , tripotent N , jE7 projector, and a grade-1+grade-5 construction. The strongest single result: $Su[0] = Vss[0]$ exactly — the neutrino is the lowest vacuum state **[B]**. Theorem F shows: the tripotent property for even vacua and the Frobenius fixed-point property (Doc. 339) are the same decomposition from two independent directions **[B]**. The electron exception at the jE7 projector is the known Furey problem **[S]**.

.1 Theorem A — N_k are idempotents in characteristic 3

$N_k = -1 + (-i B_k)$, $B_k^2 = -1$ (Witt pair):

$$N_k^2 = 1 + 2i B_k + (-1)(-1) = 2 + 2i B_k \equiv -1 + (-i B_k) \pmod{3} = N_k. \quad \mathbf{[B]}$$

The sum loses its scalar part since $-3 \equiv 0$: $N_1 + N_2 + N_3 \equiv (-i) \sum B_k = N$.

.2 Theorem B — N is a tripotent

Cross-terms $N_k N_j$ ($k \neq j$) are pure grade-4 elements; hence $N^3 = N$ **[B]**. Doug's output: $N**3 = N$, $N**4 = N**2$.

.3 Theorem C — $Su[0] = Vss[0]$: neutrino is the vacuum

$Su[0] = Vss[0]$: True and $(1+N)*(1+jE7) = Vss[0]$: True **[B]**.

FFGFT correspondence [B]. In FFGFT, v_1 lies in orbit $O_1 = \{1, 3, 9\}$, the orbit of the generator of $GF(27)^*$ — the algebraically simplest object. Both frameworks independently place the neutrino at the algebraically simplest position: convergence of two independent frameworks.

.4 Theorems D–F

D — sigma [B]. $e_1 \rightarrow e_2 \rightarrow e_4 \rightarrow e_1$, $e_3 \rightarrow e_6 \rightarrow e_5 \rightarrow e_3$; $\{1:2, 2:4, 4:1, 3:6, 6:5, 5:3\}$, two 3-cycles, $\sigma^3 = \text{id}$. Doc. 341 contains this formula; Doug's `all_g1_g5good` is the σ -invariant sum.

E — $jE7 \equiv N_k$ as projectors [B]; electron exception [S]. $F \cdot jE7 = +F$ for all Su (8/8); $F \cdot jE7 = -F$ for 7/8 of Sd (Sd[7]=electron fails). Same for N_k . Known Furey problem. FFGFT: electron in Dirac layer $\{f_3, f_4\}$, neutrino in Majorana layer $\{f_1, f_2\}$ (Doc. 343, Theorem D'''). Common root open [S].

F — Tripotent for even vacua [B]. $(V_k + P)^2 = -V_k$, $(V_k + P)^3 = V_k$ for $k \in \{0, 2, 4\}$; no collapse for $k \in \{1, 3, 5\}$ (length 32). Identical to the Frobenius fixed-point/orbit split of Doc. 339 — two independent derivations of the same decomposition.

.5 Theorem H — The Galois algebra forces Dirac neutrinos

Theorem .5.1 (Dirac constraint [B] under assumption [S]). *If the sector pairing $k \mapsto -k$ in $GF(27)^*$ corresponds to the physical charge conjugation C , then **all** neutrinos are Dirac particles.*

Proof. A Majorana particle is a fixed point of charge conjugation: $\nu = \bar{\nu}$, i.e. class f_i must map to itself under C .

Under the sector pairing $k \mapsto -k \pmod{26}$:

$$\begin{aligned} f_1 = \{1, 3, 9\} &\mapsto \{25, 23, 17\} = f_2 \neq f_1 \\ f_2 = \{17, 23, 25\} &\mapsto f_1 \neq f_2 \\ f_3 = \{5, 15, 19\} &\mapsto f_4 \neq f_3 \\ f_4 = \{7, 11, 21\} &\mapsto f_3 \neq f_4 \end{aligned}$$

No primitive class is a fixed point of the sector pairing [B]. A Majorana fixed point would require $f_i \mapsto f_i$ — this does not exist in $GF(27)^*$. The anti-particle of the neutrino f_1 is f_2 : $\nu_1 \neq \bar{\nu}_1$. $\square$

Assumption [S]. The argument holds if $k \mapsto -k$ in $GF(27)^*$ corresponds to physical charge conjugation C . In Doc. 336 the Frobenius involution $x \mapsto x^3$ is identified with the sector separation; the identification with C is physically motivated there but not algebraically proved.

Consistency [B]. In GALG/Furey (Doc. 344, Theorem C): $Sd[0] = Vss[1] \neq Vss[0] = Su[0]$, i.e. $\bar{\nu} \neq \nu$ — fully consistent with this theorem.

Plain language: What is m_{ee} and why does it matter?

The process. A radioactive nucleus converts two neutrons into two protons simultaneously, emitting two electrons. In ordinary double beta decay ($2\nu\beta\beta$) two anti-neutrinos also come out — the energy is shared among four particles. In *neutrinoless* decay ($0\nu\beta\beta$) no anti-neutrinos come out.

Why this requires Majorana. The anti-neutrino produced at the first nuclear vertex must be absorbed as a neutrino at the second vertex. This is only possible if neutrino and anti-neutrino are the same particle ($\nu = \bar{\nu}$, Majorana condition). For Dirac neutrinos ($\nu \neq \bar{\nu}$) the process is exactly forbidden.

What m_{ee} measures. The half-life of $0\nu\beta\beta$ decay is proportional to $1/m_{ee}^2$, where

$$m_{ee} = |m_1 U_{e1}^2 + m_2 U_{e2}^2 + m_3 U_{e3}^2|$$

combines all three neutrino masses weighted by PMNS mixing elements and Majorana phases.

Why interference matters. The three terms can add (constructively) or partially cancel (destructively) depending on the Majorana phases. With FFGFT masses from Doc. 340 and

phases ≈ 0 : constructive gives $m_{ee} \approx 7.1$ meV; destructive gives $m_{ee} \approx 0.12$ meV; Dirac gives $m_{ee} = 0$ exactly.

What nEXO can measure. nEXO targets sensitivity ~ 5 meV — directly in the decision range. A signal at ~ 7 meV confirms Majorana and refutes Theorem H; no signal down to 5 meV excludes the constructive Majorana prediction and is consistent with Dirac or destructive interference.

Consequence: $m_{ee} = 0$ and no neutrinoless double beta decay

For Dirac neutrinos, neutrinoless double beta decay ($0\nu\beta\beta$) is **exactly forbidden**:

$$m_{ee} = \left| \sum_i m_i U_{ei}^2 \right| \equiv 0.$$

Table 1: Predictions: Majorana (Doc. 340) vs. Dirac (Theorem H)

	Majorana ν_1, ν_2 [S]	Dirac all [B]
m_{ee}	≈ 7.1 meV	$= 0$
$0\nu\beta\beta$	allowed, ~ 10 meV	exactly forbidden
nEXO signal	~ 7 meV at phases ≈ 0	no signal
KamLAND-Zen	consistent (< 36 meV)	consistent

Testability. nEXO (sensitivity ~ 5 meV) and KamLAND-Zen (Phase III, ~ 20 meV) probe exactly this range. A null result at nEXO down to 5 meV excludes Majorana with Doc. 340 masses (phases ≈ 0) and is consistent with Theorem H. A signal at ~ 7 meV refutes Theorem H and confirms the Majorana interpretation **[B]/[S]** of Doc. 340.

.6 Theorem G — Anti-neutrinos: Dirac or Majorana?

The tension. In GALG/Furey (Theorem C) the neutrino is $Su[0] = Vss[0]$ and the anti-neutrino is $Sd[0] = cc(Su[0]) = Vss[1] \neq Vss[0]$. Hence neutrinos in Furey's construction are **Dirac particles** ($\nu \neq \bar{\nu}$).

In FFGFT (Doc. 343, Theorem D'''), ν_1 and ν_2 lie in orbits O_1 and O_3 , which are self-inverse under $k \mapsto k^{-1}$ — the *algebraic* Majorana criterion.

Theorem .6.1 (Clarification of Majorana concept **[B]/[S]**). *Two distinct Majorana concepts:*

1. **Algebraic [B]:** Orbit O_k is self-inverse under $k \mapsto k^{-1} \bmod 13$ — a statement about $GF(27)^*$ structure, not about spin.
2. **Physical [S]:** $\nu = \bar{\nu}$ in the Lorentz sense (Majorana condition). This requires $Sd[0] = Su[0]$, which does not hold in GALG/Furey.

The Galois algebra fixes the algebraic property **[B]**; the physical Majorana condition is an additional assumption **[S]**.

Position of anti-neutrinos [B]/[S]. GALG: $\bar{\nu} = Sd[0] = Vss[1]$ in the odd vacuum (sigma-orbit, other sector, Doc. 339). FFGFT: the anti-neutrino would lie in the negation $f_1 \leftrightarrow g_1$ (Doc. 343, Theorem D'') — order-13 class g_1 , other sector. Both frameworks place the anti-neutrino in the structurally "other" sector.

Consequence for m_{ee} . The prediction $m_{ee} \approx 7.1$ meV from Doc. 340 assumes physical Majorana neutrinos $\nu_{1,2}$. If ν_1, ν_2 are instead Dirac particles (as in Furey), m_{ee} vanishes entirely. The m_{ee} prediction is conditioned on the Majorana assumption **[S]**.

Open question [S]. Common algebraic root of the electron inconsistency in GALG/Furey and the Majorana/Dirac layer in FFGFT.

Empirical decision. nEXO (sensitivity ~ 5 meV) decides: signal at ~ 7 meV $\Rightarrow$ Majorana (Doc. 340 confirmed, Theorem H refuted); no signal $\Rightarrow$ consistent with Dirac (Theorem H confirmed or destructive interference).

.7 Summary

Table 2: Results Doc. 344

Theorem	Content	Status
A	$N_k^2 = N_k$, characteristic-3 proof	[B]
B	$N^3 = N$, tripotent	[B]
C	$Su[0]=Vss[0]$: neutrino=vacuum; FFGFT correspondence	[B]
D	$\sigma = \{1 \rightarrow 2, 2 \rightarrow 4, 4 \rightarrow 1, 3 \rightarrow 6, 6 \rightarrow 5, 5 \rightarrow 3\}$	[B]
E	$jE7 \equiv N_k$; electron exception (Furey problem)	[B]/[S]
F	Tripotent even vacua = Frobenius fixed points (Doc. 339)	[B]
G	Anti-neutrino $Sd[0]=Vss[1] \neq Su[0]$; algebraic Majorana $\neq$ physical Majorana	[B]/[S]
H	Sector pairing $k \mapsto -k$ has no fixed point: all neutrinos Dirac; $m_{ee} = 0$; $0\nu\beta\beta$ forbidden	[B]/[S]

Open question [S]: Common algebraic root of the electron inconsistency in GALG/Furey and the Majorana/Dirac layer separation in FFGFT (Doc. 343, Theorem D''').

Verification script

pruef_344_furey_galg.py — 6 theorems, all assertions passed.

Bibliography

- [1] C. Furey, *Standard Model Physics from an Algebra?* (2016).
- [2] D. Matzke, IPI mail 3 September 2026.
- [3] J. Pascher, *Doc. 339: Frobenius separation*, FFGFT corpus (Aug. 2026).
- [4] J. Pascher, *Doc. 341: GF(27) in GALG and FFGFT*, FFGFT corpus (Sept. 2026).
- [5] J. Pascher, *Doc. 343: Zeta function of T^4/Z_3* , FFGFT corpus (Sept. 2026).